



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

On the impact of NMC data on NLO and NNLO parton distributions and Higgs production at the Tevatron and the LHC

Citation for published version:

Ball, RD, Bertone, V, Cerutti, F, Del Debbio, L, Forte, S, Guffanti, A, Latorre, JI, Rojo, J & Ubiali, M 2011, 'On the impact of NMC data on NLO and NNLO parton distributions and Higgs production at the Tevatron and the LHC', *Physics Letters B*, vol. 704, no. 1-2, pp. 36-42. <https://doi.org/10.1016/j.physletb.2011.08.055>

Digital Object Identifier (DOI):

[10.1016/j.physletb.2011.08.055](https://doi.org/10.1016/j.physletb.2011.08.055)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Early version, also known as pre-print

Published In:

Physics Letters B

Publisher Rights Statement:

Author's Pre-print: author can archive pre-print (ie pre-refereeing)

Author's Post-print: author can archive post-print (ie final draft post-refereeing)

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Edinburgh 2011/08
IFUM-973-FT
FR-PHENO-2010-003
RWTH TTK-11-04

On the Impact of NMC Data on NLO and NNLO Parton Distributions and Higgs Production at the Tevatron and the LHC

The NNPDF Collaboration:

Richard D. Ball¹, Valerio Bertone³, Francesco Cerutti⁴, Luigi Del Debbio¹,
Stefano Forte², Alberto Guffanti³, José I. Latorre⁴, Juan Rojo² and Maria Ubiali⁵.

¹ *School of Physics and Astronomy, University of Edinburgh,
JCMB, KB, Mayfield Rd, Edinburgh EH9 3JZ, Scotland*

² *Dipartimento di Fisica, Università di Milano and INFN, Sezione di Milano,
Via Celoria 16, I-20133 Milano, Italy*

³ *Physikalisches Institut, Albert-Ludwigs-Universität Freiburg,
Hermann-Herder-Straße 3, D-79104 Freiburg i. B., Germany*

⁴ *Departament d'Estructura i Constituents de la Matèria, Universitat de Barcelona,
Diagonal 647, E-08028 Barcelona, Spain*

⁵ *Institut für Theoretische Teilchenphysik und Kosmologie, RWTH Aachen University,
D-52056 Aachen, Germany*

Abstract:

We discuss the impact of the treatment of NMC structure function data on parton distributions in the context of the NNPDF2.1 global PDF determination at NLO and NNLO. We show that the way these data are treated, and even their complete removal, has no effect on parton distributions at NLO, and at NNLO has an effect which is below one sigma. In particular, the Higgs production cross-section in the gluon fusion channel is very stable.

Fixed target deep-inelastic scattering data provide important constraints on parton distributions (PDFs) and are routinely included in PDF determinations. It has been recently suggested [1] that the results of current PDF determinations depend strongly on the treatment of the fixed target DIS data obtained by the NMC collaboration [2, 3]: in particular, according to whether data for cross-sections or structure functions are used in the fit. The substantial changes in the gluon distribution and $\alpha_s(M_Z)$ found in Ref. [1] lead to a large shift in the Higgs production cross-section, which would, if correct, have very significant implications for Higgs searches at the Tevatron and LHC. This claim has generated an ongoing discussion on the adequacy of current Higgs mass limits [4, 5]; besides its interest in this context, the issue is relevant for the understanding of the comparative merits of PDF determinations based on a wider dataset (which contain more information but might be less consistent) and those based on a more limited but more consistent set of data.

In this note we examine this issue within the context of the NNPDF2.1 NLO [6] and NNLO [7] PDF determinations. In contrast to the ABKM [8] determination, on which the results of Ref. [1] are based, NNPDF2.1 depends on a rather broader dataset, and uses the especially flexible NNPDF methodology (for a review see e.g. Ref. [9]), making it less vulnerable to parametrization bias. Related results (consistent with our findings) have been presented recently in the context of the MSTW [5] and CTEQ [10] PDF determinations.

The kinematic coverage of the NMC data is compared in Fig. 1 to that of other datasets used to determine NNPDF2.1 PDFs: the other fixed target DIS data, the HERA collider data, the fixed target Drell-Yan and Tevatron weak vector boson production data and the Tevatron inclusive jet data. We will now consider variants of NNPDF2.1 in which the NMC data are treated in different ways. In all other respects, we adopt the default settings of NNPDF2.1 as discussed in Refs. [6, 7]. In particular, we take a fixed value for the strong coupling in both the NLO and NNLO fits, $\alpha_s(M_Z) = 0.119$, close to the PDG average [13]; sets with variable $\alpha_s(M_Z)$ are also available [6, 7, 14, 15], from which combined PDF+ α_s uncertainties can be computed [16, 17].

The NMC collaboration has measured the neutral current deep-inelastic muon-nucleon cross-section

$$\frac{d^2\sigma^{\text{NC}}}{dx dQ^2}(x, y, Q^2) = \frac{2\pi\alpha^2}{xQ^4} [Y_+ F_2^{\text{NC}}(x, Q^2) \mp Y_- xF_3^{\text{NC}}(x, Q^2) - y^2 F_L^{\text{NC}}(x, Q^2)] , \quad (1)$$

where $Y_{\pm} = 1 \pm (1 - y)^2$. For NMC $Q^2 \ll M_W^2$ so the parity-violating structure function xF_3 can be neglected and only the electromagnetic components of F_2 and F_L are relevant. It is convenient to define a reduced cross-section

$$\tilde{\sigma}^{\text{NC}}(x, y, Q^2) = \left[\frac{2\pi\alpha^2}{xQ^4} Y_+ \right]^{-1} \frac{d^2\sigma^{\text{NC}}}{dx dQ^2}(x, y, Q^2) = F_2^{\text{NC}}(x, Q^2) \left(2 - 2y + \frac{y^2}{1 + R(x, Q^2)} \right) , \quad (2)$$

where

$$R(x, Q^2) = F_L(x, Q^2) / (F_2(x, Q^2) - F_L(x, Q^2)) . \quad (3)$$

Equation (2) was used by the NMC collaboration [2, 3] to extract F_2^{p} from the measured cross-section Eq. (1), using for $x \leq 0.12$ a determination of $R(x, Q^2)$ from their own data,




Figure 1: Kinematic coverage of the NMC data, compared to those of the other datasets in the NNPDF2.1 global analysis: the non-NMC fixed target DIS data, the HERA collider data, the fixed target Drell-Yan and Tevatron weak vector boson production data and the Tevatron inclusive jet data.

and for $x \geq 0.12$ a parametrization R_{1990} of R [11] obtained from a global fit to SLAC structure function data [12].

In all NLO NNPDF parton determinations [6, 15, 18–20] the NMC structure function data was used, both for the proton structure function F_2^p and the ratio of deuteron to proton structure functions, F_2^d/F_2^p . It may be reasonably argued however that data for the reduced cross-section, which is closer to what is measured experimentally, should be used instead. Note that the distinction is only relevant for the F_2^p data: since the isotriplet component of $F_L(x, Q^2)$ is very small, $R^p(x, Q^2) \approx R^d(x, Q^2)$, so

$$\frac{\tilde{\sigma}^d(x, y, Q^2)}{\tilde{\sigma}^p(x, y, Q^2)} \approx \frac{F_2^d(x, Q^2)}{F_2^p(x, Q^2)}. \quad (4)$$

In Fig. 2 (to be compared to Fig. (1) of Ref. [8]) the form of $R(x, Q^2)$ used by NMC (shown as black dots) in both regions is compared to the prediction obtained using NNPDF2.1 NLO and NNLO PDF sets. The parametrization R_{1990} does not come with an uncertainty, however the typical size of the uncertainty on it can be inferred by comparing it to the data of Ref. [12] on which it is based, also shown in Fig. 2. Note that the SLAC data are concentrated at low Q^2 , hence in most of the NMC kinematic region this parametrization is an extrapolation and thus subject to very large uncertainties. It is clear from Fig. 2 that (as emphasized in Ref. [8]) the R values used by NMC at low x do not agree well with the prediction from the use of modern PDF sets, while instead the parametrization R_{1990} is in good agreement with the NNPDF prediction within the large




Figure 2: The NNPDF2.1 NLO and NNLO results for $R(x, Q^2)$ Eq. (3) at $x = 0.11$ (left) and $x = 0.14$ (right), compared to the values of R used by NMC in Ref. [2], and the SLAC data of Ref. [12] on which the parametrization [11] used by NMC for $x > 0.12$ is based.




Figure 3: The NNPDF2.1 NLO and NNLO predictions for the NMC reduced cross-sections for $x = 0.05$ (left), $x = 0.11$ (center) and $x = 0.225$ (right), compared to the NMC data [2].

uncertainty of the data on which it is based, especially if NNLO theory is used. Thus the use of NMC cross-sections instead of structure functions (which rely on these partly inadequate assumptions on R) does indeed appear to be in principle more advisable.

However, it is unclear whether in practice the effect of this replacement may be significant, especially in view of the fact that the NMC data are known to have internal consistency problems, as shown long ago in Ref. [22]. To illustrate this, in Fig. 3 the NMC reduced cross-section data are compared to NLO and NNLO predictions obtained using the corresponding NNPDF2.1 PDF sets. It is clear that the data show larger point-by-point fluctuations than one would expect from their nominal uncertainties, thereby suggesting that the effect of the treatment of the relatively small R -dependent correction might be moderate on the scale of these fluctuations.

In order to settle the issue quantitatively, we construct and compare, both at NLO and at NNLO, three PDF sets: one in which NMC data for the proton structure function F_2^p are used, one in which data for the proton reduced cross-section are used (supplemented, in both cases, by data for the ratio F_2^d/F_2^p), and one in which NMC data (both for proton and the deuteron/proton ratio) are removed altogether from the global dataset. In all cases sets of $N_{\text{rep}} = 100$ replicas have been produced. Note that the published (default) NNPDF2.1 sets [6, 7] use NMC structure functions at NLO and NMC cross-sections at

	NNPDF2.1 NLO			NNPDF2.1 NNLO		
	str. fctn.	xsec.	noNMC	str. fctn.	xsec.	noNMC
Total	1.16	1.14	1.09	1.16	1.16	1.12
NMC-pd	0.97	0.98	-	0.93	0.93	-
<i>NMC_p</i>	<i>1.73</i>	<i>1.67</i>	-	<i>1.69</i>	<i>1.63</i>	-
SLAC	1.27	1.27	1.28	1.05	1.01	1.00
BCDMS	1.24	1.23	1.18	1.29	1.32	1.27
HERAI-AV	1.07	1.05	1.07	1.12	1.10	1.08
CHORUS	1.15	1.11	1.07	1.12	1.12	1.12
FLH108	1.37	1.34	1.38	1.27	1.26	1.29
NTVDMN	0.47	0.51	0.42	0.50	0.49	0.50
ZEUS-H2	1.29	1.23	1.24	1.32	1.31	1.30
ZEUSF2C	0.78	0.74	0.72	0.88	0.88	0.89
H1F2C	1.51	1.48	1.49	1.47	1.56	1.52
DYE605	0.85	0.93	0.88	0.81	0.81	0.81
DYE866	1.27	1.40	1.34	1.31	1.32	1.34
CDFWASY	1.85	1.87	1.60	1.55	1.65	1.41
CDFZRAP	1.62	1.76	1.64	2.16	2.12	2.18
D0ZRAP	0.60	0.57	0.56	0.67	0.67	0.67
CDFR2KT	0.97	0.73	0.81	0.79	0.74	0.80
D0R2CON	0.84	0.90	0.96	0.84	0.82	0.84

Table 1: The χ^2 of NNPDF2.1 PDFs at NLO and NNLO when NMC data are included as structure functions, reduced cross-sections, or not included.

NNLO.

In Table 1 we compare the χ^2 values obtained in these three fits, both for the global fit and individual experiments. The quality of the global fit is unchanged at NNLO and improves slightly at NLO when the structure function data are replaced by cross-section data, and in both cases the quality of the fit to NMC data improves slightly, with the quality of the fit to other data unchanged. This suggests that the use of cross-section data is indeed somewhat more consistent for NMC, but also that this has little or no effect on other experiments. When the NMC data is removed altogether, the global fit quality improves, due to the fact that χ_{NMC}^2 is rather poor in view of the aforementioned inconsistencies, regardless of how NMC data are treated. In particular, the fit to the BCDMS data, which measure the same structure functions as NMC in a partly overlapping region, improves somewhat when the NMC data are removed.

We now compare the PDFs obtained in the various cases. In Fig. 4 (NLO) and Fig. 5 (NNLO) we show the distances (as defined in Ref. [15]), computed both for central values and uncertainties, between PDFs in the sets with NMC cross-section vs. structure function data, and a set without NMC data vs. the the default NNPDF2.1 set. Recall that $d \sim 1$ corresponds to statistically indistinguishable results, while, for sets of 100 replicas, $d \sim 7$ corresponds to a shift by one sigma (i.e. results are statistically distinguishable, but compatible within uncertainties).

These plots show that at NLO the replacement of structure functions with cross-sections is at the level of statistical fluctuations. At NNLO a small, statistically significant, shift in central values and uncertainties at the level of at most a third of a sigma but mostly lower is seen in some PDFs (specifically the quark singlet and isospin triplet and the gluon). The effect of removing NMC data altogether at NLO is again almost indistinguishable from




Figure 4: Distances (defined as in Ref. [15]) between NLO PDF sets with NMC structure functions and NMC cross-sections (top) and PDF sets with NMC structure functions and PDF sets without NMC data (bottom). All distances have been computed using sets of $N_{\text{rep}} = 100$ replicas.

a statistical fluctuation with the possible exception of the quark singlet for $0.02 \lesssim x \lesssim 0.5$ which shows a shift by little more than a quarter of standard deviation (though this could be a statistical fluctuation due to the size of the replica sample). At NNLO instead the effect of removing the NMC data altogether is clearly statistically significant on the isospin triplet and gluon, corresponding to a shift in central values at the level of almost one sigma for the gluon and more than half sigma for the isospin triplet. A one sigma change of the triplet uncertainty is also observed.

Some of these PDFs at NLO and NNLO are compared in Figs. 6 and 7 respectively, at a typical electroweak scale $Q^2 = 10^4 \text{ GeV}^2$. Differences at higher scale are somewhat reduced because of asymptotic freedom, but the general pattern observed in the distance plots is clearly reproduced: at NLO replacing NMC structure functions with cross-sections has no effect, while at NNLO it has an effect which is above the threshold of statistical significance, though smaller than the change that would be observed if the data changed by an amount compatible with their uncertainties. The effect of removing NMC data altogether, both at NLO and NNLO, is qualitatively similar but quantitatively somewhat larger.

We conclude that at NLO replacing structure functions with cross-sections or even




Figure 5: Same as Fig. 4 but at NNLO. Note that in this case the PDF set with NMC cross-section data (NNLO default) is used for the comparison with the PDFs with no NMC data.

removing NMC data altogether has no effect, while at NNLO replacing structure functions with cross-sections is just above the threshold of statistical significance, and removing them altogether statistically significant, though in all cases below the effect of a one sigma change of the data: this can be viewed as an upper bound on the possible impact of the treatment of this dataset.

The main implication of the study of Ref. [1], and the reason for the the ensuing debate, was that the Higgs production cross-section via gluon-gluon fusion may change as a consequence of the treatment of NMC data by an amount which may invalidate current Higgs exclusion limits. To verify what happens in our case, we have recomputed the Higgs production cross-section using the various PDF sets discussed here, using the code of Refs. [23, 24], for a range of Higgs masses between 100 and 400 GeV. We show results for the Tevatron and the LHC 7 TeV in Fig. 8; all uncertainties shown are 68% confidence levels. We see that the replacement of NMC structure functions by cross-sections has no impact on the Higgs production cross-section, and that even removing all NMC data leads to a shift much smaller than the nominal PDF uncertainties, with a slight increase of these uncertainties. Again, this can be viewed as a (conservative) estimate of the differences arising from the different treatments of the NMC data.

Let us finally compare our results with those of Ref. [1]. In that reference, the value




Figure 6: NLO PDFs determined using NMC cross-section data (long dashes) and no NMC data (short dashes) shown as ratio to the default NNPDF2.1 set (determined using NMC structure function data) at $Q^2 = 10^4$ GeV²: singlet $\Sigma(x, Q^2)$ (top, left), gluon $g(x, Q^2)$ (top, right), triplet $T_3(x, Q^2)$ (bottom, left) and total valence $V(x, Q^2)$ (bottom, right).

of α_s was determined together with the PDFs, and the best-fit α_s was found to change significantly according to the treatment of the NMC data. In particular, the change of the best-fit α_s value was found to be of order of 1.5 sigma at NLO and 2.3 sigma at NNLO, with an increase of the Higgs cross-section at the LHC by 4% (i.e. about one sigma) at NLO and 9% (i.e. about 2.7 sigma) at NNLO when the NMC cross-section data are replaced by structure function data. In order to assess quantitatively how much of this change in Higgs cross-section is just due to the different value of α_s a comparison between ABKM sets with fixed value of α_s but different treatment of NMC data would be necessary. These sets are at present not available. However, a simple estimate (which at NLO is in fact quite accurate [14]) can be obtained by noting that, based on the size of the NLO and NNLO K -factors one expects that a percentage change $\Delta\alpha$ in the value of α_s , if everything else is kept fixed, leads to a percentage shift of the Higgs cross-section $\Delta\sigma \approx 2.5\Delta\alpha$ at NLO and $\Delta\sigma \approx 2.8\Delta\alpha$ at NNLO. Based on this, one would estimate that about 90% of the cross-section increase seen in Ref. [1] at NLO when structure function data replace cross-section data and about 80% of the increase at NNLO, is just due to the change in value of α_s . The residual change, due to the PDFs, is still perhaps somewhat larger than that which we observe, but qualitatively more in line with it.

We conclude that we do not support the conclusion that the treatment of NMC data




Figure 7: NNLO PDFs determined using NMC structure function data (long dashes) and no NMC data (short dashes) shown as ratio to the default NNPDF2.1 set (determined using NMC cross-section data) at $Q^2 = 10^4 \text{ GeV}^2$: up $u(x, Q^2)$ (top, left), gluon $g(x, Q^2)$ (top, right), antineutrino $\bar{d}(x, Q^2)$ (bottom, left) and strange $s(x, Q^2)$ (bottom, right).

may affect the Higgs cross-section and thus exclusion limits in any significant way. Of course, if the value of α_s is varied by a very large amount, then the cross-section and ensuing limits are significantly affected. In this respect, it should be noticed that the best-fit $\alpha_s(M_z) = 0.1135$ value of Refs. [1, 8] at NNLO differs by more than 7 sigma from the PDG value $\alpha_s(M_z) = 0.1184$ [13], in units of the latter's uncertainty $\Delta\alpha_s = 0.0007$. The Higgs working group [17] following PDF4LHC [25], recommends to use a more conservative $\Delta\alpha_s = 0.0012$, but even so the value of Refs. [1, 8] differs by more than four sigma from the PDG average. A NLO determination of α_s based on the NNPDF2.1 PDF fit [26] leads to values of α_s which are in good agreement with the PDG value, both when the global dataset ($\alpha_s(M_z) = 0.1191$) and deep-inelastic data only ($\alpha_s(M_z) = 0.1177$) are used. Given that, as we have just shown, the treatment of NMC data has no statistically significant impact on the NLO analysis, it is exceedingly unlikely that the NNPDF NLO determination of α_s might depend on how the NMC data are treated. It will be interesting to repeat the analysis of Ref. [26] at NNLO. The stability of NNPDF2.1 PDFs when going from NLO to NNLO [7] suggests that results should not change dramatically.

In summary, we find that the effect of the treatment of NMC data on NNPDF2.1 PDFs is of no statistical significance at NLO, and just about statistically significant at NNLO though by at least a factor three smaller than the typical PDF uncertainty due to propagated data uncertainties. The effect on the Higgs production cross-section is accord-




Figure 8: The Higgs cross-section in gluon fusion at the Tevatron Run II (top) and at the LHC 7 TeV (bottom). Left: the reference NNPDF2.1 NLO (NMC structure functions) compared to the NLO fits with NMC cross-sections and no NMC data, shown as a ratio to the reference. Right: the reference NNPDF2.1 NNLO (NMC cross-sections) compared to the NNLO fits with NMC structure functions and no NMC data, shown as a ratio to the reference. All uncertainties shown are one sigma.

ingly negligible. Even removing NMC data altogether has a moderate effect on NNPDF2.1 PDFs, which even at NNLO remains below one sigma. The considerable stability of the NNPDF2.1 results is due both to the use of a very wide dataset which includes DIS, Drell-Yan, weak vector boson production and inclusive jet data, which reduces the dependence of our results on any particular dataset, and to the extremely flexible neural network parametrization which eliminates the parametrization bias which might otherwise lead to instabilities on small shifts in input data.

Acknowledgments

We thank S. Alekhin for correspondence and discussions on the matter of this paper. M.U. is supported by the Bundesministerium für Bildung und Forschung (BmBF) of the Federal Republic of Germany (project code 05H09PAE). This work was partly supported by the Spanish MEC FIS2007-60350 grant. We would like to acknowledge the use of the computing resources provided by the Black Forest Grid Initiative in Freiburg.

References

- [1] S. Alekhin, J. Blumlein and S. Moch, (2011), arXiv:1101.5261.
- [2] New Muon Collaboration, M. Arneodo et al., Nucl. Phys. B483 (1997) 3, hep-ph/9610231.
- [3] New Muon Collaboration, M. Arneodo et al., Nucl. Phys. B487 (1997) 3, hep-ex/9611022.
- [4] J. Baglio, A. Djouadi and R. Godbole, (2011), arXiv:1107.0281.
- [5] R. S. Thorne and G. Watt, (2011), arXiv:1106.5789.
- [6] The NNPDF Collaboration, R.D. Ball et al., Nucl. Phys. B849 (2011) 296, arXiv:1101.1300.
- [7] R. D. Ball *et al.* [The NNPDF Collaboration], (2011) arXiv:1107.2652.
- [8] S. Alekhin et al., Phys. Rev. D81 (2010) 014032, arXiv:0908.2766.
- [9] S. Forte, Acta Phys. Polon. B41 (2010) 2859, arXiv:1011.5247.
- [10] J. Huston, talk at the PDF4LHC workshop, CERN, March 2011
<http://indico.cern.ch/getFile.py/access?contribId=2&resId=1&materialId=slides&confId=127425>
- [11] L.W. Whitlow et al., Phys. Lett. B282 (1992) 475.
- [12] L.W. Whitlow et al, Phys. Lett. 250B (1990) 193.
- [13] S. Bethke, Eur. Phys. J. C64 (2009) 689, arXiv:0908.1135.
- [14] F. Demartin et al., Phys. Rev. D82 (2010) 014002, arXiv:1004.0962.
- [15] The NNPDF collaboration, R.D. Ball et al., Nucl. Phys. B838 (2010) 136, 1002.4407.
- [16] S. Alekhin et al., (2011), arXiv:1101.0536.
- [17] LHC Higgs Cross Section Working Group, S. Dittmaier et al., (2011), arXiv:1101.0593.
- [18] The NNPDF collaboration, L. Del Debbio et al., JHEP 03 (2007) 039, hep-ph/0701127.
- [19] The NNPDF Collaboration, R.D. Ball et al., Nucl. Phys. B809 (2009) 1, arXiv:0808.1231.
- [20] The NNPDF collaboration, R.D. Ball et al., Nucl. Phys. B823 (2009) 195, arXiv:0906.1958.
- [21] The NNPDF collaboration, R.D. Ball et al., JHEP 05 (2010) 075, arXiv:0912.2276.
- [22] S. Forte et al., JHEP 05 (2002) 062, hep-ph/0204232.

- [23] R. Bonciani, G. Degrassi and A. Vicini, JHEP 11 (2007) 095, arXiv:0709.4227.
- [24] U. Aglietti et al., JHEP 01 (2007) 021, hep-ph/0611266.
- [25] M. Botje *et al.*, (2011) 1101.0538.
- [26] S. Lionetti et al., Phys. Lett. B701 (2011) 346, arXiv:1103.2369.