



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Optimizing mating schemes in fish breeding

Citation for published version:

D'Agaro, E, Woolliams, J, Haley, C & Lanari, D 2007, 'Optimizing mating schemes in fish breeding' Italian Journal of Animal Science, ???volume??? 6, ???pages??? 795-796., 10.4081/ijas.2007.1s.795

Digital Object Identifier (DOI):

[10.4081/ijas.2007.1s.795](https://doi.org/10.4081/ijas.2007.1s.795)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher final version (usually the publisher pdf)

Published In:

Italian Journal of Animal Science

Publisher Rights Statement:

Copyright 2007 PAGEPress®

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Optimizing mating schemes in fish breeding

E. D'Agaro¹, J. A. Woolliams², C. S. Haley², D. Lanari¹

¹ Dipartimento di Scienze Animali. Università di Udine, Italy

² Roslin Institute. Edinburgh, UK

Corresponding author: Edo D'Agaro. Dipartimento di Scienze Animali, Università di Udine, Via S. Mauro 2, Pagnacco (UD) - Tel. +39 0432650110 - Fax: +39 0432660614 - Email: dagaro@dspa.uniud.it

ABSTRACT: The optimal contribution selection method and the simulated annealing algorithm were used to maximize the genetic gain and reduce inbreeding in fish breeding populations. This study considered the following mating designs: full factorial (3 x 3); full factorial (6 x 6) and nested (6 males x 18 females). A quantitative trait based on a strictly additive and polygenic model was simulated. Two levels for the number of genotyped offspring (360 or 720) and three levels of heritability (0.1; 0.3; 0.5) were assumed. The best results in terms of ΔF and ΔG were obtained with the full factorial design (6 x 6) and considering a trait with a high heritability. The optimal family size was found at 20 fish per mating.

Key words: Simulation, Fish, optimum contribution, Annealing algorithm.

INTRODUCTION – The design of optimal mating schemes is a mean to improve fish performance worldwide. During the last two decades, breeding strategies and techniques addressing both genetic improvement has been well documented and applied in several countries (Sonesson *et al.*, 2005). Nevertheless, in Italy, selective breeding has not been widely practiced in fish farming. Recent advances in fish breeding involve comparisons and optimization of mating designs using advanced analytical models as the optimum contribution selection (Meuwissen, 1997) and the simulated annealing algorithm (Press *et al.*, 1989). The high reproductive potential in fish species allows high genetic gains by applying high selection intensities. This means that a very small number of individuals are used to generate successive generations and hence the rate of inbreeding can be high (Sonesson *et al.*, 2005). The detrimental effects of inbreeding are well documented in fish. In recent years, many selection and mating strategies have been proposed to restrict inbreeding in selection programmes (Sonesson *et al.*, 2005). In this study, a stochastic simulation was used to optimize mating schemes using different genetic parameters and under restricted inbreeding in fish breeding.

MATERIAL AND METHODS – A fish population under artificial selection was modelled by stochastic (Monte Carlo) simulation using Matlab software. Selection was applied for a single trait measured on both sexes and based on estimated breeding values (EBVs) using the ASREML2 statistical package. Generation were discrete (equal number of sires and dams were selected at each generation). The trait under selection was assumed to be determined by an infinite number of unlinked additive loci, each with an infinitesimal effect. The trait was considered to be standardized, so the initial phenotypic variance is unity. Phenotypes of unrelated base population animals (generation 0) were generated as the sum of a normally distributed environmental and genetic effects. Phenotypic values of the offspring born every generation were generated as: $P = \mu + (\sigma_A \text{RND}(0,1)_S + \sigma_A \text{RND}(0,1)_D) / 2 + \sigma_E \text{RND}(0,1) + (0.5(1 - (F_s + F_d) / 2))^{1/2} \sigma_A \text{RND}(0,1)$ where $\sigma_{A(t)} = \sigma_{A(0)} / (1 + kh^2_{(t)})$ and $k = (0.5)(k_m + k_f)$; $k_y = i_y(i_y - x_y)$ $y = \text{male or female}$. Selection was directional upwards and by truncation. Total number of offspring born per generation and numbers of selected males and females were constant (10 or 20 offspring per mating) over generations and varied according to the mating schemes. The simulated breeding schemes are described in Table 1.

Table 1. Parameters of the closed nucleus scheme.

Number of selection candidates per generation	0 or 180 or 360 or 720
Number of generations	10
Number of replicated simulations	100
Mating schemes	factorial 3 x 3; 6 x 6; nested 6 (males) x 18 (females)
Heritability	0.1 or 0.3 or 0.5

The method of Meuwissen (1997) was used to select animals. This method maximises the genetic level of the next generation of animals $G_{t+1} = c'EBV_t$ where c_t is a vector of genetic contributions of selected candidates to generation t_{+1} and EBVs is a vector of best linear unbiased prediction (BLUP) estimates of candidates in generation t . Rates of inbreeding are controlled by constraining the average coancestry to $C_{t+1} = cA_t c_t / 2$ where A_t is a $(n \times n)$ relationship matrix among n selection candidates $C_{t+1} = 1 - (1 - \Delta F)^t$ and ΔF_d is the desired rate of inbreeding. For optimizing of c_t , it was also necessary to constrain the sum the contributions of males (females) to $1/2$. In order to obtain the optimal c_t that maximize G_{t+1} Lagrangian multipliers were used. An additional restriction was to select only one full sibs per family. The minimum coancestry mating (reduce the average relationship of sires and dams and therefore also the inbreeding of their progeny is minimized) was obtained by applying the simulated annealing algorithm according to Press *et al.* (1989)

RESULTS AND CONCLUSIONS – The rate of inbreeding (ΔF) and genetic gain (ΔG) for the three mating designs are reported in Table 2.

Table 2. Rate of inbreeding (ΔF)(x100) and genetic gain (DG))(σ_p) for different mating schemes and genetic parameters.

		N. of offsprings per family				
		10			20	
Heritability	0.1	0.3	0.5	0.1	0.3	0.5
Full factorial (3 x 3)						
ΔF	3.96	5.00	4.55	4.50	4.70	4.65
ΔG	0.29	0.60	0.75	0.37	0.60	0.97
Full factorial (6 x 6)						
ΔF	2.25	2.20	2.10	2.05	1.80	1.85
ΔG	0.38	0.64	0.76	0.47	0.69	0.94
Nested (6males x 18 females)						
ΔF	2.60	2.25	2.36	2.06	2.90	2.05
ΔG	0.43	0.72	0.86	0.57	0.66	0.89

The full factorial design gives the best results in terms of ΔF and ΔG (1.85 and 0.94 or 1.80 and 0.69) for a maximal number of sires and dams (6 x 6), higher size of families per mating used for selection (720 offsprings) and higher heritability (0.3 or 0.5). According to Sorensen *et al.* (2005) the superiority of the factorial mating compared to hierarchical scheme can be explained in terms of the different genetic structure of populations obtained showing, in the factorial design, small full-sibs families, more paternal half-sibs and a group of maternal half-sibs. At a lower heritability (0.1) the nested design become competitive with the full factorial mating (6 x 6). In conclusion, the optimal design to estimate additive effects in fish seems to be a full factorial mating with full-sibs families of 20 animals.

REFERENCES – Meuwissen, T.H.E., 1997. Maximizing the response of selection. with a predefined rate of inbreeding. *J. Anim.Sci.* 75:924-940. Press, W.H., Flannery B.P., Teukolsky, S.A., Vetterling, W.T., 1989. Minimization or maximization of function. In: Numerical recipes. The act of scientific computing. University of Cambridge Press, pp. 274-334. Sonesson, A.K., Gjerde, B., Meuwissen, T.H.E., 2005. Truncation selection of BLUP-EBV and phenotypic values in fish breeding schemes. *Aquaculture* 243:61-68. Sorensen, A.C., Berg, P., Woolliams, J.A., 2005. The advantage of factorial mating under selection is uncovered by deterministically predicted rates of inbreeding. *Genet. Sel. Evol.* 37:57-81.