



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Changes in photoperiod or temperature alter the functional relationships between phytochromes and reveal roles for phyD and phyE

Citation for published version:

Halliday, KJ & Whitelam, GC 2003, 'Changes in photoperiod or temperature alter the functional relationships between phytochromes and reveal roles for phyD and phyE' *Plant physiology*, vol 131, no. 4, pp. 1913-20., 10.1104/pp.102.018135

Digital Object Identifier (DOI):

[10.1104/pp.102.018135](https://doi.org/10.1104/pp.102.018135)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher final version (usually the publisher pdf)

Published In:

Plant physiology

Publisher Rights Statement:

RoMEO green

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Changes in Photoperiod or Temperature Alter the Functional Relationships between Phytochromes and Reveal Roles for phyD and phyE¹

Karen J. Halliday* and Garry C. Whitelam

School of Biological Sciences, University of Bristol, Bristol BS8 1UG, United Kingdom (K.J.H.); and Biology Department, University of Leicester, Leicester LE1 7RH, United Kingdom (G.C.W.)

The phytochromes are one of the means via which plants obtain information about their immediate environment and the changing seasons. Phytochromes have important roles in developmental events such as the switch to flowering, the timing of which can be crucial for the reproductive success of the plant. Analysis of *phyB* mutants has revealed that *phyB* plays a major role in this process. We have recently shown, however, that the flowering phenotype of the *phyB* monogenic mutant is temperature dependent. A modest reduction in temperature to 16°C was sufficient to abolish the *phyB* mutant early-flowering phenotype present at 22°C. Using mutants null for one or more phytochrome species, we have now shown that *phyA*, *phyD*, and *phyE*, play greater roles with respect to *phyB* in the control of flowering under cooler conditions. This change in the relative contributions of individual phytochromes appears to be important for maintaining control of flowering in response to modest alterations in ambient temperature. We demonstrate that changes in ambient temperature or photoperiod can alter the hierarchy and/or the functional relationships between phytochrome species. These experiments reveal new roles for *phyD* and *phyE* and provide valuable insights into how the phytochromes help to maintain development in the natural environment.

Plant growth and development is intimately linked to external cues that signal changes in the environment. Alterations in light quality, quantity, and duration provide the plant with information that accurately reflects changes in both local environment and the changing seasons. To detect and respond to these different light signals, plants have evolved a series of highly specialized photoreceptors. This photoreceptor system includes the red (R) and far-red (FR) light-absorbing phytochromes and the blue/UV-A light-absorbing cryptochromes and phototropins (Whitelam et al., 1998).

The *Arabidopsis* phytochromes comprise the products of a family of five closely related genes, designated *PHYA* through *PHYE* (Mathews and Sharrock, 1997). The photosensory activity of the phytochromes resides in their unique capacity for reversible light-induced interconversion between a R light-absorbing Pr form and a FR light-absorbing Pfr form. Light-triggered Pfr formation also induces cytosolic to nuclear translocation and the activation of signaling via molecular interaction (Kircher et al., 2002; Quail, 2002). In the nucleus, *phyA* and *phyB* interact directly with PIF3 and *phyB* interacts with PIF4 to regulate transcription (Ni et al., 1998, 1999; Martinez-Garcia et al., 2000; Huq and Quail, 2002). Direct in-

teraction with ZTL/ADO1, ELF3, and COP1 provides a means for *phyB* to connect with the circadian clock and activate the de-etiolation switch (Jarillo et al., 2001; Liu et al., 2001; Yang et al., 2001). *phyA* and *phyB* interact with PKS1 and *phyA* with NDPK2 in the cytosol (Choi et al., 1999; Fankhauser et al., 1999). Furthermore, interactions have been demonstrated between *phyA* and *phyB* with *cry1* and *cry2*, respectively (Ahmad et al., 1998; Mas et al., 2000). This may be the means via which at least some of the reported physiological interactions between *phyA/phyB* and *cry1/cry2* occur (Casal and Mazzella, 1998; Neff and Chory, 1998; Mockler et al., 1999).

It is now well established that individual photoreceptors do not act in isolation, but as an interconnected network (Casal, 2002; Nagy and Schafer, 2002). Analysis of mutants null for one or more photoreceptors grown under specific conditions has provided valuable insights into how the photoreceptor network operates within the natural environment. Complex interactions that involve *phyA*, *phyB*, *phyD*, *cry1*, and *cry2* have been described for de-etiolation (Casal, 1995; Casal and Boccacandro, 1995; Casal and Mazzella, 1998; Neff and Chory, 1998; Hennig et al., 1999, 2001; Mazzella et al., 2001). The impact of the *cry1* and *cry2* mutations on *Lhcb*2* promoter-*gusA* expression was shown to be markedly affected by the absence of *phyA* and *phyB* (Mazzella et al., 2001). Furthermore, functional interaction between *phyA*, *phyB* and *cry1* was shown for accumulation of chlorophyll and anthocyanin (Neff and Chory, 1998; Hennig et al., 2001). Flowering is also

¹ This work was supported by the Biotechnology and Biological Science Research Council (UK).

* Corresponding author; e-mail k.j.halliday@bristol.ac.uk; fax 44-117-925-7374.

Article, publication date, and citation information can be found at www.plantphysiol.org/cgi/doi/10.1104/pp.102.018135.

subject to strong regulatory control by the photoreceptors. We now have evidence that *phyA*, *phyB*, *phyD*, *phyE*, *cry1*, and *cry2* regulate flowering through an interconnected network (Devlin et al., 1998, 1999; Mockler et al., 1999).

Temperature is also an important environmental cue in the regulation of flowering. Many plants have adopted a reproductive strategy that requires long periods of cold (1°C–10°C) to promote flowering. This strategy ensures that flowering does not occur in winter months but instead in the more favorable spring climate (Simpson and Dean, 2002). We have recently demonstrated that ambient temperature is a significant modulator of photoreceptor action in the control of flowering (Halliday et al., 2002). A modest reduction in growth temperature, from 22°C to 16°C, completely abolished the *phyB* mutant early-flowering phenotype frequently observed at higher temperatures. Thus, small changes in ambient temperature can have a large impact on photoreceptor action. These light- and temperature-controlled flowering pathways appear to regulate expression of *FT*, a known convergence point for the photoperiod and vernalization pathways (Halliday et al., 2002; Hepworth et al., 2002; Simpson and Dean, 2002; Yanovsky and Kay, 2002; Izawa et al., 2002). Therefore, *FT* (together with *LFY* and *SOC1/AGL20*) is an important integration point for multiple flowering pathways.

Studies to date have demonstrated roles for *phyD* and *phyE* in a range of developmental processes including germination, seedling establishment, elongation, and flowering responses to end-of-day-FR and low R/FR ratio light (Aukerman et al., 1997; Devlin et al., 1998, 1999; Hennig et al., 1999, 2002). For many of these responses, *phyD* and *phyE* have been shown to have redundant roles. However, our earlier studies suggest that in some instances, redundancy of action for an individual phytochrome may simply reflect suboptimal conditions for the particular phytochrome-mediated response. We have conducted a series of experiments that illustrate that changes in the photoperiod and temperature, important environmental cues, change the hierarchy of phytochrome action, revealing prominent roles for *phyD* and *phyE* in the natural environment. These experiments also highlight important changes in the functional relationships between the phytochromes that underlie developmental plasticity.

RESULTS

In SDs the *phyE* Monogenic Mutant Is Early Flowering

Consistent with earlier studies, when grown under 8-h photoperiods (SDs), the vegetative morphology of the monogenic *phyE* mutant was similar to that of the wild type (Devlin et al., 1998). However, under our growth conditions (photon irradiance 180 $\mu\text{mol m}^{-2} \text{s}^{-1}$), the *phyE* mutant flowered consistently ear-

lier than the wild type, both in terms of rosette leaf number and time to bolting (Fig. 1A; data not shown). The statistical significances for the wild type versus *phyE* and all other pairwise genotype comparisons were calculated using the Bonferroni multiple comparisons test (Fig. 1, B and D). As previously reported, the *phyA* mutant flowered slightly later than the wild type (Johnson et al., 1994; Neff and Chory, 1998). However, in SDs, plants null for both *phyA* and *phyE* flowered earlier than the monogenic *phyE* mutant (Fig. 1, A and B). This suggests an interaction of *phyA*- and *phyE*-mediated signaling in the control of flowering under SDs. The *phyD* mutant produced very slightly fewer rosette leaves than the wild type at bolting, whereas the *phyAphyD* double mutant flowered earlier than the *phyD* monogenic mutant. As for *phyE*, this suggests an interaction between the *phyA* and *phyD* mutations under SDs.

At 16°C *phyB* Acts Redundantly to Control Flowering

When grown at 16°C, the growth of wild-type plants is slower compared with plants maintained at 22°C (Halliday et al., 2002). However, the vegetative developmental phase is only slightly extended because these plants consistently produce only about four more leaves under these cooler conditions (Fig. 1A). Under SDs, the early-flowering phenotype of *phyE* was maintained at both 22°C and 16°C, although its severity was slightly reduced at the cooler temperature (Fig. 1, A and B). Likewise, the flowering responses of *phyA*, *phyD*, *phyA phyD*, and *phyA*




Figure 1. Flowering time in SDs and LDs. A, Plants were grown in SDs at either 22°C or 16°C. C, Plants were grown in LDs at 16°C. Rosette leaf number was determined at bolting (photon irradiance, 400–700 nm, 180 $\mu\text{mol m}^{-2} \text{s}^{-1}$). Bars represent the SE. WT, Laer wild type; A⁻, *phyA*; D⁻, *phyD*; and E⁻, *phyE* null mutations. B and D, Statistical significance of differences in flowering time. Pairwise comparisons for genotypes were undertaken using the Bonferroni multiple comparisons test.

phyE mutants relative to the wild type were similar under both temperature regimes. Thus, the changes in flowering time imposed by *phyE*, *phyA*, and *phyD* mutations were not markedly altered in the 16°C to 22°C temperature range. We have recently demonstrated that the *phyB* mutant flowers at the same time as the wild type under 16°C (Halliday et al., 2002). Collectively, these results suggest more prominent roles for *phyE*, *phyA*, and *phyD* in the regulation of flowering under cooler conditions. Although the monogenic *phyB* mutant is not early flowering when grown at 16°C, the *phyB* null allele does lead to accelerated flowering in the *phyE* or *phyD* backgrounds in both 16-h photoperiods (LDs) and SDs (Figs. 1, C and D, and 3). These data demonstrate synergistic interactions between *phyD* and *phyE* with *phyB*. This suggests that although *phyB* has a more minor role in repressing flowering at 16°C than it does at 22°C, it still exerts a degree of control on flowering at the cooler temperatures via synergistic interactions with other phytochromes.

Under LDs at 16°C, the monogenic *phyE* and *phyD* mutants flowered with a similar number of rosette leaves to the wild type (Fig. 1, C and D). In contrast, the late-flowering phenotype of the *phyA* mutant was retained under these conditions. This suggests that under cool LDs, the hierarchy changes such that *phyA* has a more prominent role, with respect to *phyE* and *phyD* in the control of flowering.

In LDs *phyE* Is Epistatic to *phyA* in the Control of Flowering Time

When grown under LDs, the *phyAphyD* double mutant flowered significantly earlier than the wild type (Fig. 1C). Monogenic *phyD* flowered at the same time, and monogenic *phyA* flowered later than the wild type, suggesting a functional interaction between *phyA* and *phyD* in the control of flowering. A similar relationship for *phyA* and *phyD* and for *phyA* and *phyE* was observed under SDs (Fig. 1A; see above). In contrast, under LDs, impact of the *phyA* mutation in a *phyE* background was negligible at both 22°C (data not shown) and 16°C (Fig. 1, C and D). Under LDs, plants carrying the *phyA* and *phyE* mutations flowered at the same time as the *phyE* mutant. Furthermore, the *phyAphyBphyE* and *phyAphyBphyDphyE* mutants flowered at the same times as *phyBphyE* and *phyBphyDphyE*, respectively. These data suggest that under LDs *phyE* is required for the *phyA* mutant phenotype.

The Monogenic *phyD* Mutant Has Reduced Leaf Size in LDs

Like *phyE*, *phyD* has been shown to act redundantly with *phyB* to control leaf shape (Devlin et al., 1998, 1999). However, we have shown that small adjustments in temperature reveal a striking leaf phe-

notype in the monogenic *phyD* mutant. When grown in LDs under cooler conditions (16°C), *phyD* produced markedly smaller leaves than the wild type, revealing a new role for *phyD* in the promotion of leaf expansion (Fig. 2, A and B). Removal of *phyA* in addition to *phyD* restored much of the wild-type phenotype, suggesting that *phyA* was required for the monogenic *phyD* mutant phenotype. This phenotype is not only temperature conditional, it is also photoperiod dependent. When grown under SDs, the rosette diameter of *phyD* was very similar to the wild type. Under these conditions, *phyD* leaf area was slightly smaller than the wild type; however, the removal of *phyA* in addition to *phyD* completely restored the wild-type phenotype (Fig. 2, A and B). Taken together, these data suggest that the role of *phyD* in controlling leaf development is photoperiod and temperature conditional. Furthermore, *phyA* appears to have a role in moderating *phyD* action in this response.

phyD Slows Rosette Leaf Formation Rate

Earlier work established a prominent role for *phyB* in controlling the rate of rosette leaf production (Mazzella et al., 2001; Halliday et al., 2002). Analysis




Figure 2. Basal rosette diameter in LDs and SDs. A, Laer WT, and *phyD*, *phyAphyD* mutants grown in LDs or SDs for 28 d. B, Basal rosette leaf diameter (centimeters) was determined for 28-d-old plants grown in LDs or SDs (photon irradiance, 400–700 nm, 180 $\mu\text{mol m}^{-2} \text{s}^{-1}$) at 16°C. Bars represent the SE. WT, Laer wild type; A⁻, *phyA*; and D⁻, *phyD*.

of the monogenic *phyD* mutant revealed a role for *phyD* in the control of rosette leaf production rate throughout vegetative development, but one that gains prominence in the second half of the vegetative phase. When grown in SDs at 16°C, the first seven to eight leaves were produced at a similar rate in the *phyD* mutant and the wild type, thereafter in *phyD*, leaf production slowed (Fig. 3A). A further slowing of leaf production was observed during the final third of the developmental phase. These data are consistent with our recent analysis of the *phyAphyB-phyD* mutant that suggested this role for *phyD* in the second half of the vegetative developmental phase (Halliday et al., 2002). As for the *phyB* mutant, the *phyD* phenotype was seen at both 22°C and 16°C (Halliday et al., 2002; data not shown). This phenotype contrasts with that of *phyA* and *phyE*, both of which produce leaves at a wild-type rate (Halliday et al., 2002; data not shown). Removal of *phyB* in addition to *phyD* slowed leaf production further (Fig. 3, A and B). Leaf production of mutants null for *phyB*, *phyD*, and *phyE* was very severely retarded. On occasion, growth was more severely disrupted in *phyBphyDphyE* mutants, these plants appeared pale and sickly and developed necrotic lesions (data not shown). We have not observed these phenotypic traits in our *phyAphyBphyD* or *phyAphyBphyE* triple mutants, which may reflect the relative importance of *phyB*, *phyD* and *phyE* for normal vegetative development.




Figure 3. Rosette leaf production rate in 16°C SDs. A, Rosette leaf number was counted at time intervals (days) until flowering time in plants grown at 16°C in SDs (photon irradiance, 400–700 nm, 180 $\mu\text{mol m}^{-2} \text{s}^{-1}$). Bars represent the SE. WT, Laer wild type; B⁻, *phyB*; D⁻, *phyD*; and E⁻, *phyE* null mutations. B, Laer WT, and *phyBphyD*, *phyBphyDphyE* mutants grown in SDs for 46 d.




Figure 4. Temperature-dependent internode elongation. A, The *phyAphyBphyE* triple mutant, grown at 16°C (left) and 21°C (right) at photon irradiance, 400 to 700 nm, 100 $\mu\text{mol m}^{-2} \text{s}^{-1}$. B, Rosette internode length (millimeters) of 21°C-grown wild type and *phyA-phyBphyE*. Plants were grown in SDs for these experiments; bars represent the SE. C, Wild-type seedlings grown in the dark at 16°C and 26°C. Fresh and electron micrograph images were taken of seedlings grown on 3% (w/v) Suc for 3 weeks.

The Elongated Internode Phenotype of *phyAphyBphyE* Is Temperature Dependent

The identification of phytochrome controlled internode elongation, and flowering responses in the *phyAphyB* double mutant provided the basis of a screen that identified the *phyE* null mutation (Devlin et al., 1996, 1998). The constitutively early flowering and elongated internode phenotype of the *phyAphyB-phyE* mutant provided evidence for the role of *phyE* in these aspects of photomorphogenesis. However, in a similar fashion to mutants lacking *phyA*, *phyB*, and *cry1*, the elongated internode phenotype of the *phyA-phyBphyE* mutant is only evident if plants are grown at an inductive temperature (Mazzella et al., 2000). When *phyAphyBphyE* plants were grown under SDs at 16°C, the mutant exhibited a normal rosette habit, whereas growth at or above 22°C resulted in the *phyAphyBphyE* mutant producing distinct internodes (Fig. 4, A and B; Devlin et al., 1998). The elongated internode phenotype was not observed in double mutant combinations of *phyA*, *phyB*, or *phyE* under these conditions. We therefore reasoned that under warmer growth conditions, internode elongation may be the default situation and that *phyE* (together with *phyA* and *phyB*) inhibit this elongation to maintain the rosette growth habit. To test this, wild-type seedlings were grown in darkness on vertically oriented Suc-containing plates at either 16°C or 26°C. Suc availability in the aerial part of the plant is

known to promote seedling development in the absence of photoreceptor action (Roldan et al., 1999). Although the seedlings exhibited an elongated growth habit under both temperature regimes, internodes were only elongated in seedlings grown at the warmer temperature (Fig. 4C). This suggests that at permissive temperature, internode elongation is the default position and that phyE, phyB and phyA are important for maintaining the compact rosette habit under these conditions.

DISCUSSION

We set out to gain further insights into how the phytochrome network controls development in the natural environment. By growing plants deficient in one or more phytochrome species under different photoperiods and temperatures, we have been able to establish new roles for phyD and phyE. We have also demonstrated that changes in photoperiod and temperature dramatically alter functional relationships between phytochrome species. A growing body of evidence suggests that this is the means via which the photoreceptor system manipulates development in response to changed environmental conditions. We show this also provides a mechanism for photoreceptors to maintain developmental stability under different ambient temperatures.

The Early-Flowering Phenotype of *phyE* Is Specific to SDs

PhyB has been shown to be an important regulator of flowering time in response to light quality and photoperiod (Whitelam et al., 1998; Salome et al., 2002). We have shown that in SDs, like *phyB*, the monogenic *phyE* mutant also flowers early. Furthermore, we did not observe this phenotype under LDs, which suggests the early-flowering phenotype of *phyE* is specific to SDs. Thus it appears that under SDs, phyB and phyE play major roles in regulating flowering time. Devlin and co-workers (1998) previously described the phyE phenotype as wild type, however, this apparent contradiction may simply reflect the comparatively high-light levels used in our experiments. The apparent specificity of phyE action to SDs may occur as an indirect consequence of the short photoperiod. Alternatively, this may represent a mechanism via which light interacts with the circadian system to delay flowering under SDs.

The *phyD* mutant flowered slightly earlier than the wild type under SDs. Again like *phyE*, this effect was not observed under LDs. The enhanced effect of the *phyE* and to a lesser extent the monogenic *phyD* mutations under SDs may reflect more influential roles for phyD and phyE under shorter photoperiods in the inhibition of flowering. Although the effects of the monogenic *phyD* and *phyE* mutations were not severe, in a *phyA* background, they had a larger

impact. In SDs, we have shown the genetic interactions of *phyD* and *phyE* with *phyA* are synergistic. This genetic relationship of phytochrome genes enables specific modification of flowering when phyA signaling is perturbed in addition to phyD or phyE. Because phyA, phyD, and phyE are differentially regulated by light and exhibit different action kinetics, this may be a means for the plant to distinguish and respond to a simultaneous change in two or more parameters in the light environment (Eichenberg et al., 2000; Kircher et al., 2002). This type of mechanism may facilitate acceleration of flowering in response to neighboring vegetation. Under these circumstances, fluence rates are high (degrading phyA), but the light reflected from the potential competitors is FR-enhanced, lowering the proportion of active phyD or phyE. This type of signaling provides plants with a means to interpret and process complex changes in the light environment.

Photoperiod Affects the Functional Relationship of *phyA* and *phyE* in the Control of Flowering

The synergistic relationship of *phyA* and *phyE* in the control of flowering observed in SDs was not observed under LDs. In LDs, we have shown that *phyE* is epistatic to *phyA* in this response. These data suggest that the length of the photoperiod has a significant impact on how the phyA and phyE pathways interact. Under SDs, the phyA- and phyE-signaling pathways are functionally distinct, whereas under LDs phyE is necessary for phyA action. One could speculate that altering the functional relationships of phyA and phyE in this way provides one route via which flowering can be adjusted in response to the prevailing photoperiod. For example, in LDs, the absence of both phyA and phyE had practically no effect on flowering time, whereas in SDs, *phyAphyE* was early flowering. Thus, the combined action of the phyA and phyE appears to be inhibitory under SDs, conditions that delay flowering in the wild type. Conditional synergism has previously been demonstrated for *cry1* and *phyB* in the control of hypocotyl length (Casal and Mazzella, 1998). They demonstrated that in saturating light conditions, phyB and *cry1* acted independently, but under conditions that were non-saturating for either phyB or *cry1* action, they acted synergistically. These types of experiments illustrate how changes in the light environment can dramatically change the functional relationship between photoreceptors. Our data suggest that photoperiod-mediated changes in the functional relationship between phyA and phyE may contribute to the changes in flowering time observed in different photoperiods.

At 16°C, phyE and phyD Have More Prominent Roles in the Control of Flowering

The early-flowering phenotype of the monogenic *phyB* mutant is well known (Whitelam et al., 1998).

However, we have recently shown that this phenotype is abolished when plants are grown at 16°C, a typical summertime temperature in a range of northern latitudes (Halliday et al., 2002). The data in this paper demonstrate that *phyB* does have a role in the control of flowering under cooler conditions, but its role is redundant in the presence of *phyD* and *phyE*. Under LDs at 16°C, the monogenic *phyD* and *phyE* mutations had no impact on flowering time, however, loss of *phyB* in addition to *phyD* or *phyE* accelerated flowering. This suggests that under LDs at 16°C, *phyD*, *phyE*, and *phyB* have largely redundant roles in the control of flowering, however, the interaction between *phyB* and *phyD* or *phyE* was synergistic. Under SDs at 16°C, a redundant role was still observed for *phyB*, however, the *phyE* and, to a lesser extent, the monogenic *phyD* mutations accelerated flowering. Thus, at lower temperatures, *phyE* and *phyD* have more prominent roles in the control of flowering under SDs.

We observed that the late-flowering phenotype of the monogenic *phyA* mutant was retained at 16°C under both SDs and LDs. Thus, *phyA* appears to have a more prominent role in LDs, but shares prominence with *phyE* and *phyD* in SDs at cooler temperatures. Our recent work demonstrated that wild-type plants display a normal early-flowering response to low R/FR ratio at 16°C (Halliday et al., 2002). This, together with our current findings suggests that *phyB* takes the principle role under warmer conditions, however, the action of *phyA*, *phyE*, and *phyD* gain importance under cooler conditions. This change in the hierarchy of phytochrome action at 16°C maintains phytochrome control of flowering under these conditions. This type of accommodative action or “developmental canalization” has been proposed for *phyA*, *phyB*, *cry1*, and *cry2* in the control of seedling de-etiolation (Mazzella et al., 2001). This type of complex, highly connected, and yet plastic network is thought to be essential for normal development as it buffers both environmental change and genetic variation (Stearns, 2002). Our observations are interesting in context with recent findings that a drop in temperature from 23°C to 16°C enhanced the late flowering phenotype of *cry2* considerably (Blazquez et al., 2003). Therefore, like *phyB*, the *cry2* phenotype is also very sensitive to changes in temperature. However, in contrast to *phyB*, *cry2* action appears to be enhanced under cooler conditions.

Photoperiod and Temperature Affect the Role of *phyD* in the Control of Leaf Expansion

When grown in LDs at 16°C, the *phyD* mutant rosette leaves were notably smaller than those of the wild type. However, monogenic *phyD* mutant rosettes had a wild-type appearance under SDs and warmer LD conditions (Aukerman et al., 1997; Devlin et al., 1999). These data suggest that this rather striking

phyD phenotype is dependent upon both photoperiod and temperature. Furthermore, while under permissive conditions, the *phyD* mutation inhibits leaf expansion; the removal of *phyA* in addition to *phyD* greatly attenuates this response. This suggests that *phyA* is required for the *phyD* small rosette phenotype. Recent work has demonstrated that *phyD* acts redundantly with *phyB* in the inhibition of leaf elongation when plants are grown in either LDs or SDs under warmer conditions (Aukerman et al., 1997; Devlin et al., 1999). In contrast, under cool LDs, *phyD* appears to be important for promotion of leaf blade expansion. The ecological significance of this finding is not clear, however, under these conditions, *phyD* appears to have an opposing action to *phyB* in the control of leaf shape.

phyD Controls the Rate of Rosette Leaf Formation

Recent reports have shown that the *phyB* mutation severely affects the rate of rosette leaf production (Mazzella et al., 2001; Halliday et al., 2002). Our recent analysis of the *phyB*, *phyAphyB*, and *phyAphyBphyD* suggested that *phyD* also regulated leaf production rate, but only in the second half of the vegetative phase (Halliday et al., 2002). Analysis of the monogenic *phyD* mutant revealed that *phyD* contributes to the control of rosette leaf production throughout development. However, its role was greatest in the final third of the vegetative phase. Thus, both *phyB* and *phyD* control the rate of rosette leaf formation, but their relative contributions are dependent on the developmental phase. These phytochrome-mediated effects are clearly a means of adjusting leaf production to suit the prevailing light environment. Such a strategy may be important when resources are limited, for example, under conditions of heavy vegetation shade.

The Elongated Phenotype of the *phyAphyBphyE* Mutant Is Temperature Dependent

Earlier work by Mazzella et al. (2000) demonstrated that the elongated internode phenotype of *phyB*, *phyAphyB*, *phyBcry1*, and *phyAphyBcry1* mutants grown in continuous white light was a temperature-dependent phenomenon. Our experiments provide evidence that the elongated internode phenotype *phyAphyBphyE* is also temperature dependent. When grown under SDs at 16°C, *phyAphyBphyE* grew with a compact rosette, whereas at 22°C, internodes were clearly visible. Because double mutant combinations of *phyA*, *phyB*, and *phyE* did not produce internodes under our conditions, it appears that *phyA*, *phyB*, and *phyE* act redundantly to maintain the basal rosette during development. These data are consistent with previous data that demonstrate roles for *phyA*, *phyB*, and *cry1* in this respect (Mazzella et al., 2001). Because multiple photoreceptors appear to suppress

internode formation, we were interested to establish whether elongation was the default condition at warmer temperatures. To do this, we grew wild-type seedlings on Suc at 16°C and 26°C in darkness. These seedlings developed internodes at 26°C but not 16°C. These data are consistent with internode elongation being the default situation under warmer temperatures. When seedlings are grown in the light phyE, phyA, phyB, and cry1 act collectively to preserve the rosette growth habit.

Continually surveying their surroundings, the light receptors act as an integrated signaling network keeping development in tune with the environment. This complex task requires a flexible network that can both respond to and accommodate environmental change. The data presented in this paper provide a window into the complex light-signaling network that finely tunes development. Changes in the functional relationship between photoreceptors appear to be crucial for adjusting development in response to environmental cues such as photoperiod. However, they are also necessary for maintaining responses under varied environmental conditions. Changes in the hierarchy of phytochrome action under different ambient temperatures appear to be an important mechanism for maintaining control of flowering in the natural environment where temperatures fluctuate. Such accommodative behavior is an acknowledged characteristic of highly interconnected networks that act to buffer the effect of environmental or genetic perturbations (Casal, 2002; Stearns, 2002). Understanding the mechanisms that control both responsive and accommodative photoreceptor action will be one of our future challenges.

MATERIALS AND METHODS

Plant Material and Growth Conditions

In all of our experiments, we used *Arabidopsis* ecotype Landsberg *erecta* (Laer). Phytochrome mutant alleles were *phyA-2* (Whitelam et al., 1993), *phyB-1* (Koorneef et al., 1980), *phyD-1* (Aukerman et al., 1997), and *phyE-1* (Devlin et al., 1998). The *phyD-1* mutation is a naturally occurring allele found in the Wassilewskija ecotype, therefore, near-isogenic Laer *phyD-1* mutant lines were created by introgression of the *phyD-1* mutation into the Laer ecotype (Aukerman et al., 1997).

In each of the experiments, seeds were sown on 0.8% (w/v) Lehle medium (Lehle Seeds, Round Rock, TX), and stratified in darkness at 4°C for 5 d before transfer to SDs or LDs or at 16°C or 22°C. After a further 5 d, uniformly sized seedlings were transplanted to 5 × 5 × 5-cm pots containing a 3:1 compost:horticultural silver sand mix. Light was provided by L65/80W/30 warm-white fluorescent tubes (photon irradiance 400 to 700 nm, 180 μmol m⁻² s⁻¹; Osram Ltd., St. Helens, UK).

Seedlings in the dark internode elongation experiments were stratified and germinated as above, and then grown on 3% (w/v) Suc Murashige and Skoog medium in complete darkness for 3 weeks.

Fixation and Scanning of Tissue

A scanning electron microscope was used to obtain the close-up views of internodes. Samples were fixed in the fixing buffer 2% (w/v) glutaraldehyde in 30 mM sodium-cacodylate for 24 h. After three 10-min washes in fixing buffer, a secondary fix (1% [w/v] osmium in fixing buffer) was applied for a further 24 h followed again by three 10-min washes. Samples were then

dehydrated via 15-min soaks in each of the acetone series (v/v): 30%, 50%, 70%, 90%, and 100% × 3. After four 15-min exchanges through liquid CO₂, the samples were dried using a Balzers Critical Point Drier CPD030. Samples were mounted on aluminum stubs and sputter coated with gold/palladium to an approximate thickness of 673A in a Polaron SC7640. Images were collected by a scanning electron microscope (S-3000H, Hitachi, Tokyo).

Plant Growth Assays

For plants grown under SDs, rosette leaf counts were carried out twice a week. Leaves were counted only when the petiole was visible to the naked eye. Flowering time was recorded as primary rosette leaf number at inflorescence production. Rosette leaves were distinguished from axillary leaves on the basis of morphological differences. Rosette diameter was measured at the widest point with a ruler. For quantification of internode length, images were taken with a digital camera, and measurements were made using Sigma Scan software (SPSS Science Software UK Ltd., Woking, Surrey, UK).

Statistical Analysis

Statistical analysis was performed using ANOVA and the Bonferroni multiple comparisons test. For each experiment, pairwise comparisons were made between all relevant genotypes, a subset of which is shown in Figure 1, B and D.

ACKNOWLEDGMENT

We thank Wendy Stoddart for technical assistance.

Received November 26, 2002; returned for revision January 2, 2003; accepted January 2, 2003.

LITERATURE CITED

- Ahmad M, Jarillo JA, Smirnova O, Cashmore AR (1998) The CRY1 blue light photoreceptor of *Arabidopsis* interacts with phytochrome A in vitro. *Mol Cell* 1: 939–948
- Aukerman MJ, Hirschfeld M, Wester L, Weaver M, Clack T, Amasino RM, Sharrock RA (1997) A deletion in the PHYD gene of the *Arabidopsis* Wassilewskija ecotype defines a role for phytochrome D in red/far-red light sensing. *Plant Cell* 9: 1317–1326
- Blazquez MA, Ahn JH, Weigel D (2003) A thermosensory pathway controlling flowering time in *Arabidopsis thaliana*. *Nat Genet* 33: 168–171
- Casal JJ (1995) Coupling of phytochrome B to the control of hypocotyl growth in *Arabidopsis*. *Planta* 196: 23–29
- Casal JJ (2002) Environmental cues affecting development. *Curr Opin Plant Biol* 5: 37–42
- Casal JJ, Boccalandro H (1995) Co-action between phytochrome B and HY4 in *Arabidopsis thaliana*. *Planta* 197: 213–218
- Casal JJ, Mazzella MA (1998) Conditional synergism between cryptochrome 1 and phytochrome B is shown by the analysis of phyA, phyB, and hy4 simple, double, and triple mutants in *Arabidopsis*. *Plant Physiol* 118: 19–25
- Choi G, Yi H, Lee J, Kwon YK, Soh MS, Shin B, Luka Z, Hahn TR, Song PS (1999) Phytochrome signalling is mediated through nucleoside diphosphate kinase 2. *Nature* 401: 610–613
- Devlin PF, Halliday KJ, Harberd NP, Whitelam GC (1996) The rosette habit of *Arabidopsis thaliana* is dependent upon phytochrome action: novel phytochromes control internode elongation and flowering time. *Plant J* 10: 1127–1134
- Devlin PF, Patel SR, Whitelam GC (1998) Phytochrome E influences internode elongation and flowering time in *Arabidopsis*. *Plant Cell* 10: 1479–1487
- Devlin PF, Robson PR, Patel SR, Goosey L, Sharrock RA, Whitelam GC (1999) Phytochrome D acts in the shade-avoidance syndrome in *Arabidopsis* by controlling elongation growth and flowering time. *Plant Physiol* 119: 909–915
- Eichenberg K, Baurle I, Paulo N, Sharrock RA, Rudiger W, Schafer E (2000) *Arabidopsis* phytochromes C and E have different spectral characteristics from those of phytochromes A and B. *FEBS Lett* 470: 107–112

- Fankhauser C, Yeh KC, Lagarias JC, Zhang H, Elich TD, Chory J (1999) PKS1, a substrate phosphorylated by phytochrome that modulates light signaling in *Arabidopsis*. *Science* **284**: 1539–1541
- Halliday KJ, Salter MG, Thingnaes E, Whitelam GC (2003) Phytochrome control of flowering is temperature sensitive and correlates with expression of the floral integrator *FT*. *Plant J* **33**: 875–888
- Hennig L, Funk M, Whitelam GC, Schafer E (1999) Functional interaction of cryptochrome 1 and phytochrome D. *Plant J* **20**: 289–294
- Hennig L, Poppe C, Sweere U, Martin A, Schafer E (2001) Negative interference of endogenous phytochrome b with phytochrome a function in *Arabidopsis*. *Plant Physiol* **125**: 1036–1044
- Hennig L, Stoddart WM, Dieterle M, Whitelam GC, Schafer E (2002) Phytochrome E controls light-induced germination of *Arabidopsis*. *Plant Physiol* **128**: 194–200
- Hepworth SR, Valverde F, Ravenscroft D, Mouradov A, Coupland G (2002) Antagonistic regulation of flowering-time gene *SOC1* by *CONSTANS* and *FLC* via separate promoter motifs. *EMBO J* **21**: 4327–4337
- Huq E, Quail PH (2002) PIF4, a phytochrome-interacting bHLH factor, functions as a negative regulator of phytochrome B signaling in *Arabidopsis*. *EMBO J* **21**: 2441–2450
- Izawa T, Oikawa T, Sugiyama N, Tanisaka T, Yano M, Shimamoto K (2002) Phytochrome mediates the external light signal to repress *FT* orthologs in photoperiodic flowering of rice. *Genes Dev* **16**: 2006–2020
- Jarillo JA, Capel J, Tang RH, Yang HQ, Alonso JM, Ecker JR, Cashmore AR (2001) An *Arabidopsis* circadian clock component interacts with both *CRY1* and *phyB*. *Nature* **410**: 487–490
- Johnson E, Bradley M, Harberd NP, Whitelam GC (1994) Photoresponses of light-grown *phyA* mutants of *Arabidopsis*: phytochrome A is required for the perception of daylength extensions. *Plant Physiol* **105**: 141–149
- Kircher S, Gil P, Kozma-Bognar L, Fejes E, Speth V, Husselstein-Muller T, Bauer D, Adam E, Schafer E, Nagy F (2002) Nucleocytoplasmic partitioning of the plant photoreceptors phytochrome A, B, C, D, and E is regulated differentially by light and exhibits a diurnal rhythm. *Plant Cell* **14**: 1541–1555
- Koornneef M, Rolf E, Spruit CJP (1980) Genetic control of light-inhibited hypocotyl elongation in *Arabidopsis thaliana* (L.) Heynh. *Z Pflanzenphysiol* **100**: 147–160
- Liu XL, Covington MF, Fankhauser C, Chory J, Wagner DR (2001) *ELF3* encodes a circadian clock-regulated nuclear protein that functions in an *Arabidopsis* PHYB signal transduction pathway. *Plant Cell* **13**: 1293–1304
- Martinez-Garcia JF, Huq E, Quail PH (2000) Direct targeting of light signals to a promoter element-bound transcription factor. *Science* **288**: 859–863
- Mas P, Devlin PF, Panda S, Kay SA (2000) Functional interaction of phytochrome B and cryptochrome 2. *Nature* **408**: 207–211
- Mathews S, Sharrock RA (1997) Phytochrome gene diversity. *Plant Cell Environ* **20**: 666–671
- Mazzella MA, Bertero D, Casal JJ (2000) Temperature-dependent internode elongation in vegetative plants of *Arabidopsis thaliana* lacking phytochrome B and cryptochrome 1. *Planta* **210**: 497–501
- Mazzella MA, Cerdan PD, Staneloni RJ, Casal JJ (2001) Hierarchical coupling of phytochromes and cryptochromes reconciles stability and light modulation of *Arabidopsis* development. *Development* **128**: 2291–2299
- Mockler TC, Guo H, Yang H, Duong H, Lin C (1999) Antagonistic actions of *Arabidopsis* cryptochromes and phytochrome B in the regulation of floral induction. *Development* **126**: 2073–2082
- Nagy F, Schafer E (2002) Phytochromes control photomorphogenesis by differentially regulated, interacting signaling pathways in higher plants. *Annu Rev Plant Physiol Plant Mol Biol* **53**: 329–355
- Neff MM, Chory J (1998) Genetic interactions between phytochrome A, phytochrome B, and cryptochrome 1 during *Arabidopsis* development. *Plant Physiol* **118**: 27–35
- Ni M, Tepperman JM, Quail PH (1998) PIF3, a phytochrome-interacting factor necessary for normal photoinduced signal transduction, is a novel basic helix-loop-helix protein. *Cell* **95**: 657–667
- Ni M, Tepperman JM, Quail PH (1999) Binding of phytochrome B to its nuclear signalling partner PIF3 is reversibly induced by light. *Nature* **400**: 781–784
- Quail PH (2002) Phytochrome photosensory signalling networks. *Nat Rev Mol Cell Biol* **3**: 85–93
- Roldan M, Gomez-Mena C, Ruiz-Garcia L, Salinas J, Martinez-Zapater JM (1999) Sucrose availability on the aerial part of the plant promotes morphogenesis and flowering of *Arabidopsis* in the dark. *Plant J* **20**: 581–590
- Salome PA, Michael TP, Kearns EV, Fett-Neto AG, Sharrock RA, McClung CR (2002) The out of phase 1 mutant defines a role for PHYB in circadian phase control in *Arabidopsis*. *Plant Physiol* **129**: 1674–1685
- Simpson GG, Dean C (2002) *Arabidopsis*, the Rosetta stone of flowering time? *Science* **296**: 285–289
- Stearns SC (2002) Progress on canalization. *Proc Natl Acad Sci USA* **99**: 10229–10230
- Whitelam GC, Johnson E, Peng J, Carol P, Cowl JS, Harberd NP (1993) Phytochrome A null mutants of *Arabidopsis* display a wild-type phenotype in white light. *Plant Cell* **5**: 757–768
- Whitelam GC, Patel S, Devlin PF (1998) Phytochromes and photomorphogenesis in *Arabidopsis*. *Philos Trans R Soc Lond B Biol Sci* **353**: 1445–1453
- Yang HQ, Tang RH, Cashmore AR (2001) The signaling mechanism of *Arabidopsis* *CRY1* involves direct interaction with *COP1*. *Plant Cell* **13**: 2573–2587
- Yanovsky MJ, Kay SA (2002) Molecular basis of seasonal time measurement in *Arabidopsis*. *Nature* **419**: 308–312