



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Enhanced carbon pump inferred from relaxation of nutrient limitation in the glacial ocean

Citation for published version:

Pichevin, LE, Reynolds, BC, Ganeshram, RS, Cacho, I, Pena, L, Keefe, K & Ellam, RM 2009, 'Enhanced carbon pump inferred from relaxation of nutrient limitation in the glacial ocean' Nature, vol. 459, no. 7250, pp. 1114-1117. DOI: 10.1038/nature08101

Digital Object Identifier (DOI):

[10.1038/nature08101](https://doi.org/10.1038/nature08101)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Nature

Publisher Rights Statement:

The final version of this work was published in Nature of the Nature Publishing Group (2009)

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



This is the author's final draft or 'post-print' version. The final version was published in Nature copyright of the Nature Publishing Group (2009) and is available online.

Cite As: Pichevin, LE, Reynolds, BC, Ganeshram, RS, Cacho, I, Pena, L, Keefe, K & Ellam, RM 2009, 'Enhanced carbon pump inferred from relaxation of nutrient limitation in the glacial ocean' *Nature*, vol 459, no. 7250, pp. 1114-1117.

DOI: 10.1038/nature08101

Enhanced Carbon Pump Inferred from Relaxation of Nutrient Limitation in the Glacial Ocean

Authors: Pichevin L.E., Reynolds B.C., Ganeshram R.S., Cacho I., Pena L., Keefe K., Ellam R.M.

Address for correspondence:

Laetitia Pichevin
School of Geosciences,
Grant Institute,
University of Edinburgh,
The King's Buildings
West Mains Road,
Edinburgh, UK
EH9 3JW

Email: laetitia.pichevin@ed.ac.uk

Enhanced Carbon Pump Inferred from Relaxation of Nutrient Limitation in the Glacial Ocean

Pichevin L. E.¹, Reynolds B. C.², Ganeshram R. S.¹, Cacho I.³, Pena L.³, Keefe K.⁴,
Ellam R. M.⁴

1. School of Geosciences, Grant Institute, University of Edinburgh, West Main Road, EH10 3JW, Edinburgh, UK, laetitia.pichevin@ed.ac.uk, Tel +44 131 650 8547, Fax +44 131 668 3184

2. IGMR, ETH Zürich, Clausiusstrasse 25, CH-8092 Zürich, Switzerland

3. GRC Geociències Marines, Facultat de Geologia, Universitat de Barcelona C/ Martí Franques s/n 08028 Barcelona, Spain

4. Scottish Universities Environment Research Centre, Rankine Avenue, Scottish Enterprise Technology Park, East Kilbride, G75 0QF, UK

The modern Eastern Equatorial Pacific (EEP) is a major oceanic source of carbon to the atmosphere¹. During glacial periods, increased deep-sea carbon export in this region supported by additional input of dust-born iron^{2, 3} could have contributed towards lower atmospheric CO₂ levels. Such a role for the EEP is supported by higher organic carbon burial rates^{4, 5} documented in underlying glacial sediments but the lower opal accumulation rates had cast doubts^{6, 7}. Here we present a new silicon isotope record that suggests the paradoxical decline in opal accumulation rate in the glacial EEP resulted from a decrease in Si:C uptake ratio of diatoms under conditions of increased Fe availability from enhanced dust inputs. Consequently, our study provides support for an invigorated biological pump in this region during the last glacial period that

could have contributed to glacial CO₂ drawdown. Additionally, using evidence from silicon and nitrogen isotope changes we infer that, in contrast to modern situation, the biological productivity in this region is not constrained by the availability of Fe, Si and N during the glacial period. Therefore, we hypothesise that an invigorated biological CO₂ pump constrained perhaps only by P limitation was a more common occurrence in low-latitude areas of the glacial ocean.

The Eastern Equatorial Pacific (EEP), an important area of biogenic opal production and burial in the ocean⁸, accounts for more than two third of the marine CO₂ efflux to the atmosphere today¹. This situation arises because primary productivity in the EEP is co-limited by silicic acid and iron availability⁹ and therefore the biological CO₂ pump is unable to compensate for the CO₂ out-gassing that occurs through upwelling of deep waters¹⁰ (Fig.1).

In figure 2, we present glacial-interglacial records of biogenic accumulation as well as stable N and Si isotope records spanning the last 35 ka from a EEP marine sediment Core ODP 1240 (Fig. 1). Organic carbon concentrations in the sediment were double during the Last Glacial Maximum (LGM) compared to the Holocene and peak during the deglacial. The ²³⁰Th-normalised organic carbon accumulation rates also retain the same trend (Fig. 2). The doubling of organic carbon accumulation during the LGM is in agreement with previous studies showing that organic carbon export in the EEP was higher during the LGM^{4, 5}, and may imply an invigorated biological CO₂ pump during this period (see supplementary material). Contradicting such an assertion, opal concentration in Core 1240 was reduced roughly by a factor of 2 during the LGM, consistent with existing opal accumulation records in the region⁷.

⁶. Such differences in biogenic accumulation records cannot be explained by local changes in upwelling alone.

The sedimentary ratio of Si_{opal} to organic carbon (Si:C) in Core 1240 increases drastically during the glacial to interglacial transition with values almost 3 times higher during the early Holocene compared to the LGM (Fig. 2). This change in the relative accumulation of opal and carbon is seen across the entire EEP and therefore cannot be simply explained by factors such as differential distribution and preservation of these biogenic components (supp. mat.).

The contrast between opal and organic carbon accumulation trends during the LGM in the EEP could be reconciled by invoking a shift in the phytoplankton communities from diatomaceous to coccolithophorid production. This would have reduced the C rain-rate ratios (the relative contributions of organic carbon and CaCO_3 to total export production) in the glacial EEP and hence the capacity of the marine C reservoir to sequester carbon¹¹. However, CaCO_3 contents and accumulation rates in Core 1240 do not show a clear glacial-interglacial pattern to support such a shift in the rain-rate ratio of particulate export (Fig. 2). The estimates of carbonate accumulation rates in the EEP either based on the ^{230}Th normalisation technique or high resolution ^{14}C stratigraphy unequivocally document reduced carbonate fluxes and production during the LGM¹². Moreover, a recent study¹³ of carbonate accumulation corrected for sedimentary dissolution has even suggested that primary production in the EEP shifted from CaCO_3 to opal production during the LGM compared to the Holocene. Thus, sedimentary opal records from the EEP, which show halved accumulation rates during the LGM compared to the Holocene epoch⁷, implying a less efficient biological CO_2 pump, are in contradiction to the inferences based on enhanced organic carbon production and has until now remained a paradox. Resolving this

paradox has important implications in understanding the role of EEP in glacial CO₂ drawdown and the impact of enhanced Fe delivery on nutrient limitation and biological CO₂ pump during the glacial periods.

Availability of silicic acid in the surface water of the EEP is considered as the major control on opal production¹⁰ - a view that stems from modern observations of Si-limitation in this region (Fig. 1). However, LGM conditions could have been drastically different for diatom growth when dust-borne iron delivery to the surface EEP was twice as much as today, as recorded by sediment cores in this area^{2, 3} (Fig.2). The impact of this increased iron delivery on opal production in the Fe-limited low latitude Pacific ocean^{2, 3} has never been fully explored. Results from *in situ* Fe-fertilisation experiments^{14, 15} reveal that the alleviation of Fe limitation in the EEP causes a 2 to 3 fold decrease in Si:C uptake ratios of siliceous producers due to decreased cell silicification^{14, 15}. If this were to happen during the Fe-replete LGM this may have resulted in lower Si uptake during diatom growth and consequently reduced opal accumulation. In addition, given the large decline in Si:C uptake ratios documented in these experiments, silicic acid may have been available in surplus during diatom production. Thus, the reconstruction of glacial silicic acid utilisation in the EEP could be the missing link to unravel the opal accumulation “paradox” and to address carbon mass balance during the LGM.

We evaluate the possibility of lowered Si utilisation during diatom growth under Fe-replete glacial periods using a record of $\delta^{30}\text{Si}$ of opal (Fig.2). During the Holocene $\delta^{30}\text{Si}$ values are $\sim +1.3\%$, a value close to the isotopic composition of silicic acid in intermediate water masses of the tropical Pacific^{16, 17}. This indicates near complete utilisation of the available silicic acid, consistent with modern observations of its limitation in the EEP. In contrast, the LGM values were much lower ($<0.9\%$). This

lower $\delta^{30}\text{Si}$ signature may reflect a decrease in the relative utilisation of silicic acid in the EEP during glacial period provided this shift is not affected by glacial change in the nutrient source.

During the glacial, silicic acid leakage from the Southern Ocean, whereby enhanced dust-borne iron input to the Southern Ocean resulted in the conservation of silicic acid over other nutrients leading to its export to the low latitude^{18, 19}, is one mechanism that could have potentially altered silicic acid supply to the EEP and its isotopic signature. It has been estimated that during the glacial, low latitude Pacific could have received 3 times more silicic acid through the sub-Antarctic mode waters, with a $\delta^{30}\text{Si}$ signature as high as +3‰²⁰. Complete diatom utilisation of this additional supply of heavy $\text{Si}(\text{OH})_4$ in the EEP should have increased glacial opal burial accompanied by heavier shift in $\delta^{30}\text{Si}$. To the contrary, Core 1240 records a glacial decline in opal accumulation with lighter isotopic values (Fig.2).

The reason for the apparent contrast becomes clear if one considers how glacial increase in dust-borne Fe inputs would have impacted silicic acid utilisation locally within the EEP. The increased Fe input and the resultant decline in Si:C uptake ratios, as was proposed for the Southern Ocean, should have also operated in the glacial EEP leading to the conservation of silicic acid^{14, 18}. This would explain the decline in opal accumulation and the lighter $\delta^{30}\text{Si}$ values in the glacial intervals of Core 1240. Paradoxically, the increased silicic acid supply from the Southern Ocean appears to have occurred at the time the demand for this nutrient declined locally within the EEP -a contention that is supported by the similar glacial-interglacial histories of dust input in these two regions³. Thus, this new result documenting excess silicic acid in the glacial EEP strongly implies that the decline in opal accumulation in LGM sediments is not controlled by the availability of silicic acid and therefore could

be attributed to the consequence of Fe-fertilisation and lowered Si:C uptake ratios during diatom growth. Importantly, the glacial decrease in opal accumulation observed across the EEP^{6, 7} does not reflect a decline in C rain-rate ratio and the biological CO₂ pump as previously suggested⁶.

Our results suggest that the glacial scenario of nutrient limitation and the constraints on biological production in the EEP were drastically different from the modern situation depicted in Figure 1. In contrast to modern conditions, enhanced Fe delivery and the resultant switch to lower Si:C uptake ratio by diatoms led to conditions where Fe and Si are no longer limiting factors for biological productivity. Based on glacial $\delta^{30}\text{Si}$ values in Core 1240 and applying a steady state model¹⁷ we estimate relative silicic acid utilisation during the LGM declined by about 25% from a near complete utilisation during the Holocene. This calculation assumes a constant Si(OH)₄ source of around +1.2‰ in agreement with low latitude Pacific intermediate water values^{16, 17}. This estimate of decreased Si(OH)₄ utilisation would be even greater if the leakage of isotopically heavy silicic acid from the sub-Antarctic is taken into account²⁰. Similarly, the N-isotope record allows us to assess the potential for glacial nitrate limitation. The lack of Fe and Si co-limitation during the glacial should have led to increased nitrate utilisation and heavier glacial $\delta^{15}\text{N}$. In contrast, the N-isotope profile (Fig. 2) trends towards lower $\delta^{15}\text{N}$ values during the LGM relative to the Holocene, consistent with other similar studies in this region²¹. However, given that nitrate utilisation is incomplete in the modern EEP a further $\delta^{15}\text{N}$ decrease (>3‰) during the glacial period cannot be simply explained by changes in nitrate uptake by biota alone as it would entail a very large reduction in utilisation during this period. Therefore, we suggest that the low LGM $\delta^{15}\text{N}$ values relative to the Holocene reflect additional

supply of isotopically lighter N due to reduced denitrification in the oxygen minimum zones (OMZ) off Peru and Mexico^{22, 23}. Today the prevalence of denitrification in these margins bordering the EEP makes the source waters supplied to the EEP thermocline depleted in nitrate relative to phosphate (sub-redfieldian N:P) and with a relatively heavy N-isotope signature (+6.5‰ as opposed to the +4.8‰ ocean average) due to mixing with partially-denitrified heavy nitrate²⁴. Thus, reduced denitrification during glacial periods would enhance the nitrate inputs into the EEP by increasing the N/P ratio of upwelling waters and provide nitrate that is isotopically lighter. In combination both these factors would lead to a shift to lighter sedimentary N-isotope signatures irrespective of any glacial change in local upwelling^{25, 26}. Thus, the lighter N-isotope values provide evidence that nitrate is also not a limiting nutrient in the glacial EEP- a condition comparable to the modern situation. Therefore the data strongly suggest simultaneous removal of constraints imposed by Fe, Si and N limitation on biological production in the EEP during the LGM. This argues for a more invigorated biological CO₂ pump in the glacial EEP perhaps ultimately constrained by the availability of dissolved phosphate. The resulting increase in carbon export and rain-rate ratio as documented in the glacial sediments of the EEP (supp. mat.) should have reduced CO₂ evasion. Such an assertion is also consistent with the timing of the first phase of the atmospheric CO₂ rise during the last glacial termination (~18 ka) which is contemporaneous, within dating uncertainties, to the decline in dust delivery and the increases in sedimentary Si:C and $\delta^{30}\text{Si}$ in EEP records.

Finally, we suggest that our results have much wider ramifications because the EEP provides an illustration of the constraints imposed on the biological CO₂ pump in

low-latitude oceans of the glacial periods. In currently Fe limited open ocean High Nitrate Low Chlorophyll regions, such as the EEP and Southern Ocean, the switch to Fe-replete LGM conditions led to the generation of excess silicic acid during diatom growth, which in turn is subject to dispersal through surface and subsurface ocean circulation increasing the availability of silicic acid over much wider area of the glacial ocean^{18, 19}. This additional silicic acid supply occurs at a time when its demand by biota is already reduced due to Fe-fertilisation, which should have caused drastic reductions in Si limitation over large areas of the glacial ocean.

Also, the documented glacial decline in denitrification in the world's OMZ is expected to have increased the availability of nitrate^{27, 23}. For instance, NO₃:PO₄ ratio at the surface of the modern EEP is about 12.5, i.e. lower than N:P Redfield ratio of 16. In contrast, a 30% increase in nitrate inventory during the last glacial period compared to today as suggested by modelling work²⁸ could have been sufficient to cause the N:P ratio in the EEP to exceed the Redfield ratio. Such changes occurring more widely would have resulted in an invigorated CO₂ pump in low latitude oceans ultimately constrained by the availability of dissolved phosphate as suggested by the case study reported here and predicted by recent model results²⁹. Therefore, we hypothesise that P limitation was much more widespread during the glacial periods - a situation fundamentally different to that of the modern ocean.

Method Summary

Core ODP 202 1240 was retrieved from Cocos ridge (00°01.31N; 86°27.76W, 2,921 m depth) in the EEP. The age model is based on 13 AMS¹⁴C dates on planktonic foraminifers²⁵. Determination of the opal content (%) was performed by molybdate-blue spectrophotometry on alkaline extracts. Organic C (%), total N (%) and δ¹⁵N (‰)

were determined on bulk sediments using a Carlo Erba elemental analyzer coupled to a VG Prism III mass spectrometer at the University of Edinburgh. ^{230}Th normalization was performed using acid digestion on bulk samples and column chemistry followed by MC-ICPMS analyses.

Purification of the diatom samples for silicon isotope measurement has been performed by chemical leaching of the carbonate and organic fractions, sieving and differential settling. Silicon isotope determination has been conducted in ETH Zürich on the Nu1700 high-resolution MC-ICPMS³⁰.

References

1. Takahashi, T. et al. Global sea-air CO₂ flux based on climatological surface ocean pCO₂, and seasonal biological and temperature effects. *Deep-Sea Res. art II-Top. Stud. Oceanogr.* **49**, 1601-1622 (2002).
2. McGee, D., Marcantonio, F. & Lynch-Stieglitz, J. Deglacial changes in dust flux in the eastern equatorial Pacific. *Earth Planet. Sci. Lett.* **257**, 215-230 (2007).
3. Winckler, G., Anderson, R. F., Fleisher, M. Q., McGee, D. & Mahowald, N. Covariant glacial-interglacial dust fluxes in the equatorial Pacific and Antarctica. *Science* **320**, 93-96 (2008).
4. Pedersen, T. F. Increased Productivity in the Eastern Equatorial Pacific During the Last Glacial Maximum (19,000 to 14,000 Yr Bp). *Geology* **11**, 16-19 (1983).
5. Lyle, M. Climatically Forced Organic-Carbon Burial in Equatorial Atlantic and Pacific Oceans. *Nature* **335**, 529-532 (1988).

6. Kienast, S. S., Kienast, M., Jaccard, S., Calvert, S. E. & Francois, R. Testing the silica leakage hypothesis with sedimentary opal records from the eastern equatorial Pacific over the last 150 kyrs. *Geophys. Res. Lett.* **33**, doi: 10.1029/2006GL026651 (2006).
7. Bradtmiller, L. I., Anderson, R. F., Fleisher, M. Q. & Burckle, L. H. Diatom productivity in the equatorial Pacific Ocean from the last glacial period to the present: A test of the silicic acid leakage hypothesis. *Paleoceanography* **21** doi: 10.1029/2006PA001282 (2006).
8. Lisitzin, A. P. Distribution of Siliceous Micro Fossils in Suspension and in Bottom Sediments. The Micropalaeontology of Oceans. Symposium. 828p. Illus. Maps. Cambridge University Press: New York, N.Y., U.S.A, 173-195 (1971).
9. Moore, J. K., Doney, S. C. & Lindsay, K. Upper ocean ecosystem dynamics and iron cycling in a global three-dimensional model. *Glob. Biogeochem. Cycles* **18**, doi: 10.1029/2004GB002220 (2004).
10. Dugdale, R. C. & Wilkerson, F. P. Silicate regulation of new production in the equatorial Pacific upwelling. *Nature* **391**, 270-273 (1998).
11. Archer, D. & Maier-reimer, E. Effect of Deep-Sea Sedimentary Calcite Preservation on Atmospheric CO₂ Concentration. *Nature* **367**, 260-263 (1994).
12. Loubere, P. & Richaud, M. Some reconciliation of glacial-interglacial calcite flux reconstructions for the eastern equatorial Pacific. *Geochem. Geophys. Geosyst.* **8**, doi: 10.1029/2006GC001367 (2007).

13. Richaud, M., Loubere, P., Pichat, S. & Francois, R. Changes in opal flux and the rain ratio during the last 50,000 years in the equatorial Pacific. *Deep-Sea Res. Prt II- Top. Stud. Oceanogr.* **54**, 762-771 (2007).
14. Takeda, S. Influence of iron availability on nutrient consumption ratio of diatoms in oceanic waters. *Nature* **393**, 774-777 (1998).
15. Hutchins, D. A. & Bruland, K. W. Iron-limited diatom growth and Si : N uptake ratios in a coastal upwelling regime. *Nature* **393**, 561-564 (1998).
16. Reynolds, B. C., Frank, M. & Halliday, A. N. Silicon isotope fractionation during nutrient utilization in the North Pacific. *Earth Planet. Sci. Lett.* **244**, 431-443 (2006).
17. Beucher, C. P., Brzezinski, M. A. & Jones, J. L. Sources and biological fractionation of Silicon isotopes in the Eastern Equatorial Pacific. *Geochim. Cosmochim. Acta* **72**, 3063-3073 (2008).
18. Brzezinski, M. A. et al. A switch from Si(OH)₄ to NO₃⁻ depletion in the glacial Southern Ocean. *Geophys. Res. Lett.* **29**, doi: 10.1029/2001GL014349 (2002).
19. Matsumoto, K., Sarmiento, J. L. & Brzezinski, M. A. Silicic acid leakage from the Southern Ocean: A possible explanation for glacial atmospheric pCO₂. *Glob. Biogeochem. Cycles* **16**, doi: 10.1029/2001GB001442 (2002).
20. Beucher, C. P., Brzezinski, M. A. & Crosta, X. Silicic acid dynamics in the glacial sub-Antarctic: Implications for the silicic acid leakage hypothesis. *Glob. Biogeochem. Cycles* **21**, doi: 10.1029/2006GB002746 (2007).
21. Farrell, J. W., Pedersen, T. F., Calvert, S. E. & Nielsen, B. Glacial Interglacial Changes in Nutrient Utilization in the Equatorial Pacific-Ocean. *Nature* **377**, 514-517 (1995).

22. Toggweiler, J. R. & Carson, S. What are upwelling systems contributing to the ocean's carbon and nutrient budgets (ed. C.P. Summerhayes, K.-C. E., M.V. Angel, R.L. Smith, and B. Zeitzschel), 337-360 (John Wiley & Sons, 1995).
23. Ganeshram, R. S., Pedersen, T. F., Calvert, S. E., McNeill, G. W. & Fontugne, M. R. Glacial-interglacial variability in denitrification in the world's oceans: Causes and consequences. *Paleoceanography* **15**, 361-376 (2000).
24. Altabet, M. A. Nitrogen isotopic evidence for micronutrient control of fractional NO₃⁻ utilization in the equatorial Pacific. *Limnol. Oceanogr.* **46**, 368-380 (2001).
25. Pena, L. D., Cacho, I., Ferretti, P. & Hall, a. M. A. El Niño–Southern Oscillation–like variability during glacial terminations and interlatitudinal teleconnections. *Paleoceanography*, **23**, doi:10.1029/2008PA001620 (2008).
26. Koutavas, A., Lynch-Stieglitz, J., Marchitto, T. M. & Sachs, J. P. El Nino-like pattern in ice age tropical Pacific sea surface temperature. *Science* **297**, 226-230 (2002).
27. Ganeshram, R. S., Pedersen, T. F., Calvert, S. E. & Francois, R. Reduced nitrogen fixation in the glacial ocean inferred from changes in marine nitrogen and phosphorus inventories. *Nature* **415**, 156-159 (2002).
28. Deutsch, C., Sigman, D. M., Thunell, R. C., Meckler, A. N. & Haug, G. H. Isotopic constraints on glacial/interglacial changes in the oceanic nitrogen budget. *Glob. Biogeochem. Cycles* **18**, doi: 10.1029/2003GB002189 (2004).
29. Moore, J. K. & Doney, S. C. Iron availability limits the ocean nitrogen inventory stabilizing feedbacks between marine denitrification and nitrogen fixation. *Glob. Biogeochem. Cycles* **21**, doi: 10.1029/2006GB002762 (2007).

30. Georg, R. B., Reynolds, B. C., Frank, M. & Halliday, A. N. New sample preparation techniques for the determination of Si isotopic compositions using MC-ICPMS. *Chemical Geology* **235**, 95-104 (2006).
31. Monnin, E. et al. Atmospheric CO₂ concentrations over the last glacial termination. *Science* **291**, 112-114 (2001).

Supplementary Information is linked to the online version of the paper at www.nature.com/nature

Acknowledgements

We thank R. Francois, S. Kienast, W. Geibert, E. Galbraith and G. Leduc for thought-provoking discussions. Funding for this project was provided by the Scottish Alliance for Geoscience Environment Society (SAGES), Natural Environment Research Council (NERC) through Standard NERC grant to R.S.G and L.E.P and a Marie Curie Intra-European fellowship to L.E.P.

Author Contributions:

L.E.P. and R.S.G. initiated the project. L.E.P. and B.C.R. measured silicon isotopes in ETH Zurich, L.E.P. measured elemental composition and nitrogen isotopes, L.E.P., K.K. and R.E. made the ²³⁰Thorium measurements. I.C. and L.P. provided the age model. L.E.P. and R.S.G. wrote the paper with the participation of B.R.C. All authors participated to the discussions on the results and commented on the manuscript.

Figure captions

Figure 1: Annually averaged CO₂ partial pressure difference between the atmosphere and the ocean for year 1995 (modified from¹; <http://www.ldeo.columbia.edu/res/pi/CO2/carbondioxide/>). Also depicted is the hierarchy of nutrient limitation for diatom growth across the low latitude Pacific Ocean. Black lines delimit the area where the most severe limitation is imposed by iron, dotted lines delimit the area of silicic acid limitation. Beyond these lines the ocean is firstly nitrate limiting (modified from⁹). Note the geographical correspondence between CO₂ efflux maxima (warm colours) and iron and silicic acid limitation areas. Core site ODP 202 1240 is shown with a white dot.

Figure 2: Sedimentary records from Core 1240 plotted versus calendar ages B.P; (a) $\delta^{15}\text{N}$ signal of bulk material (bold line = 2 points average), (b) the $\delta^{30}\text{Si}$ signal of diatoms (errors bars are 1 sigma error of the mean), (c) Si:C ratio (bold line = 2 points average), elemental concentrations (solid line) and ²³⁰thorium normalized accumulation rates (dotted lines) of (d) organic carbon (wt.%), (e) opal (wt.%), (f) carbonate (wt.%), (g) dust fluxes in Core ODP 138-849A and 138-850A², and (h) atmospheric PCO₂ record from EPICA dome C ice core³¹. T1 and LGM stand for Termination 1 and Last Glacial Maximum, respectively.

Methods

Core ODP 202 1240 (00°01.31N; 86°27.76W, 2,921 m depth) was retrieved on Cocos ridge (see ODP report http://www-odp.tamu.edu/publications/202_IR/chap_11/chap_11.htm).

The age model is based on 13 AMS¹⁴C dates on planktonic foraminifers. Determination of the opal content (%) was performed by molybdate-blue

spectrophotometry on alkaline extracts according to the method published in³². Organic Carbon (%) and N (%) contents were determined by using a Carlo-Erba elemental analyser on 10 mg of freeze-dried and powdered bulk sediment. The isotopic composition of sedimentary nitrogen ($\delta^{15}\text{N}$) was measured with a VG Prism III mass spectrometer on freeze-dried and powdered bulk sediments. Thorium 230 normalization was used to account for the effect of sediment focusing³³ and performed using acid digestion on bulk samples and column chemistry followed by multi collector ICP-MS analyses at the Scottish Universities Environment Research Centre ³⁴(supp. mat.).

Purification of the diatom samples for silicon isotope measurement has been performed by chemical leaching of the carbonate and organic fractions, sieving and differential settling following a cleaning method published recently³⁵. Silicon isotope determination has been conducted in ETH Zürich on the Nu1700 high-resolution MC-ICPMS⁶. A small amount of biogenic opal (0.5 mg) was dried down with concentrated perchloric acid at $\sim 180^\circ\text{C}$ in Teflon[©] vials, and then dissolved in 100 μl of 1 M NaOH, before being diluted to 5 ml with 0.01 M HCl after 24 hours. Equivalent to 10 μg of opal was loaded onto a pre-cleaned 1.8ml DOWEX 50W-X12 cation exchange resin bed (in H^+ form) and eluted with 5 ml of purified water (Milli-Q element $18.2 \text{ M}\Omega\cdot\text{cm}^{-1}$). The Si isotope composition was determined on the diluted solution (0.6ppm Si) on the Nu1700 high-resolution MC-ICPMS at ETH Zürich, using a standard-sample-standard bracketing protocol. All results in this study were calculated using the $\delta^{30}\text{Si}$ notation for deviations of the measured $^{30}\text{Si}/^{28}\text{Si}$ from the international Si standard NBS28 in parts per thousand (‰). The long-term reproducibility was better than 0.07‰ $\delta^{30}\text{Si}$ (1 s.d.)³⁶. Samples were measured at

least 5 times, which resulted in a 95 % confidence level below 0.08%. Error bars on the $\delta^{30}\text{Si}$ plot are calculated as 1 sigma error of the mean (Fig. 2).

References

32. Mortlock, R. A. & Froelich, P. N. A Simple Method for the Rapid-Determination of Biogenic Opal in Pelagic Marine-Sediments. *Deep-Sea Res. Part a-Oceanogr. Res. Papers* **36**, 1415-1426 (1989).
33. Francois, R., Frank, M., van der Loeff, M. M. R. & Bacon, M. P. Th-230 normalization: An essential tool for interpreting sedimentary fluxes during the late Quaternary. *Paleoceanography* **19**, doi: 10.1029/2003PA000939 (2004).
34. Ellam, R. M. & Keefe, K. MC-ICP-MS analysis of non-natural U isotope ratios using a Th-229/Th-232 external mass bias correction. *J. Analytic. Atom. Spectrometry* **22**, 147-152 (2007).
35. Morley, D. W. et al. Cleaning of lake sediment samples for diatom oxygen isotope analysis. *J. Paleolimno.* **31**, 391-401 (2004).
36. Reynolds, B. C., Georg, R. B., Oberli, F., Wiechert, U. H. & Halliday, A. N. Re-assessment of silicon isotope reference materials using high-resolution multi-collector ICP-MS. *J. Analytic. Atom. Spectrometry* **21**, 266-269 (2006).

Sea-Air PCO_2 difference (μatm)



