



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

A novel, resistance-linked ovine PrP variant and its equivalent mouse variant modulate the in vitro cell-free conversion of rPrP to PrPres

Citation for published version:

Kirby, L, Goldmann, W, Houston, F, Gill, AC & Manson, JC 2006, 'A novel, resistance-linked ovine PrP variant and its equivalent mouse variant modulate the in vitro cell-free conversion of rPrP to PrPres' *Journal of General Virology*, vol 87, no. 12, pp. 3747-3751., 10.1099/vir.0.82086-0

Digital Object Identifier (DOI):

[10.1099/vir.0.82086-0](https://doi.org/10.1099/vir.0.82086-0)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Author final version (often known as postprint)

Published In:

Journal of General Virology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



1 **A novel resistance-linked ovine PrP variant and its equivalent mouse variant**
2 **modulate the *in vitro* cell-free conversion of rPrP to PrP^{res}**

3

4 **Louise Kirby¹, Wilfred Goldmann², Fiona Houston¹ Andrew C. Gill¹ and Jean**
5 **C. Manson²**

6

7 ¹Institute for Animal Health, Compton Laboratories, Newbury, Berkshire, RG20 7NN,
8 UK and ²Neuropathogenesis Unit, Institute for Animal Health, Ogston Building, West
9 Mains Road, Edinburgh, EH9 3JF, UK.

10

11 **Author for correspondence:** Louise Kirby

12 Tel: [+44] 01635 577259

13 Fax: [+44] 01635 577263

14 e-mail: louise.kirby@bbsrc.ac.uk

15

16 Running title: Cell-free conversion of ovine PrP variant L¹⁶⁸

17 Number or words in summary: 150

18 Number of words in main text: 2491

19 Number of figures and tables: 3

20

21 **Summary**

22 **Prion diseases are associated with the conversion of the normal cellular**
23 **prion protein, PrP^c, to the abnormal disease associated PrP^{Sc}. This conversion**
24 **can be mimicked *in vitro* using the cell-free conversion assay. We have**
25 **recently shown that this assay can be modified to use bacterial recombinant**

26 **PrP as substrate and mimic the *in vivo* transmission characteristics of rodent**
27 **scrapie. Here we demonstrate that the assay replicates the ovine**
28 **polymorphism barriers of scrapie transmission. In addition, the recently**
29 **identified ovine PrP variant ARL^{168Q}, which is associated with survival of**
30 **sheep to experimental BSE, modulates the cell-free conversion of ovine**
31 **recombinant PrP to PrP^{res} by 3 different types of PrP^{Sc}, reducing conversion**
32 **efficiencies to levels similar to the ovine resistance-associated ARR variant.**
33 **Also, the equivalent variant in mice (L¹⁶⁴) is resistant to conversion by 87V**
34 **scrapie. Together these results suggest a significant role for this position**
35 **and/or amino acid in conversion.**

36

37 Transmissible spongiform encephalopathies (TSEs) are fatal
38 neurodegenerative diseases including Creutzfeldt-Jakob disease in humans, scrapie
39 in sheep and goats and bovine spongiform encephalopathy (BSE) in cattle. The
40 causative agent responsible for TSEs or prion diseases has yet to be fully defined.
41 However, a fundamental event in disease is the conversion of the normal, proteinase
42 K (PK) sensitive isoform of the prion protein, PrP^c, to an abnormal, partially PK
43 resistant isoform, PrP^{Sc}, and the accumulation of this abnormal isoform in the central
44 nervous system of infected animals (Hope *et al.*, 1986; Meyer *et al.*, 1986; Oesch *et*
45 *al.*, 1985). Although the mechanism of conversion is unknown, interaction between
46 PrP^c and PrP^{Sc} is critically implicated by *in vitro* studies (Caughey & Chesebro, 1997)
47 and features in the current models of replication (Prusiner, 1991; Jarrett & Lansbury,
48 1993).

49 Polymorphisms in PrP are associated with susceptibility to and pathology of
50 TSEs. The major determinants controlling the susceptibility of sheep to scrapie are

51 polymorphisms at PrP amino acid codons 136, 154 and 171 (Hunter *et al.*, 1997).
52 The PrP ARR allele (amino acids, in single letter code, at positions 136, 154 and 171
53 respectively) is associated with resistance to classical scrapie. The PrP ARQ and
54 VRQ alleles are associated with susceptibility to disease. Novel ovine PrP
55 polymorphisms are regularly being identified in sheep genotyping programs. Most of
56 these novel polymorphisms occur with low frequency and their association with
57 disease susceptibility is not known (Baylis & Goldman, 2004). It is important to
58 assess the disease association of such variants in the hope of identifying additional
59 scrapie resistant alleles, as not all breeds or populations have significant frequencies
60 of the known resistant ARR allele and it has been reported that sheep homozygous
61 for the ARR allele may be susceptible to atypical scrapie (Bushman *et al.*, 2004)
62 and intracerebral (ic) inoculation with BSE (Houston *et al.*, 2003). The identification
63 of other resistant alleles may also help to protect against novel strains of scrapie or
64 adaptation to a particular genotype.

65 Due to the long incubation time and high cost of animal experiments, the *in*
66 *vitro* cell-free conversion assay has provided a quick, well defined system in which to
67 assess the disease association of such alleles (Kocisko *et al.*, 1994). However, for
68 natural infection, dose, route, strain of agent, influence of second allele and breed of
69 sheep are all likely to play a role in TSE susceptibility and additional evidence from
70 experimental challenge would be required to support any association of particular
71 alleles with resistance to TSEs. In the cell-free conversion assay PrP^{Sc}, isolated from
72 the brains of scrapie infected animals, induces the conversion of radiolabelled
73 recombinant PrP, to a PK resistant isoform, PrP^{res} (Kocisko *et al.*, 1994). The assay
74 has been shown to replicate *in vivo* species specificity, strain properties and
75 polymorphism barriers (Bessen *et al.*, 1995; Bossers *et al.*, 1997; Bossers *et al.*,

76 2000; Iniguez *et al.*, 2000; Kocisko *et al.*, 1995; Zhang *et al.*, 2002; Raymond *et al.*,
77 1997; Horiuchi *et al.*, 2000) and has been used to study many aspects of molecular
78 conversion. As yet, however, no *in vitro* cell-free generated PrP^{res} has been shown to
79 be infectious (Hill *et al.*, 1999). Recently, we reported the use, as substrate, of
80 mouse and hamster PrP, biochemically purified from recombinant bacteria and
81 demonstrated that the assay replicated several characteristics of *in vivo* disease
82 (Kirby *et al.*, 2003).

83 Evidence is presented here that bacterial recombinant ovine PrP can be
84 converted to PrP^{res} in the cell-free conversion assay and the sheep polymorphism
85 barriers of scrapie transmission are replicated. Full length ovine PrP of the ARR,
86 ARQ and VRQ genotypes, with the N-terminal signal sequence replaced with
87 methionine and the C-terminal signal sequence removed, corresponding to amino
88 acids 25-233, were PCR amplified from genomic DNA using the 5' and 3' primers, 5'-
89 GGATCCATCATGAAGAAGCGACCAAAACCTGGC-3' and 5'-
90 CCGAATTCTCATGCCCCCCTTTGGTAATAA-3', respectively. Plasmid pTrcHis B
91 (Invitrogen) was digested with restriction enzymes *Nco*I and *Eco*RI to remove the 6-
92 histidine tag. PCR fragments were digested with restriction enzymes *Eco*RI and *Rca*I
93 and ligated into the modified pTrcHis B plasmid. Therefore, the vector encodes full-
94 length, untagged ovine PrP. Calcium chloride competent *Escherichia coli*, strain
95 1B392 (Wright *et al.*, 1986), were transformed with the recombinant vectors. Ovine
96 PrP variants were expressed, radiolabelled, purified, refolded and characterized by
97 mass spectrometry and circular dichroism, as described in Kirby *et al.*, (2003). A
98 representative autoradiograph is shown in Figure 1, lanes 1-3. PrP^{Sc} was purified
99 from the brain stems of VRQ homozygous sheep clinically infected with scrapie
100 (SSBP/1 source), from ARQ homozygous sheep clinically infected with BSE and

101 from BSE infected cows, based on a method described by Hope *et al.*, (1986). Cell-
102 free conversion assays were carried out using the 3 radiolabelled ovine PrP variants
103 (^{35}S -rARRPrP, ^{35}S -rARQPrP, and ^{35}S -rVRQPrP) as substrates and the 3 different
104 PrP^{Sc} types as seeds, as described previously (Kirby *et al.*, 2003). Briefly, 1 μg of
105 PrP^{Sc} was incubated with 200 ng of ^{35}S -rPrP for 24 hours at 37°C, in a non-
106 guanidine containing conversion buffer. Following incubation, 1/20th of the reaction
107 was treated with 60 μg PK ml⁻¹ for 1 hour at 37°C. PK digestion was stopped by
108 adding Pefabloc to 1 mM. All samples were methanol precipitated and analysed by
109 SDS-PAGE and autoradiography. Autoradiographs were quantified using Phoretix
110 Gel Analysis Software. A typical autoradiograph is shown in Figure 1A. The
111 experiment was repeated 3 times, efficiencies of conversion determined by
112 densitometric analysis and the mean conversion efficiencies calculated ($\pm\text{SE}$) by
113 densitometric analysis of labeled PrP before and after PK treatment (Figure 1B).

114 Figure 1A shows that bacterial recombinant ovine PrP is converted into a PK
115 resistant form in the cell-free conversion assay. The conversion efficiencies obtained
116 using bacterial recombinant ovine PrP were low (Figure 1B) in comparison to the
117 efficiencies obtained using bacterial recombinant rodent PrP (Kirby *et al.*, 2003).
118 Conversion efficiency may be increased by the addition of guanidine to the
119 conversion buffer (Bossers *et al.*, 1997). Indeed, it has recently been reported that
120 the addition of guanidine to the assay buffer is essential to obtain any conversion of
121 sheep PrP (Piening *et al.*, 2006). However, this reduces the physiological relevance
122 of the assay as a model of conversion. In addition, the conversion efficiencies
123 between sets of experiments (a set is defined as the 5 types of ^{35}S -rOvPrP and 1
124 type of PrP^{Sc}) varied enormously (Figure 1B). This has been reported by others
125 using a similar cell-free conversion assay (Bossers *et al.*, 1997). The reason for this

126 is unknown, but breed of sheep or the area of brain selected for purification of PrP^{Sc}
127 may contribute. However, the relative conversion efficiencies within a set were
128 similar each time the experiment was repeated. Of the 3 frequently occurring ovine
129 PrP variants, ARR, ARQ and VRQ, ³⁵S-rARRPrP, the variant associated with
130 resistance to classical scrapie, consistently produced the lowest amount of PrP^{res}
131 when used as substrate in the cell-free conversion reaction (Figure 1A, lanes 6, 11
132 and 16). ³⁵S-rARQPrP and ³⁵S-rVRQPrP, variants associated with susceptibility to
133 scrapie and short incubation times, converted with higher efficiencies (Figure 1A,
134 lanes 7, 8, 12, 13, 17 and 18). A switch in the convertibility of ³⁵S-rARQPrP and ³⁵S-
135 rVRQPrP is evident with homologous and heterologous PrP^{Sc}. The homologous
136 conversion reactions produced the greatest amount of PrP^{res} (Figure 1A, lanes 8 and
137 17). These results indicate that the known *in vivo* polymorphism barriers of scrapie
138 transmission can be mimicked in the cell-free conversion assay, using bacterial
139 recombinant ovine PrP.

140 Recently, Goldmann *et al.*, (2005) have identified two novel ovine PrP alleles.
141 Sheep carrying a PrP variant with a proline to leucine polymorphism at amino acid
142 168 (L¹⁶⁸) were shown to have increased survival time to experimental BSE in 2
143 independent experiments (Goldmann, **this reference refers to Wilfred Goldmann's
144 paper to be published alongside my paper). ARL¹⁶⁸Q occurs at low frequency and
145 although it may be linked with resistance to experimental BSE no data exists on its
146 resistance to scrapie infection. The other variant ARQE¹⁷⁵, with a change from a
147 glutamine to a glutamic acid at amino acid position 175 (E¹⁷⁵) is also rare and not yet
148 associated with scrapie or BSE susceptibility. Therefore, the cell-free conversion
149 assay was used to predict whether these ovine variants are associated with
150 resistance to scrapie infection. The 168 and 175 amino acid positions are of interest

151 as they are located close to the putative factor-X binding site (Telling *et al.*, 1995;
152 Kaneko *et al.*, 1997) and the resistance-associated R¹⁷¹ position.

153 In a further experiment ³⁵S-rARL¹⁶⁸QPrP and ³⁵S-rARQE¹⁷⁵PrP were
154 produced (Figure 1A, lanes 4 and 5), as described above for the other ovine
155 variants, and used as substrate in the cell-free conversion assay incubating with the
156 3 different PrP^{Sc} types. ³⁵S-rARL¹⁶⁸QPrP converted with low efficiency using all 3
157 types of PrP^{Sc} (Figure 1A, lanes 9, 14 and 19), indicating that the ARL¹⁶⁸Q effect on
158 conversion is significant for scrapie as well as BSE. ³⁵S-rARQE¹⁷⁵PrP converted with
159 an efficiency similar to ³⁵S-rARQPrP with the 3 different types of PrP^{Sc} (Figure 1A,
160 lanes 10, 15 and 20), suggesting that this amino acid change does not affect
161 conversion. Table 1 provides a rank order of conversion efficiencies of the ³⁵S-
162 rOvPrP variants with the 3 different PrP^{Sc} types.

163 Due to the low conversion efficiencies and to strengthen the data showing the
164 reduced conversion efficiency effect of the ovine ARL¹⁶⁸QPrP variant, it was
165 determined whether the equivalent site in mouse PrP would have the same effect
166 and reduce conversion efficiency. The mouse cell-free conversion assay gives
167 higher conversion efficiencies and therefore the effect on conversion can be more
168 easily seen. Also, polymorphisms associated with resistance to TSEs in both sheep
169 (R¹⁷¹) and humans (K²¹⁹) have been engineered on a mouse background (R¹⁶⁷ and
170 K²¹⁸, respectively) and the recombinant versions shown not to convert when
171 expressed in ScN2a cells (Kaneko *et al.*, 1997), suggesting that mouse PrP can be
172 used to model the effects of PrP polymorphisms in other species.

173 The murine equivalent of ovine amino acid 168 is 164. Full length mouse PrP
174 (amino acids 23-230) of the *Prn-p^a* genotype with a proline to leucine mutation at
175 amino acid 164 was constructed by site directed mutagenesis (QuickChange II,

176 Stratagene) using the full length mouse PrP clone, the production of which has been
177 described previously (Kirby *et al.*, 2003), as a template and the following primers; 5'-
178 CCAAGTGTACTACAGGCTAGTGGATCAGTACAGC-3' and 5'-
179 GCTGTACTGATCCACTAGCCTGTAGTACACTTGG-3'. Rosetta *Escherichia coli*
180 (Novagen), which over expresses the rare leucine tRNA, were transformed with
181 pTrcMoL¹⁶⁴PrP. MoL¹⁶⁴PrP and MoPrP were expressed, radiolabelled, purified and
182 characterized as described for the ovine variants. PrP^{Sc} was purified from the brains
183 of terminally ill 87V-infected VM mice, based on a method described by Hope *et al.*,
184 (1986). Cell-free conversion assays, using the 2 mouse PrP variants as substrates
185 and 87V PrP^{Sc} as seed, were carried out in the absence of guanidine and analysed
186 as described previously (Kirby *et al.*, 2003). The experiment was repeated at least 3
187 times and a typical autoradiograph is shown in Figure 2.

188 Figure 2 shows that ³⁵S-rMoPrP is efficiently converted to PrP^{res} in the
189 presence of 87V PrP^{Sc} (Figure 2, lane 2). However, ³⁵S-rMoL¹⁶⁴ PrP is not converted
190 to PrP^{res} in the presence of 87V PrP^{Sc} (Figure 2, lane 3), suggesting that this amino
191 acid/position may be a significant site in the conversion process in other species.

192 It is not understood how substitution of different amino acids at certain
193 positions within PrP has such a profound effect on susceptibility. It has been
194 suggested that mutations can modulate the stability of PrP^c, PrP^{Sc}, or both, or can
195 affect the binding of PrP to effector molecules. In the case of a proline to leucine
196 mutation, the amino acids share similar hydrophobicity, but are structurally diverse.
197 Proline, the only imino acid, has backbone torsion angles that are tightly controlled
198 as a result of its cyclic structure. Because of this, it results in turns in protein
199 backbones and often occurs at the end of β -sheets. A change from proline to
200 leucine, which has a greater range of flexibility in its backbone angles, may reduce

201 the propensity of PrP to form β -sheets, and explain the significant protective effect of
202 this mutation. Alternatively, leucine is an amino acid capable of involvement in many
203 types of secondary structure, including α -helices, and a change to a leucine may aid
204 the stabilisation of PrP^c. Interestingly, the same mutation, proline to leucine, is
205 associated with apparent spontaneous disease in humans when it occurs at position
206 102 in human PrP, the reverse of what we find for ovine position 168 and its
207 equivalent mouse position. This is almost undoubtedly a reflection of the very
208 different tertiary structures in the different parts of the molecule and the different
209 involvement of these areas in conversion of PrP^c to PrP^{Sc}. In addition, extensive
210 gene-targeted transgenic mouse experiments show that L¹⁰¹ mice have altered
211 susceptibility to a range of TSE isolates compared to wildtype mice (Barron *et al.*,
212 2001), further complicating interpretation. The neutral phenotype of the ovine PrP
213 E¹⁷⁵ polymorphism in our assay suggests that not every change in this region of PrP
214 will affect susceptibility and indicates that the underlying mechanism may be highly
215 positional and residue specific.

216 To determine the molecular mechanisms responsible for the protective effect
217 of the L¹⁶⁸ mutation, substitution of a range of different amino acids at position 168
218 would be required and their affect on conversion assessed. Such experiments are
219 currently underway in our laboratory using the murine cell-free conversion assay as
220 a model. In addition, the ovine and murine version of the cell-free conversion assay
221 using bacterial recombinant PrP can be used to assess the link between novel ovine
222 polymorphisms, as they are identified, to classical scrapie, atypical scrapie, such as
223 NOR98, and to BSE.

224

225 **Acknowledgements**

226 The authors would like to thank the USDA and BBRSC for funding, Ian Sylvester for
227 providing the ARR and VRQ PrP expressing clones and James Graham for CD
228 analysis of recombinant mouse L¹⁶⁴ PrP.

229

230 **Figure 1 legend:**

231 **A)** An autoradiograph of a cell-free conversion reaction using 5 ovine recombinant
232 variants (³⁵S-rOvPrP) as substrate and PrP^{Sc}, isolated from scrapie infected
233 VRQ/VRQ homozygous sheep brains (lanes 6-10), BSE infected cow brains (lanes
234 11-15) and BSE infected ARQ/ARQ homozygous sheep brains (lanes 16-20) as
235 seeds. Lanes 6-20 have been exposed for a longer time interval than lanes 1-5 in
236 order to detect PrP^{res}. Molecular mass markers are indicated on the left in kDa. (PK)
237 proteinase K. Boxed area indicates PrP^{res}. **B)** Average conversion efficiencies (±SE)
238 for each set of conversion reactions (5 types of ³⁵S-rOvPrP and 1 type of PrP^{Sc}).

239

240 **Table 1 legend:**

241 A summary table of sheep conversion assay data. Each set of conversions (5 ³⁵S-
242 rOvPrP variants converted by 1 type of PrP^{Sc}) are ranked in order of decreasing
243 conversion efficiency, predicting the susceptibility of the ovine variants to scrapie and
244 BSE.

245

246 **Figure 2 legend:**

247 An autoradiograph of a cell-free conversion reaction using mouse PrP with and
248 without a proline to leucine mutation at amino acid position 164 (equivalent to ovine
249 position 168) as substrates and 87V PrP^{Sc} as seed. Molecular mass markers are
250 indicated on the left in kDa. (PK) proteinase K. Boxed area indicates PrP^{res}.

251

252 **References**

253 **Barron, R. M., Thomson, V., Jamieson, E., Melton, D. W., Ironside, J., Will, R. &**
254 **Manson, J. C. (2001).** Changing a single amino acid in the N-terminus of murine PrP
255 alters TSE incubation time across three species barriers. *Embo J* **20**, 5070-8.

256

257 **Baylis, M. and Goldman, W. (2004).** The Genetics of scrapie in sheep and goats.
258 *Current Molecular Medicine* **4**, 385- 396

259

260 **Bessen, R. A., Kocisko, D. A., Raymond, G. J., Nandan, S., Lansbury, P. T. &**
261 **Caughey, B. (1995).** Non-genetic propagation of strain-specific properties of scrapie
262 prion protein. *Nature* **375**, 698-700.

263

264 **Bossers, A., Belt, P., Raymond, G. J., Caughey, B., de Vries, R. & Smits, M. A.**
265 **(1997).** Scrapie susceptibility-linked polymorphisms modulate the in vitro conversion
266 of sheep prion protein to protease-resistant forms. *Proc Natl Acad Sci U S A* **94**,
267 4931-6.

268

269 **Bossers, A., de Vries, R. & Smits, M. A. (2000).** Susceptibility of sheep for scrapie
270 as assessed by in vitro conversion of nine naturally occurring variants of PrP. *J Virol*
271 **74**, 1407-14.

272

273 **Buschmann A., Luhken G., Schultz J., Erhardt G., Groschup M. H.(2004)**
274 Neuronal accumulation of abnormal prion protein in sheep carrying a scrapie-
275 resistant genotype (PrP ARR/ARR). *J. Gen. Virol.* **85**, 2727-2733

276

277 **Caughey, B. & Chesebro, B. (1997).** Prion protein and the transmissible spongiform
278 encephalopathies. *Trends in Cell Biology* **7**, 56-62.

279

280 **Goldmann, W., Baylis, M., Chihota, C., Stevenson, E. & Hunter, N. (2005).**
281 Frequencies of *PrP* gene haplotypes in British sheep flocks and the implications for
282 breeding programs. *J App Microbiol*, **98**, 1294-1302

283

284 **Hill, A. F., Antoniou, M. & Collinge, J. (1999).** Protease-resistant prion protein
285 produced in vitro lacks detectable infectivity. *J Gen Virol* **80 (Pt 1)**, 11-4.

286

287 **Hope, J., Morton, L. J., Farquhar, C. F., Multhaup, G., Beyreuther, K. &**
288 **Kimberlin, R. H. (1986).** The major polypeptide of scrapie-associated fibrils (SAF)
289 has the same size, charge distribution and N-terminal protein sequence as predicted
290 for the normal brain protein (PrP). *Embo J* **5**, 2591-7.

291

292 **Horiuchi, M., Priola, S. A., Chabry, J. & Caughey, B. (2000).** Interactions between
293 heterologous forms of prion protein: binding, inhibition of conversion, and species
294 barriers. *Proc Natl Acad Sci U S A* **97**, 5836-41.

295

296 **Houston, F., Goldmann, W., Chong, A., Jeffrey, M., González, L., Foster, J.,**
297 **Parnham, D. & Hunter, N. (2003).** Transmission of BSE to sheep thought to be
298 genetically resistant to infection. *Nature*, **423**, 498

299

300 **Hunter, N., Goldmann, W., Foster, J. D., Cairns, D. & Smith, G. (1997).** Natural

301 scrapie and PrP genotype: case-control studies in British sheep. *Vet Rec* **141**, 137-
302 40.

303

304 **Iniguez, V., McKenzie, D., Mirwald, J. & Aiken, J. (2000).** Strain-specific
305 propagation of PrP(Sc) properties into baculovirus-expressed hamster PrP(C). *J Gen*
306 *Viro* **81**, 2565-71.

307

308 **Jarrett, J. T. & Lansbury, P. T. (1993).** Seeding One-Dimensional Crystallization of
309 Amyloid - a Pathogenic Mechanism in Alzheimers-Disease and Scrapie. *Cell* **73**,
310 1055-1058.

311

312 **Kaneko, K., Zulianello, L., Scott, M., Cooper, C. M., Wallace, A. C., James, T. L.,**
313 **Cohen, F. E. & Prusiner, S. B. (1997).** Evidence for protein X binding to a
314 discontinuous epitope on the cellular prion protein during scrapie prion propagation.
315 *Proc Natl Acad Sci U S A* **94**, 10069-74.

316

317 **Kirby, L., Birkett, C. R., Rudyk, H., Gilbert, I. H. & Hope, J. (2003).** In vitro cell-free
318 conversion of bacterial recombinant PrP to PrP(res) as a model for conversion. *J*
319 *Gen Viro* **84**, 1013-20.

320

321 **Kocisko, D. A., Come, J. H., Priola, S. A., Chesebro, B., Raymond, G. J.,**
322 **Lansbury, P. T. & Caughey, B. (1994).** Cell-free formation of protease-resistant
323 prion protein. *Nature* **370**, 471-4.

324

325 **Kocisko, D. A., Priola, S. A., Raymond, G. J., Chesebro, B., Lansbury, P. T., Jr.**

326 **& Caughey, B. (1995).** Species specificity in the cell-free conversion of prion protein
327 to protease-resistant forms: a model for the scrapie species barrier. *Proc Natl Acad*
328 *Sci U S A* **92**, 3923-7.

329

330 **Meyer, R. K., McKinley, M. P., Bowman, K. A., Braunfeld, M. B., Barry, R. A. &**
331 **Prusiner, S. B. (1986).** Separation and properties of cellular and scrapie prion
332 proteins. *Proc Natl Acad Sci U S A* **83**, 2310-4.

333

334 **Oesch, B., Westaway, D., Walchli, M., McKinley, M. P., Kent, S. B., Aebersold,**
335 **R., Barry, R. A., Tempst, P., Teplow, D. B., Hood, L. E. & et al. (1985).** A cellular
336 gene encodes scrapie PrP 27-30 protein. *Cell* **40**, 735-46.

337

338 **Piening, N., Nonno, R., Di Bari, M., Walter, S., Windl, O., Agrimi, U.,**
339 **Kretzschmar, Hans A. & Bertsch, U. (2006).** Conversion efficiency of bank vole
340 prion protein in vitro is determined by residues 155 and 170, but does not correlate
341 with the high susceptibility of bank voles to sheep scrapie in vivo. *Journal of*
342 *Biological Chemistry* **182**, 9373-9384.

343

344 **Prusiner, S. B. (1991).** Molecular-Biology of Prion Diseases. *Science* **252**, 1515-
345 1522.

346

347 **Raymond, G. J., Hope, J., Kocisko, D. A., Priola, S. A., Raymond, L. D.,**
348 **Bossers, A., Ironside, J., Will, R. G., Chen, S. G., Petersen, R. B., Gambetti, P.,**
349 **Rubenstein, R., Smits, M. A., Lansbury, P. T., Jr. & Caughey, B. (1997).**

350 Molecular assessment of the potential transmissibilities of BSE and scrapie to

351 humans. *Nature* **388**, 285-8.

352

353 **Telling, G. C., Scott, M., Mastrianni, J., Gabizon, R., Torchia, M., Cohen, F. E.,**

354 **DeArmond, S. J. & Prusiner, S. B. (1995).** Prion propagation in mice expressing

355 human and chimeric PrP transgenes implicates the interaction of cellular PrP with

356 another protein. *Cell* **83**, 79-90.

357

358 **Wright, E.M & Humphreys et al., (1986).** Dual-origin plasmids containing an

359 amplifiable ColE1 ori: temperature-controlled expression of cloned genes. *Gene* **49**,

360 311-321.

361


362 **Zhang, F. P., Zhang, J., Zhou, W., Zhang, B. Y., Hung, T. & Dong, X. P. (2002).**

363 Expression of PrP(C) as HIS-fusion form in a baculovirus system and conversion of

364 expressed PrP-sen to PrP-res in a cell-free system. *Virus Res* **87**, 145-53.

Figure 1

A



B





Figure 2



367

Table 1

368

369

370

PrP ^{Sc}	Conversion efficiency of ³⁵ S-rOvPrP				
	High				Low
VRQ/SSBP1	VRQ	ARQE ¹⁷⁵	ARQ	ARR	ARL ¹⁶⁸ Q
Bovine/BSE	ARQE ¹⁷⁵	VRQ	ARR	ARQ	ARL ¹⁶⁸ Q
ARQ/BSE	ARQE ¹⁷⁵	ARQ	VRQ	ARR	ARL ¹⁶⁸ Q
consensus	ARQ/ARQE ¹⁷⁵		VRQ	ARR/ARL ¹⁶⁸ Q	

371