



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Overcoming reduced glucocorticoid sensitivity in airway disease

Citation for published version:

Marwick, JA, Adcock, IM & Chung, KF 2010, 'Overcoming reduced glucocorticoid sensitivity in airway disease: molecular mechanisms and therapeutic approaches' *Drugs*, vol 70, no. 8, pp. 929-48., 10.2165/10898520-000000000-00000

Digital Object Identifier (DOI):

[10.2165/10898520-000000000-00000](https://doi.org/10.2165/10898520-000000000-00000)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Preprint (usually an early version)

Published In:

Drugs

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Overcoming Reduced Glucocorticoid Sensitivity in Airway Disease

Molecular Mechanisms and Therapeutic Approaches

John A. Marwick,¹ Ian M. Adcock² and Kian Fan Chung²

- 1 MRC Centre for Inflammation Research, Queens Medical Research Institute, University of Edinburgh Medical School, Edinburgh, UK
- 2 Section of Airways Disease, National Heart & Lung Institute, Imperial College London, London, UK

Abstract

There is a considerable and growing unmet medical need in respiratory disease concerning effective anti-inflammatory therapies for conditions such as severe asthma, chronic obstructive pulmonary disease and cystic fibrosis. These diseases share a predominant characteristic of an enhanced and uncontrolled inflammatory response in the lungs, which contributes to disease progression, hospitalization and mortality. These diseases are poorly controlled by current anti-inflammatory therapies including glucocorticoids, which are otherwise effective in many other inflammatory conditions or in milder disease such as asthma. The exact cause of this apparent impairment of glucocorticoid function remains largely unclear; however, recent studies have now implicated a number of possible mechanisms. Central among these is an elevation of the oxidant burden in the lungs and the resulting reduction in the activity of histone deacetylase (HDAC)-2. This contributes to both the enhancement of proinflammatory mediator expression and the impaired ability of the glucocorticoid receptor (GR)- α to repress proinflammatory gene expression. The oxidant-mediated reduction in HDAC-2 activity is, in part, a result of an elevation in the phosphoinositol 3-kinase (PI3K) δ /Akt signalling pathway. Blockade of the PI3K δ pathway restores glucocorticoid function in both *in vitro* and *in vivo* models, and in primary cells from disease. In addition, inhibition of the PI3K δ and PI3K γ isoforms is anti-inflammatory in both innate and adaptive immune responses. Consequently, selective inhibition of this pathway may provide a therapeutic strategy both as a novel anti-inflammatory and in combination therapy with glucocorticoids to restore their function. However, a number of other oxidant-related and -unrelated mechanisms, including altered kinase signalling and expression of the dominant negative GR β , may also play a role in the development of glucocorticoid insensitivity. Further elucidation of these mechanisms and pathways will enable novel therapeutic targeting for alternative anti-inflammatory drugs or combination therapies providing restoration for the anti-inflammatory action of glucocorticoids.

1. Overview of Glucocorticoid Insensitivity in Airway Diseases: Medical Need

Glucocorticoids are effective anti-inflammatory drugs that are used to control both acute and chronic inflammatory responses in a wide range of diseases. However, in some diseases the anti-inflammatory actions of glucocorticoids are impaired or ineffective. These include depression, inflammatory bowel disease, multiple sclerosis, rheumatoid arthritis, acute lymphoblastic leukaemia, cystic fibrosis (CF), idiopathic pulmonary fibrosis, severe asthma and chronic obstructive pulmonary disease (COPD).^[1-7] Respiratory diseases are predominant amongst these and their incidence, particularly severe asthma and COPD, are increasing.^[8,9] The problem of relative glucocorticoid insensitivity in these diseases is compounded by the fact that the precise immunological and molecular mechanisms of these diseases remain complex and unclear.^[10] Current anti-inflammatory therapies for severe asthma, COPD and CF are inadequate, which highlights a substantial unmet medical need with significant disease management problems and cost burden for healthcare systems worldwide. Recent attempts to find effective anti-inflammatory alternatives, such as anti-tumour necrosis factor (TNF)- α agents for COPD, have proven unsuccessful and place some doubt on the reliability of 'single target' therapies in complex and poorly understood inflammatory diseases.^[11] In addition, the incidence of a relative glucocorticoid insensitivity is seen in a wide range of diseases, yet many of these harbour 'similar' characteristics including oxidative stress and innate immunity. Therefore, the likelihood of at least some degree of mechanistic commonality is large; this remains unaddressed but would be important in developing a therapeutic strategy with a broad disease application.

In this review we aim to introduce the reader to glucocorticoids, their mechanism of action and the basic pathophysiology of COPD, severe asthma and CF. In addition, we aim to describe the phenomenon of relative glucocorticoid insensitivity in these diseases, including proposed mech-

anisms and thereafter potential anti-inflammatory therapeutic strategies.

2. Glucocorticoids: Mechanism of Action

Glucocorticoids mediate a broad-ranging immunosuppression through binding and activation of the glucocorticoid receptor (GR)- α . Non-glucocorticoid bound GR α is sequestered in the cytosol by a complex of chaperone proteins including heat shock proteins 70 and 90, which prevent non-ligand-mediated GR α activity.^[12,13] Association with this complex enables GR α to adopt the correct conformation to allow glucocorticoid binding.^[13] Upon glucocorticoid binding, GR α dissociates from this complex and translocates to the nucleus where it mediates both repression and induction of gene expression (figure 1).^[14,15] The exact contributions of GR α -mediated gene repression (transrepression) and gene activation (transactivation) to the anti-inflammatory actions of glucocorticoids remain unresolved and controversial, but may be stimulus dependent.^[16] There is also increasing evidence for rapid 'non-genomic' glucocorticoid actions that may arise from (i) specific interactions with as yet unidentified cell surface receptors or (ii) direct action in the cell lipid membrane.^[17,18] These non-genomic actions are proposed to lead to rapid signalling through the generation of secondary signalling messengers and thereafter influencing signalling pathways.^[17] However, investigation into these non-genomic mechanisms are in their infancy and their role, if any, in either the anti-inflammatory actions of glucocorticoids or in disease is unknown. Therefore, these are not discussed in this manuscript.

2.1 Glucocorticoid-Mediated Transrepression

GR α monomers are able to associate directly with promoter-bound transcription factors such as nuclear factor (NF)- κ B and activator protein (AP)-1.^[19] This association allows GR α to be targeted directly to the activated promoter region of proinflammatory genes controlled by these transcription factors. Thereafter, GR α facilitates silencing of gene transcription by modulation of critical covalent modifications on the core histone proteins




Fig. 1. Diagrammatic representation of the basic mechanism by which glucocorticoid (GC) receptor- α (GR α) [a] represses transcription factor-driven expression of proinflammatory genes (transrepression) and (b) activates gene expression from GC response element (GRE) promoter sites. Nuclear factor- κ B is represented by a heterodimer of its p50/p65 subunits. HAT = histone acetyltransferase; HDAC = histone deacetylase; RNA Pol II = RNA polymerase II; TBP = TATA binding protein.

such as acetylation and methylation.^[20-22] These modifications are central in mediating conformational changes that regulate access for the transcriptional machinery to the promoter region.^[20] GR α itself has no intrinsic enzymatic capacity to directly modulate these histone modifications, but recruits co-repressor complexes such as mammalian Sin3a (mSin3a) and nucleosome remodelling and deacetylase/Mi2 (NuRD/Mi2) complexes.^[23,24] These co-repressor complexes contain DNA and chromatin-modifying enzymes, including methyltransferases and histone deacetylases (HDACs), which ‘turn off’ gene transcription by altering the covalent modifications on the core histone proteins and DNA^[21,25] (figure 1). The recruitment of HDAC-2 appears to be of particular importance for GR α -mediated transrepression. Specific reduction of HDAC-2 expression using small interfering RNA (siRNA) abolishes glucocorticoid-GR α -

mediated repression of proinflammatory gene expression.^[26] HDAC-2 induces gene silencing in a nonspecific manner by removing the acetyl groups from the amino terminal tails (deacetylation) of the core histone proteins.^[20] This enables DNA to re-associate with the core histone proteins forming a compact or ‘condensed’ tertiary chromatin structure which dislodges the transcriptional machinery and prevents further binding.^[20,27] HDAC-2 also deacetylates GR α itself, which serves to enhance its association to NF- κ B.^[26]

2.2 Glucocorticoid-Mediated Glucocorticoid Receptor (GR)- α Transactivation

In addition to repression of gene expression, GR α can also promote gene expression (transactivation) by forming homodimers and directly binding to the glucocorticoid response element (GRE)

promoter (figure 1). GR α -GRE binding induces the expression of a number of anti-inflammatory mediators including interleukin (IL)-10, secretory leukocyte protease inhibitor and mitogen-activated protein kinase (MAPK) phosphatase 1 (MKP-1).^[28,29] In addition, glucocorticoid repression of inflammatory genes under the control of NF- κ B may also be partly facilitated through GR α -mediated induction of inhibitory protein κ B α (I κ B α) from the GRE. The resulting elevated I κ B expression sequesters NF- κ B to the cytosol where it is held in an inactive cytoplasmic complex.^[30]

Particular attention has been paid to the glucocorticoid-dependent induction of MKP-1. Induction of the potent proinflammatory p38 MAPK/NF- κ B signalling pathway can regulate GR α function (via p38 MAPK phosphorylation; discussed in sections 2.4 and 4.3), which is in turn regulated by GR α through induction of MKP-1 expression.^[31,32] Adenoviral overexpression of MKP-1 reduced p38 MAPK and NF- κ B activation, and the expression of CXC chemokine ligand 8 (CXCL8).^[31] In addition, siRNA-mediated knockdown of MKP-1 partially reverses dexamethasone repression of TNF α -mediated activation of p38 MAPK. TNF α -induced NF- κ B-mediated proinflammatory gene expression is repressed by dexamethasone in the presence of MAPK kinase 6 (MKK6), a p38 MAPK inhibitor; however, blocking MKP-1 by addition of MKP-1 siRNA diminishes this effect.^[31] These studies provide evidence for the direct importance of GR α transactivation in mediating the anti-inflammatory actions of glucocorticoids.

However, GR α -mediated gene expression is also thought to be responsible for many of the unacceptable side effects seen with high doses of glucocorticoids and is a common reason for the withdrawal of high-dose oral glucocorticoids as an anti-inflammatory therapy.

2.3 Relative Contribution of GR α Transrepression and Transactivation in Glucocorticoid Inflammatory Mediator Suppression

GR α transactivation is widely associated with the unacceptable side effects seen at higher doses

of glucocorticoids. Therefore, the relative contributions of both transrepression and transactivation may be therapeutically important if these mechanisms can be selectively targeted. Transgenic mice expressing a mutant GR α , which is unable to dimerise and thereby bind to the DNA (GR α^{dim} mice) are able to repress inflammatory responses induced by both local and systemic lipopolysaccharide exposure comparable to wild-type animals.^[16] Therefore, it was proposed that GR α DNA binding and consequent induction of gene expression (transactivation) was non-essential for the anti-inflammatory actions of GR α . However, further investigations using these mice in a model of contact allergy demonstrated that the anti-inflammatory response of glucocorticoids was impaired in the GR α^{dim} mice.^[33] This suggests that the relative contributions of GR α transrepression and transactivation may depend on the inflammatory stimulus.

To date, more importance has been placed on GR α transrepression in glucocorticoid-mediated inflammatory repression. This is principally due to greater experimental investigation into GR α -mediated transrepression and concern over the transactivational-related adverse effects rather than substantial experimental evidence showing a secondary/minor role for GR-mediated transactivation in the glucocorticoid-mediated anti-inflammatory action. Therefore, further, more direct investigation is needed to decipher the real contribution of GR α -mediated transactivation in a range of inflammatory responses. Critically, this must also be done using human cells and tissues as murine cells do not express the GR β isoform, which may play an important role (discussed in sections 2.5 and 4.4). Investigation in primary human cells and tissues using well characterized dissociated corticosteroids may resolve this issue in the near future.

2.4 Regulation of GR α Function by Phosphorylation

Phosphorylation plays an important role in the regulation of GR α function.^[34,35] Human GR α is phosphorylated on at least five serine residues (S113, S141, S203, S211 and S226), which

are positioned within the major transactivational domain (AF1).^[34] Although the exact functional effects of phosphorylation at these sites are unclear, a number of studies have shown how they may determine the activity of the AF1 domain on GR α . Phosphorylation of S211 and S226 is thought to be important for the transcriptional activity of GR α , whilst S203 may be involved in GR α translocation.^[35,36] Phosphorylation of S211 and S226 may also enhance the interaction of GR α with co-regulators, including the vitamin D₃ receptor-interacting protein/thyroid receptor-associated protein (DRIP/TRAP) complex and the vitamin D receptor interacting protein 150 (MED14), as well as potentially influencing GR α activity in a gene-selective manner.^[35]

An array of kinases facilitates phosphorylation of GR α at distinct sites, thereby integrating GR α activation in cell-signalling pathways. These include extracellular-regulated kinase (ERK)-2, p38 MAPK, cyclin-dependent kinases (CDKs) 1, 2 and 5, glycogen synthase kinase 3 (GSK3) and c-Jun N-terminal kinase (JNK).^[36-38] These kinases mediate phosphorylation that may be either activational (CDK1, 2 and 5) or inhibitory (JNK, GSK3). In addition, certain kinases such as p38 MAPK appear to facilitate both activation and inhibition of GR α activity in a cell-specific manner.^[36,39]

2.5 A Role for GR β ?

The human glucocorticoid gene encodes two isoforms of the GR receptor generated by splice variants resulting in expression of the GR α and GR β isoforms.^[40] In contrast to the canonical GR α , GR β is an orphan receptor which is localized to the nuclear compartment and is devoid of the ability to bind glucocorticoids.^[41] The sole function of GR β was thought to be as a dominant negative regulator of GR α through competition for co-regulators including the GR interacting protein 1 (GRIP-1).^[40,42] However, although the GR β isoform lacks the ability to bind to glucocorticoids, recent evidence suggests that, in addition to its dominant negative regulatory role for GR α function, GR β may also facilitate direct transcriptional effects of its own. Although no

endogenous ligands for GR β have been identified, Cidlowski and colleagues^[41] demonstrated that GR β is also localized in the cytosol using COS-1 and U-2 OS cells and has the ability to bind the synthetic GR α receptor antagonist mifepristone (RU486), which mediated nuclear translocation of GR β . Interestingly, in the absence of GR α , microarray analysis showed that GR β was capable of regulating gene transcription, which was diminished upon binding of mifepristone.^[41] Additional studies using Hela cells have also shown that GR β has intrinsic gene-specific transcriptional activity (both negative and positive), where the majority of affected genes were distinct from those regulated by GR α .^[43] These recent findings clearly demonstrate that the role of GR β may be more complex and important than its perceived single role as a dominant negative regulator of GR α . The further complication of additional splice variants with potentially distinct functions also needs further examination.^[44] However, these are novel discoveries and further investigations, particularly in primary human cells, are needed to elucidate the extent and role of these proposed additional GR β actions as well as any cellular or tissue specificity.

3. Relative Glucocorticoid Insensitivity in Respiratory Disease

3.1 Severe Asthma

Allergic responses in the airways of patients with asthma may induce narrowing of the large airways that is largely reversible by bronchodilator drugs such as short- and long-acting β_2 agonists. The allergic response in asthma is characterized by an adaptive/T helper (T_H) type 2 cell-mediated response that induces an isotype switch in B cells for the production of immunoglobulin (Ig) E. This in turn initiates mast cell degranulation through activation of high-affinity IgE receptors expressed on the mast cells, and also through the induction of eosinophilic activation and responses.^[10] Low doses of inhaled glucocorticoids effectively control the inflammatory response in most patients with asthma

and are considered a first-line treatment.^[45] However, a small proportion of asthmatic patients have a severe form of asthma that is not adequately controlled by existing asthma medications, including glucocorticoids, even when used at high doses.

The inflammatory response in the lungs of patients with severe (relatively glucocorticoid-'insensitive' or -'unresponsive') asthma is not different to that observed in the lungs of patients with mild/moderate (relatively glucocorticoid-'sensitive' or -'responsive') asthma, as demonstrated by an elevation in both eosinophil and leukocyte numbers in bronchial biopsies from patients with severe asthma.^[46] It is clear that patients with severe asthma are treated with much higher doses of glucocorticoids, which therefore gives rise to the idea of relative glucocorticoid insensitivity.^[47] This relative glucocorticoid insensitivity is supported by observations that glucocorticoids fail to repress the release of pro-inflammatory mediators from both circulating inflammatory cells (peripheral blood mononuclear cells [PBMCs]) and lung macrophages (obtained from bronchoalveolar lavage) induced by lipopolysaccharide from patients with severe asthma as compared with patients with mild/moderate asthma, including CXCL8, TNF α and granulocyte macrophage-colony stimulating factor (GM-CSF).^[48,49] The ratio of matrix metalloprotease (MMP)-9 to the tissue inhibitor of MMP (TIMP)-1 is higher in the lungs of patients with severe asthma.^[50] Glucocorticoids elevate TIMP1 expression but this response is absent in alveolar macrophages from patients with severe asthma, which may contribute to the abnormal tissue remodelling in the airways of severe asthma leading to a reduction in lung function and β -agonist reversibility in these patients.^[2,50] Indeed, Macedo et al.^[46] have recently shown that there is a greater degree of subepithelial fibrosis and airway smooth muscle in patients with severe asthma as compared with non-severe asthma.

Interestingly, the inflammatory component of severe asthma may have some similarities to the inflammatory response seen in COPD.^[10] This is particularly apparent with the presence of an innate/T_h1 response in addition to the T_h2 cells that

predominate in mild/moderate asthma as well as more CD8+ T cells.^[10,51-53] In contrast to mild/moderate asthma, patients with severe asthma also have increased neutrophils in their sputum, increased CXCL8 and TNF α , and an elevated oxidant burden in the lung.^[51,54-57]

Asthmatic patients who smoke (accounting for ~25% of asthmatic patients) also have a reduced response to inhaled and oral glucocorticoids, and develop more severe asthma with an associated decline in lung function compared with asthmatic patients who do not smoke.^[58,59] These smoking asthmatic patients also have an increase in sputum neutrophils and CXCL8, which is more closely associated with severe asthma and COPD than with mild/moderate asthmatic patients who do not smoke.^[59] The evidence for this smoking-related effect is further found in COPD (discussed in section 3.2) where the major aetiological factor is cigarette smoke, which may also contribute to the development of the relative glucocorticoid-insensitivity seen in this disease.^[2,60]

3.2 Chronic Obstructive Pulmonary Disease (COPD)

COPD is characterized by airflow limitation that is progressive, poorly reversible and is associated with airway remodelling and an enhanced chronic inflammation in the lungs.^[61] Although cigarette smoke is the main aetiological factor in the development of COPD, the exact mechanism and contributing factors involved in the development of COPD are complex and not well understood. Airflow limitation is a central feature of both COPD and asthma; however, an important defining factor differentiating these two diseases is that the airflow limitation in asthma is largely reversible, whereas it is largely irreversible in COPD.^[62] A seminal study from Hogg and colleagues^[62] identified a potential cause of the airflow obstruction in COPD to be remodelling of the small airways, which correlated with disease progression.

Another defining factor between COPD and asthma is the type of underlying inflammation in the lungs. The inflammation in the lungs of COPD patients is predominantly driven by an

innate/ T_{h1} type immune response, whereas in asthma it is a predominantly adaptive/ T_{h2} immune-mediated response.^[10,60] The chronic enhanced lung inflammation in COPD is associated with an elevation in the number of macrophages, neutrophils, T_{h1} and type 1 cytotoxic T ($Tc1$) cells and $CD8+$ T cells in the lung.^[10,51,60] In addition, COPD patients have mucus hypersecretion and reduced mucociliary clearance, which is a major contributing factor in bacterial colonization and infection resulting in exacerbations as well as varying degrees of emphysema (alveolar wall destruction) and chronic bronchitis.^[60,63,64]

Unlike asthma patients, the majority of patients with COPD respond very poorly to inhaled and oral glucocorticoids, even at high doses. Glucocorticoid treatment in patients with COPD has no impact on the rate of decline in lung function, disease progression or mortality. There is evidence that glucocorticoids may have a modest beneficial effect on the number of exacerbations, although this has recently been questioned.^[65,66] The inflammatory response seen in the lungs of COPD patients is glucocorticoid insensitive. A number of studies have shown that inhaled glucocorticoids have no impact on the numbers of inflammatory cells in the lungs or on the release of proinflammatory mediators in COPD patients.^[29,67,68] In addition, glucocorticoids fail to repress the release of proinflammatory mediators from alveolar macrophages obtained from the bronchoalveolar lavage from COPD patients.^[69]

3.3 Cystic Fibrosis

CF is one of the most common lethal genetic diseases in Caucasians, and is characterized by an aggressive and predominantly innate/ T_{h1} inflammatory response with extensive destruction of the lungs.^[70] In contrast to both COPD and severe asthma, the underlying cause of CF is known. The development of CF is a result of a mutation in the gene that encodes the CF transmembrane conductance regulator (CFTR).^[70] The CFTR is expressed in epithelial cells and leukocytes, and acts mainly as a chloride channel, although it has a plethora of other key regulatory functions, including regulation of adenosine tri-

phosphate channels, intracellular vesicle transport, acidification of intracellular organelles, and inhibition of both epithelial sodium channels and endogenous calcium-activated chloride channels.^[70] The development of CF is often rapid; the lungs of children with CF appear normal at birth, but then rapidly become infected and inflamed. The influx of inflammatory polymorphonuclear cells is associated with an imbalance in pro- and anti-inflammatory mediators including an elevation in NF- κ B signalling, and increased IL-6, CXCL8, TNF α and eicosanoid expression, whilst the levels of anti-inflammatory mediators such as IL-10, lipoxin and docosahexaenoic acid are reduced.^[70-73] Despite the importance of the inflammatory response in the pathogenesis of CF, the current therapies are ineffective. Similarly to COPD and severe asthma, glucocorticoids are ineffective at controlling the aggressive inflammatory response seen in CF. Ibuprofen is commonly used as an alternative anti-inflammatory therapy as it is inexpensive, has relatively few adverse effects, and has been shown to be beneficial when given before the onset of severe inflammation and pathological changes in the lung.^[74] However, due to the aggressive nature of the inflammatory response in CF and its undoubted importance in the progression of the disease, the lack of an effective broad-ranging and powerful anti-inflammatory drug such as a functional glucocorticoid-type drug represents a major unmet medical need.

4. Underlying Mechanisms of Glucocorticoid Insensitivity

The molecular mechanisms of the relative glucocorticoid insensitivity in these respiratory diseases remain both unclear and understudied. Although there is a recognised inherent variation in glucocorticoid responsiveness between different cell types and tissues, the majority of cells that make up the lung and the infiltrating inflammatory cells are responsive to glucocorticoids.^[75] One exception to this are neutrophils, which are less responsive to glucocorticoids (discussed in detail in section 4.5). Furthermore, many other inflammatory conditions involving many of these

cell types are well controlled. It is therefore unlikely that an intrinsic relative glucocorticoid unresponsiveness of a particular cell type or of the lung itself can fully account for the reduction of glucocorticoid-insensitivity seen in CF, severe asthma and COPD.

There may also be some commonality in the mechanisms of glucocorticoid insensitivity between these diseases and indeed in other diseases that respond relatively poorly to glucocorticoids. Common immunological features of severe asthma, COPD and CF include an enhanced inflammatory response that either has components of or is predominantly driven by the innate/ T_H1 type immune response. The innate/ T_H1 response

itself is well documented as responding well to glucocorticoids; however, an 'enhanced' inflammatory response may incorporate an overriding factor. One of the more prominent theories implicated is elevated or 'overriding' oxidant burden in the lungs^[60,76] (figure 2). This concept is attractive not only from the experimental and mechanistic evidence, but also by its alignment with other proposed mechanisms. However, severe asthma, COPD and CF are often studied separately, whereas an approach to dissect the pathways governing glucocorticoid function across these diseases may prove more fruitful. Other proposed mechanisms, both distinct and aligned with oxidative stress, include genetic factors,




Fig. 2. Diagrammatic representation of the proposed mechanisms by which oxidative stress impairs glucocorticoid receptor- α ($GR\alpha$) function leading to an enhanced proinflammatory response that is resistant to glucocorticoid-mediated repression. Nuclear factor (NF)- κB is represented by a heterodimer of its p50/p65 subunits. **HDAC**=histone deacetylase; **mSin3a**=mammalian Sin3a; **RNS**=reactive nitrogen species; **ROS**=reactive oxygen species.




Fig. 3. Diagrammatic representation of the proposed mechanisms by which glucocorticoid receptor- α (GR α) function may become defective in airways disease leading to glucocorticoid insensitivity. Nuclear factor (NF)- κ B is represented by a heterodimer of its p50/p65 subunits. **GRE** = glucocorticoid response element; **HAT** = histone acetyltransferase; **HDAC** = histone deacetylase; **MKP-1** = mitogen-activated protein kinase phosphatase 1; **mSin3a** = mammalian Sin3a.

altered kinase signalling, elevated GR β expression and alteration of the histone acetylation/deacetylation balance (figure 3).

4.1 Oxidative Stress and Histone Deacetylase Activity

An elevated oxidant burden in the lungs is a major component of severe asthma, COPD and CF, which may be a central factor in the development of relative glucocorticoid insensitivity in these diseases.^[60,76] This oxidant stress may be derived from both exogenous sources such as pollution and cigarette smoke and endogenously from the respiratory burst of proinflammatory cells such as macrophages and neutrophils. Under physiological conditions, oxidants and the redox state of the cell are an integral part of cellular signalling and function, including that of GR.^[77,78] However, a significant elevation in

oxidants can overcome both intra- and extracellular antioxidant defences, and can alter signalling pathways and protein function.^[79] The role of additional exogenous oxidative stress such as cigarette smoke is highlighted by both its induction of reduced glucocorticoid insensitivity in experimental models, the development of glucocorticoid insensitivity in COPD and the development of relative glucocorticoid insensitivity in patients with mild/moderate asthma who smoke.^[58,60,80,81] Oxidative stress also plays a key role in the inflammatory processes associated with CF.^[82]

A central factor in both an oxidant-mediated enhancement of inflammatory responses and development of relative glucocorticoid insensitivity may be facilitated by a direct alteration of the acetylation-deacetylation balance of the core histones.^[83] Oxidative stress leads to a reduction of HDAC-2 activity (acute exposure) with an

additional reduction in HDAC-2 expression (seen with more chronic exposures) in *in vitro* and *in vivo* models.^[80,84,85] The reduction in both activity and expression results from covalent modifications, including hyperphosphorylation, nitration and carbonylation mediated by reactive oxygen species (e.g. reactive carbonyls, reactive nitrogen species), and by kinase signalling pathways activated by the oxidative stress.^[83,86] These modifications directly result in the reduction HDAC-2 activity and also reduce its expression by marking it out for proteasomal degradation.^[84,85,87-89] As histone acetylation is a key factor in gene transcription, this oxidant-mediated loss of HDAC activity and/or elevated histone acetyltransferase (HAT) activity is likely to contribute to the enhanced inflammatory responses in these diseases (figure 2).

HDAC-2 activity and expression is reduced in both COPD and severe asthma, with the reduction of activity in the peripheral lungs of COPD patients correlating with disease severity.^[49,90,91] Importantly, HDAC-2 is fundamental for functional GR α transrepression of proinflammatory genes, and a reduction in its activity/expression mediated either by oxidative stress or by siRNA severely impairs GR α transrepression of proinflammatory mediators.^[26] Conversely, protection or restoration of HDAC-2 activity in *in vitro* and *in vivo* models restores glucocorticoid responsiveness.^[81,92] Therefore, the reduction in HDAC-2 activity and expression seen in COPD and severe asthma is likely to be an important factor in the mechanisms of the relative glucocorticoid insensitivity seen in these diseases (figure 2).

Total HDAC activity measurement in CF has only been performed as part of one study and there was no alteration compared with healthy lungs.^[91] However, this study was performed on total lung extracts and alterations may be more local or cell-type restricted, which may be missed in a relatively crude total lung analysis. Indeed, recent studies have shown that the loss of CFTR may result in a reduction of HDAC-2 activity in airway epithelial cells.^[93] In addition, oxidative stress-induced hyperacetylation of the CXCL8 promoter in CF airway cell models and HAT activity may be elevated in the peripheral lungs

of CF patients.^[91,94] However, with so few studies in CF relating to the HDAC/HAT balance, the evidence of a role for HDAC-2 activity in the relative glucocorticoid insensitivity in CF appears promising but is not yet conclusive. This highlights a critical need for further investigations relating to glucocorticoid insensitivity in this disease.

4.2 Genetics

A number of studies have linked genetic mutations and alterations in cellular expression with the development of glucocorticoid insensitivity, as elegantly reviewed by Gross et al.^[95] Early studies identified a higher incidence of relative glucocorticoid-insensitive asthma within families, which indicated that there may be a genetic link to the development of glucocorticoid insensitivity.^[96] Genetic analysis of PBMCs have identified 11 genes that discriminate between relative glucocorticoid-insensitive and glucocorticoid-sensitive asthmatic patients, including those for several asthma-related proinflammatory signalling pathways, suggesting that there may be a genetic factor in the development of relative glucocorticoid-insensitive asthma.^[97] A link between genetic factors and the development of relative glucocorticoid insensitivity in COPD has not been directly assessed; however, there is evidence that genetic susceptibility is likely to play a role. Studies investigating the antioxidant capacity in the lungs of COPD patients and smokers suggest that subjects who develop COPD have a reduced capacity to elevate their antioxidant defences, which may, in part, account for the development of COPD in a subset of smokers (~20%).^[98-100] Furthermore, given the likely central role for oxidant stress in the development of glucocorticoid insensitivity, the reduced antioxidant capacity is also likely to contribute to the development of a relative glucocorticoid insensitivity in COPD.

The genetics of severe asthma versus mild/moderate asthma are currently being intensively investigated, but as of yet there has been no indication of genetic link with antioxidant capacity that may thereafter play a role in glucocorticoid insensitivity. In CF, the onset and level of the inflammatory response and ensuing lung

destruction is aggressive and a clear driving factor is represented by the mutation of the CFTR gene. Therefore, unlike the slow progression and development of COPD where, in addition to other factors, the oxidant imbalance may take many years to be overcome, an underlying genetic susceptibility in CF relating the antioxidant capacity is unlikely to play a central role in either the development of the disease or glucocorticoid insensitivity.

4.3 Kinase Signalling

The activation of signalling kinases is a fundamental part of physiologically controlled inflammation. However, many of these kinases such as p38 MAPK, GSK-3 β , ERK1/2 and JNK also regulate GR α activity and may also influence its gene specificity.^[101,102] The activity of ERK1/2, JNK and p38 MAPK are elevated in severe asthma compared with mild/moderate asthma and the elevation of p38 MAPK has also been linked to elevated phosphorylation of GR α leading to reduced ligand-binding affinity.^[103,104] COPD patients also display altered kinase signalling with an elevation in the activity of both p38 MAPK and JNK compared with healthy smokers.^[105,106] Elevated p38 MAPK may also be directly responsible for reduced GR α function in severe asthma through a reduction in the expression of MKP-1.^[49] Altered kinase signalling may also diminish GR α function indirectly through impairment of HDAC-2 activity by elevated PI3K δ signalling in COPD.^[81,107]

In CF, the absence of the CFTR induces an intrinsic activation of proinflammatory transcription factors including AP-1 and NF- κ B by mechanisms involving ERK1/2 and IKK signalling pathways.^[108] This activation is then consequently sustained via autocrine stimulation by proinflammatory mediators such as IL-1 β .^[108] Both ERK1/2 and p38 MAPK may also be involved in the enhanced proinflammatory response in CFTR defective cells.^[109] Further studies are needed to directly investigate the activation of these kinases in CF in relation to altered GR α function and glucocorticoid insensitivity, and the negative feedback regulatory mechanisms utilised by GR α such as MKP-1 expression.

4.4 GR β Expression

Classically, GR β is perceived as a negative regulator of GR α and its level of expression would thereby dictate the relative functional effectiveness of GR α . Consequently, the expression of GR β correlates with effective glucocorticoid function, particularly in cells/tissues.^[75,110] It is therefore likely that a high GR β to GR α ratio in disease would impair GR α function contributing to glucocorticoid insensitivity. However, the level of GR β in disease is a controversial area. Glucocorticoid insensitivity in severe asthma was initially found to be independent of GR α and GR β expression.^[111] Other studies have shown a relationship between GR β expression and severe asthma.^[112] In COPD, the expression of GR α is reduced with no elevation of GR β levels.^[81,113] No studies have looked at the expression of GR α or GR β in CF. Therefore, with so few studies assessing the relative expressions of GR α and GR β in these diseases it is difficult to assign a role for GR β in the development of glucocorticoid insensitivity with any confidence. More studies are therefore needed to clarify their relative expression in disease.

A role for GR β may also be found in light of recent evidence that it has an intrinsic transcriptional activity and may also be able to bind ligands.^[41,43] In light of these novel and important observations, the role of GR β in primary human cells and tissues, including those from disease, must be established. Thereafter, any potential impact on glucocorticoid insensitivity may be assessed with more confidence than a simplistic dominant negative regulation of GR α .

4.5 Neutrophils and Neutrophilic Inflammation

The majority of cells involved in the pathogenesis of lung diseases are responsive to glucocorticoids. However, neutrophils and neutrophilic inflammation are relatively unresponsive to glucocorticoid-mediated immunosuppression compared with other cell types.^[81,114,115] Studies have shown a relatively higher expression of GR β and a further induction of GR β expression upon stimulation, which could thereafter impair the functional

effectiveness of GR α and contribute to a relative reduction in glucocorticoid sensitivity.^[110] However, other studies have shown low expression of GR β and therefore the precise relative expression levels of both GR α and GR β in neutrophils remains both unresolved and controversial.^[116]

Importantly, neutrophil-associated exacerbations of severe asthma and COPD are also poorly controlled by glucocorticoids.^[114,115] However, it has not been determined whether this is an extension of the existing glucocorticoid insensitivity or if there is an additional unresponsiveness mediated by the acute elevation in neutrophils during the exacerbation. It is therefore feasible that the presence of neutrophils in both the stable and acute inflammatory responses in severe asthma, COPD and CF may contribute, in part, to their overall reduced responsiveness to glucocorticoid treatment.

Despite an apparent lack of glucocorticoid-mediated immunosuppression in neutrophils and neutrophilic inflammation, glucocorticoid treatment of neutrophils elicits GR α -dependent responses, most notably a delay in their constitutive apoptosis.^[117,118] These studies demonstrate that glucocorticoids are able to mediate important GR α -specific responses in neutrophils. Induction of these responses may have important clinical implications, particularly in severe asthma and COPD, which despite being relatively glucocorticoid insensitive, is still treated with high levels of glucocorticoids. Further investigations are needed to depict the relevance and possible roles of these GR α -dependent responses when treating a neutrophil-associated inflammatory response with glucocorticoids.

4.6 Impairment of GR α Transrepression or Transactivation, or Both?

Most functional studies concerning glucocorticoid insensitivity in models mimicking aspects of airway diseases such as asthma and COPD look at the ability of glucocorticoids to repress the pro-inflammatory genes, co-factors concerned with GR α transrepression or the inflammatory cell influx.^[26,49,80,81,90,92] A failure of glucocorticoids to repress these inflammatory factors is seen as glucocorticoid insensitivity. However, alterations in GR α transactivation in patients with relative gluco-

corticoid insensitivity has not been extensively assessed and may also be important, particularly as the potential contribution of GR α transactivation to the full anti-inflammatory actions of glucocorticoid in responsive airway diseases has not been established. Of course, the major reason that patients have limited or are withdrawn from glucocorticoid treatment is unacceptable side effects, which in turn are classically attributed to GR transactivation. Therefore, this would suggest that GR α transactivation is functional in these relative glucocorticoid-insensitive patients. However, recent evidence suggests that the regulation of specific genes by GR α may be dependent on GR α phosphorylation at specific sites.^[40,101,102] It could therefore be speculated that the phosphorylation status of GR α is altered in relative glucocorticoid-insensitive patients in such a manner that transcription of genes associated with its anti-inflammatory action are impaired but not those largely associated with mediating glucocorticoid-mediated adverse effects. Furthermore, studies using the GR α^{drim} mice, which are essentially devoid of GR α transactivation, show that these mice still have adverse effects from glucocorticoid treatment.^[119] Therefore, we cannot rule out both a substantial contribution of GR α transactivation to glucocorticoid-mediated immunosuppression in glucocorticoid-responsive airway diseases and thereafter any impairment of GR α transactivational immunosuppression in glucocorticoid-insensitive airway diseases. It is therefore essential that further studies attempt to depict the role of GR α transactivation in airway diseases as this may then influence future therapeutic strategies. Investigation using recently described dissociative glucocorticoids in patients with severe asthma and COPD may resolve many of these issues.^[120]

5. Therapeutic Approaches to Overcome Glucocorticoid Insensitivity in COPD and Asthma: Alternative Treatments and Restorative Approaches

5.1 Alternative Anti-Inflammatory Therapeutic Targets

Selective inhibition of single inflammatory mediators that are thought to be important in

COPD and asthma have not proven successful.^[11] This is likely to be due to the complex nature of the inflammatory responses in these diseases where removal of proinflammatory signalling derived from a single proinflammatory mediator may simply be compensated by the multitude of downstream intracellular signalling networks activated by the overall inflammatory response. Therefore, an effective alternative anti-inflammatory strategy must be directed against a target that has broad-ranging anti-inflammatory effects such as those seen with glucocorticoids. One attractive strategy that is being investigated is the inhibition of selective protein kinases, which act as signalling messengers for multiple inflammatory stimuli. These kinases are a pivotal part of multiple interconnecting inflammatory pathways and, therefore, regulate the expression of a plethora of different proinflammatory mediators.^[121,122] An obvious hurdle with targeting these kinases is that many are ubiquitously expressed and in addition to inflammatory signalling are also involved in critical physiological cellular processes. However, specific inhibition of selected isoforms and the relatively unique opportunity to deliver therapies locally to the lung may provide an avenue to overcome possible toxic effects. Two such protein kinase targets, phosphoinositol 3-kinase (PI3K) and p38 MAPK kinase, are discussed in the following sections.

5.1.1 p38 Mitogen-Activated Protein Kinase

There are four p38 MAPK isoforms, α , β , γ and δ , which are encoded by separate genes and are expressed in a tissue-dependent manner. p38 MAPK is a central mediator in many of the inflammatory signalling pathways, including activation of NF- κ B, regulation of GR α and marking out proinflammatory genes for early gene transcription through phosphorylation of histone 3 at serine 10.^[123] The specific roles of the individual p38 MAPK isoforms in the inflammatory response are unknown due to a lack of selective pharmacological tools; however, selective inhibition of the p38 MAPK α isoform potently inhibits the release of many proinflammatory cytokines including TNF α , IL-6, CXCL-8 and GM-CSF.^[124,125]

Many of these pathways and genes are prominent in the relatively glucocorticoid-insensitive inflammatory responses seen in severe asthma, COPD and CF.^[51] Selective inhibition of p38 MAPK also reduced the inflammatory response in nasal biopsies from CF patients, including CXCL8 release, COX-2 upregulation and neutrophil migration, although the contribution of the individual p38 MAPK isoforms has yet to be assessed.^[126] Therefore, selective inhibition of p38 MAPK may also prove to be an effective anti-inflammatory therapeutic strategy in CF.

As oxidative stress and oxidant-driven inflammation is such a prominent component of these diseases, which can not only alter drug targeting but may also induce glucocorticoid insensitivity, it is critical to assess if these inhibitors are able to suppress oxidant-driven inflammatory responses where glucocorticoids fail.^[69,127] Encouragingly, inhibition of p38 MAPK α also reduces the inflammatory response in cigarette smoke-mediated glucocorticoid-insensitive models *in vivo*.^[128] In addition, inhibition of p38 MAPK may also improve glucocorticoid function in PBMCs from COPD patients.^[129]

Consequently, direct inhibition of p38 MAPK has been proposed as a potential novel anti-inflammatory strategy and a number of selective p38 MAPK α inhibitors are currently under development for respiratory disease.^[122]

5.1.2 PI3K δ/γ

The lipid kinases PI3K δ and PI3K γ have a relatively restricted expression to mast cells and leukocytes, and are central in mediating both the innate and adaptive inflammatory responses.^[130] Selective inhibition of these PI3K isoforms has proven particularly successful in models of allergic inflammation by reducing airway hyperresponsiveness, eosinophilia and mast cell degranulation, as well as B- and T-cell function.^[130-135] Consequently, a number of small-molecule inhibitors of these isoforms are currently under development for asthma.^[136]

In contrast to allergy, the roles of these isoforms in diseases that are driven by innate (T_H1) immunity such as COPD and CF, or that have a T_H1 component such as severe asthma, have not

been directly studied. However, the impact of these isoforms on the key cells, including neutrophils and macrophages, involved in these diseases has been investigated. Neutrophil migration, directional chemotaxis and respiratory burst are dependent on PI3K γ and δ signalling.^[134,137,138] Monocyte recruitment into the lungs is also dependant on PI3K δ signalling.^[139] In addition, T_h1 and Tc1 cells are now proposed to be important in the orchestration of chronic inflammatory responses. PI3K signalling also appears to be important in T-cell trafficking and the retention of specific T cell types, such as Tc1 cells.^[140]

Therefore, the role of PI3K γ and δ in the regulation of the innate immune response and the ability to selectively target these isoforms presents an attractive and novel anti-inflammatory therapeutic strategy for the treatment of severe asthma, COPD and CF.

5.2 Therapeutic Targets for the Restoration of Glucocorticoid Function

In addition to novel anti-inflammatory targets, another strategy being pursued is the pharmacological restoration of glucocorticoid function, which would have the benefit of re-establishing an effective anti-inflammatory therapy that was well profiled in terms of application, drug interactions and adverse effects. Restoration of glucocorticoid function has been achieved both in *in vitro* and *in vivo* glucocorticoid-insensitive models and in primary cells from patients with disease through selective inhibition of the PI3K δ /Akt pathway.^[81,107] These targets along with additional strategies are discussed in the following sections.

5.2.1 Antioxidants

As oxidant stress is proposed to play a central role in not only the development and progression of COPD, severe asthma and CF but also in the mechanisms of relative glucocorticoid insensitivity, it would be obvious to directly target this oxidant stress with antioxidants. A number of studies have suggested that use of antioxidant supplements such as N-acetyl-cysteine (NAC) or

an increase in dietary antioxidants may be beneficial in COPD.^[141,142] This is an attractive therapeutic strategy; however, one of the major hurdles is that antioxidants have a poor bioavailability and attempts to deliver an effective concentration to the lungs has proven difficult, with little clinical benefit seen in COPD to date.^[143,144] If bioavailability and delivery hurdles are overcome and antioxidant drugs such as NAC are able to be delivered into the lung at an effective concentration, then it is likely that they would have a beneficial impact on the function of inhaled glucocorticoids.

5.2.2 Long-Acting β_2 Agonists

Long-acting β_2 agonists (LABAs) are bronchodilator drugs used in both asthma and COPD. Although their use as a monotherapy is controversial, their use as a combination therapy with inhaled glucocorticoids has been shown to have several beneficial effects, including improved glucocorticoid function.^[45,145] When inhaled glucocorticoid therapy alone fails to control asthma, addition of a LABA as a combination therapy improves symptom score, improves lung function, reduces the use of rapid-acting β_2 agonists and reduces the number of exacerbations.^[45] Several large-scale studies have demonstrated that use of LABAs in combination with inhaled glucocorticoids in patients with moderate-to-severe COPD improved lung function (although it does not influence its long-term deterioration), exacerbations, breathlessness and health status.^[145]

The exact molecular mechanisms of this improved glucocorticoid function remain unclear, but it is thought that LABAs may improve the penetration of inhaled glucocorticoids into lung cells. Although the evidence is clear that a combination therapy of LABA-inhaled glucocorticoid improves symptoms and lung function in both asthma and COPD compared with inhaled glucocorticoids alone,^[146] there is little evidence to suggest that their use results in a substantial improvement in glucocorticoid immunosuppression and ultimate disease progression. Therefore, the additional use of an effective anti-inflammatory is still required in these diseases.

5.2.3 Theophylline

Several studies have shown that the first-generation bronchodilator theophylline used at low concentrations restored glucocorticoid function in an HDAC-2-dependent manner.^[92,127,147,148] Enhancement of glucocorticoid function by theophylline has been seen clinically for a number of years where it has been used in patients with moderate to severe asthma.^[147,149] However, although theophylline may act through a number of pathways including that of PI3K δ /Akt (discussed in section 5.2.4), the exact target(s) by which theophylline acts at these low concentrations has not yet been established. Interestingly, the binding profile of theophylline is altered under conditions of oxidative stress, which may explain why it has no apparent impact on HDAC-2 activity in non-oxidant stressed cells.^[127] Importantly, this finding may also have wider implications in terms of drug binding and targeting in diseases with a prominent component of oxidative stress and should be taken into consideration in the development of therapies for these diseases.

5.2.4 PI3K δ

In models of oxidant-mediated glucocorticoid insensitivity, inhibition of the PI3K/Akt pathway also restores glucocorticoid function in an HDAC-2-dependent manner.^[81] Our studies using selective small molecule inhibitors and transgenic mice in oxidant-mediated glucocorticoid insensitivity have revealed that it is the PI3K δ isoform that is responsible for this action.^[81] Furthermore, selective inhibition of PI3K δ restores glucocorticoid function in primary cells from COPD patients as compared with age-matched smoking controls.^[107] The precise mechanisms by which inhibition of PI3K δ signalling restores glucocorticoid function is not clear but appears to involve the protection of HDAC-2 activity by reducing oxidant-mediated hyperphosphorylation and nitration.^[81] In addition, abolition of PI3K δ and γ signalling also protects the expression of other key components of the HDAC-2-containing co-repressor complexes (including Mi2 expression), which are recruited by GR α to mediate gene repression.

However, PI3K δ signalling is an early event occurring at the inner phospholipid layer of the cell surface membrane and is therefore not likely to directly interact with HDAC-2 or other co-repressors. PI3K δ signalling is activated by oxidative stress and is the major PI3K isoform responsible for activation of Akt.^[107,150] Akt may thereafter act directly or on signalling kinases further downstream, although extensive investigations are still needed to elucidate the signalling pathways that link oxidant activation of PI3K δ /Akt and their protective effects on HDAC-2/co-repressor expression and activity.

Therefore, as small-molecule inhibitors of PI3K δ and dual inhibitors of PI3K δ / γ are currently being considered for allergic disease, these findings represent a significant development in a possible therapeutic strategy for the restoration of glucocorticoid function in COPD, which may also be applicable to severe asthma and CF.

6. Conclusions

Relative glucocorticoid-insensitive inflammation is an unmet medical need in a number of diseases, including severe asthma, COPD and CF. The clinical significance of this relative glucocorticoid insensitivity in these diseases, which include disease management problems and cost burden, has led to increasing investigation into new effective anti-inflammatory strategies in what has been an understudied area.

Development of inhibitors against novel targets with broad-ranging anti-inflammatory functions such as p38 MAPK α and PI3K δ / γ , which may also act to restore the function of glucocorticoids, may provide an effective therapeutic strategy against the enhanced glucocorticoid-insensitive inflammation seen in these diseases. However, despite the identification of the importance of oxidant-mediated alterations in kinase signalling and co-repressor expression/activity, the precise mechanism(s) of the relative glucocorticoid insensitivity in these diseases still remain unknown. Due to the complexity of the inflammatory responses in these diseases, which include components of both the innate and adaptive immune systems, oxidant stress and

protease imbalance, it is likely that the mechanism(s) by which glucocorticoid function is impaired will be complex and multifaceted. However, given the similarities in aspects of the inflammation present in the lungs of patients with severe asthma, COPD and CF, there may be common mechanisms (such as those mediated by oxidant stress) that may provide a viable therapeutic strategy applicable across these diseases. It is therefore important that all of the potential mechanisms of relative glucocorticoid insensitivity in these diseases are elucidated to provide a clearer picture from which effective alternative anti-inflammatory and/or glucocorticoid-restorative therapies may be developed.

Acknowledgements

JAM holds funding from the European Respiratory Society and Medical Research Scotland (fellowship number 87). IMA and KFC hold funding from the EU, Medical Research Council and the Wellcome Trust. IMA has received honoraria for attending Advisory Board meetings or for consultancy work for Chiesi, GSK and Novartis with regard to treatment of asthma and COPD; he has also received research grants from GSK, AstraZeneca and Pfizer. KFC has received honoraria for attending Advisory Board meetings or for consultancy work for Gilead, GSK, Novartis, Boehringer Ingelheim and Merck with regard to treatment of asthma and COPD; he has also received research grants from GSK.

References

- Pujols L, Xaubert A, Ramirez J, et al. Expression of glucocorticoid receptors α and β in steroid sensitive and steroid insensitive interstitial lung diseases. *Thorax* 2004; 59: 687-93
- Barnes PJ, Adcock IM. Glucocorticoid resistance in inflammatory diseases. *Lancet* 2009; 373: 1905-17
- Beesley AH, Firth MJ, Ford J, et al. Glucocorticoid resistance in T-lineage acute lymphoblastic leukaemia is associated with a proliferative metabolism. *Br J Cancer* 2009; 100: 1926-36
- Pace TW, Hu F, Miller AH. Cytokine-effects on glucocorticoid receptor function: relevance to glucocorticoid resistance and the pathophysiology and treatment of major depression. *Brain Behav Immun* 2007; 21: 9-19
- Onda K, Rimbara E, Hirano T, et al. Role of mRNA expression of transcription factors in glucocorticoid sensitivity of peripheral blood mononuclear cells and disease state in rheumatoid arthritis. *J Rheumatol* 2004; 31: 464-9
- Langhoff E, Pedersen PS, Koch C. Methylprednisolone resistance of cystic fibrosis lymphocytes. *Pediatr Res* 1984; 18: 488-9
- Matysiak M, Makosa B, Walczak A, et al. Patients with multiple sclerosis resisted to glucocorticoid therapy: abnormal expression of heat-shock protein 90 in glucocorticoid receptor complex. *Mult Scler* 2008; 14: 919-26
- Mannino DM, Buist AS. Global burden of COPD: risk factors, prevalence, and future trends. *Lancet* 2007; 370: 765-73
- Chung F, Barnes N, Allen M, et al. Assessing the burden of respiratory disease in the UK. *Respir Med* 2002; 96: 963-75
- Barnes PJ. Immunology of asthma and chronic obstructive pulmonary disease. *Nat Rev Immunol* 2008; 8: 183-92
- Rennard SI, Fogarty C, Kelsen S, et al., on behalf of the COPD Investigators. The safety and efficacy of infliximab in moderate to severe chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2007; 175: 926-34
- Pratt WB, Toft DO. Steroid receptor interactions with heat shock protein and immunophilin chaperones. *Endocr Rev* 1997; 18: 306-60
- Murphy PJM, Morishima Y, Chen H, et al. Visualization and mechanism of assembly of a glucocorticoid receptor-Hsp70 complex that is primed for subsequent hsp90-dependent opening of the steroid binding cleft. *J Biol Chem* 2003; 278: 34764-73
- Kovacs JJ, Murphy PJM, Gaillard S, et al. HDAC6 regulates Hsp90 acetylation and chaperone-dependent activation of glucocorticoid receptor. *Mol Cell* 2005; 18: 601-7
- Rhen T, Cidlowski JA. Antiinflammatory action of glucocorticoids: new mechanisms for old drugs. *N Engl J Med* 2005; 353: 1711-23
- Reichardt HM, Tuckermann JP, Gottlicher M, et al. Repression of inflammatory responses in the absence of DNA binding by the glucocorticoid receptor. *EMBO J* 2001; 20: 7168-73
- Haller J, Mikics E, Makara GB. The effects of non-genomic glucocorticoid mechanisms on bodily functions and the central neural system: a critical evaluation of findings. *Front Neuroendocrinol* 2008; 29: 273-91
- Bartholome B, Spies C, Gaber T, et al. Membrane glucocorticoid receptors (mGCR) are expressed in normal human peripheral blood mononuclear cells and up-regulated after in vitro stimulation and in patients with rheumatoid arthritis. *FASEB J* 2004; 18: 70-80
- Scheinman RI, Gualberto A, Jewell CM, et al. Characterization of mechanisms involved in transrepression of NF-kappa B by activated glucocorticoid receptors. *Mol Cell Biol* 1995; 15: 943-53
- Li B, Carey M, Workman JL. The role of chromatin during transcription. *Cell* 2007; 128: 707-19
- Li J, Lin Q, Wang W, et al. Specific targeting and constitutive association of histone deacetylase complex during transcriptional repression. *Genes Dev* 2002; 16: 687-92
- John S, Sabo PJ, Johnson TA, et al. Interaction of the glucocorticoid receptor with the chromatin landscape. *Mol Cell* 2008; 29: 611-24
- Knoepfler PS, Eisenman RN. Sin meets NuRD and other tails of repression. *Cell* 1999; 99: 447-50
- Silverstein RA, Ekwall K. Sin3: a flexible regulator of global gene expression and genome stability. *Curr Genetics* 2005; 47: 1-17
- Denslow SA, Wade PA. The human Mi-2/NuRD complex and gene regulation. *Oncogene* 2007; 26: 5433-8

26. Ito K, Yamamura S, Essilfie-Quaye S, et al. Histone deacetylase 2-mediated deacetylation of the glucocorticoid receptor enables NF- κ B suppression. *J Exp Med* 2006; 203: 7-13
27. Allfrey VG, Mirsky AE. Structural modifications of histones and their possible role in the regulation of RNA synthesis. *Science* 1964; 144: 599
28. Usmani OS, Ito K, Manechotesuwan K, et al. Glucocorticoid receptor nuclear translocation in airway cells after inhaled combination therapy. *Am J Respir Crit Care Med* 2005; 172: 704-12
29. Culpitt SV, Maziak W, Loukidis S, et al. Effect of high dose inhaled steroid on cells, cytokines, and proteases in induced sputum in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1999; 160: 1635-9
30. Auphan N, DiDonato JA, Rosette C, et al. Immunosuppression by glucocorticoids: inhibition of NF-kappaB activity through induction of I kappa B synthase. *Science* 1995; 270: 286-90
31. King EM, Holden NS, Gong W, et al. Inhibition of NF- κ B-dependent Transcription by MKP-1. *J Biol Chem* 2009; 284: 26803-15
32. Abraham SM, Lawrence T, Kleiman A, et al. Antiinflammatory effects of dexamethasone are partly dependent on induction of dual specificity phosphatase 1. *J Exp Med* 2006; 203: 1883-9
33. Tuckermann JP, Kleiman A, Moriggl R, et al. Macrophages and neutrophils are the targets for immune suppression by glucocorticoids in contact allergy. *J Clin Invest* 2007; 117: 1381-90
34. Kumar R, Calhoun WJ. Differential regulation of the transcriptional activity of the glucocorticoid receptor through site-specific phosphorylation. *Biologics* 2008; 2: 845-54
35. Chen W, Dang T, Blind RD, et al. Glucocorticoid receptor phosphorylation differentially affects target gene expression. *Mol Endocrinol* 2008; 22: 1754-66
36. Miller AL, Webb MS, Copik AJ, et al. p38 Mitogen-activated protein kinase (MAPK) is a key mediator in glucocorticoid-induced apoptosis of lymphoid cells: correlation between p38 MAPK activation and site-specific phosphorylation of the human glucocorticoid receptor at serine 211. *Mol Endocrinol* 2005; 19: 1569-83
37. Krstic MD, Rogatsky I, Yamamoto KR, et al. Mitogen-activated and cyclin-dependent protein kinases selectively and differentially modulate transcriptional enhancement by the glucocorticoid receptor. *Mol Cell Biol* 1997; 17: 3947-54
38. Rogatsky I, Waase C, Garabedian MJ. Phosphorylation and inhibition of rat glucocorticoid receptor transcriptional activation by glycogen synthase kinase-3 (GSK-3). *J Biol Chem* 1998; 273: 14315-21
39. Szatmari Z, Garabedian MJ, Vil-iek J. Inhibition of glucocorticoid receptor-mediated transcriptional activation by p38 mitogen-activated protein (MAP) kinase. *J Biol Chem* 2004; 279: 43708-15
40. Duma D, Jewell CM, Cidlowski JA. Multiple glucocorticoid receptor isoforms and mechanisms of post-translational modification. *J Steroid Biochem Mol Biol* 2006; 102: 11-21
41. Lewis-Tuffin LJ, Jewell CM, Bienstock RJ, et al. Human glucocorticoid receptor β binds RU-486 and is transcriptionally active. *Mol Cell Biol* 2007; 27: 2266-82
42. Charmandari E, Chrousos GP, Ichijo T, et al. The human glucocorticoid receptor (hGR) β isoform suppresses the transcriptional activity of hGR α by interfering with formation of active coactivator complexes. *Mol Endocrinol* 2005; 19: 52-64
43. Kino T, Manoli I, Kelkar S, et al. Glucocorticoid receptor (GR) β has intrinsic, GR α -independent transcriptional activity. *Biochem Biophys Res Commun* 2009; 381: 671-5
44. Lu N, Cidlowski J. The origin and functions of multiple human glucocorticoid receptor isoforms. *Ann N Y Acad Sci* 2004; 1024: 102-23
45. Bateman ED, Hurd SS, Barnes PJ, et al. Global strategy for asthma management and prevention: GINA executive summary. *Eur Respir J* 2008; 31: 143-78
46. Macedo P, Hew M, Torrego A, et al. Inflammatory biomarkers in airways of patients with severe asthma compared with non-severe asthma. *Clin Exp Allergy* 2009; 39: 1668-76
47. Moore WC, Bleecker ER, Curran-Everett D, et al. Characterization of the severe asthma phenotype by the National Heart, Lung, and Blood Institute's Severe Asthma Research Program. *J Allergy Clin Immunol* 2007; 119: 405-13
48. Hew M, Bhavsar P, Torrego A, et al., for the National Heart Lung and Blood Institute's Severe Asthma Research Program. Relative corticosteroid insensitivity of peripheral blood mononuclear cells in severe asthma. *Am J Respir Crit Care Med* 2006; 174: 134-41
49. Bhavsar P, Hew M, Khorasani N, et al. Relative corticosteroid insensitivity of alveolar macrophages in severe asthma compared with non-severe asthma. *Thorax* 2008; 63: 784-90
50. Goleva E, Hauk PJ, Boguniewicz J, et al. Airway remodeling and lack of bronchodilator response in steroid-resistant asthma. *J Allergy Clin Immunol* 2007; 120: 1065-72
51. Barnes PJ. The cytokine network in asthma and chronic obstructive pulmonary disease. *J Clin Invest* 2008; 118: 3546-56
52. Kurashima K, Fujimura M, Ishiura Y, et al. Asthma severity is associated with an increase in both blood CXCR3+ and CCR4+ T cells. *Respirology* 2006; 11: 152-7
53. van Rensen ELJ, Sont JK, Evertse CE, et al., AMPUL Study Group. Bronchial CD8 cell infiltrate and lung function decline in asthma. *Am J Respir Crit Care Med* 2005; 172: 837-41
54. Jatakanon A, Uasuf C, Maziak W, et al. Neutrophilic inflammation in severe persistent asthma. *Am J Respir Crit Care Med* 1999; 160: 1532-9
55. Cho SH, Stanciu LA, Holgate ST, et al. Increased interleukin-4, interleukin-5, and interferon- γ in airway CD4+ and CD8+ T cells in atopic asthma. *Am J Respir Crit Care Med* 2005; 171: 224-30
56. Wood LG, Gibson PG. Reduced circulating antioxidant defences are associated with airway hyper-responsiveness, poor control and severe disease pattern in asthma. *Br J Nutr* 2010; 103 (5): 735-41
57. Fitzpatrick A, Brown L, Holguin F, et al. Levels of nitric oxide oxidation products are increased in the epithelial lining fluid of children with persistent asthma. *J Allergy Clin Immunol* 2009; 124: 990-6. e1-9

58. Tomlinson JEM, McMahon AD, Chaudhuri R, et al. Efficacy of low and high dose inhaled corticosteroid in smokers versus non-smokers with mild asthma. *Thorax* 2005; 60: 282-7
59. Thomson NC, Chaudhuri R, Livingston E. Asthma and cigarette smoking. *Eur Respir J* 2004; 24: 822-33
60. Chung KF, Adcock IM. Multifaceted mechanisms in COPD: inflammation, immunity, and tissue repair and destruction. *Eur Respir J* 2008; 31: 1334-56
61. Global Initiative for Chronic Obstructive Lung Disease. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease. NHLBI/WHO workshop report. Bethesda (MD): National Heart, Lung, and Blood Institute, 2008 Jan 1: 1-100. NIH publication no.: 2701
62. Hogg JC, Chu F, Utokaparch S, et al. The nature of small-airway obstruction in chronic obstructive pulmonary disease. *N Engl J Med* 2004; 350: 2645-53
63. Patel I, Seemungal T, Wilks M, et al. Relationship between bacterial colonisation and the frequency, character, and severity of COPD exacerbations. *Thorax* 2002; 57: 759-64
64. Sethi S. Bacterial infection and the pathogenesis of COPD. *Chest* 2000; 117: 286S-91S
65. Yang I, Fong K, Sim E, et al. Inhaled corticosteroids for stable chronic obstructive disease. *Cochrane Database Syst Rev* 2007; (2): CD002991
66. Suissa S, Ernst P, Vandemheen KL, et al. Methodological issues in therapeutic trials of COPD. *Eur Respir J* 2008; 31: 927-33
67. Keatings VM, Jatakanon A, Worsdell YM, et al. Effects of inhaled and oral glucocorticoids on inflammatory indices in asthma and COPD. *Am J Respir Crit Care Med* 1997; 155: 542-8
68. Bourbeau J, Christodouloupoulos P, Maltais F, et al. Effect of salmeterol/fluticasone propionate on airway inflammation in COPD: a randomised controlled trial. *Thorax* 2007; 62: 938-43
69. Culpitt SV, Rogers DF, Shah P, et al. Impaired inhibition by dexamethasone of cytokine release by alveolar macrophages from patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2003; 167: 24-31
70. O'Sullivan BP, Freedman SD. Cystic fibrosis. *Lancet* 2009; 373: 1891-904
71. Karp CL, Flick LM, Park KW, et al. Defective lipoxin-mediated anti-inflammatory activity in the cystic fibrosis airway. *Nat Immunol* 2004; 5: 388-92
72. Carrabino S, Carpani D, Livraghi A, et al. Dysregulated interleukin-8 secretion and NF- κ B activity in human cystic fibrosis nasal epithelial cells. *J Cyst Fibros* 2006; 5: 113-9
73. Freedman SD, Blanco PG, Zaman MM, et al. Association of cystic fibrosis with abnormalities in fatty acid metabolism. *N Engl J Med* 2004; 350: 560-9
74. Konstan MW. Ibuprofen therapy for cystic fibrosis lung disease: revisited. *Curr Opin Pulm Med* 2008; 14 (6): 567-73
75. Gross KL, Cidlowski JA. Tissue-specific glucocorticoid action: a family affair. *Trends Endocrinol Metab* 2008; 19: 331-9
76. Ciencewicz J, Trivedi S, Kleeberger SR. Oxidants and the pathogenesis of lung diseases. *J Allergy Clin Immunol* 2008; 122: 456-68
77. Hwang NR, Yim S, Kim YM, et al. Oxidative modifications of glyceraldehyde-3-phosphate dehydrogenase play a key role in its multiple cellular functions. *Biochem J* 2009; 423: 253-64
78. Kitagawa H, Yamaoka I, Akimoto C, et al. A reduction state potentiates the glucocorticoid response through receptor protein stabilization. *Genes Cells* 2007; 12: 1281-7
79. Wright VP, Reiser PJ, Clanton TL. Redox modulation of global phosphatase activity and protein phosphorylation in intact skeletal muscle. *J Physiol* 2009; 587: 5767-81
80. Ito K, Lim S, Caramori G, et al. Cigarette smoking reduces histone deacetylase 2 expression, enhances cytokine expression, and inhibits glucocorticoid actions in alveolar macrophages. *FASEB J* 2001; 15 (6): 1110-2
81. Marwick JA, Caramori G, Stevenson CS, et al. Inhibition of PI3K δ restores glucocorticoid function in smoking-induced airway inflammation in mice. *Am J Respir Crit Care Med* 2009; 179: 542-8
82. Rottner M, Freyssinet JM, Martinez MC. Mechanisms of the noxious inflammatory cycle in cystic fibrosis. *Respir Res* 2009; 10: 23
83. Rahman I, Marwick J, Kirkham P. Redox modulation of chromatin remodeling: impact on histone acetylation and deacetylation, NF- κ B and pro-inflammatory gene expression. *Biochem Pharmacol* 2004; 68: 1255-67
84. Marwick JA, Kirkham PA, Stevenson CS, et al. Cigarette smoke alters chromatin remodeling and induces proinflammatory genes in rat lungs. *Am J Respir Cell Mol Biol* 2004; 31: 633-42
85. Meja KK, Rajendrasozhan S, Adenuga D, et al. Curcumin restores corticosteroid function in monocytes exposed to oxidants by maintaining HDAC2. *Am J Respir Cell Mol Biol* 2008; 39: 312-23
86. Marwick JA, Ito K, Adcock IM, et al. Oxidative stress and steroid resistance in asthma and COPD: pharmacological manipulation of HDAC-2 as a therapeutic strategy. *Expert Opin Ther Targets* 2007; 11: 745-55
87. Ito K, Hanazawa T, Tomita K, et al. Oxidative stress reduces histone deacetylase 2 activity and enhances IL-8 gene expression: role of tyrosine nitration. *Biochem Biophys Res Commun* 2004; 315: 240-5
88. Osoata GO, Yamamura S, Ito M, et al. Nitration of distinct tyrosine residues causes inactivation of histone deacetylase 2. *Biochem Biophys Res Commun* 2009; 384: 366-71
89. Galasinski SC, Resing KA, Goodrich JA, et al. Phosphatase inhibition leads to histone deacetylases 1 and 2 phosphorylation and disruption of corepressor interactions. *J Biol Chem* 2002; 277: 19618-26
90. Cosio BG, Mann B, Ito K, et al. Histone acetylase and deacetylase activity in alveolar macrophages and blood monocytes in asthma. *Am J Respir Crit Care Med* 2004; 170: 141-7
91. Ito K, Ito M, Elliott WM, et al. Decreased histone deacetylase activity in chronic obstructive pulmonary disease. *N Engl J Med* 2005; 352: 1967-76

92. Cosio BG, Tsaprouni L, Ito K, et al. Theophylline restores histone deacetylase activity and steroid responses in COPD macrophages. *J Exp Med* 2004; 200: 689-95
93. Bartling TR, Drumm ML. Loss of CFTR results in reduction of histone deacetylase 2 in airway epithelial cells. *Am J Physiol Lung Cell Mol Physiol* 2009; 297: L35-43
94. Bartling TR, Drumm ML. Oxidative stress causes IL8 promoter hyperacetylation in cystic fibrosis airway cell models. *Am J Respir Cell Mol Biol* 2009; 40: 58-65
95. Gross KL, Lu NZ, Cidlowski JA. Molecular mechanisms regulating glucocorticoid sensitivity and resistance. *Mol Cell Endocrinol* 2009; 300: 7-16
96. Carmichael J, Paterson I, Diaz P, et al. Corticosteroid resistance in asthma. *BMJ* 1981; 282: 1419-22
97. Hakonarson H, Bjornsdottir US, Halapi E, et al. Profiling of genes expressed in peripheral blood mononuclear cells predicts glucocorticoid sensitivity in asthma patients. *Proc Natl Acad Sci U S A* 2005; 102: 14789-94
98. Smolonska J, Wijmenga C, Postma DS, et al. Meta-analyses on suspected chronic obstructive pulmonary disease genes: a summary of 20 years' research. *Am J Respir Crit Care Med* 2009; 180: 618-31
99. Young RP, Hopkins R, Black PN, et al. Functional variants of antioxidant genes in smokers with COPD and in those with normal lung function. *Thorax* 2006; 61: 394-9
100. Mak JCW, Ho SP, Yu WC, et al., on behalf of the Hong Kong Thoracic Society Chronic Obstructive Pulmonary Disease Study Group. Polymorphisms and functional activity in superoxide dismutase and catalase genes in smokers with COPD. *Eur Respir J* 2007; 30: 684-90
101. Galliher-Beckley AJ, Cidlowski JA. Emerging roles of glucocorticoid receptor phosphorylation in modulating glucocorticoid hormone action in health and disease. *IUBMB Life* 2009; 61: 979-86
102. Galliher-Beckley AJ, Williams JG, Collins JB, et al. Glycogen synthase kinase 3 β -mediated serine phosphorylation of the human glucocorticoid receptor redirects gene expression profiles. *Mol Cell Biol* 2008; 28: 7309-22
103. Irusen E, Matthews JG, Takahashi A, et al. p38 Mitogen-activated protein kinase-induced glucocorticoid receptor phosphorylation reduces its activity: role in steroid-insensitive asthma. *J Allergy Clin Immunol* 2002; 109: 649-57
104. Liu W, Liang Q, Balzar S, et al. Cell-specific activation profile of extracellular signal-regulated kinase 1/2, Jun N-terminal kinase, and p38 mitogen-activated protein kinases in asthmatic airways. *J Allergy Clin Immunol* 2008; 121: 893-902
105. Renda T, Baraldo S, Pelaia G, et al. Increased activation of p38 MAPK in COPD. *Eur Respir J* 2008; 31: 62-9
106. Rumora L, Milevoj L, Popovic-Grle S, et al. Levels changes of blood leukocytes and intracellular signalling pathways in COPD patients with respect to smoking attitude. *Clin Biochem* 2008; 41: 387-94
107. Marwick J, Caramori G, Adcock IM, et al. PI3Kdelta expression is increased in peripheral lung macrophages in COPD patients [abstract]. *Eur Respir J* 2009; 34 Suppl. 53: P917
108. Verhaeghe C, Remouchamps C, Hennuy B, et al. Role of IKK and ERK pathways in intrinsic inflammation of cystic fibrosis airways. *Biochem Pharmacol* 2007; 73: 1982-94
109. Muselet-Charlier C, Roque T, Boncoeur E, et al. Enhanced IL-1 β -induced IL-8 production in cystic fibrosis lung epithelial cells is dependent of both mitogen-activated protein kinases and NF- κ B signaling. *Biochem Biophys Res Commun* 2007; 357: 402-7
110. Strickland I, Kisich K, Hauk PJ, et al. High constitutive glucocorticoid receptor β in human neutrophils enables them to reduce their spontaneous rate of cell death in response to corticosteroids. *J Exp Med* 2001; 193: 585-94
111. Gagliardo R, Chanez P, Vignola AM, et al. Glucocorticoid receptor alpha and beta in glucocorticoid dependent asthma. *Am J Respir Crit Care Med* 2000; 162: 7-13
112. Bergeron C, Fukakusa M, Olivenstein R, et al. Increased glucocorticoid receptor- β expression, but not decreased histone deacetylase 2, in severe asthma. *J Allergy Clin Immunol* 2006; 117: 703-5
113. Korn SH, Thunnissen FB, Wesseling GJ, et al. Glucocorticoid receptor mRNA levels in bronchial epithelial cells of patients with COPD: influence of glucocorticoids. *Respir Med* 1998; 92: 1102-9
114. Ito K, Herbert C, Siegle JS, et al. Steroid-resistant neutrophilic inflammation in a mouse model of an acute exacerbation of asthma. *Am J Respir Cell Mol Biol* 2008; 39: 543-50
115. Barnes PJ. New molecular targets for the treatment of neutrophilic diseases. *J Allergy Clin Immunol* 2007; 119: 1055-62
116. Pujols L, Mulla J, Roca-Ferrer J, et al. Expression of glucocorticoid receptor alpha- and beta-isoforms in human cells and tissues. *Am J Physiol Cell Physiol* 2002; 283: C1324-31
117. Meagher LC, Cousin JM, Seckl JR, et al. Opposing effects of glucocorticoids on the rate of apoptosis in neutrophilic and eosinophilic granulocytes. *J Immunol* 1996; 156: 4422-8
118. Ward C, Chilvers ER, Lawson MF, et al. NF- κ B activation is a critical regulator of human granulocyte apoptosis in vitro. *J Biol Chem* 1999; 274: 4309-18
119. Kleiman A, Tuckermann JP. Glucocorticoid receptor action in beneficial and side effects of steroid therapy: lessons from conditional knockout mice. *Mol Cell Endocrinol* 2007; 275: 98-108
120. Schacke H, Berger M, Rehwinkel H, et al. Selective glucocorticoid receptor agonists (SEGRAs): novel ligands with an improved therapeutic index. *Mol Cell Endocrinol* 2007; 275: 109-17
121. Adcock I, Chung K, Caramori G, et al. Kinase inhibitors in airway inflammation. *Eur J Pharmacol* 2006; 533: 118-32
122. Kumar S, Boehm J, Lee JC. p38 MAP kinases: key signalling molecules as therapeutic targets for inflammatory diseases. *Nat Rev Drug Discov* 2003; 2: 717-26
123. Saccani S, Pantano S, Natoli G. p38-dependent marking of inflammatory genes for increased NF- κ B recruitment. *Nat Immunol* 2002; 3: 69-75
124. Tudhope SJ, Finney-Hayward TK, Nicholson AG, et al. Different mitogen-activated protein kinase-dependent cytokine responses in cells of the monocyte lineage. *J Pharmacol Exp Ther* 2008; 324: 306-12
125. Smith S, Fenwick P, Nicholson A, et al. Inhibitory effect of p38 mitogen-activated protein kinase inhibitors on cytokine release from human macrophages. *Br J Pharmacol* 2006; 149: 393-404

126. Raia V, Maiuri L, Ciacci C, et al. Inhibition of p38 mitogen activated protein kinase controls airway inflammation in cystic fibrosis. *Thorax* 2005; 60: 773-80
127. Marwick JA, Wallis G, Meja K, et al. Oxidative stress modulates theophylline effects on steroid responsiveness. *Biochem Biophys Res Commun* 2008; 377: 797-802
128. Medicherla S, Fitzgerald MF, Spicer D, et al. p38 α -Selective mitogen-activated protein kinase inhibitor sd-282 reduces inflammation in a subchronic model of tobacco smoke-induced airway inflammation. *J Pharmacol Exp Ther* 2008; 324: 921-9
129. Bhavsar PK, Khorasani N, Johnson M, et al. Reversal of relative corticosteroid insensitivity in PBMCs from patients with COPD by p38 MAPK inhibition [abstract]. *Am J Respir Crit Care Med* 2009; 179: A6187
130. Rommel C, Camps M, Ji H. PI3K δ and PI3K γ : partners in crime in inflammation in rheumatoid arthritis and beyond? *Nat Rev Immunol* 2007; 7: 191-201
131. Lee KS, Lee HK, Hayflick JS, et al. Inhibition of phosphoinositide 3-kinase δ attenuates allergic airway inflammation and hyperresponsiveness in murine asthma model. *FASEB J* 2006; 20: 455-65
132. Lee KS, Park SJ, Kim SR, et al. Phosphoinositide 3-kinase- δ inhibitor reduces vascular permeability in a murine model of asthma. *J Allergy Clin Immunol* 2006; 118: 403-9
133. Ali K, Bilancio A, Thomas M, et al. Essential role for the p110 δ phosphoinositide 3-kinase in the allergic response. *Nature* 2004; 431: 1007-11
134. Sasaki T, Irie-Sasaki J, Jones RG, et al. Function of PI3K in thymocyte development, T cell activation, and neutrophil migration. *Science* 2000; 287: 1040-6
135. Laffargue M, Calvez R, Finan P, et al. Phosphoinositide 3-kinase gamma is an essential amplifier of mast cell function. *Immunity* 2002; 16: 441-51
136. Marone R, Cmiljanovic V, Giese B, et al. Targeting phosphoinositide 3-kinase: moving towards therapy. *Biochim Biophys Acta* 2008; 1784: 159-85
137. Condliffe AM, Davidson K, Anderson KE, et al. Sequential activation of class IB and class IA PI3K is important for the primed respiratory burst of human but not murine neutrophils. *Blood* 2005; 106: 1432-40
138. Sadhu C, Masinovsky B, Dick K, et al. Essential role of phosphoinositide 3-kinase δ in neutrophil directional movement. *J Immunol* 2003; 170: 2647-54
139. Ferreira AM, Isaacs H, Hayflick JS, et al. The p110 δ isoform of PI3K differentially regulates β 1 and β 2 integrin-mediated monocyte adhesion and spreading and modulates diapedesis. *Microcirculation* 2006; 13: 439-56
140. Ward SG, Marelli-berg FM. Mechanisms of chemokine and antigen-dependent T-lymphocyte navigation. *Biochem J* 2009; 418: 13-27
141. Cai S, Chen P, Zhang C, et al. Oral N-acetylcysteine attenuates pulmonary emphysema and alveolar septal cell apoptosis in smoking-induced COPD in rats. *Respirology* 2009; 14: 354-9
142. Rahman I. Antioxidant therapeutic advances in COPD. *Ther Adv Respir Dis* 2008; 2: 351-74
143. Schermer T, Chavannes N, Dekhuijzen J, et al. Fluticasone and N-acetylcysteine in primary care patients with COPD or chronic bronchitis. *Respir Med* 2009; 103: 542
144. Stav D, Raz M. Effect of N-Acetylcysteine on air trapping in COPD. *Chest* 2009; 136: 381-6
145. Calverley PMA, Anderson JA, Celli B, et al. Salmeterol and fluticasone propionate and survival in chronic obstructive pulmonary disease. *N Engl J Med* 2007; 356: 775-89
146. Chung KF, Caramori C, Adcock IM. Inhaled corticosteroids as combination therapy with β -adrenergic agonists in airways disease: present and future. *Eur J Clin Pharmacol* 2009; 65: 835-71
147. Evans DJ, Taylor DA, Zetterstrom O, et al. Theophylline plus low dose inhaled steroid as effective as high dose steroid in the control of asthma. *N Engl J Med* 1997; 337: 1412-8
148. Ito K, Lim S, Caramori G, et al. A molecular mechanism of action of theophylline: induction of histone deacetylase activity to decrease inflammatory gene expression. *Proc Natl Acad Sci U S A* 2002; 99: 8921-6
149. Ukena D, Harnest U, Sakalauskas R, et al. Comparison of addition of theophylline to inhaled steroid with doubling of the dose of inhaled steroid in asthma. *Eur Respir J* 1997; 10: 2754-60
150. Molnarfi N, Brandt KJ, Gruaz L, et al. Differential regulation of cytokine production by PI3K δ in human monocytes upon acute and chronic inflammatory conditions. *Mol Immunol* 2008; 45: 3419-27

Correspondence: Dr John Marwick, MRC Centre for Inflammation Research, Queens Medical Research Institute, University of Edinburgh Medical School, 47 Little France Crescent, Edinburgh, EH16 4TJ, UK.
E-mail: John.Marwick@ed.ac.uk