



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Short communication: Methicillin-resistant *Staphylococcus aureus* detection in US bulk tank milk

Citation for published version:

Virgin, JE, Van Slyke, TM, Lombard, JE & Zadoks, RN 2009, 'Short communication: Methicillin-resistant *Staphylococcus aureus* detection in US bulk tank milk' *Journal of Dairy Science*, vol. 92, no. 10, pp. 4988-4991. DOI: 10.3168/jds.2009-2290

Digital Object Identifier (DOI):

[10.3168/jds.2009-2290](https://doi.org/10.3168/jds.2009-2290)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Journal of Dairy Science

Publisher Rights Statement:

© American Dairy Science Association, 2009

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Short communication: Methicillin-resistant *Staphylococcus aureus* detection in US bulk tank milk

J. E. Virgin,* T. M. Van Slyke,† J. E. Lombard,*¹ and R. N. Zadoks†²

*USDA:APHIS:VS, Centers for Epidemiology and Animal Health, 2150 Centre Ave., Bldg B, Fort Collins, CO 80526-8117

†Quality Milk Production Services, College of Veterinary Medicine, Cornell University, Ithaca, NY 14850-1263

ABSTRACT

Staphylococcus aureus is a major cause of mastitis in dairy cattle. This study estimated the herd prevalence of methicillin-resistant *Staph. aureus* (MRSA) among US dairy herds by testing bulk tank milk (BTM) samples using genotypic and phenotypic methods. A nationally representative sample of 542 operations had BTM cultured for *Staph. aureus*, and 218 BTM samples were positive upon initial culture. After 4 wk to 4 mo of frozen storage, 87% of 218 samples (n = 190) were still culture positive for *Staph. aureus* on blood agar, but none were positive for MRSA on the selective indicator medium CHROMagar MRSA. A duplex PCR was used to detect the *Staph. aureus*-specific *nuc* gene and the methicillin resistance gene, *mecA*, in mixed staphylococcal isolates from the 190 BTM samples that were positive for *Staph. aureus* after storage. Seven samples tested positive for *nuc* and *mecA*, and 2 samples tested positive for *mecA* only. *MecA*-positive *Staphylococcus* spp., but not MRSA, were subsequently isolated from 5 samples, whereas neither *mecA*-positive *Staphylococcus* spp. nor MRSA was isolated from the remaining 4 samples. Presence of methicillin-resistant, coagulase-negative *Staphylococcus* spp. may complicate the detection of MRSA by means of PCR on BTM. Bulk tank milk in the United States is not a common source of MRSA.

Key words: methicillin-resistant *Staphylococcus aureus* (MRSA), dairy cow, bulk tank milk, NAHMS

Methicillin-resistant *Staphylococcus aureus* (MRSA) is a major cause of nosocomial infections in humans. In addition, community-acquired MRSA has now become a major concern (Otter and French, 2008). New evidence also suggests that domestic animals, including food animals, are capable of serving as reservoirs and

shedders of MRSA and that transmission between host species may be possible (Loo et al., 2007). Methicillin-resistant *Staph. aureus* has been detected in the milk of cows with mastitis, dogs, cats, horses, pigs, sheep, goats, rabbits, chickens, and exotic species (Lee, 2003; Leonard and Markey, 2006). Recent studies have shown genetic similarity between MRSA isolates from food animals, including dairy cows, and those in humans, suggesting a mode of transmission between the 2 species (Juhász-Kaszanyitzky et al., 2007; Moon et al., 2007). Infection of dairy cattle with MRSA has been attributed to human-to-animal transfer, but the directionality of transmission is not always known (Devriese and Hommez, 1975; Juhász-Kaszanyitzky et al., 2007). Although *Staph. aureus* is a common mastitis pathogen, there has been little evidence to suggest that MRSA is common in milk. Furthermore, the risk of dairy product-borne transmission of MRSA is easily minimized by pasteurization, and most reported instances of foodborne MRSA have occurred through contamination by infected food handlers rather than the food itself (Kluytmans et al., 1995; Jones et al., 2002).

Previous research indicates that MRSA is rarely found in milk from dairy cattle in the United States. Analysis of 2,978 *Staph. aureus* isolates from milk samples submitted to diagnostic laboratories in Michigan and Wisconsin between 1994 and 2001 showed that 1.4% of isolates were resistant to oxacillin (Erskine et al., 2002; Makovec and Ruegg, 2003). Resistance to oxacillin was not detected among 357 *Staph. aureus* isolates recovered from milk in North Carolina (Anderson et al., 2006). A survey of milk samples from cows with mastitis from 153 dairies in Korea, however, yielded 21 (2.5%) MRSA positives and 19 (2.4%) methicillin-resistant CNS (Moon et al., 2007). Further studies are needed in the United States to ascertain the prevalence of MRSA in food animals, especially dairy cattle. The objective of this study was to determine the prevalence of MRSA detected in bulk tank milk (BTM) samples that were collected as part of the National Animal Health Monitoring System's (NAHMS) Dairy 2007 study and represented the diversity of dairy operations in the United States. Bulk tank milk samples were sub-

Received April 13, 2009.

Accepted June 30, 2009.

¹Corresponding author: jason.e.lombard@aphis.usda.gov

²Current address: Royal (Dick) School of Veterinary Studies, The University of Edinburgh, Easter Bush Veterinary Centre, Roslin, Scotland, UK.

mitted to Quality Milk Production Services (QMPS, Ithaca, NY) for detection of mastitis pathogens and MRSA. We hypothesized that *Staphylococcus aureus* would be isolated from BTM samples but the prevalence of MRSA would be low.

The NAHMS Dairy 2007 study was conducted in 17 of the top US dairy-producing states. The regions (states) included West (California, Idaho, New Mexico, Texas, and Washington) and East (Indiana, Iowa, Kentucky, Michigan, Minnesota, Missouri, New York, Ohio, Pennsylvania, Vermont, Virginia, and Wisconsin). Herds were categorized by the number of milk cows into small (1–99 cows), medium (100–499 cows), and large (500+ cows). Phase I of the study covered the population of dairy operations with at least one milk cow in the 17 participating states. As of January 1, 2007, these operations accounted for 82.5% of milk cows and 79.5% of dairy operations in the United States. Collection of BTM occurred during phase II of the study and included those operations in phase I that elected to continue participating and had at least 30 milk cows. All respondent data were statistically weighted to reflect the population of dairy operations with 30 or more cows in the 17 participating states. The inverse of the probability of selection for each operation was the initial selection weight. The selection weight was adjusted for nonresponse within each state and size group to allow for inferences back to the original sample population. Weights were again adjusted to account for nonrespondents in phase II. Of the 3,554 operations in the original sample, 2,194 completed phase I and of the 1,073 eligible operations, 582 completed phase II. Between February 28 and August 30, 2007, approximately 100 mL of bulk tank milk was aseptically collected from 542 of the 582 operations that completed phase II. Samples were packed in ice and shipped overnight to QMPS.

The BTM samples were received by the QMPS bacteriology laboratory and cultured for the presence of mastitis pathogens including *Staph. aureus* using routine methods (Hogan et al., 1999; Zadoks et al., 2004). The samples then were stored at -20°C for 4 wk to 4 mo until further processing. To detect MRSA in BTM, phenotypic and genotypic methods were used. Phenotypic detection was based on plating on a selective indicator medium, BBL CHROMagar MRSA (BD Diagnostic Systems, Sparks, MD; Han et al., 2007). Samples that tested positive for *Staph. aureus* upon initial routine culture ($n = 218$) were thawed at room temperature and streaked onto CHROMagar MRSA using sterile cotton swabs. As positive controls, MRSA strain ATCC 700698 and strain QMP S1-027, a MRSA isolate obtained from heifer milk and confirmed by *nuc* and *mecA* PCR (gift of John Barlow, University of Ver-

mont, Burlington), were used. Plates were incubated aerobically at 35 to 37°C and evaluated after 24 ± 4 h. Presence and appearance of colonies were recorded, and special attention was paid to colony color. Mauve colonies on CHROMagar MRSA were observed for both control strains, but not for any of the BTM samples tested. Blue colonies were observed on a large proportion of CHROMagar MRSA plates ($n = 145$, or 67% of BTM samples tested), and pale pink colonies or blue colonies with a pink halo were observed on 11 plates (5% of samples tested). Based on microscopic evaluation of colony morphology, catalase and coagulase testing, all pink colonies, all blue colonies with a pink halo, and an arbitrarily selected subset of blue colonies were identified to the genus level. Colonies belonged to the genera *Streptococcus* (blue colonies with or without pink halo), *Corynebacterium* (pale pink colonies), or *Staphylococcus* (pale pink colonies; all coagulase negative). A large proportion of BTM samples contain *Streptococcus* spp., *Staphylococcus* spp., and *Corynebacterium* spp. (Zadoks et al., 2004). *Staphylococcus* and *Corynebacterium* species are known to give occasional false-positive results on CHROMagar MRSA (Han et al., 2007). *Streptococcus* spp., *Staphylococcus* spp., and *Corynebacterium* spp. were phenotypically distinct from MRSA isolates in the current study, but growth of such colonies on CHROMagar MRSA complicates interpretation of culture results and may lead to false-positive readings. Thus, CHROMagar MRSA, which was developed for detection of MRSA in nasal swabs, may not be suitable for routine detection of MRSA in BTM.

In a parallel assay, the thawed milk samples were plated on trypticase soy agar with 5% sheep blood and 0.1 esculin (TSA-BE; PML Microbiologicals, Mississauga, Ontario, Canada) to obtain staphylococcal colonies for subsequent genotypic detection of MRSA based on PCR using *Staph. aureus* and *mecA*-specific primer sets (Brakstad et al., 1992; Martineau et al., 2000). A mixture of staphylococcal colonies was used to represent the mixed staphylococcal flora that can be present in bulk milk samples. For samples that tested positive for *nuc* and *mecA*, PCR was repeated on subcultures of isolated staphylococcal colonies to determine whether the 2 genes were present in the same isolate or in different isolates. This stepwise approach enabled us to process a large number of samples at low cost. For screening of the BTM samples, mixed staphylococcal colonies from TSA-BE were suspended in $1 \times$ Tris-EDTA buffer (Promega, Madison, WI). An aliquot of the suspension was used as DNA template in PCR reactions and the remainder was stored at 4°C . The *nuc* primers (NUC1 = 5'-GCG ATT GAT GGT GAT ACG GTT-3' and NUC2 = 5'-AGC CAA GCC TTG ACG AAC TAA AGC-3') were used to detect the gene encoding the thermostable

nuclease of *Staph. aureus* (Brakstad et al., 1992). The *mecA* primers (*mecA174* = 5'-AAC AGG TGA ATT ATT AGC ACT TGT AAG-3' and *mecA174B* = 5'-ATT GCT GTT AAT ATT TTT TGA GTT GAA-3') were used to detect the methicillin resistance gene (Martineau et al., 2000). Reactions were run in a total volume of 25 μ L consisting of 2 μ L of template, 12.5 μ L of GoTaq Green (Promega), 0.125 μ L of each primer (100 mM), and 10 μ L of nuclease-free dH₂O, using an iCycler (BioRad Laboratories, Hercules, CA) and the following run parameters: initial denaturation at 94°C for 15 min, followed by 37 cycles at 94°C for 60 s, 55°C for 30 s, and 72°C for 60 s, and final extension at 72°C for 10 min. In each PCR assay, the 2 MRSA control strains and 2 negative controls were included, one for lysate preparation and one for the PCR reaction itself. Presence of *nuc* and *mecA* PCR products, which are 270 and 174 bp long, respectively, was determined through electrophoresis in 1.5% agarose gels. Of the 218 BTM samples that had been positive for *Staph. aureus* upon initial culture (i.e., before frozen storage of the milk samples), 190 (87.2%) tested positive for *Staph. aureus* using the PCR method on a mixture of staphylococcal colonies from the poststorage culture. The failure to detect *Staph. aureus* in 28 (12.8%) of samples after frozen storage may be the result of loss of viability during storage, random variability in detection of *Staph. aureus* in samples with low numbers of colony-forming units, or lack of sensitivity of the culture and PCR-based methods used after frozen storage.

Nine colony suspensions tested positive for *mecA*. In 7 samples, *nuc* and *mecA* were detected, and in 2 samples *mecA* but not *nuc* was detected. Detection of *mecA* in the absence of *nuc* can be explained by the presence of methicillin-resistant *Staphylococcus* spp. other than *Staph. aureus*. Methicillin resistance is more common in non-aureus staphylococci from milk samples than in *Staph. aureus* (Makovec and Ruegg, 2003). Given that detection of *mecA* can be due to presence of *Staphylococcus* spp., there was a need to determine whether *nuc*-positive, *mecA*-positive samples contained MRSA or a mixture of methicillin-resistant *Staphylococcus* spp. and methicillin-susceptible *Staph. aureus* (MSSA). To this end, the stored portion of the colony suspensions that had been used to set up PCR reactions was plated on TSA-BE and incubated overnight at 37°C for samples that had yielded *mecA* amplicons. After incubation, individual staphylococcal colonies were streaked for isolation onto fresh TSA-BE to obtain pure cultures, which were subsequently tested by *nuc-mecA* PCR to determine whether MSSA, MRSA, or methicillin-resistant *Staphylococcus* spp. had been present in the suspension. Methicillin-resistant *Staphylococcus* spp. but not MRSA were obtained from 5 BTM samples. No

mecA-positive colonies were isolated from the remaining 4 samples. One of those samples had tested negative for *nuc*, suggesting that methicillin-resistant *Staphylococcus* spp. was the source of the *mecA* gene. For the remaining 3 samples, we could not determine whether the *mecA* gene originated from *Staphylococcus* spp. or *Staph. aureus*. Testing of isolated rather than mixed colonies from BTM samples would preclude the need to determine whether *mecA*-positive results were due to MRSA or *Staphylococcus* spp. Given the high prevalence of *Staph. aureus* in BTM and the low prevalence of *mecA*, such an approach might be cost-prohibitive in routine veterinary diagnostic laboratories.

On the basis of our results we conclude that 1) MRSA could not be detected in nationally representative BTM samples from the NAHMS Dairy 2007 study using phenotypic (CHOMagar MRSA) and genotypic (*nuc-mecA* PCR) methods, suggesting that BTM is not a common source of MRSA in the United States; 2) the *mecA* gene that encodes methicillin resistance does occur in staphylococci isolated from BTM; 3) presence of *mecA* in BTM isolates often is due to presence of methicillin-resistant *Staphylococcus* spp. other than *Staph. aureus*; and 4) there is a need for validated, cheap, and fast tools to detect MRSA in BTM routinely and to differentiate it from other methicillin-resistant *Staphylococcus* spp. that may be present in BTM. Such a tool could contribute to routine monitoring of milk quality and safety.

REFERENCES

- Anderson, K. L., R. L. Lyman, S. M. Bodeis-Jones, and D. G. White. 2006. Genetic diversity and antimicrobial susceptibility profiles among mastitis-causing *Staphylococcus aureus* isolated from bovine milk samples. *Am. J. Vet. Res.* 67:1185–1191.
- Brakstad, O. G., K. Aasbakk, and J. A. Maeland. 1992. Detection of *Staphylococcus aureus* by polymerase chain reaction amplification of the *nuc* gene. *J. Clin. Microbiol.* 30:1654–1660.
- Devriese, L. A., and J. Hommez. 1975. Epidemiology of methicillin-resistant *Staphylococcus aureus* in dairy herds. *Res. Vet. Sci.* 19:23–27.
- Erskine, R. J., R. D. Walker, C. A. Bolin, P. C. Bartlett, and D. G. White. 2002. Trends in antibacterial susceptibility of mastitis pathogens during a seven-year period. *J. Dairy Sci.* 85:1111–1118.
- Han, Z., E. Lautenbach, N. Fishman, and I. Nachamkin. 2007. Evaluation of mannitol salt agar, CHROMagar *Staph aureus* and CHROMagar MRSA for detection of methicillin-resistant *Staphylococcus aureus* from nasal swab specimens. *J. Med. Microbiol.* 56:43–46.
- Hogan, J. S., R. N. González, R. J. Harmon, S. C. Nickerson, S. P. Oliver, J. W. Pankey, and K. L. Smith. 1999. Laboratory Handbook on Bovine Mastitis. Rev. ed. National Mastitis Council, Madison, WI.
- Jones, T. F., M. Kellum, S. Porter, M. Bell, and W. Schaffner. 2002. An outbreak of community-acquired foodborne illness caused by methicillin-resistant *Staphylococcus aureus*. *Emerg. Infect. Dis.* 8:82–84.
- Juhász-Kaszanyitzky, E., S. Janosi, P. Somogyi, A. Dan, L. Van der Graaf-van Bloois, E. Van Duijkeren, and J. A. Wagenaar. 2007.

- MRSA transmission between cows and humans. *Emerg. Infect. Dis.* 13:630–632.
- Kluytmans, J., W. van Leeuwen, W. Goessens, R. Hollis, S. Messer, L. Herwaldt, H. Bruining, M. Heck, J. Rost, N. van Leeuwen, A. Van Belkum, and H. Verbrugh. 1995. Food-initiated outbreak of methicillin resistant *Staphylococcus aureus* analyzed by pheno- and genotyping. *J. Clin. Microbiol.* 33:1121–1128.
- Lee, J. H. 2003. Methicillin (oxacillin)-resistant *Staphylococcus aureus* strains isolated from major food animals and their potential transmission to humans. *Appl. Environ. Microbiol.* 69:6489–6494.
- Leonard, F. C., and B. K. Markey. 2006. Methicillin-resistant *Staphylococcus aureus* in animals: A review. *Vet. J.* 175:17–36.
- Loo, I., X. Huijsdens, E. Tiermersma, A. Neeling, N. van de Sande-Bruinsma, D. Beaujean, A. Voss, and J. Klutmans. 2007. Emergence of methicillin-resistant *Staphylococcus aureus* of animal origin in humans. *Emerg. Infect. Dis.* 13:1834–1839.
- Makovec, J. A., and P. L. Ruegg. 2003. Antimicrobial resistance of bacteria isolated from dairy cow milk samples submitted for bacterial culture: 8,905 samples (1994–2001). *J. Am. Vet. Med. Assoc.* 222:1582–1589.
- Martineau, F., F. J. Picard, N. Lansac, C. Menard, P. H. Roy, M. Ouellette, and M. G. Bergeron. 2000. Correlation between the resistance genotype determined by multiplex PCR assays and the antibiotic susceptibility patterns of *Staphylococcus aureus* and *Staphylococcus epidermidis*. *Antimicrob. Agents Chemother.* 44:231–238.
- Moon, J. S., A. R. Lee, H. M. Kang, E. S. Lee, M. N. Kim, Y. H. Paik, Y. H. Park, Y. S. Joo, and H. C. Koo. 2007. Phenotypic and genetic antibiogram of methicillin-resistant *Staphylococci* isolated from bovine mastitis in Korea. *J. Dairy Sci.* 90:1176–1185.
- Otter, J. A., and G. L. French. 2008. The emergence of community-associated methicillin-resistant *Staphylococcus aureus* at a London teaching hospital, 2000–2006. *Clin. Microbiol. Infect.* 14:670–676.
- Zadoks, R. N., R. N. González, K. J. Boor, and Y. H. Schukken. 2004. Mastitis-causing streptococci are important contributors to bacterial counts in raw bulk tank milk. *J. Food Prot.* 67:2644–2650.