



# THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### Entrainment to feeding but not to light: Circadian phenotype of VPAC(2) receptor-null mice

**Citation for published version:**

Sheward, WJ, Maywood, ES, French, KL, Horn, JM, Hastings, MH, Seckl, JR, Holmes, MC & Harmar, AJ 2007, 'Entrainment to feeding but not to light: Circadian phenotype of VPAC(2) receptor-null mice' *Journal of Neuroscience*, vol. 27, no. 16, pp. 4351-4358. DOI: 10.1523/JNEUROSCI.4843-06.2007

**Digital Object Identifier (DOI):**

[10.1523/JNEUROSCI.4843-06.2007](https://doi.org/10.1523/JNEUROSCI.4843-06.2007)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Publisher's PDF, also known as Version of record

**Published In:**

*Journal of Neuroscience*

**Publisher Rights Statement:**

Copyright © 2013 by the Society for Neuroscience

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



# Entrainment to Feeding but Not to Light: Circadian Phenotype of VPAC<sub>2</sub> Receptor-Null Mice

W. John Sheward,<sup>1,2</sup> Elizabeth S. Maywood,<sup>3</sup> Karen L. French,<sup>2</sup> Jacqueline M. Horn,<sup>1,2</sup> Michael H. Hastings,<sup>3</sup> Jonathan R. Seckl,<sup>2</sup> Megan C. Holmes,<sup>1,2</sup> and Anthony J. Harmar<sup>1,2</sup>

<sup>1</sup>Centre for Neuroscience Research, School of Biomedical Sciences, University of Edinburgh, Edinburgh, EH8 9JZ, United Kingdom, <sup>2</sup>Centre for Cardiovascular Science, University of Edinburgh, Queen's Medical Research Institute, Edinburgh EH16 4TJ, United Kingdom, and <sup>3</sup>Medical Research Council Laboratory of Molecular Biology, Cambridge CB2 2QH, United Kingdom

The master clock driving mammalian circadian rhythms is located in the suprachiasmatic nuclei (SCN) of the hypothalamus and entrained by daily light/dark cycles. SCN lesions abolish circadian rhythms of behavior and result in a loss of synchronized circadian rhythms of clock gene expression in peripheral organs (e.g., the liver) and of hormone secretion (e.g., corticosterone). We examined rhythms of behavior, hepatic clock gene expression, and corticosterone secretion in VPAC<sub>2</sub> receptor-null (*Vipr2*<sup>-/-</sup>) mice, which lack a functional SCN clock. Unexpectedly, although *Vipr2*<sup>-/-</sup> mice lacked robust circadian rhythms of wheel-running activity and corticosterone secretion, hepatic clock gene expression was strongly rhythmic, but advanced in phase compared with that in wild-type mice. The timing of food availability is thought to be an important entrainment signal for circadian clocks outside the SCN. *Vipr2*<sup>-/-</sup> mice consumed food significantly earlier in the 24 h cycle than wild-type mice, consistent with the observed timing of peripheral rhythms of circadian gene expression. When restricted to feeding only during the daytime (RF), mice develop rhythms of activity and of corticosterone secretion in anticipation of feeding time, thought to be driven by a food-entrainable circadian oscillator, located outside the SCN. Under RF, mice of both genotypes developed food-anticipatory rhythms of activity and corticosterone secretion, and hepatic gene expression rhythms also became synchronized to the RF stimulus. Thus, food intake is an effective zeitgeber capable of coordinating circadian rhythms of behavior, peripheral clock gene expression, and hormone secretion, even in the absence of a functional SCN clock.

**Key words:** circadian rhythms; corticosterone; entrainment; feeding; HPA axis; liver; VIP

## Introduction

The master clock driving mammalian circadian rhythms resides within the suprachiasmatic nuclei (SCN) of the hypothalamus. However, many peripheral tissues also contain endogenous circadian clocks that are thought to play critical roles in the local control of physiology and metabolism (Balsalobre et al., 1998; Zylka et al., 1998; Damiola et al., 2000; Yamazaki et al., 2000; McNamara et al., 2001; Abe et al., 2002; Yoo et al., 2004). For example, in the liver, there are circadian rhythms in the expression of genes involved in nutrient metabolism, heme and glutamine biosynthesis, and drug detoxification (Kornmann et al., 2001; Reddy et al., 2006). The signals that entrain peripheral circadian clocks to the environment are unknown and are the subject of intense current investigation.

If rats and mice, which are nocturnal animals, are fed exclusively during the daytime [restricted feeding (RF)], the phase of circadian gene expression in some peripheral organs such as liver, kidney, heart, and pancreas becomes uncoupled from the SCN

master pacemaker (Damiola et al., 2000; Stokkan et al., 2001; Horikawa et al., 2005). Animals then display two periods of activity in each 24 h, one during the subjective night and another during the day, in anticipation of feeding [food-anticipatory activity (FAA)] (Mistlberger, 1994; Davidson et al., 2003; Dudley et al., 2003; Horikawa et al., 2005). RF also induces food-anticipatory components of other daily rhythms, including body temperature, plasma corticosterone, and drinking. These feeding-entrained rhythms persist for several days if feedings are omitted. Daily rhythms of FAA and of gastrointestinal and metabolic functions can be entrained by RF in SCN-lesioned rodents. These studies (for review, see Stephan, 2002) have led to the proposal that there is a food-entrainable circadian oscillator (FEO) that drives the increase in locomotor activity in anticipation of scheduled feeding and is distinct from the light-entrainable oscillator in the SCN.

Vasoactive intestinal peptide (VIP), acting through the VPAC<sub>2</sub> receptor (encoded by the *Vipr2* gene), plays a pivotal role in the control of circadian activity in the SCN, promoting rhythmicity and synchronizing pacemaking neurons (Aton et al., 2005; Maywood et al., 2006). Both *Vipr2*<sup>-/-</sup> and VIP-deficient mice display a severely disrupted circadian phenotype (Harmar et al., 2002; Colwell et al., 2003), showing markedly attenuated or abolished behavioral rhythms in constant darkness and running wheel behavior that is determined predominantly by the prevail-

Received Nov. 7, 2006; revised Feb. 28, 2007; accepted March 5, 2007.

This work was supported by the Medical Research Council. We thank Dr. Marty Straume for assistance with the use of COSOPT software.

Correspondence should be addressed to Anthony J. Harmar, Centre for Neuroscience Research, School of Biomedical Sciences, University of Edinburgh, 1 George Square, Edinburgh EH8 9JZ, UK. E-mail: tony.harmar@ed.ac.uk.

DOI:10.1523/JNEUROSCI.4843-06.2007

Copyright © 2007 Society for Neuroscience 0270-6474/07/274351-08\$15.00/0

ing lighting conditions (masking) rather than by the SCN clock. Unlike most other circadian mutant mice, which have alterations of the clock genes, *Vipr2*<sup>-/-</sup> mice have an intact molecular circadian oscillator and are therefore potentially able to sustain rhythms in peripheral organs. In this study, we provide the first characterization of the control of peripheral circadian rhythms in *Vipr2*<sup>-/-</sup> mice.

There were robust rhythms of clock gene expression in the liver of *Vipr2*<sup>-/-</sup> mice; these were phase advanced relative to wild type, consistent with an abnormal timing of food intake. Rhythms of hepatic clock gene expression and food-anticipatory rhythms of activity and corticosterone secretion could be entrained by RF, demonstrating their independence from the SCN.


## Materials and Methods

**Animals.** Adult male mice (wild type or *Vipr2*<sup>-/-</sup> on a C57BL/6J background; 4–12 months old at the start of study) were housed individually in polycarbonate cages and maintained under controlled conditions of lighting and temperature. For behavioral experiments (see below), the cages were also equipped with running wheels. Powdered diet was provided in custom 60 ml glass jars (Unifab, Kalamazoo, MI). Each jar was covered with a stainless steel lid containing a hole 22 mm in diameter. A stainless steel disc was placed under the lid on top of the diet; the disc contains six circular openings (each 11 mm in diameter) to allow food access while minimizing spillage. The design of the hoppers and the use of powdered diet allowed access to the food but prevented animals from removing and storing excess food in the cage. When the animals were housed in a light/dark (LD) cycle, the lighting schedule was 12 h of bright white light and 12 h of “darkness” of dim red light (<7 lux at cage level), lights on between 7:00 A.M. and 7:00 P.M. The dim red light was permanently on to facilitate animal care and to enable behavioral studies to be performed during the dark period. DD denotes constant lighting with dim red light. All experimentation was conducted in accordance with the United Kingdom Animals (Scientific Procedures) Act, 1986.

**Food restriction studies.** The RF regimen was introduced progressively: food availability was reduced over a 3 d period from 8 to 4 h/d (Marchant and Mistlberger, 1997). During RF, food hoppers were introduced into the cages daily at zeitgeber time or circadian time 4 (ZT4/CT4) and taken out at ZT8/CT8 (where, by convention, ZT0 is defined as the time of lights on in an LD cycle, and CT0 is time of lights on in the previous LD cycle). Mice had *ad libitum* access to water throughout all of the experiments. Body weight and food intake were closely monitored during periods of RF, and animals failing to feed or showing a >20% loss in weight were returned to normal feeding and excluded from the study or humanely killed.

**Behavior and feeding studies.** Running wheel activity was recorded continuously throughout the experiment. During the *ad libitum* study, food intake was measured at intervals of 4 h over a 48 h period in both LD and DD conditions. For DD conditions, these measurements were commenced on the second complete circadian cycle (35 h) after transfer to DD, in accordance with our previous studies of activity (Harmar et al., 2002).

**Data analyses.** Food intake in the *ad libitum* and RF studies was calculated from the weight of food hoppers on entering and directly after



**Figure 1.** Rhythms of food intake and wheel running in wild-type and *Vipr2*<sup>-/-</sup> mice fed *ad libitum*. **A, B**, Representative profiles of locomotor activity in wild-type (**A**) and *Vipr2*<sup>-/-</sup> (**B**) mice are presented in double-plotted format. Periods of darkness are shaded. For the first 16 d, animals were exposed to a 12 h LD cycle (darkness from 7:00 P.M. to 7:00 A.M.). From day 17, animals were maintained in constant darkness. **C–F**, Cumulative wheel running counts (**C, D**) and food intake (**E, F**) were monitored in wild-type (black bars) and *Vipr2*<sup>-/-</sup> (gray bars) mice over 4 h periods for 48 h in LD (**C, E**) or 2 d after transfer into DD (**D, F**). Values represent mean  $\pm$  SEM;  $n = 8$ . The bars at the top indicate the dark period in black and the light period in white (**C, E**) or subjective night in black and subjective day in gray (**D, F**). \* $p < 0.05$ ; \*\* $p < 0.01$ ; \*\*\* $p < 0.001$  compared with *Vipr2*<sup>-/-</sup>.

removing them from the cages. The mean and SEM of the weight of food consumed was calculated for every 4 h period for both wild-type (WT) and *Vipr2*<sup>-/-</sup> mice. In a few cases, the animals had spread large amounts of bedding material into the hoppers, which made accurate assessment of food intake impossible. In these cases, the individual animal's food intake during the 4 h period was excluded from calculations of mean and SEM for the period.


Clocklab software (Actimetrics, Wilmette, IL) was used to determine the time of activity onset, least-squares fit of activity onset, total activity indices, and  $\chi^2$  periodograms of each of the animals in both the *ad libitum* and the RF studies. Circadian gene expression was determined using the cosine wave-fitting algorithm COSOPT (Straume, 2004), to identify statistically significant rhythms and corresponding phase optima.

**Plasma hormone and liver gene expression studies.** In a second series of experiments, WT or *Vipr2*<sup>-/-</sup> mice were housed individually in LD or DD conditions and either fed *ad libitum* or entrained to a restricted feeding regimen as described above. Mice were killed rapidly by decapitation at 4 h intervals (at ZT/CT 4, 8, 12, 16, 20, and 24), and trunk blood was collected into heparinized tubes. The brains (for a separate study) and samples of liver tissue were removed, quickly frozen on dry ice, and subsequently stored at  $-80^\circ\text{C}$  until required. Blood samples were centrifuged, and plasma was stored at  $-20^\circ\text{C}$  before determination of plasma concentrations of corticosterone by radioimmunoassay (Holmes et al., 1997). Total RNA was extracted from the liver samples using TRIzol reagent (Invitrogen, Carlsbad, CA) according to the manufacturer's instructions. TaqMan reverse transcriptase (RT)-PCR was performed using primers and probes for *Per1* (Sangoram et al., 1998), *Per2* (Panda et al., 2002), *Bmal1*, and *Dbp* (Bunger et al., 2000). Relative mRNA abundance was determined by use of the comparative  $\delta$ -CT method using 18S

**Table 1.** Mean  $\pm$  SEM acrophase and phase differences of running wheel activity and feeding behavior in wild-type and *Vipr2*<sup>-/-</sup> mice in LD and 2 d after transfer into DD conditions and with *ad libitum* access to food

		Running	Feeding
LD	Wild type	Acrophase	0.48 $\pm$ 0.81
	<i>Vipr2</i> <sup>-/-</sup>	Acrophase	22.20 $\pm$ 1.10
		Phase difference (h)	2.28
DD	Wild type	Acrophase	23.80 $\pm$ 1.20
	<i>Vipr2</i> <sup>-/-</sup>	Acrophase	16.20 $\pm$ 1.00
		Phase difference (h)	7.60
			Feeding
LD	Wild type	Acrophase	1.10 $\pm$ 1.40
	<i>Vipr2</i> <sup>-/-</sup>	Acrophase	19.75 $\pm$ 0.85
		Phase difference (h)	5.35
DD	Wild type	Acrophase	23.85 $\pm$ 0.71
	<i>Vipr2</i> <sup>-/-</sup>	Acrophase	16.72 $\pm$ 0.96
		Phase difference (h)	7.13

The phase difference is shown as the phase advance in *Vipr2*<sup>-/-</sup> mice.



**Figure 2.** Rhythms of clock gene expression in the liver of wild-type and *Vipr2*<sup>-/-</sup> mice. Values represent mean  $\pm$  SEM;  $n = 5–8$ . **A–D**, Rhythms of *Per1* (**A, B**) and *Per2* (**C, D**) gene expression in the liver of wild-type (filled squares) and *Vipr2*<sup>-/-</sup> (open squares) mice in LD (**A, C**) or 2 d after transfer into DD (**B, D**) under conditions of *ad libitum* feeding are shown. Data were obtained by TaqMan RT-PCR. In each panel, one 24 h set of data has been plotted twice to better show the pattern of variation with time. The bars at the top indicate the dark period in black and the light period in white (**A, C**) or subjective night in black and subjective day in gray (**B, D**). \* $p < 0.05$ ; \*\* $p < 0.01$ ; \*\*\* $p < 0.001$  compared with *Vipr2*<sup>-/-</sup>.

ribosomal RNA as internal control. *In situ* hybridization was conducted and quantified as described previously (Akhtar et al., 2002), using tissues from mice killed at 6 h intervals in LD (at ZT 6, 12, 18, and 24) and at 5 h intervals in DD (at CT 0, 5, 10, 15, and 20).

**Statistical analysis.** The significance of the differences between means was determined by two-way ANOVA followed by Bonferroni's *post hoc* test or by unpaired Student's *t* test, as appropriate.  $p < 0.05$  was considered statistically significant.

## Results

### Timing of food intake

The wheel-running activity of both wild-type and *Vipr2*<sup>-/-</sup> mice housed in a 12 h LD cycle was almost entirely confined to the dark period (99% in wild type and 97% in knock-out), consistent with a strong masking effect of light (Fig. 1A–C). There was a diurnal rhythm of food intake (Fig. 1E), but this was much less tightly confined to the dark period (70% in wild type and 55% in knock-out), showing less evidence of masking by light. *Vipr2*<sup>-/-</sup> mice consumed food significantly earlier in the 24 h cycle than wild-type mice (Fig. 1E, Table 1). Two days after transfer into DD, the rhythm of food intake in wild-type animals was similar to that in LD, whereas the knock-out animals' rhythm was advanced by  $\sim 7$  h (Fig. 1F, Table 1).

### Circadian rhythmicity of clock gene expression in *Vipr2*<sup>-/-</sup> mice

There were robust rhythms of clock gene expression in the liver of *Vipr2*<sup>-/-</sup> mice ( $p < 0.001$ ) (Fig. 2). These rhythms were abnormal in phase, consistent with the abnormal phasing of food intake: in *Vipr2*<sup>-/-</sup> mice exposed to an LD cycle, the times of peak expression of the clock genes examined (*Per1*, *Per2*, *Dbp*, and *Bmal1*) were advanced (by 4–6 h) compared with wild-type mice (Fig. 2A, C, Table 2). Two days after transfer into DD, even greater

phase advances (7–11 h) were seen (Fig. 2B, D, Table 2). To determine the generality and extent of this altered peripheral synchronization in *Vipr2*<sup>-/-</sup> mice, *in situ* hybridization was used to examine cycles of expression of the core clock genes *Per2*, *Bmal1*, *Rev-erb $\alpha$* , and *Cry1* and the clock-regulated genes *Dbp*, *Wee1*, *Pnp*, and *Nfix* in both liver and heart. In both organs, all of these circadian genes were phase advanced in mutant mice under a light/dark cycle (Fig. 3), and even more so 2 d after transfer into DD (Fig. 4). In the absence of a competent SCN, therefore, the circadian transcriptional program in the periphery remained functional but underwent a marked phase shift.

The temporal distribution of circadian gene expression was therefore advanced in direct correspondence to the altered rhythms of food intake in *Vipr2*<sup>-/-</sup> mice. This showed that *Vipr2*<sup>-/-</sup> mice retain some capacity for circadian coordination in the periphery and that feeding-related cues may be important in internal coordination.

### Effects of RF on wheel-running activity

To test whether an FEO was functional in *Vipr2*<sup>-/-</sup> mice, we compared the ability of wild-type and *Vipr2*<sup>-/-</sup> mice to entrain to an RF schedule (Fig. 5). When subjected to 4 h of RF during the light phase of an LD cycle, both wild-type and *Vipr2*<sup>-/-</sup> mice developed two daily peaks of wheel-running activity, the first (food-anticipatory activity) during the daytime immediately preceding food presentation and the second at the onset of darkness (Fig. 5A–C). When transferred to DD, both wild-type and *Vipr2*<sup>-/-</sup> mice continued to display daily bouts of food-anticipatory wheel running of increased intensity (Fig. 5A, B, D) ( $p < 0.05$ ; unpaired Student's *t* test). There was, however, a marked difference in behavior between the genotypes. Whereas wild-type mice continued to display a second daily bout of activity at the onset of subjective night, consistent with the activity of the SCN circadian clock, *Vipr2*<sup>-/-</sup> mice displayed only a single daily peak of activity in anticipation of food presentation (i.e., the SCN-regulated activity bout was not evident). In mice of both genotypes, daily bouts of food-anticipatory activity did not persist when *ad libitum* access to food was restored. In wild-type animals, circadian bouts of activity at the onset of subjective night continued, again consistent with a functional SCN, whereas *Vipr2*<sup>-/-</sup> mice displayed a broad peak of weakly rhythmic activity, with no clear time of onset and with a maximum in the light phase of the previous LD cycle. The timing of this activity was similar to that seen in *Vipr2*<sup>-/-</sup> mice that had not been subjected to food restriction. These data show that a functional FEO was retained in the SCN-compromised *Vipr2*<sup>-/-</sup> mice.


### Effects of RF on hepatic gene expression rhythms

To test the impact of RF on peripheral gene expression, *Vipr2*<sup>-/-</sup> and wild-type mice were entrained to an RF schedule, and liver


**Table 2.** Mean ± SEM acrophase and phase differences in hepatic gene expression rhythms of wild-type and *Vipr2*<sup>-/-</sup> mice in LD and 2 d after transfer into DD conditions and with *ad libitum* access to food

		<i>Per1</i>	<i>Per2</i>	<i>Bmal1</i>	<i>Dbp</i>	
LD	Wild type	Acrophase	16.56 ± 0.19	19.80 ± 1.20	4.34 ± 0.79	15.10 ± 1.1
	<i>Vipr2</i> <sup>-/-</sup>	Acrophase	12.40 ± 1.70	14.13 ± 0.79	20.50 ± 1.40	8.66 ± 0.97
		Phase difference (h)	4.26	5.67	7.84	6.44
DD	Wild type	Acrophase	17.35 ± 0.99	23.00 ± 1.60	5.60 ± 1.70	15.55 ± 0.82
	<i>Vipr2</i> <sup>-/-</sup>	Acrophase	10.4 ± 3.40	12.10 ± 1.80	22.10 ± 3.40	5.90 ± 2.10
		Phase difference (h)	6.95	10.90	7.50	9.65

The phase difference is shown as the phase advance in *Vipr2*<sup>-/-</sup> mice.




**Figure 3.** *a–h*, Rhythms of clock gene expression in the heart (*a–d*) and liver (*e–h*) of wild-type and *Vipr2*<sup>-/-</sup> mice in LD. Rhythms of *Per2* (*a, e*), *Bmal1* (*b, f*), *Dbp* (*c, g*), and *RevErbα* (*d, h*) gene expression in wild-type (filled squares) and *Vipr2*<sup>-/-</sup> (open squares) mice are shown. Data were obtained by *in situ* hybridization. The bars at the top indicate the dark period in black and the light period in white. Values represent mean ± SEM.



**Figure 4.** *a–h*, Rhythms of clock gene expression in the heart (*a–d*) and liver (*e–h*) of wild-type and *Vipr2*<sup>-/-</sup> mice 2 d after transfer into DD. Rhythms of *Per2* (*a, e*), *Bmal1* (*b, f*), *Dbp* (*c, g*), and *RevErbα* (*d, h*) gene expression in wild-type (filled squares) and *Vipr2*<sup>-/-</sup> (open squares) mice are shown. Data were obtained by *in situ* hybridization. The bars at the top indicate subjective night in black and subjective day in gray. Values represent mean ± SEM.

gene expression rhythms were examined. In WT mice, the time of peak expression of the *Per1* gene was coincident with the onset of food presentation (at ZT/CT 4) under both LD and DD conditions (Fig. 5*E,F*). This same pattern was evident in *Vipr2*<sup>-/-</sup> mice: there was no difference between genotypes in the effect of RF on *Per1* rhythms in the liver. *Dbp* showed a similar pattern to

*Per1*, whereas maximal expression of *Per2* and *Bmal1* occurred 2 and 12 h later, respectively (data not shown). In contrast to the phase differences seen in mice fed *ad libitum*, there were no significant differences in the profiles of gene expression of any of the clock genes measured between wild-type and *Vipr2*<sup>-/-</sup> mice un-



**Figure 5.** Entrainment of behavior and clock gene expression to an RF schedule in *Vipr2*<sup>-/-</sup> mice. **A, B**, Representative profiles of locomotor activity in wild-type (**A**) and *Vipr2*<sup>-/-</sup> (**B**) mice are presented in double-plotted format. After an initial period of entrainment to an LD cycle with *ad libitum* access to food (LD; days 1–4), food availability was reduced to 4 h/d (**A, B**, boxed regions; from 11:00 A.M. to 3:00 P.M.), initially in an LD cycle (LD-RF; days 5–21) and subsequently in DD (DD-RF; days 22–34). *Ad libitum* access to food was then restored (DD; days 35–43). **C, D**, Cumulative wheel running counts were monitored under an RF schedule in wild-type (black bars) and *Vipr2*<sup>-/-</sup> (gray bars) mice over 48 h in LD (**C**) or 2 d after transfer into DD (**D**). **E, F**, Rhythms of *Per1* gene expression in the liver of wild-type (filled squares) and *Vipr2*<sup>-/-</sup> (open squares) mice under an RF schedule were measured by TaqMan RT-PCR in LD (**E**) or 2 d after transfer into DD (**F**). One 24 h set of data has been plotted twice to better show the pattern of variation with time. **C–F**, The bars at the top indicate the dark period in black and the light period in white (**C, E**) or subjective night in black and subjective day in gray (**D, F**). The period of food availability in **C–F** is indicated by a striped bar. Values represent mean  $\pm$  SEM.

der a restricted feeding regimen in either LD or DD conditions. It is therefore evident that the RF was able to coordinate circadian gene expression in the periphery as effectively in *Vipr2*<sup>-/-</sup> mice as it does in WT mice and that it can therefore act independently of the SCN.

#### Circadian rhythms of corticosterone secretion

To obtain a second, independent report on the competence of the FEO, plasma corticosterone levels were assayed in both genotypes under *ad libitum* and restricted feeding. In agreement with previous reports, there was a significant circadian rhythm of plasma corticosterone ( $p < 0.001$ ) in wild-type mice, with a peak coincident with the onset of the activity period (Fig. 6A,B). Unexpectedly, there was no significant rhythmicity in corticosterone secretion in *ad libitum* fed *Vipr2*<sup>-/-</sup> mice, even in conditions (under an LD cycle) in which both behavior and peripheral circadian gene expression were strongly rhythmic, albeit phase ad-


vanced in the latter case (Fig. 6A). This demonstrates that rhythmic corticosterone secretion is regulated independently of activity/rest cycles: it is not a passive consequence of behavioral activity. Circadian gene expression in the liver (and heart) is not dependent on rhythmic corticosterone secretion.

In RF conditions, the profile of corticosterone secretion was altered in both wild-type and *Vipr2*<sup>-/-</sup> mice. In wild-type mice entrained to an RF schedule, in both LD and DD conditions, two daily peaks of glucocorticoid secretion were observed, one resembling that seen in animals fed *ad libitum*, reaching maximal values just before the day–night transition (or subjective night for DD mice) and the second occurring at the beginning of the period of food availability (Fig. 6C,D). In contrast, in *Vipr2*<sup>-/-</sup> mice in LD, there was just a single peak of corticosterone secretion coincident with the onset of food availability. Moreover, in DD conditions, the plasma corticosterone concentrations of *Vipr2*<sup>-/-</sup> mice were high for most of the 24 h period, again reaching a peak immediately before feeding and declining to a minimum at the end of feeding before showing a progressive increase to peak before the next period of food presentation. These data independently confirm the existence of a competent FEO in the *Vipr2*<sup>-/-</sup> mice and demonstrate the potency of food-related cues in establishing internal circadian synchrony.

#### Discussion

Our data show that there is a functional FEO in *Vipr2*<sup>-/-</sup> mice. Moreover, they clearly show that under a variety of experimental conditions, the entrainment of the liver circadian clock and the secretion of corticosterone are linked to the timing of feeding and support the hypothesis (Stokkan et al., 2001) that the liver and other peripheral oscillators are not directly coupled to the SCN but controlled indirectly, through rhythmic feeding behavior.

In contrast to our findings with *Vipr2*<sup>-/-</sup> mice, previous studies in groups of SCN lesioned mice (Hara et al., 2001; Akhtar et al., 2002; Iijima et al., 2002; Terazono et al., 2003; Kudo et al., 2004; Guo et al., 2005), including two studies performed with a similar lighting protocol to ours (Akhtar et al., 2002; Guo et al., 2005), have reported a loss of circadian rhythms of clock gene expression in the liver and other peripheral organs. It is not clear that ability of light to suppress activity (masking) was intact after SCN lesion in any of these studies, and in some, the absence of masking by light was used as a criterion for the success of SCN lesion. The role of the SCN in masking of locomotor activity by light is controversial (Redlin and Mrosovsky, 1999; Li et al., 2005), but it seems that masking is lost in most SCN lesioned mice, either because of collateral damage to extra-SCN pathways



**Figure 6.** Rhythms of corticosterone secretion in wild-type and *Vipr2*<sup>-/-</sup> mice. Values represent mean  $\pm$  SEM;  $n = 5$ –8. **A–D**, Peripheral blood was sampled at 4 h intervals from wild-type (filled squares) and *Vipr2*<sup>-/-</sup> (open squares) mice housed in a 12/12 h LD cycle (**A, C**) or 2 d after transfer into constant darkness (**B, D**) under conditions of *ad libitum* feeding (**A, B**) or with RF (**C, D**). In each panel, one 24 h set of data has been plotted twice to better show the pattern of variation with time. The bars at the top indicate the dark period in black and the light period in white (**A, C**) or subjective night in black and subjective day in gray (**B, D**). **C, D**, The period of food availability is indicated by a striped bar. \* $p < 0.05$ ; \*\* $p < 0.01$ ; \*\*\* $p < 0.001$  compared with *Vipr2*<sup>-/-</sup>.

required for masking or because an intact (albeit arrhythmic) SCN may be required for the masking response (Hara et al., 2001; Iijima et al., 2002; Terazono et al., 2003; Kudo et al., 2004). Masking responses to light are preserved when circadian rhythmicity is lost in mice with mutations in core clock genes (Vitaterna et al., 1999; Bunger et al., 2000; Zheng et al., 2001). In SCN lesioned animals, we would predict that feeding behavior would be arrhythmic, leading to a loss of synchronized rhythms of clock gene expression in peripheral tissues.

A masking response to light was intact in *Vipr2*<sup>-/-</sup> mice, so that 97% of the wheel-running activity of *Vipr2*<sup>-/-</sup> mice housed in an LD cycle took place in the dark period. Food intake was also rhythmic, although less tightly confined to the dark period. Wheel running and food intake were also significantly rhythmic in the mutant animals 2–3 d after transfer from an LD cycle into DD. In both LD and DD, *Vipr2*<sup>-/-</sup> mice consumed food significantly earlier in the 24 h cycle than wild-type mice, and there were corresponding differences in the timing of clock gene expression in the liver. We conclude that there is a functional hierarchy in the control of peripheral circadian rhythms, in which, even in the absence of a functional SCN clock, the daily patterns of activity imposed by an LD cycle are able to maintain feeding cycles and thereby entrain a FEO and rhythms of gene expression in liver and elsewhere. This rhythm persists, at least for a few days, when animals are transferred into constant conditions. It may not be appropriate to use wheel-running behavior as an index of the activity of the FEO (as has been done in many previous studies), because under some conditions, there is dissociation between the timing of feeding and of wheel-running behavior.

*Vipr2*<sup>-/-</sup> mice developed food-anticipatory behavior when subjected to a daily RF schedule, further consistent with the presence of an FEO independent of the SCN. SCN lesioned mice (Marchant and Mistlberger, 1997; Hara et al., 2001) and homozy-

gous Clock mutant (*Clk/Clk*) mice (Pitts et al., 2003; Horikawa et al., 2005) also display food-anticipatory behavior that persists after temporal feeding cues are removed for several cycles, indicating that the FEO is a circadian timer located outside the SCN and not dependent on a functional Clock gene.

In nocturnal rodents, there is a circadian rhythm of corticosterone secretion from the adrenal cortex, which peaks during the late subjective day. The SCN is thought to drive this rhythm, which is abolished by SCN lesions (Moore and Eichler, 1972; Abe et al., 1979; Filipowski et al., 2004), through neural connections to the corticotrophin-releasing factor and arginine-vasopressin neurons of the paraventricular nuclei of the hypothalamus (Buijs et al., 1993) and probably also through the sympathetic innervation of the adrenal gland (Jasper and Engeland, 1994). Although the activity of the SCN is the predominant pathway influencing the hypothalamo-pituitary-adrenal (HPA) axis under conditions of *ad libitum* feeding, which occurs mainly at night, restricted daytime feeding establishes a bimodal temporal pattern of corticosterone secretion in rodents (Krieger, 1974; Honma et al., 1984; Holmes et al., 1997; Balsalobre et al., 2000; Le Minh et al., 2001). One peak of glucocorticoid secretion resembles that seen in animals fed *ad libitum*, reaching maximal values just before the light/dark transition and the period of activity. A second peak of glucocorticoid secretion, seen only in animals fed during the day, occurs just before the period of food availability, when the HPA axis is normally quiescent. It is possible that the same mechanism responsible for FAA (the putative FEO) may underlie the anticipatory rise in plasma corticosterone observed in animals entrained to an RF schedule, because this is suppressed by *ad libitum* feeding but reappears at the phase predicted by the previous RF schedule when animals are fasted (Honma et al., 1996) and can only be entrained by feeding at intervals within the circadian range (Honma et al., 1984).

Our data on corticosterone are consistent with a model in which the circadian activity of both the SCN and of a separate FEO influences rhythms of hormone secretion. There was no significant rhythmicity in corticosterone secretion in *Vipr2*<sup>-/-</sup> mice fed *ad libitum*, suggesting that circadian rhythms of corticosterone secretion are “hard wired” to the SCN clock and are directly controlled by events downstream of oscillations of clock gene expression in the SCN. Our findings accord with an earlier study (Scarborough et al., 1996) showing that antisense antagonism of VIP mRNA in the SCN temporarily abolished the circadian rhythm of corticosterone secretion. In both wild-type and *Vipr2*<sup>-/-</sup> mice entrained to an RF schedule, a peak of corticosterone secretion preceding the period of food presentation was observed. Thus, *Vipr2*<sup>-/-</sup> mice develop anticipatory rhythms of glucocorticoid secretion when entrained to an RF schedule. In *Vipr2*<sup>-/-</sup> mice entrained to an RF schedule in DD conditions, plasma concentrations of corticosterone were low immediately after feeding and then increased progressively to reach maximal levels before feeding. The reasons for the hypersecretion of corticosterone under these conditions is not known; additional studies are necessary to examine possible differences in the HPA axis and in stress responses between wild-type and *Vipr2*<sup>-/-</sup> mice.

We conclude from these studies that (1) the SCN clock is not required for the expression of circadian rhythms in peripheral tissues; (2) *Vipr2*<sup>-/-</sup> mice, which lack a robust SCN circadian clock, nevertheless possess food-anticipatory rhythms of behavior and corticosterone secretion, consistent with the presence of an intact FEO; (3) the phasing of the circadian clock in the liver is abnormal in *Vipr2*<sup>-/-</sup> mice, probably because their feeding behavior is timed differently from wild-type animals as a conse-

quence of the defective SCN clock; (4) the SCN may form part of an anatomical pathway essential for the masking of behavior by light; and (5) the circadian rhythm of corticosterone secretion seen in *ad libitum* fed animals is driven by the SCN clock. These studies establish that the *Vipr2*<sup>-/-</sup> mouse is likely to be an important animal model in which to dissect the mechanisms by which peripheral circadian rhythms are controlled.

## References

- Abe K, Kroning J, Greer MA, Critchlow V (1979) Effects of destruction of the suprachiasmatic nuclei on the circadian rhythms in plasma corticosterone, body temperature, feeding and plasma thyrotropin. *Neuroendocrinology* 29:119–131.
- Abe M, Herzog ED, Yamazaki S, Straume M, Tei H, Sakaki Y, Menaker M, Block GD (2002) Circadian rhythms in isolated brain regions. *J Neurosci* 22:350–356.
- Akhtar RA, Reddy AB, Maywood ES, Clayton JD, King VM, Smith AG, Gant TW, Hastings MH, Kyriacou CP (2002) Circadian cycling of the mouse liver transcriptome, as revealed by cDNA microarray, is driven by the suprachiasmatic nucleus. *Curr Biol* 12:540–550.
- Aton SJ, Colwell CS, Harmar AJ, Waschek J, Herzog ED (2005) Vasoactive intestinal polypeptide mediates circadian rhythmicity and synchrony in mammalian clock neurons. *Nat Neurosci* 8:476–483.
- Balsalobre A, Damiola F, Schibler U (1998) A serum shock induces circadian gene expression in mammalian tissue culture cells. *Cell* 93:929–937.
- Balsalobre A, Brown SA, Marcacci L, Tronche F, Kellendonk C, Reichardt HM, Schutz G, Schibler U (2000) Resetting of circadian time in peripheral tissues by glucocorticoid signaling. *Science* 289:2344–2347.
- Buijs RM, Markman M, Nunes-Cardoso B, Hou YX, Shinn S (1993) Projections of the suprachiasmatic nucleus to stress-related areas in the rat hypothalamus: a light and electron microscopic study. *J Comp Neurol* 335:42–54.
- Bunger MK, Wilsbacher LD, Moran SM, Clendenin C, Radcliffe LA, Hogenesch JB, Simon MC, Takahashi JS, Bradfield CA (2000) *Mop3* is an essential component of the master circadian pacemaker in mammals. *Cell* 103:1009–1017.
- Colwell CS, Michel S, Itri J, Rodriguez W, Tam J, Lelievre V, Hu Z, Liu X, Waschek JA (2003) Disrupted circadian rhythms in VIP- and PHI-deficient mice. *Am J Physiol Regul Integr Comp Physiol* 285:R939–R949.
- Damiola F, Le Minh N, Preitner N, Kornmann B, Fleury-Olela F, Schibler U (2000) Restricted feeding uncouples circadian oscillators in peripheral tissues from the central pacemaker in the suprachiasmatic nucleus. *Genes Dev* 14:2950–2961.
- Davidson AJ, Poole AS, Yamazaki S, Menaker M (2003) Is the food-entrainable circadian oscillator in the digestive system? *Genes Brain Behav* 2:32–39.
- Dudley CA, Erbel-Sieler C, Estill SJ, Reick M, Franken P, Pitts S, McKnight SL (2003) Altered patterns of sleep and behavioral adaptability in NPAS2-deficient mice. *Science* 301:379–383.
- Filipski E, King VM, Etienne MC, Li X, Claustrat B, Granda TG, Milano G, Hastings MH, Levi F (2004) Persistent twenty-four hour changes in liver and bone marrow despite suprachiasmatic nuclei ablation in mice. *Am J Physiol Regul Integr Comp Physiol* 287:R844–R851.
- Guo H, Brewer JM, Champhekar A, Harris RB, Bittman EL (2005) Differential control of peripheral circadian rhythms by suprachiasmatic-dependent neural signals. *Proc Natl Acad Sci USA* 102:3111–3116.
- Hara R, Wan K, Wakamatsu H, Aida R, Moriya T, Akiyama M, Shibata S (2001) Restricted feeding entrains liver clock without participation of the suprachiasmatic nucleus. *Genes Cells* 6:269–278.
- Harmar AJ, Marston HM, Shen S, Spratt C, West KM, Sheward WJ, Morrison CF, Dorin JR, Piggins HD, Reubi J-C, Kelly JS, Maywood ES, Hastings MH (2002) The VPAC<sub>2</sub> receptor is essential for circadian function in the mouse suprachiasmatic nuclei. *Cell* 109:497–508.
- Holmes MC, French KL, Seckl JR (1997) Dysregulation of diurnal rhythms of serotonin 5-HT<sub>2C</sub> and corticosteroid receptor gene expression in the hippocampus with food restriction and glucocorticoids. *J Neurosci* 17:4056–4065.
- Honma KI, Honma S, Hiroshige T (1984) Feeding-associated corticosterone peak in rats under various feeding cycles. *Am J Physiol* 246:R721–R726.
- Honma S, Katsuno Y, Abe H, Honma K (1996) Aging affects development and persistence of feeding-associated circadian rhythm in rat plasma corticosterone. *Am J Physiol* 271:R1514–R1520.
- Horikawa K, Minami Y, Iijima M, Akiyama M, Shibata S (2005) Rapid damping of food-entrained circadian rhythm of clock gene expression in clock-defective peripheral tissues under fasting conditions. *Neuroscience* 134:335–343.
- Iijima M, Nikaido T, Akiyama M, Moriya T, Shibata S (2002) Methamphetamine-induced, suprachiasmatic nucleus-independent circadian rhythms of activity and mPer gene expression in the striatum of the mouse. *Eur J Neurosci* 16:921–929.
- Jasper MS, Engeland WC (1994) Splanchnic neural activity modulates ultradian and circadian rhythms in adrenocortical secretion in awake rats. *Neuroendocrinology* 59:97–109.
- Kornmann B, Preitner N, Rifat D, Fleury-Olela F, Schibler U (2001) Analysis of circadian liver gene expression by ADDER, a highly sensitive method for the display of differentially expressed mRNAs. *Nucleic Acids Res* 29:E51–E51.
- Krieger DT (1974) Food and water restriction shifts corticosterone, temperature, activity and brain amine periodicity. *Endocrinology* 95:1195–1201.
- Kudo T, Nakayama E, Suzuki S, Akiyama M, Shibata S (2004) Cholesterol diet enhances daily rhythm of *Pai-1* mRNA in the mouse liver. *Am J Physiol Endocrinol Metab* 287:E644–E651.
- Le Minh N, Damiola F, Tronche F, Schutz G, Schibler U (2001) Glucocorticoid hormones inhibit food-induced phase-shifting of peripheral circadian oscillators. *EMBO J* 20:7128–7136.
- Li X, Gilbert J, Davis FC (2005) Disruption of masking by hypothalamic lesions in Syrian hamsters. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* 191:23–30.
- Marchant EG, Mistlberger RE (1997) Anticipation and entrainment to feeding time in intact and SCN-ablated C57BL/6j mice. *Brain Res* 765:273–282.
- Maywood ES, Reddy AB, Wong GK, O'Neill JS, O'Brien JA, McMahon DG, Harmar AJ, Okamura H, Hastings MH (2006) Synchronization and maintenance of timekeeping in suprachiasmatic circadian clock cells by neuropeptidergic signaling. *Curr Biol* 16:599–605.
- McNamara P, Seo SP, Rudic RD, Sehgal A, Chakravarti D, FitzGerald GA (2001) Regulation of CLOCK and MOP4 by nuclear hormone receptors in the vasculature: a humoral mechanism to reset a peripheral clock. *Cell* 105:877–889.
- Mistlberger RE (1994) Circadian food-anticipatory activity: formal models and physiological mechanisms. *Neurosci Biobehav Rev* 18:171–195.
- Moore RY, Eichler VB (1972) Loss of a circadian adrenal corticosterone rhythm following suprachiasmatic lesions in the rat. *Brain Res* 42:201–206.
- Panda S, Antoch MP, Miller BH, Su AI, Schook AB, Straume M, Schultz PG, Kay SA, Takahashi JS, Hogenesch JB (2002) Coordinated transcription of key pathways in the mouse by the circadian clock. *Cell* 109:307–320.
- Pitts S, Perone E, Silver R (2003) Food-entrained circadian rhythms are sustained in arrhythmic *Clk/Clk* mutant mice. *Am J Physiol Regul Integr Comp Physiol* 285:R57–R67.
- Reddy AB, Karp NA, Maywood ES, Sage EA, Deery M, O'Neill JS, Wong GK, Chesham J, Odell M, Lilley KS, Kyriacou CP, Hastings MH (2006) Circadian orchestration of the hepatic proteome. *Curr Biol* 16:1107–1115.
- Redlin U, Mrosovsky N (1999) Masking by light in hamsters with SCN lesions. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol* 184:439–448.
- Sangoram AM, Saez L, Antoch MP, Gekakis N, Staknis D, Whiteley A, Fruechte EM, Vitaterna MH, Shimomura K, King DP, Young MW, Weitz CJ, Takahashi JS (1998) Mammalian circadian autoregulatory loop: a timeless ortholog and mPer1 interact and negatively regulate CLOCK-BMAL1-induced transcription. *Neuron* 21:1101–1113.
- Scarborough K, Harney JP, Rosewell KL, Wise PM (1996) Acute effects of antisense antagonism of a single peptide neurotransmitter in the circadian clock. *Am J Physiol* 270:R283–R288.
- Stephan FK (2002) The “other” circadian system: food as a Zeitgeber. *J Biol Rhythms* 17:284–292.
- Stokkan KA, Yamazaki S, Tei H, Sakaki Y, Menaker M (2001) Entrainment of the circadian clock in the liver by feeding. *Science* 291:490–493.
- Straume M (2004) DNA microarray time series analysis: automated statistical assessment of circadian rhythms in gene expression patterning. *Methods Enzymol* 383:149–166.
- Terazono H, Mutoh T, Yamaguchi S, Kobayashi M, Akiyama M, Udo R,



- Ohdo S, Okamura H, Shibata S (2003) Adrenergic regulation of clock gene expression in mouse liver. *Proc Natl Acad Sci USA* 100:6795–6800.
- Vitaterna MH, Selby CP, Todo T, Niwa H, Thompson C, Fruechte EM, Hitomi K, Thresher RJ, Ishikawa T, Miyazaki J, Takahashi JS, Sancar A (1999) Differential regulation of mammalian period genes and circadian rhythmicity by cryptochromes 1 and 2. *Proc Natl Acad Sci USA* 96:12114–12119.
- Yamazaki S, Numano R, Abe M, Hida A, Takahashi R, Ueda M, Block GD, Sakaki Y, Menaker M, Tei H (2000) Resetting central and peripheral circadian oscillators in transgenic rats. *Science* 288:682–685.
- Yoo SH, Yamazaki S, Lowrey PL, Shimomura K, Ko CH, Buhr ED, Siepka SM, Hong HK, Oh WJ, Yoo OJ, Menaker M, Takahashi JS (2004) PERIOD2::LUCIFERASE real-time reporting of circadian dynamics reveals persistent circadian oscillations in mouse peripheral tissues. *Proc Natl Acad Sci USA* 101:5339–5346.
- Zheng B, Albrecht U, Kaasik K, Sage M, Lu W, Vaishnav S, Li Q, Sun ZS, Eichele G, Bradley A, Lee CC (2001) Nonredundant roles of the mPer1 and mPer2 genes in the mammalian circadian clock. *Cell* 105:683–694.
- Zylka MJ, Shearman LP, Weaver DR, Reppert SM (1998) Three period homologs in mammals: differential light responses in the suprachiasmatic circadian clock and oscillating transcripts outside of brain. *Neuron* 20: 1103–1110.