



# THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### What memory binding functions is the hippocampus responsible for?

**Citation for published version:**

Fabi, K, Parra, M, Luzzi, S, Cubelli, R & Della Sala, S 2010, 'What memory binding functions is the hippocampus responsible for?' XLI Congress of the Italian Society of Neurology, Catania, Italy, 23/10/10 - 27/10/10, .

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Author final version (often known as postprint)

**Publisher Rights Statement:**

© Fabi, K., Parra, M., Luzzi, S., Cubelli, R., & Della Sala, S. (2010). What memory binding functions is the hippocampus responsible for?. Poster session presented at XLI Congress of the Italian Society of Neurology, Catania, Italy.

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



## **What memory binding functions is the hippocampus responsible for?**

**Katia Fabi <sup>a</sup>, Mario A. Parra <sup>b,c</sup>, Simona Luzzi <sup>a</sup>, Roberto Cubelli <sup>c</sup>, and Sergio Della Sala <sup>b</sup>**

a: Clinical Neurology, Polytechnic University of Marche, Ancona, Italy

b: Human Cognitive Neuroscience, Psychology, University of Edinburgh, UK

c: Department of Cognitive Sciences and Education, University of Trento, Italy

### **Abstract**

The role of the hippocampus in binding information in working memory (WM) is little understood. When complex experiences comprise associations between different pieces of information such as objects and locations (relational binding), the function of the hippocampus is required to hold them in WM (Mitchell et al., 2000; 2006; Piekema, 2006). However, recent evidence suggests that if the to-be-associated information leads to the formation of integrated objects such as coloured shapes (conjunctive binding), the hippocampus is less involved in holding temporary representations of these complex events in WM (Baddeley et al., 2010; Piekema, 2006). We investigated the relational and conjunctive binding hypotheses of the hippocampal functions in a patient with right hippocampal damage. The patient and controls were asked to study visual arrays of stimuli which consisted of shape-colour relations (shape-colour pairs) or shape-colour conjunctions (coloured shapes). After the study array, they were presented with a new screen consisting of one set of shapes (line drawings) and one set of colours. They were asked to reconstruct the bindings by selecting the shapes and their corresponding colours. As compared to healthy controls, the patient was impaired in holding relations of shapes and colours in WM whereas he could retain the conjunctions similarly to controls. These results lend support to the role of the hippocampus in supporting memory for inter-item associations but not memory for conjunctions of features which define objects' identity.