



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

An emerging role for NAADP-mediated Ca²⁺ signaling in the pancreatic beta-cell

Citation for published version:

Arredouani, A, Evans, AM, Ma, J, Parrington, J, Zhu, MX & Galione, A 2010, 'An emerging role for NAADP-mediated Ca²⁺ signaling in the pancreatic beta-cell' *Islets*, vol 2, no. 5, pp. 323-330., 10.4161/isl.2.5.12747

Digital Object Identifier (DOI):

[10.4161/isl.2.5.12747](https://doi.org/10.4161/isl.2.5.12747)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Preprint (usually an early version)

Published In:

Islets

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



An emerging role for NAADP-mediated Ca²⁺ signaling in the pancreatic β -cell

Abdelilah Arredouani,¹ A. Mark Evans,² Jianjie Ma,³ John Parrington,¹ Michael X. Zhu⁴ and Antony Galione^{1,*}

¹Department of Pharmacology; University of Oxford; Oxford, UK; ²Centre for Integrative Physiology; College of Medicine and Veterinary Medicine; University of Edinburgh; Edinburgh, Scotland UK; ³Department of Physiology and Biophysics; UMDNJ-Robert Wood Johnson Medical School; Piscataway, NJ USA;

⁴Department of Integrative Biology and Pharmacology; The University of Texas Health Science Center in Houston; Houston, TX USA

Several recent reports, including one in this journal, have reignited the debate about whether the calcium-mobilizing messenger, nicotinic adenine nucleotide diphosphate (NAADP) plays a central role in the regulation of calcium signaling in pancreatic β -cells.¹⁻⁵ These studies have highlighted a role for NAADP-induced Ca²⁺ mobilization not only in mediating the effects of the incretin, GLP-1 and the autocrine proliferative effects of insulin, but also possibly a fundamental role in glucose-mediated insulin secretion in the pancreatic β -cell.

Introduction

The K_{ATP} channel-dependent hypothesis for stimulus-secretion coupling in pancreatic β -cells has gained prominence in recent years as the major pathway by which glucose triggers signaling events to bring about membrane depolarization and the activation of voltage-dependent calcium channels (VDCCs) and Ca²⁺ influx, the major regulator of insulin granule exocytosis.⁶ However, several shortcomings of this hypothesis have been articulated.^{7,8} First, glucose may still evoke Ca²⁺ signals and insulin secretion after knockout of K_{ATP} channel components,⁹⁻¹¹ and secondly K_{ATP} channel closure alone is not sufficient to depolarize the membrane to cause activation of VDCCs. A background current, for example, an inward cation current is required, together with the closure of the K_{ATP} channel which itself increases the membrane resistance to cause sufficient depolarization to activate VDCCs.

The Ca²⁺ mobilizing messenger, NAADP is gaining prominence as a universal trigger for events at both the ER and at the plasma membrane.¹²⁻¹⁵ It does this by causing local Ca²⁺ release from acidic stores, which may then be amplified by recruiting Ca²⁺ release channels on the endoplasmic reticulum (ER) membrane or activating Ca²⁺-dependent channels at the plasma membrane. Recent studies have demonstrated the presence of NAADP-sensitive acidic Ca²⁺ stores in pancreatic β -cells,^{1,16-18} and their role in triggering signaling events in β -cells is gathering momentum.

The most recent reports highlight the role of NAADP in signal transduction in response to activation of autocrine insulin receptors.^{4,5} Another highlights the role of NAADP in mediating the effects of the incretin, GLP-1.¹ Furthermore, a recent important development describes a new selective membrane-permeant NAADP antagonist, Ned-19 which inhibited glucose-evoked calcium spiking in mouse pancreatic β -cells in a concentration-dependent manner.² Finally, a groundbreaking discovery has been made recently with the identification of the two pore channels (TPCs) as NAADP-gated calcium release channels on endo-lysosomal membranes and the demonstration that NAADP evoked oscillatory non-selective cation currents in mouse pancreatic β -cells, this effect being abolished in cells prepared from *Tpc2*^{-/-} mice.³ These studies represent the re-emergence of an interest in NAADP signaling in pancreatic β -cells that occurred some years ago and cement earlier observations on a possible role for

Key words: pancreatic β -cell, calcium lysosome, two-pore channel (TPC), NAADP

Submitted: 05/30/10

Accepted: 06/21/10

Previously published online:
www.landesbioscience.com/journals/islets/article/12747

DOI: 10.4161/isl.2.5.12747

*Correspondence to: Antony Galione;
 Email: antony.galione@pharm.ox.ac.uk




Figure 1. Structure of NAADP. NAADP differs from β -NADP only in that the base nicotinic acid is substituted for nicotinamide.

NAADP in glucose-induced calcium signaling in these cells.¹⁹ These earlier studies suggested a role for NAADP in mediating the autocrine effects of insulin in human islets,²⁰ the release of Ca^{2+} from acidic stores in β -cell lines,¹⁶ and the role of NAADP-evoked calcium release as a critical mediator of Ca^{2+} signaling responses to glucose in MIN6 cells.^{17,21} With these new studies, NAADP is emerging as an important player in the regulation of β -cell Ca^{2+} signaling.

NAADP as a Calcium Mobilizing Messenger

NAADP is the most potent of the major Ca^{2+} mobilizing messengers described, with activity reported at low nanomolar intracellular concentrations.²² It differs in structure from the more familiar coenzyme NADP by the substitution of the nicotinamide moiety by nicotinic acid (Fig. 1). Its actions were first reported in sea urchin eggs,²² but NAADP is now known to have widespread actions in most cell types studied.^{13,23} However, it is fair to say that in terms of its synthesis, regulation and mechanism of action, NAADP signaling is poorly understood. NAADP is thought to be synthesized from NADP as a precursor, and the only mammalian enzymes that have been demonstrated to catalyze this reaction, at least in vitro, are ADP-ribosyl cyclases (ARCs) such as CD38.^{24,25} Interestingly, CD38 and its alternate Ca^{2+} mobilizing product, cADPR, had previously been linked to stimulus-secretion coupling in β -cells^{26,27} as discussed below. Various stimuli, including agonists acting at a variety of cell surface receptors, have been demonstrated to

increase cellular NAADP levels, adding to growing evidence that it is an intracellular messenger.²³ The concentration-response relationship between NAADP and Ca^{2+} release in β -cells,^{20,21} in common with other mammalian cell types, is described as bell-shaped, with maximal Ca^{2+} release evoked by around 100 nM NAADP.²⁸ Higher concentrations are less effective; with micromolar concentrations causing apparent desensitization of the NAADP-sensitive Ca^{2+} release mechanism in the absence of measurable Ca^{2+} release.^{1,20,21} In many cell types, NAADP appears to target a separate Ca^{2+} store, which may be acidic in nature such as lysosomes, rather than the long-established ER Ca^{2+} store,²⁹ although Ca^{2+} release from acidic stores can trigger subsequent Ca^{2+} -induced Ca^{2+} release (CICR) from the ER.^{14,30,31} Thus NAADP has been proposed to serve as a key triggering Ca^{2+} mobilizing messenger, not only releasing Ca^{2+} by directly targeting NAADP-sensitive stores, but also acting to coordinate Ca^{2+} release from the ER via IP_3R and RyR through CICR-dependent mechanisms.^{12,14}

The Molecular Identity of NAADP Receptors

The molecular targets for NAADP have been a matter of debate for some time.³² Pharmacological studies have generally supported the idea that NAADP targets a mechanism distinct from the two major Ca^{2+} release channels, IP_3 or RyRs. Of the known lysosome resident channels, mucolipin 1 was the first to be suggested as the target receptor channel for NAADP³³ but this protein had previously been ruled out by others.³⁴ Ryanodine receptors have also

been suggested in some cells to be directly targeted by NAADP,^{35,36} although this is not universally accepted.³⁷ However, the recent identification of two-pore channels (TPCs) as NAADP receptors,³ this finding being subsequently confirmed by others,^{38,39} perhaps provides the most convincing candidates, as they mirror many of the properties previously ascribed to NAADP-gated Ca^{2+} release mechanisms including endo-lysosomal localization.^{14,40}

TPCs are poorly characterized members of the voltage-gated superfamily of cation channels. They are an ancient family, and a TPC homolog functions as a vacuolar channel in plants.⁴¹ They consist of two-linked 6 transmembrane domain units and thus have a total of 12 transmembrane alpha helices. It is likely that the functional channel is a dimer. There are three non-allelic TPC genes in most animal species, but only TPC1 and TPC2 are found in humans and rodents. Expression of HsTPCs in HEK293 cells showed them to be localized to the endolysosomal system, with TPC2 predominantly expressed in lysosomes while TPC1 was endosomal. In cells overexpressing TPC2, NAADP evoked a biphasic Ca^{2+} release, the first phase from acidic stores, whilst the second phase was due to amplification by the recruitment of ER IP_3Rs , presumably by CICR.^{3,40} The concentration-dependence of NAADP action showed the characteristic bell-shaped concentration-response curve. siRNA TPC2 probes abolished the sensitivity to NAADP, and stimulation of Ca^{2+} activated cation currents in mouse pancreatic β -cells were absent in cells from *Tpc2*^{-/-} mice. In addition, expression of TPC2 in HEK293 cells was associated with increased [³²P]NAADP binding,³ and immunopurified endogenous sea urchin TPC1 and TPC3 complexes bind [³²P]NAADP with nanomolar affinity.⁴⁰

Taken together, TPCs represent strong candidates as NAADP receptors^{14,15,42} and provide for the first time a molecular handle with which to investigate NAADP signaling.

Calcium Stores in Pancreatic β -cells

The traditional model for glucose-evoked Ca^{2+} signaling has emphasized the role of

Ca²⁺ influx through L-type voltage-gated Ca²⁺ channels, with a minor role for the ER in shaping the stereotypic Ca²⁺ spikes. Indeed, the earliest Ca²⁺ response observed to be initiated by glucose is a decrease ascribed to enhancement of Ca²⁺ uptake by the ER by increase in ATP synthesis⁴³ probably mediated by SERCA2b. Pharmacological ablation of SERCA-mediated pumping of Ca²⁺ into the ER with thapsigargin or genetic ablation of SERCA2b and 3 isoforms in mouse β-cells generally enhances amplitudes of glucose-evoked Ca²⁺ oscillations.⁴⁴ Thus the ER plays a role in Ca²⁺ buffering, although cycles of uptake and release may be important in glucose-mediated fast Ca²⁺ spiking⁴⁵ and play a prominent role in the modulation of Ca²⁺ spiking by activation of incretin receptors coupled to IP₃ production.⁴⁶ The finding that NAADP targets acidic Ca²⁺ stores²⁹ has prompted the investigation of the role of acidic stores in Ca²⁺ handling in pancreatic β-cells.

In contrast to the ER, little is known of the Ca²⁺ uptake mechanisms in acidic stores, nor how Ca²⁺ is stored in these organelles, although polyanions have been suggested to play a role.⁴⁷ However, Ca²⁺ uptake and storage seem to be dependent on the maintenance of the pH gradient across their membranes. Two classes of pharmacological agents have been principally used to abrogate Ca²⁺ storage by these organelles. The first is bafilomycin and related compounds such as concanamycin, which inhibit the vacuolar H⁺-ATPase expressed widely in acidic organelles including the endolysosomal system and secretory granules.⁴⁸ The ability of these compounds to discharge Ca²⁺ depends on the leakiness of stores to both protons and Ca²⁺ and often requires prolonged incubations. Both nigericin, a protonophore and NAADP have been found to accelerate Ca²⁺ leak by these organelles.²⁹ A second compound, glycyphenylalanine-2-naphthylamide (GPN), causes osmotic lysis of acidic stores based on the presence of cathepsin C, found principally within lysosomes, and is accompanied by bursts of Ca²⁺ release.²⁹ In MIN6 cells in which the bioluminescent Ca²⁺ reporter aequorin was targeted to the ER or insulin-containing secretory granules, it was found that NAADP released Ca²⁺ from secretory

granules but not the ER, an effect that was not blocked by either ryanodine or dantrolene, a RyR1 inhibitor.¹⁶ It was found that photolysis of caged NAADP evoked a Ca²⁺ response that was largely abolished by pretreatment with bafilomycin, but thapsigargin did not affect the amplitude, although it reduced the duration of the response.¹⁷ A further study showed in mouse primary β-cells that NAADP-evoked Ca²⁺ release was blocked by GPN and partially inhibited by thapsigargin but not by dantrolene.¹⁸ Acidic Ca²⁺ pools were shown to be primarily responsible for the delayed plateau of Ca²⁺ release after prolonged depolarization of β-cells termed the “Ca²⁺ hump”. The apparent large size of this Ca²⁺ pool may indicate that Ca²⁺ release from the small number of lysosomes in β-cells alone cannot account for this phenomenon, but might include secretory granules or other acidic organelles in addition.¹⁸ Additionally, in one of the recent studies, NAADP was firmly shown to mobilize Ca²⁺ from acidic stores but not the ER in mouse primary β-cells.¹ However, in permeabilized mouse β-cells, NAADP was found to be ineffective at mobilizing Ca²⁺.⁴⁹ There are few reports of NAADP efficacy in broken cell systems.²⁸ One possible explanation is that stores are labile, small and coupling to CICR amplification mechanisms required for detection are broken.

NAADP Synthesis and Regulation

An impetus for studying the role of NAADP in β-cells followed on from the work of Okamoto and colleagues. In 1993, they proposed a key role for cADPR in stimulus-secretion coupling.²⁶ cADPR was proposed to enhance insulin secretion in pancreatic β-cells by either mobilizing Ca²⁺ from the ER through activating RyRs²⁶ or at higher concentrations by enhancing Ca²⁺ influx through activating plasma membrane TRPM2 channels.⁵⁰ Inhibition or genetic ablation of CD38 function has been correlated with reduced Ca²⁺ responses and insulin secretion in response to elevated glucose concentrations,⁵¹ and even linked to Type 2 diabetes.^{27,52-55} Subsequent studies showed that CD38, which cyclizes NAD to produce

cADPR, played a key role in insulin secretion.⁵¹ However, given that cADPR targets RyRs mainly on the ER, and this organelle shapes, rather than has a triggering role in evoking Ca²⁺ signals in β-cells,⁵⁶ and that a role for cADPR in β-cell stimulus-secretion coupling has not been universally confirmed,^{57,58} the role of NAADP was investigated since it was shown that CD38 and ARCs could catalyze NAADP formation in cell-free systems²⁴ and NAADP mobilizes Ca²⁺ largely from a store separate from the ER.

While a number of studies have provided evidence for the role of ARCs such as CD38 in the synthesis of cADPR,⁵⁹ evidence for their role in the synthesis of endogenous NAADP remains conflicting, with some studies supporting such a role,^{60,61} others arguing against.⁶² In a recent study in β-cells, CD38 was shown to contribute to NAADP synthesis stimulated by GLP-1, but this appeared not to be the only mechanism for generating NAADP.¹

CD38 was initially considered an ectoenzyme⁵⁹ but it has now been shown that ARCs may also be localized inside cells, where they are more appropriately sited for production of intracellular messengers.⁶³ There is growing evidence that ARCs, including CD38, may be present in endosomes,⁶⁴ secretory granules⁶³ and even lysosomes.¹ In an interesting study, internalization of plasma membrane CD38 was induced by the reducing agent, L-oxothiazolidine-4-carboxylic acid, a pro-drug of cysteine, which like glucose enhances dimerization of CD38, enhances glucose-mediated Ca²⁺ signaling and induces antidiabetic effects in ameliorating glucose intolerance in db/db mice.⁶⁴ The localization of messenger-synthesising enzymes within organelles presents a topological problem in that substrates have to be transported in, and products out to their targets. However, evidence has been presented in the sea urchin egg for such processes.⁶³ Moreover, the luminal acidic pH may favour enzyme activities especially for NAADP generation and since a base-exchange mechanism has been proposed for this messenger,^{24,25} nicotinic acid may accumulate there at sufficient concentrations for NAADP synthesis by this mechanism.

Although NAADP has been found to increase in islets and β -cells in response to glucose, GLP-1 and insulin, stimulus-coupling to intra-organellar enzymes again presents a topological problem. At present these mechanisms are unclear. However, in β -cells, ATP,⁶⁵ cGMP-dependent kinases, PKC⁴ and cAMP (perhaps via EPAC)¹ have all been proposed. Our understanding of the way in which β -cell agonists couple to the enzymatic synthesis of NAADP and cADPR is in urgent need of further study in common with the situation in other systems, as is the identification of other possible NAADP synthases.

NAADP and Receptor-mediated Ca^{2+} Signaling

Incretins such as GLP-1 do not trigger insulin secretion per se but potentiate the actions of glucose.⁶⁶ They act at cell surface receptors and activate signal transduction processes. A recent report has demonstrated that GLP-1 is coupled to both NAADP and cADPR synthesis and Ca^{2+} release from acidic and ER stores.¹ Unusually in this study, NAADP was shown to increase Ca^{2+} in β -cells in response to extracellular application of NAADP at concentrations similar to those required for responses in studies employing whole cell patch clamp application.²

GLP1 (10 nM) was found to evoke Ca^{2+} signals only in the presence of elevated glucose.^{1,67} Ca^{2+} signals were maintained with the initial phase ascribed to NAADP whilst thapsigargin and cADPR antagonists (but not IP_3 R blockers) caused the abolition or abbreviation of the second maintained phase. Both desensitizing concentrations of NAADP or pretreatment with bafilomycin abolished both phases of the GLP-1 response. Thus the initial transient evoked by NAADP is required for the second phase of Ca^{2+} release evoked by cADPR. This is similar to models of NAADP trigger action in other systems¹² such as the fertilization of the sea urchin egg, where NAADP can fire a Ca^{2+} wave first by evoking Ca^{2+} release from acidic stores and then by recruiting ER CICR mechanisms.^{68,69}

Experiments with *Cd38*^{-/-} islets showed a partial dependence on CD38 for both NAADP and cADPR synthesis,

suggesting other enzymes may also be involved in their production.¹

Another receptor on β -cells linked to Ca^{2+} signals dependent on NAADP-sensitive Ca^{2+} release is the insulin receptor. Paracrine or autocrine effects of insulin receptors have been shown to evoke Ca^{2+} signals in β -cells.^{70,71} In the first report of NAADP effects in human β -cells, microinjection of NAADP at different concentrations displayed the hallmark bell-shaped curve for Ca^{2+} release with maximum effects at 100 nM. Injecting a desensitizing concentration of 10 mM, insulin-evoked Ca^{2+} signals were abolished. Furthermore, the insulin concentration-response curve was bell-shaped for Ca^{2+} release, mirroring that for NAADP. As for GLP-1, the initial phase of Ca^{2+} release evoked by insulin was insensitive to SERCA pump inhibitors suggesting release from a non-ER store. The second phase of insulin-evoked Ca^{2+} release was partially blocked by the IP_3 R antagonist, xestospongine but not by ryanodine. Insulin stimulation of islets lead to increased insulin content but not secretion, thus implicating a role for NAADP-mediated Ca^{2+} signaling in insulin regulation of insulin gene expression. Interestingly, as we describe below, while NAADP may depolarize β -cell membranes (Fig. 2), insulin apparently does not.⁷⁰

As recently reported in this journal, the role of NAADP in the mediation of insulin responses has been reinvestigated. In this study, NAADP was again implicated in initiating the Ca^{2+} mobilizing effects of insulin, with a sequential role for IP_3 and cADPR consistent with the triggering role for NAADP. Here a role for insulin-mediated NAADP-dependent signaling was implicated in the control of β -cell proliferation.⁴ Additionally, insulin was shown to increase islet NAADP levels, in contrast to the previous report from the same group.¹

A Triggering Role for NAADP in Nutrient-Mediated Calcium Signaling

Perhaps one of the most interesting new aspects in this area is that a key role for NAADP for the actual triggering of

glucose-mediated responses has now been reported in several studies. The first study to suggest this role used the ability of high concentrations of NAADP to desensitize NAADP-evoked Ca^{2+} release²¹ in MIN6 cells. Using microinjected caged NAADP it was found that photolysis of low amounts evoked a large Ca^{2+} release, which was largely resistant to inhibition by thapsigargin. However, with photolysis of higher amounts of caged NAADP, no Ca^{2+} release was seen. Having established conditions for desensitization of NAADP-evoked Ca^{2+} release, cells were then challenged with 20 mM glucose. Glucose-evoked Ca^{2+} oscillations were greatly suppressed, suggesting a role of NAADP in the actual triggering processes in β -cell stimulus-secretion coupling. In addition, high affinity binding sites for NAADP were found in islets, and importantly glucose was able to evoke an increase in cellular NAADP levels.²¹ Since NAADP was reported to mobilize Ca^{2+} from acidic stores,²⁹ the effects of disrupting Ca^{2+} storage in acidic organelles upon glucose-evoked Ca^{2+} signaling was examined.¹⁷ Bafilomycin (2 μM) abolished glucose responses but not those to acetylcholine, an incretin acting on β -cell muscarinic receptors which are linked to IP_3 production and release of Ca^{2+} from ER stores. Conversely, thapsigargin abolished acetylcholine but not glucose responses.¹⁷ Taken together, these data are suggestive of a role for NAADP-mobilization from acidic stores in the glucose-mediated triggering response. The employment of a newly developed membrane-permeant selective NAADP antagonist, Ned-19 has confirmed these findings. Ned-19 is weakly fluorescent and labels acidic stores in pancreatic β cells, an effect that is reduced by prior treatment with the membrane-permeant NAADP analogue, NAADP-AM.² Ned-19 abolished glucose-evoked Ca^{2+} responses in a concentration-dependent manner (Fig. 3),² although it did not affect activation of voltage-gated Ca^{2+} channels by potassium depolarization or mitochondrial metabolism. Since the large glucose-evoked Ca^{2+} signals are dependent on extracellular Ca^{2+} as they rely on depolarization-evoked openings of VDCCs, what then is the role of NAADP-mediated Ca^{2+} release from acidic stores?

One clue has come from intracellular perfusion of β -cells with NAADP. NAADP evokes a series of oscillatory depolarizing cation currents, which are abolished by Ned-19.² The channels carrying these currents are unknown but intriguingly they are blocked by TRPM4/5 blockers (unpublished observations), and Ca^{2+} -activated TRPM4 and TRPM5 channels in the plasma membrane have been suggested to play an important role in shaping Ca^{2+} signals⁷² and controlling insulin secretion in β -cells.⁷³⁻⁷⁵ Thus NAADP-evoked Ca^{2+} release via TPCs from acidic stores just under the plasma membrane could comprise a signaling module with Ca^{2+} -activated TRPM4/5 channels mediating plasma membrane depolarization by NAADP (Fig. 4). Interestingly, this depolarizing effect seems unique to NAADP as a Ca^{2+} mobilizing messenger, since IP_3 has been shown to hyperpolarize the β -cell membrane.⁷⁶ One possibility then is that NAADP signaling mechanisms contribute together with the closure of K_{ATP} channels to the depolarization of the β -cell membrane to a threshold for activation of VDCCs (Fig. 2). The operation of this triggering mechanism may explain how glucose still evokes Ca^{2+} transients in β -cells from knockout mice with defects in Kir6.2 or SUR1, the components of the K_{ATP} channel. Since NAADP production and Ca^{2+} release may be localized to regions just under the plasma membrane such Ca^{2+} microdomains, as previously hypothesized,⁷⁷ may play a key role in β -cell excitability. The currents generated by this mechanism in response to glucose may be small and transient, but sufficient to depolarize the membrane especially when the membrane resistance is increased by closure of K_{ATP} channels. Thus, although this mechanism is strictly

K_{ATP} -independent, it may synergize with K_{ATP} -dependent mechanisms in the triggering phase for glucose action. A major breakthrough in our understanding of NAADP-mediated Ca^{2+} release has come from the recent discovery of two pore channels (TPCs) as NAADP-gated Ca^{2+} release channels of endolysosomal compartments.^{3,14} Importantly, as described above, in β -cells from *Tpc2*^{-/-} mice, NAADP no longer evokes the oscillatory depolarizing currents. The study of

glucose-mediated Ca^{2+} signaling in TPC knockout mice will be important in our understanding of the role of NAADP in β -cell Ca^{2+} signaling. Preliminary results already reported suggest that glucose-evoked Ca^{2+} spiking and membrane depolarization is abolished or greatly attenuated in *Tpc2*^{-/-} β -cells.⁷⁸ Further studies of *Tpc2*^{-/-}, *Tpc1*^{-/-} and possibly *Tpc1/Tpc2* double knockout β -cells that are ongoing may provide exciting new information about a possible new triggering pathway of




Figure 2. NAADP-evoked Ca^{2+} -activated cation currents in mouse pancreatic β -cells. (A) Ca^{2+} -dependent current traces are from mouse β cells that were voltage-clamped with a patch pipette containing 100 nM NAADP in the presence or absence of the membrane permeant NAADP antagonist Ned-19 (100 μM) in the bathing solution. (B) Cation currents at -70 mV evoked by intracellular dialysis of 100 nM NAADP via patch pipette in whole cell mode in pancreatic β -cells isolated from wild-type (top) and *Tpc2* knockout (bottom) mice. The left hand parts show a single representative trace, whilst the right hand part shows records from several cells [Taken from ref. 2 and 3].




Figure 3. The NAADP antagonist blocks glucose-evoked Ca^{2+} signaling in mouse primary pancreatic beta cells. Ned-19 reduces glucose-induced Ca^{2+} increases in a concentration-dependent manner. Islets were pre-incubated with the indicated concentration of Ned-19 for 30 min before the addition of 15 mM glucose [Taken from ref. 2].




Figure 4. Scheme for how NAADP-regulated TPC2 and K_{ATP} channels synergize to trigger glucose-evoked insulin secretion. NAADP-induced Ca^{2+} release from acidic stores (AS) synergizes with the K_{ATP} -dependent pathway to depolarize the plasma membrane and activate VDCCs. The ATP-mediated closure of K_{ATP} channels increases membrane resistance, which together with Ca^{2+} -dependent depolarizing (non-selective cation) currents (possibly mediated by Ca^{2+} -activated TRPM4 or TRPM5 channels), activated by NAADP-induced Ca^{2+} -release via TPC2 expressed on acidic stores, may depolarize the plasma membrane to threshold for VDCC activation. Ca^{2+} influx via VDCCs then triggers exocytosis of insulin granules. AS: acidic stores; NSCC: non-selective cation channel; R_m : membrane resistance; VDCC: voltage-dependent Ca^{2+} channels; Glc: glucose.

NAADP for stimulus-secretion coupling in the pancreatic β -cell.

Conclusions

There are now a growing number of reports that NAADP mobilizes Ca^{2+} in pancreatic β -cells. The effects of this messenger appear to be more robust and less controversial compared to those of cADPR. The generation of selective pharmacological inhibitors for NAADP-evoked Ca^{2+} release and the identification of TPCs as target NAADP-gated Ca^{2+} release channels provide new tools for the study of NAADP-mediated Ca^{2+} signaling in pancreatic β -cells. NAADP appears to have multiple roles, which may depend on the agonist that generates its production. Whilst insulin stimulated NAADP production may control insulin synthesis and β -cell proliferation, GLP-1-mediated NAADP production may potentiate glucose-mediated Ca^{2+} signaling and insulin secretion. Perhaps most intriguing of all, NAADP-mediated Ca^{2+} release from acidic stores appears to play a key role in triggering glucose-mediated Ca^{2+} signaling. The mechanism is unclear but seems to involve the activation of calcium-activated plasma membrane cation currents which, with closure of K_{ATP} channels, may be an important component of the triggering pathway in stimulus-secretion

coupling in the pancreatic β -cell. The characterization of these currents, and the elucidation of the mechanisms coupling glucose metabolism to NAADP production are urgently required.

Acknowledgements

We would like to thank Dr. Anthony Morgan for drawing the scheme in Figure 4.

References

- Kim BJ, Park KH, Yim CY, Takasawa S, Okamoto H, Im MJ, et al. Generation of NAADP and cADPR by glucagon-like peptide-1 evokes Ca^{2+} signal that is essential for insulin secretion in mouse pancreatic islets. *Diabetes* 2008; 57:868-78.
- Naylor E, Lewis AM, Arredouani A, Parkesh R, Vasudevan SR, Galione A, et al. Discovery of a drug-like, small-molecule antagonist of the second messenger NAADP enabled by accessible virtual screening. *Nat Chem Biol* 2009; 5:220-6.
- Calcraft PJ, Ruas M, Pan Z, Cheng X, Arredouani A, Hao X, et al. NAADP mobilizes calcium from acidic organelles through two-pore channels. *Nature* 2009; 459:596-600.
- Shawl AI, Park K-H, Kim U-H. Insulin receptor signaling for the proliferation of pancreatic beta cells. Involvement of Ca^{2+} mobilizing messengers, IP_3 , NAADP and cADPR. *Islets* 2009; 1:1-8.
- Alejandro EU, Kalynyak TB, Taghizadeh F, Gwiazda KS, Rawstron EK, Jacob KJ, et al. Acute insulin signaling in pancreatic beta-cells is mediated by multiple Raf-1 dependent pathways. *Endocrinology* 2010; 151:502-12.
- Henquin JC. Triggering and amplifying pathways of regulation of insulin secretion by glucose. *Diabetes* 2000; 49:1751-60.
- Henquin JC, Nenquin M, Ravier MA, Szollosi A. Shortcomings of current models of glucose-induced insulin secretion. *Diabetes Obes Metab* 2009; 11:168-79.
- Islam MS. Calcium signaling in the islets. In: Islam MS, ed. *The Islets of Langerhans (Advances in Experimental Medicine and Biology)*: Springer 2010; 235-59.
- Miki T, Nagashima K, Tashiro F, Kotake K, Yoshitomi H, Tamamoto A, et al. Defective insulin secretion and enhanced insulin action in K_{ATP} channel-deficient mice. *Proc Natl Acad Sci USA* 1998; 95:10402-6.
- Seghers V, Nakazaki M, DeMayo F, Aguilar-Bryan L, Bryan J. *Surl* knockout mice. A model for K_{ATP} channel-independent regulation of insulin secretion. *J Biol Chem* 2000; 275:9270-7.
- Szollosi A, Nenquin M, Aguilar-Bryan L, Bryan J, Henquin JC. Glucose stimulates Ca^{2+} influx and insulin secretion in 2-week-old beta-cells lacking ATP-sensitive K^+ channels. *J Biol Chem* 2007; 282:1747-56.
- Patel S, Churchill GC, Galione A. Coordination of Ca^{2+} signalling by NAADP. *Trends Biochem Sci* 2001; 26:482-9.
- Guse AH, Lee HC. NAADP: a universal Ca^{2+} trigger. *Sci Signal* 2008; 1:10.
- Galione A, Evans AM, Ma J, Parrington J, Arredouani A, Cheng X, et al. The acid test: the discovery of two-pore channels (TPCs) as NAADP-gated endolysosomal Ca^{2+} release channels. *Pflugers Arch* 2009; 458:869-76.
- Zhu M, Ma J, Parrington J, Calcraft PJ, Galione A, Evans AM. Calcium signaling via two-pore channels: local or global, that is the question. *Am J Physiol Cell Physiol* 2010; 298: C430-C441.
- Mitchell KJ, Lai FA, Rutter GA. Ryanodine receptor type I and nicotinic acid adenine dinucleotide phosphate receptors mediate Ca^{2+} release from insulin-containing vesicles in living pancreatic beta-cells (MIN6). *J Biol Chem* 2003; 278:11057-64.
- Yamasaki M, Masgrau R, Morgan AJ, Churchill GC, Patel S, Ashcroft SJ, et al. Organelle selection determines agonist-specific Ca^{2+} signals in pancreatic acinar and beta cells. *J Biol Chem* 2004; 279:7234-40.
- Duman JG, Chen L, Palmer AE, Hille B. Contributions of intracellular compartments to calcium dynamics: Implicating an acidic store. *Traffic* 2006; 7:859-72.
- Patel S. NAADP on the up in pancreatic beta cells—a sweet message? *Bioessays* 2003; 25:430-3.

20. Johnson JD, Misler S. Nicotinic acid-adenine dinucleotide phosphate-sensitive calcium stores initiate insulin signaling in human beta cells. *Proc Natl Acad Sci USA* 2002; 99:14566-71.
21. Masgrau R, Churchill GC, Morgan AJ, Ashcroft SJ, Galione A. NAADP: a new second messenger for glucose-induced Ca²⁺ responses in clonal pancreatic beta cells. *Curr Biol* 2003; 13:247-51.
22. Lee HC, Aarhus R. A derivative of NADP mobilizes calcium stores insensitive to inositol trisphosphate and cyclic ADP-ribose. *J Biol Chem* 1995; 270:2152-7.
23. Galione A. NAADP, a new intracellular messenger that mobilizes Ca²⁺ from acidic stores. *Biochem Soc Trans* 2006; 34:922-6.
24. Aarhus R, Graeff RM, Dickey DM, Walseth TF, Lee HC. ADP-ribosyl cyclase and CD38 catalyze the synthesis of a calcium-mobilizing metabolite from NADP. *J Biol Chem* 1995; 270:30327-33.
25. Palade PT. The hunt for an alternate way to generate NAADP. *Am J Physiol Cell Physiol* 2006; 292:C4-C7.
26. Takasawa S, Nata K, Yonekura H, Okamoto H. Cyclic ADP-ribose in insulin secretion from pancreatic beta cells. *Science* 1993; 259:370-3.
27. Okamoto H, Takasawa S. Recent advances in the okamoto model: The CD38-cyclic ADP-ribose signal system and the regenerating gene protein (Reg)-reg receptor system in beta-cells. *Diabetes* 2002; 51:462-73.
28. Morgan AJ, Galione A. Investigating cADPR and NAADP in intact and broken cell preparations. *Methods* 2008; 46:194-203.
29. Churchill GC, Okada Y, Thomas JM, Genazzani AA, Patel S, Galione A. NAADP mobilizes Ca²⁺ from reserve granules, lysosome-related organelles, in sea urchin eggs. *Cell* 2002; 111:703-8.
30. Cancela JM, Churchill GC, Galione A. Coordination of agonist-induced Ca²⁺-signalling patterns by NAADP in pancreatic acinar cells. *Nature* 1999; 398:74-6.
31. Galione A, Churchill GC. Interactions between calcium release pathways: multiple messengers and multiple stores. *Cell Calcium* 2002; 32:343-54.
32. Galione A, Petersen OH. The NAADP receptor: New receptors or new regulation? *Mol Interv* 2005; 5:73-9.
33. Zhang F, Li PL. Reconstitution and characterization of a nicotinic acid adenine dinucleotide phosphate (NAADP)-sensitive Ca²⁺ release channel from liver lysosomes of rats. *J Biol Chem* 2007; 282:25259-69.
34. Pryor PR, Reimann F, Gribble FM, Luzzio JP. Mucolipin-1 Is a lysosomal membrane protein required for intracellular lactosylceramide traffic. *Traffic* 2006; 7:1388-98.
35. Gerasimenko JV, Maruyama Y, Yano K, Dolman NJ, Tepikin AV, Petersen OH, et al. NAADP mobilizes Ca²⁺ from a thapsigargin-sensitive store in the nuclear envelope by activating ryanodine receptors. *J Cell Biol* 2003; 163:271-82.
36. Dammermann W, Zhang B, Nebel M, Cordiner C, Odoardi F, Kirchberger T, et al. NAADP-mediated Ca²⁺ signaling via type 1 ryanodine receptor in T cells revealed by a synthetic NAADP antagonist. *Proc Natl Acad Sci USA* 2009; 106:10678-83.
37. Copello JA, Qi Y, Jeyakumar LH, Ogunbunmi E, Fleischer S. Lack of effect of cADP-ribose and NAADP on the activity of skeletal muscle and heart ryanodine receptors. *Cell Calcium* 2001; 30:269-84.
38. Zang X, Schieder M, Cuny H, Fenske S, Gruner C, Rotzer K, et al. The two-pore channel TPCN2 mediates NAADP-dependent Ca²⁺-release from lysosomal stores. *Pflügers Arch* 2009; 458:891-9.
39. Brailoiu E, Churamani D, Cai X, Schrlau MG, Brailoiu GC, Gao X, et al. Essential requirement for two-pore channel 1 in NAADP-mediated calcium signaling. *J Cell Biol* 2009; 186:201-9.
40. Ruas M, Rietdorf K, Arredouani A, Davis LC, Lloyd-Evans E, Koegel H, et al. Purified TPC isoforms form NAADP receptors with distinct roles for Ca²⁺ signaling and endo-lysosomal trafficking. *Curr Biol* 2010; 20:703-709.
41. Peiter E, Maathuis FJ, Mills LN, Knight H, Pelloux J, Hetherington AM, et al. The vacuolar Ca²⁺-activated channel TPC1 regulates germination and stomatal movement. *Nature* 2005; 434:404-8.
42. Guse AH. Second messenger signaling: multiple receptors for NAADP. *Curr Biol* 2009; 19:521-3.
43. Tengholm A, Hellman B, Gylfe E. The endoplasmic reticulum is a glucose-modulated high-affinity sink for Ca²⁺ in mouse pancreatic b-cells. *J Physiol* 2001; 530:533-40.
44. Arredouani A, Henquin JC, Gilon P. Contribution of the endoplasmic reticulum to the glucose-induced [Ca²⁺]_i response in mouse pancreatic islets. *Am J Physiol* 2002; 282:982-91.
45. Grapengiesser E, Gylfe E, Hellman B. Three types of cytoplasmic Ca²⁺ oscillations in stimulated pancreatic beta-cells. *Arch Biochem Biophys* 1989; 268:404-7.
46. Ahren B, Gromada J, Schmitz O. Incretin hormones and insulin secretion. *Horm Metab Res* 2004; 36:733-4.
47. Docampo R, de Souza W, Miranda K, Rohloff P, Moreno SN. Acidocalcisomes—conserved from bacteria to man. *Nat Rev Microbiol* 2005; 3:251-61.
48. Drose S, Altendorf K. Bafilomycins and concanamycins as inhibitors of V-ATPases and P-ATPases. *J Exp Biol* 1997; 200:1-8.
49. Tengholm A, Hellman B, Gylfe E. Mobilization of Ca²⁺ stores in individual pancreatic beta-cells permeabilized or not with digitonin or alpha-toxin. *Cell Calcium* 2000; 27:43-51.
50. Togashi K, Hara Y, Tominaga T, Higashi T, Konishi Y, Mori Y, et al. TRPM2 activation by cyclic ADP-ribose at body temperature is involved in insulin secretion. *EMBO J* 2006; 25:1804-1815.
51. Kato I, Yamamoto Y, Fujimura M, Noguchi N, Takasawa S, Okamoto H. CD38 disruption impairs glucose-induced increases in cyclic ADP-ribose, [Ca²⁺]_i, and insulin secretion. *J Biol Chem* 1999; 274:1869-72.
52. Ikehata F, Satoh J, Nata K, Tohgo A, Nakazawa T, Kato I, et al. Autoantibodies against CD38 (ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase) that impair glucose-induced insulin secretion in noninsulin-dependent diabetes patients. *J Clin Invest* 1998; 102:395-401.
53. Yagui K, Shimada F, Mimura M, Hashimoto N, Suzuki Y, Tokuyama Y, et al. A missense mutation in the CD38 gene, a novel factor for insulin secretion: association with Type II diabetes mellitus in Japanese subjects and evidence of abnormal function when expressed *in vitro*. *Diabetologia* 1998; 41:1024-8.
54. Pupilli C, Giannini S, Marchetti P, Lupi R, Antonelli A, Malavasi F, et al. Autoantibodies to CD38 (ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase) in Caucasian patients with diabetes: effects on insulin release from human islets. *Diabetes* 1999; 48:2309-15.
55. An NH, Han MK, Um C, Park BH, Park BJ, Kim HK, et al. Significance of ecto-cyclase activity of cd38 in insulin secretion of mouse pancreatic islet cells. *Biochem Biophys Res Commun* 2001; 282:781-6.
56. Henquin JC, Ravier MA, Nenquin M, Jonas JC, Gilon P. Hierarchy of the beta-cell signals controlling insulin secretion. *Eur J Clin Invest* 2003; 33:742-50.
57. Rutter GA, Theler JM, Li G, Wollheim CB. Ca²⁺ stores in insulin-secreting cells: lack of effect of cADP-ribose. *Cell Calcium* 1994; 16:71-80.
58. Willmott NJ, Galione A, Smith PA. A cADP-ribose antagonist does not inhibit secretagogue-, caffeine- and nitric oxide-induced Ca²⁺ responses in rat pancreatic beta-cells. *Cell Calcium* 1995; 18:411-9.
59. Malavasi F, Deaglio S, Funaro A, Ferrero E, Horenstein AL, Ortolan E, et al. Evolution and function of the ADP-ribosyl cyclase/CD38 gene family in physiology and pathology. *Physiol Rev* 2008; 88:841-86.
60. Kim SY, Cho BH, Kim UH. CD38-mediated Ca²⁺ signaling contributes to angiotensin II-induced activation of hepatic stellate cells: attenuation of hepatic fibrosis by CD38 ablation. *J Biol Chem* 2009; 285:576-82.
61. Rah SY, Mushtaq M, Nam TS, Kim SH, Kim UH. Generation of cyclic ADP-Ribose and nicotinic acid adenine dinucleotide phosphate by CD38 for Ca²⁺ signaling in interleukin-8-treated lymphokine-activated killer cells. *J Biol Chem* 2010; 286:2187-8.
62. Soares S, Thompson M, White T, Isbell A, Yamasaki M, Prakash Y, et al. NAADP as a second messenger: neither CD38 nor base-exchange reaction are necessary for *in vivo* generation of NAADP in myometrial cells. *Am J Physiol Cell Physiol* 2007; 292:227-39.
63. Davis LC, Morgan AJ, Ruas M, Wong JL, Graeff RM, Poustkka AJ, et al. Ca²⁺ signaling occurs via second messenger release from intraorganelle synthesis sites. *Curr Biol* 2008; 18:1612-8.
64. Han MK, Kim SJ, Park YR, Shin YM, Park HJ, Park KJ, et al. Antidiabetic effect of a prodrug of cysteine, L-2-oxothiazolidine-4-carboxylic acid, through CD38 dimerization and internalization. *J Biol Chem* 2002; 277:5315-21.
65. Takasawa S, Tohgo A, Noguchi N, Koguma T, Nata K, Sugimoto T, et al. Synthesis and hydrolysis of cyclic ADP-ribose by human leukocyte antigen CD38 and inhibition of the hydrolysis by ATP. *J Biol Chem* 1993; 268:26052-4.
66. Drucker DJ. The biology of incretin hormones. *Cell Metab* 2006; 3:153-65.
67. Meier JJ. The contribution of incretin hormones to the pathogenesis of type 2 diabetes. *Best Pract Res Clin Endocrinol Metab* 2009; 23:433-41.
68. Churchill GC, Galione A. Spatial control of Ca²⁺ signaling by nicotinic acid adenine dinucleotide phosphate diffusion and gradients [In Process Citation]. *J Biol Chem* 2000; 275:36867-92.
69. Churchill GC, Galione A. NAADP induces Ca²⁺ oscillations via a two-pool mechanism by priming IP₃- and cADPR-sensitive Ca²⁺ stores. *EMBO J* 2001; 20:2666-71.
70. Aspinwall CA, Lakey JR, Kennedy RT. Insulin-stimulated insulin secretion in single pancreatic beta cells. *J Biol Chem* 1999; 274:6360-5.
71. Aspinwall CA, Qian WJ, Roper MG, Kulkarni RN, Kahn CR, Kennedy RT. Roles of insulin receptor substrate-1, phosphatidylinositol 3-kinase and release of intracellular Ca²⁺ stores in insulin-stimulated insulin secretion in beta cells. *J Biol Chem* 2000; 275:22331-8.
72. Colsout B, Schraenen A, Lemaire K, Quintens R, Van Lommel L, Segal A, et al. Loss of high-frequency glucose-induced Ca²⁺ oscillations in pancreatic islets correlates with impaired glucose tolerance in *Trpm5*^{-/-} mice. *Proc Natl Acad Sci USA* 2010; 107:5208-521.
73. Cheng H, Beck A, Launay P, Gross SA, Stokes AJ, Kinet JP, et al. TRPM4 controls insulin secretion in pancreatic beta-cells. *Cell Calcium* 2007; 41:51-61.
74. Marigo V, Courville K, Hsu WH, Feng JM, Cheng H. TRPM4 impacts on Ca²⁺ signals during agonist-induced insulin secretion in pancreatic beta-cells. *Mol Cell Endocrinol* 2009; 299:194-203.
75. Brixi LR, Monteilh-Zoller MK, Ingenbrandt CS, Fleig A, Penner R, Enklaar T, et al. TRPM5 regulates glucose-stimulated insulin secretion. *Pflügers Arch* 2010; 460:69-76.
76. Ammala C, Larsson O, Berggren PO, Bokvist K, Juntti-Berggren L, Kindmark H, et al. Inositol trisphosphate-dependent periodic activation of a Ca²⁺-activated K⁺ conductance in glucose-stimulated pancreatic beta-cells. *Nature* 1991; 353:849-52.

-
77. Rutter GA, Tsuboi T, Ravier MA. Ca^{2+} microdomains and the control of insulin secretion. *Cell Calcium* 2006; 40:539-51.
 78. Arredouani A, Parkesh R, Pillinger T, Coltart G, Clough F, Shimomura K, et al. Essential role of NAADP-evoked calcium release in glucose-mediated depolarization, $[\text{Ca}^{2+}]_i$ spiking and insulin secretion in mouse pancreatic beta cell. *Diabetologica* 2009; 2009:433.