



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

The Genome Sequence of Taurine Cattle: A Window to Ruminant Biology and Evolution

Citation for published version:

Elsik, CG, Tellam, RL, Worley, KC, Gibbs, RA, Abatepaulo, ARR, Abbey, CA, Adelson, DL, Aerts, J, Ahola, V, Alexander, L, Alioto, T, Almeida, IG, Amadio, AF, Anatriello, E, Antonarakis, SE, Anzola, JM, Astashyn, A, Bahadue, SM, Baldwin, CL, Barris, W, Baxter, R, Bell, SN, Bennett, AK, Bennett, GL, Biase, FH, Boldt, CR, Bradley, DG, Brinkman, FSL, Brinkmeyer-Langford, CL, Brown, WC, Brownstein, MJ, Buhay, C, Caetano, AR, Camara, F, Carroll, JA, Carvalho, WA, Casey, T, Cervelatti, EP, Chack, J, Chacko, E, Chandrabose, MM, Chapin, JE, Chapple, CE, Chen, H-C, Chen, L, Cheng, Y, Cheng, Z, Childers, CP, Chitko-McKown, CG, Chiu, R, Choi, J, Chrast, J, Colley, AJ, Connelley, T, Cree, A, Curry, S, Dalrymple, B, Diep Dao, M, Davis, C, de Oliveira, CJF, de Miranda Santos, IKF, de Campos, TA, Deobald, H, Devinoy, E, Dickins, CM, Yan, D, Dinh, HH, De Donato, M, Donohue, KE, Donthu, R, Dovc, P, Dugan-Rocha, S, Durbin, KJ, Eberlein, A, Edgar, RC, Egan, A, Eggen, A, Eichler, EE, Elhaik, E, Ellis, SA, Elnitski, L, Ermolaeva, O, Eyras, E, Fitzsimmons, CJ, Fowler, GR, Franzin, AM, Fritz, K, Gabisi, RA, Garcia, GR, Garcia, JF, Genini, S, Gerlach, D, German, JB, Gilbert, JGR, Gill, CA, Gladney, CJ, Glass, EJ, Goodell, J, Grant, JR, Graur, D, Greaser, ML, Green, JA, Green, RD, Guan, L, Guigo, R, Hadsell, DL, Hagen, DE, Hakimov, HA, Halgren, R, Hamernik, DL, Hamilton, C, Harhay, GP, Harrow, JL, Hart, EA, Hastings, N, Havlak, P, Henrichsen, CN, Hernandez, J, Hernandez, M, Herzig, CTA, Hiendleder, SG, Hines, S, Hitchens, ME, Hlavina, W, Hobbs, M, Holder, M, Holt, RA, Hu, Z-L, Hume, J, Iivanainen, A, Ingham, A, Iso-Touru, T, Jamis, C, Jann, O, Jensen, K, Jhangiani, SN, Jiang, H, Johnson, AJ, Jones, SJM, Joshi, V, Junier, T, Kapetis, D, Kappes, SM, Kapustin, Y, Keele, JW, Kent, MP, Kerr, T, Khalil, SS, Khatib, H, Kiryutin, B, Kitts, P, Kokocinski, F, Kolbehdari, D, Kovar, CL, Kriventseva, EV, Kumar, CG, Kumar, D, Lahmers, KK, Landrum, M, Larkin, DM, Lau, LPL, Leach, R, Lee, JCM, Lee, S, Lemay, DG, Lewin, HA, Lewis, LR, Li, C, Lien, S, Liu, GE, Liu, Y, Liu, Y, Logan, KM, Lopez, J, Lozado, RJ, Lutzow, YS, Lynn, DJ, MacNeil, MD, Maglott, D, Malinverni, R, Maqbool, NJ, Marques, E, Marra, MA, Martin, WF, Martins, NF, Maruyama, SR, Matukumalli, LK, Mazza, R, McEwan, JC, McKay, SD, McLean, KL, McWilliam, S, Medrano, JF, Memili, E, Moen, C, Molenaar, AJ, Moore, SS, Moore, R, More, DD, Moreno, BT, Morgan, MB, Muntean, CT, Muzny, DM, Nandakumar, HP, Nazareth, LV, Nguyen, NB, Nicholas, FW, Nogueira, MFG, Okwuonu, GO, Olsaker, I, Pant, SD, Panzitta, F, Pastor, RCP, Patel, BM, Payne, GM, Plass, M, Poli, MA, Poslusny, N, Pruitt, K, Pu, L-L, Qin, X, Rachagani, S, Raison, JM, Ranganathan, S, Ratnakumar, A, Razpet, A, Reecy, J, Reese, JT, Ren, Y, Reymond, A, Riggs, PK, Rijnkels, M, Rincon, G, Roberts, A, Rodriguez-Osorio, N, Rodriguez-Zas, SL, Romero, NE, Rosenwald, A, Ruiz, SJ, Sabo, A, Salih, H, Sando, L, Santibanez, J, Sapojnikov, V, Schein, JE, Schmutz, SM, Schnabel, RD, Schook, L, Searle, SM, Seo, S, Shen, Y, Shen, L, Sherman, L, Skow, LC, Smith, T, Snelling, WM, Sodergren, E, Solovyev, V, Song, H, Song, J, Sonstegard, TS, Southey, BR, Souvorov, A, Spurlock, D, Steffen, D, Stone, RT, Stothard, P, Sugimoto, Y, Sweedler, JV, Takasuga, A, Tammen, I, Taniguchi, M, Taylor, JF, Telugu, BPVL, UCLA, C, Urbanski, JM, Utsunomiya, YT, Van, CP, Vattathil, S, Verschoor, CP, Waardenberg, AJ, Walker, A, Wang, Z, Ward, R, Warren, JT, Waterman, RC, Weikard, R, Weinstock, GM, Welsh, TH, Wheeler, DA, Wheeler, TT, White, SN, Whiteside, MD, Wilczek-Boney, K, Williams, JL, Williams, RL, Wilming, LG, Womack, J, Wright, RA, Wunderlich, KR, Wyss, C, Yang, MQ, Yang, J, Zdobnov, EM, Zhang, J, Zhao, F-Q, Zhu, B & Bovine Genome Sequencing & Analysis Consortium 2009, 'The Genome Sequence of Taurine Cattle: A Window to Ruminant Biology and Evolution' *Science*, vol 324, no. 5926, pp. 522-528. DOI: 10.1126/science.1169588

Digital Object Identifier (DOI):

[10.1126/science.1169588](https://doi.org/10.1126/science.1169588)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:
Science





Published in final edited form as:

Science. 2009 April 24; 324(5926): 522–528. doi:10.1126/science.1169588.

The Genome Sequence of Taurine Cattle: A window to ruminant biology and evolution

The Bovine Genome Sequencing and Analysis Consortium^{*}, Christine G. Elsik, Ross L. Tellam, and Kim C. Worley

Abstract

To understand the biology and evolution of ruminants, the cattle genome was sequenced to ~7× coverage. The cattle genome contains a minimum of 22,000 genes, with a core set of 14,345 orthologs shared among seven mammalian species of which 1,217 are absent or undetected in non-eutherian (marsupial or monotreme) genomes. Cattle-specific evolutionary breakpoint regions in chromosomes have a higher density of segmental duplications, enrichment of repetitive elements, and species-specific variations in genes associated with lactation and immune responsiveness. Genes involved in metabolism are generally highly conserved, although five metabolic genes are deleted or extensively diverged from their human orthologs. The cattle genome sequence thus provides an enabling resource for understanding mammalian evolution and accelerating livestock genetic improvement for milk and meat production.

Domesticated cattle (*Bos taurus* and *Bos taurus indicus*) provide a significant source of nutrition and livelihood to nearly 6.6 billion humans. Cattle belong to a phylogenetically distant clade compared to humans and rodents, the Cetartiodactyl order of eutherian mammals, which first appeared ~60 million years ago (1). Cattle represent the Ruminantia, which occupy diverse terrestrial environments with their ability to efficiently convert low quality forage into energy-dense fat, muscle and milk. These biological processes have been exploited by humans since domestication, which began in the Near East some 8,000-10,000 years ago (2). Since then, over 800 cattle breeds have been established representing an important world heritage and an enabling scientific resource for understanding the genetics of complex traits.

The cattle genome was assembled with methods similar to the rat and sea urchin genomes (3, 4). The most recent assemblies, Btau3.1 and Btau4.0, combined bacterial artificial chromosome (BAC) and whole genome shotgun (WGS) sequences. Btau3.1 was used for gene-specific analyses. Btau4.0, which includes finished sequence data and used different mapping methods to place the sequence on chromosomes, was used for all global analyses other than gene prediction. The contig N50 (50% of the genome is in contigs of this size or greater) is 48.7kb for both assemblies; the scaffold N50 for Btau4.0 is 1.9 Mb. In the Btau4.0 assembly, 90% of the total genome sequence was placed on the 29 autosomes and X chromosome and validated (3). Of 1.04 million expressed sequence tag (EST) sequences, 95.0% were contained in the assembled contigs. With an equivalent gene distribution in the remaining 5% of the genome, the estimated genome size is 2.87 Gbp. Comparison to 73 finished BACs and single nucleotide polymorphism (SNP) linkage data (5,6) confirmed this assembly quality with greater than 92%

Corresponding Authors: Christine G. Elsik, Department of Biology, 406 Reiss, 37th & O Streets NW, Georgetown University, Washington, DC 20057, USA, ce75@georgetown.edu, Ross L. Tellam, CSIRO Livestock Industries, 306 Carmody Rd, St Lucia, QLD 4067, Australia, ross.tellam@csiro.au, Kim C. Worley, Human Genome Sequencing Center, Department of Molecular and Human Genetics, Baylor College of Medicine, MS BCM226, One Baylor Plaza, Houston, TX 77030, USA, kworley@bcm.edu.

^{*}All authors with their affiliations and contributions are listed at the end of this paper.

genomic coverage and fewer than 0.8% of SNP were incorrectly positioned at the resolution of these maps (3,4).

We used the cattle genome to catalog protein-coding genes, microRNA genes and ruminant-specific interspersed repeats and manually annotated over 4000 genes. The consensus protein-coding gene set for Btau3.1 (OGSv1), from six predicted gene sets (4), consists of 26,835 genes with a validation rate of 82% (4). On this basis we estimate that the cattle genome contains at least 22,000 protein-coding genes. We identified 496 microRNA genes of which 135 were putative novel microRNAs (4). About half of the cattle miRNA occur in 60 genomic microRNA clusters, containing 2 to 7 microRNA genes separated by less than 10 kbp (Fig. S2). The overall GC content of the cattle genome is 41.7%, with an observed-to-expected CpG ratio of 0.234, similar to other mammals.

The cattle genome has transposable element classes similar to other mammals as well as large numbers of ruminant-specific repeats (Table S4) that comprise 27% of its genome. The consensus sequence of BovB, a non-LTR LINE retrotransposon, lacked a functional open reading frame (ORF), suggesting it was inactive (7). However, BovB repeats with intact ORF were identified in the genome and their phylogeny (Fig. S4) indicates that some are still actively expanding and evolving. Mapping chromosomal segments of high and low density ancient repeats, L2/MIR (a LINE/SINE pair) and BovB, and more recent repeats, BovB/Art2A (BovB derived SINE pair), revealed that the genome consists of ancient regions enriched for L2/MIR and recent regions enriched for BovB/Art2A (Fig. S7). Exclusion of BovB/Art2A from contiguous blocks of ancient repeats suggests that evolution of the ruminant/cattle genome experienced invasions of new repeats into regions lacking ancient repeats. Alternatively, older repeats may have been destroyed by insertion of ruminant/cattle-specific repeats. AGC trinucleotide repeats, the most common simple sequence repeat (SSR) in artiodactyls (which include cattle, pigs and sheep), are 90 and 142 fold over-represented in cattle compared to human and dog, respectively (Fig. S10). 39% of the AGC repeats in the cattle genome were associated with Bov-A2 SINE elements.

A comparative analysis examined the rate of protein evolution and the conservation of gene repertoires among orthologs in the genomes of dog, human, mouse and rat (representing placental mammals), opossum (marsupial), and platypus (monotreme). Orthology was resolved for >75% of cattle and >80% of human genes (Fig. 1A). There were 14,345 orthologous groups with representatives in human, cattle or dog, mouse or rat, and opossum or platypus, which represent 16,749 cattle and 16,177 human genes, respectively of which 12,592 are single copy orthologs. We also identified 1,217 placental mammal-specific orthologous groups with genes present in human, cattle or dog, mouse or rat, but not in opossum or platypus. About 1,000 orthologs shared between rodents and laurasiatherians (cattle and dog), many of which encode G-protein coupled receptors, appear to have been lost or may be mis-annotated in the human genome (Fig. 1B). Gene repertoire conservation among these mammals correlates with conservation at the amino-acid sequence level (Fig. 1C). The elevated rate of evolution in rodents relative to other mammals (10) was supported by the higher amino acid sequence identity between human and dog or cattle proteins relative to that between human and rodent proteins. However, maximum-likelihood analysis of amino acid substitutions in single-copy orthologs supports the accepted sister lineage relationship of primates and rodents (1) (Fig. 1D).

Alternative splicing is a major mechanism for transcript diversification (8), yet the extent of its evolutionary conservation and functional impact remain unclear. We used the cattle genome to analyze the conservation of the most common form of alternative splicing, exon skipping, defined as a triplet of exons in which the middle exon is absent in some transcripts, in a set of 1,930 exon-skipping events across human, mouse, dog and cattle (4). We examined 277 cases,

with different conservation patterns between human and mouse, in 16 different cattle tissues with RT-PCR (4). These splicing events were divided into a shared set (163 in both human and mouse) and a non-shared set (114 in human but not in mouse). Of the 277, we detected exon skipping for 188 cases in cattle (Table S5) suggesting that the majority of genes with exon skipping in human were present and regulated in cattle, and that if an event is shared between human and mouse, it was more likely to be found in cattle. It was estimated that at most 40% of exon skipping is conserved among mammals and our data agrees with the upper bound from previous analyses with human and rodents [e.g. (9)].

We constructed a cattle-human Oxford Grid (Fig. S12) (4) to conduct synteny-based chromosomal comparisons which reinforced that human genome organization is more similar to cattle than to rodents because most cattle chromosomes primarily correspond to part of one human chromosome; albeit with multiple rearrangements [e.g. (10)]. In contrast, the cattle-mouse Oxford Grid shows poorer chromosomal correspondence. Lineage-specific evolutionary breakpoints were identified for cattle, artiodactyls, and ferungulates (a group encompassing artiodactyls and carnivores, represented by cattle, pig and dog), and are shown with cattle (Fig. S11) and human sequence coordinates (Fig. 2) (4). Primate, dog, rodent, mouse, and rat lineage-specific breakpoint positions were similarly identified. A total of 124 evolutionary breakpoint regions (EBRs) were identified in the cattle lineage, of which 100 were cattle/ruminant specific and 24 were artiodactyl-specific (e.g. Fig. 2). Nine additional EBRs represent presumptive ferungulate-specific rearrangements. *Bos taurus* chromosome 16 (BTA16) is populated with four ferungulate specific EBRs, suggesting that this region was rearranged before the Artiodactyla and Carnivora divergence (Fig. 2). Such conserved regions demonstrate many inversions that occurred prior to the divergence of the carnivores and artiodactyls have probably been retained in the ancestral form within the human genome. In contrast to the cattle genome, a pig physical map identified only 77 lineage-specific EBRs. Interchromosomal rearrangements and inversions characterize most of the lineage-specific rearrangements observed in the cattle, dog, and pig genomes.

An examination of repeat families and individual transposable elements within cattle-, artiodactyl- and ferungulate-specific EBRs showed a significantly higher density of LINE-L1 elements and the ruminant-specific LINE-RTE repeat family (11) in cattle-specific EBRs relative to the remainder of the cattle genome (Table S6). In contrast, the SINE-BovA repeat family and the more ancient tRNAGlu-derived SINE repeats (12) were present in lower density in cattle-specific EBRs, similar to other LINES and SINEs (Table S7). The differences in repeat densities were generally consistent in cattle-, artiodactyl- and ferungulate-specific EBRs, with the exception of the tRNAGlu-derived and LTR-ERV1 repeats, which are at higher densities in artiodactyl EBRs compared to the rest of the genome.

The tRNAGlu (CHRS) repeats originated in the common ancestor of Suina (pigs and peccaries), Ruminantia and Cetacea (whales) (12), suggesting that tRNAGlu-derived SINEs were involved in ancestral artiodactyl chromosome rearrangements. Furthermore, the lower density of the more ancient repeat families in cattle-specific EBRs suggests that either more recently arising repeat elements were inserted into regions lacking ancient repeats or that older repeats were destroyed by this insertion (Table S7). The differing density of repeat elements in EBRs were also found in regions of homologous synteny suggesting that repeats may promote evolutionary rearrangements (see below). Differences in repeat density in cattle-specific EBRs are thus unlikely to be caused by the accumulation of repeats in EBRs after such rearrangements occur. We identified a cattle-specific EBR associated with a bidirectional promoter (Figs. S14 and S15), that may affect control of the expression of the *CYB5R4* gene which has been implicated in human diabetes and therefore may be important in the regulation of energy flow in cattle (4).

1,020 segmental duplications (SDs) corresponding to 3.1% (94.4 Mbp) of the cattle genome were identified (4). Duplications assigned to a chromosome showed a bipartite distribution with respect to length and percent identity (Fig. S16) and interchromosomal duplications were shorter (median length 2.5 kbp) and more divergent (<94% identity), relative to intrachromosomal duplications (median length 20 kbp, ~97% identity), and tended to be locally clustered (Fig. S17). Twenty-one of these duplications were >300 kbp and located in regions enriched for tandem duplications (e.g. BTA18, Fig. S18). This pattern is reminiscent of the duplication pattern of the dog, rat and mouse but different from that of primate/great-ape genomes (13,14). On average cattle SDs >10 kbp represent 11.7% of base pairs in 10 kbp intervals located within cattle-specific EBRs and 23.0% of base pairs located within the artiodactyl-specific EBRs. By contrast, in the remainder of the genome sequence assigned to chromosomes the fraction of SDs was 1.7% ($p < 1 \times 10^{-12}$). These data indicate that SDs play a role in promoting chromosome rearrangements by non-allelic homologous recombination [e.g. (15)] and suggest that either a significant fraction of the SDs observed in cattle occurred before the Ruminant-Suina split, and/or that the sites for accumulation of SDs are non-randomly distributed in artiodactyl genomes.

SDs involving genic regions may give rise to new functional paralogs. Seventy six percent (778/1,020) of the cattle SDs correspond to complete or partial gene duplications with high sequence identity (median 98.7%). This suggests that many of these gene duplications are specific to either the artiodactyla or the *Bos* lineage and tend to encode proteins that often interface with the external environment, particularly immune proteins and sensory/olfactory receptors. Several of these gene duplications are also duplicated in other mammalian lineages (e.g. cytochrome P450, sulfotransferase, ribonuclease A, defensins and pregnancy-associated glycoproteins). Paralogs located in segmental duplications present exclusively in cattle may have functional implications for the unique physiology, environment and diet of cattle.

An over-representation of genes involved in reproduction in cattle SDs (Tables S8 and S9) is associated with several gene families expressed in the ruminant placenta. These families encode the intercellular signaling proteins pregnancy associated glycoproteins (on BTA29), trophoblast Kunitz domain proteins (on BTA13) and interferon tau (*IFNT*) (on BTA8). A gene family encoding prolactin-related proteins (on BTA23) was only identified in the assembly-dependent analysis of SDs. These genes regulate ruminant-specific aspects of fetal growth, maternal adaptations to pregnancy and the coordination of parturition (16, 17). While Type I IFN genes are primarily involved in host defense (18), *IFNT* prevents regression of the corpus luteum during early pregnancy resulting in a uterine environment receptive to early conceptus development (19).

Signatures of positive selection (obtained by measurement of their rates of synonymous and nonsynonymous substitutions) identified 71 genes (4) including ten immune related genes (i.e. *IFNAR2*, *IFNG*, *CD34*, *TREM1*, *TREM1L1*, *FCER1A*, *IL23R*, *IL24*, *IL15* and *LEAP2*). As previously mentioned, immune genes are over-represented in SDs (see Table 1 and Fig S20). Examples of genes varying in cattle relative to mouse include a cluster of β -defensin genes, which encode antimicrobial peptides, the anti-microbial cathelicidin genes [which show increased sequence diversity of the mature cathelicidin peptides (20)], changes in the numbers of interferon genes (21) and the number and organization of genes involved in adaptive immune responses in cattle compared to human and mouse (4). This extensive duplication and divergence of genes involved innate immunity may be because of the substantial load of microorganisms present in the rumen of cattle which increases the risk of opportunistic infections at mucosal surfaces and positive selection for the traits that enabled stronger and more diversified innate immune responses at these locations. Another possibility is that immunity may have been under selection due to the herd structure which can promote rapid disease transmission. Also, immune function-related duplicated genes have gained non-

immune functions e.g. *IFNT* (see above), and the C-class lysozyme genes, which are involved in microbial degradation in the rumen, a fermentative foregut (see below).

There has been substantial reorganization of gene families encoding proteins present in milk. One such rearrangement affecting milk composition involves the histatherin (*HSTN*) gene within the casein gene cluster on BTA6 (Fig. S21). In the cattle genome *HSTN* is juxtaposed to a regulatory element (*BCE*) important (22) for β -casein (*CSN2*) expression, and as a probable consequence *HSTN* is regulated like the casein genes during the lactation cycle. This rearrangement that led to the juxtaposition of *HSTN* next to the *BCE* is also the probable cause of deletion of one of the two copies of α -S2-like casein genes (*CSNIS2A*) present in other mammalian genomes (23). The biological implications of this change in casein gene copy number are not yet clear.

Additionally, the cattle serum amyloid A (*SAA*) gene cluster arose from both a laurasiatherian SD and a cattle-specific EBR, resulting in two mammary gland-expressed *SAA3*-like genes, *SAA3.1* and *SAA3.2* on BTA29, and an *SAA3*-like gene on BTA15 (Fig. S21). *SAA3.2* has been shown to inhibit microbial growth (24) Two additional milk protein genes were associated with SDs: cathelicidin (*CATHL1*) and beta-2 microglobulin (*B2M*) - part of the neonatal Fc receptor (FcRn) that transfers IgG immunoglobulin across epithelial cells of many tissues including the gut and mammary gland (25,26). IgG is the predominant immunoglobulin in cow's milk compared to IgA in human milk (27), and unlike humans, which acquire passive immunity from the mother via placental transfer of immunoglobulins during pregnancy, calves acquire passive immunity via ingestion of IgG in milk (27). *B2M* is also redistributed in epithelial cells upon calving and it protects IgG from degradation (25). A genetic variant of *B2M* has negative effects on passive immune transfer (28), The additional copy of the gene encoding *B2M* may potentially be associated with the abundance of IgG in cows' milk and an increased capacity for uptake in the neonatal gut. Considering that the passive transfer of immunity to the calf is one of the important functions of milk, it is striking that lactation-related genes affected by genomic rearrangements encode immune-related proteins in milk.

Cattle metabolic pathways demonstrated a strong degree of conservation among the comprehensive set of genes involved in core mammalian metabolism (4) and permitted an examination of unique genetic events that may be related to ruminant-specific metabolic adaptations. However, among 1,032 genes examined from the human metabolic pathways, five were deleted or extensively diverged in cattle: *PLA2G4C* (phospholipase A2, group IVC), *FAAH2* (fatty acid amide hydrolase 2), *IDI2* (isopentenyl-diphosphate delta isomerase 2), *GSTT2* (glutathione S-transferase theta 2) and *TYMP* (thymidine phosphorylase), which may be adaptations that impact on fatty acid metabolism, the mevalonate pathway (synthesis of dolichols, vitamins, steroid hormones and cholesterol), detoxification, pyrimidine metabolism, respectively. Phylogenetic analysis shows that *PLA2G4C* was deleted ~87-97 Mya in the Laurasiatherian lineages (Fig. S22). Strikingly, ~20% of the sequences from two abomasum (last chamber of the cattle stomach) EST libraries (a total of 2,392 sequences) correspond to three C-type lysozyme genes. Lysozyme primarily functions in animals as an antibacterial protein suggesting they probably function in the abomasum (similar to the monogastric stomach) to degrade the cell walls of bacteria entering from the foregut (29). The cattle genome contains 10 C-type lysozyme genes (Table S14, Fig. S23) and EST evidence (Fig. S23) shows that six of the seven remaining C-type lysozyme genes are expressed primarily in the rumen and/or intestine suggesting additional roles for the encoded proteins in ruminant digestion.

In summary, the biological systems most impacted by changes in the number and organization of genes in the cattle lineage include reproduction, immunity, lactation, and digestion. We highlighted the evolutionary activity associated with chromosomal breakpoint regions and their propensity for promoting gene birth and rearrangement. These changes in the cattle lineage

probably reflect metabolic and immune adaptations due to microbial fermentation in the rumen, the herd environment and its influence on disease transmission, and the reproductive strategy of cattle. The cattle genome and associated resources will facilitate the identification of novel functions and regulatory systems of general importance in mammals and may provide an enabling tool for genetic improvement within the beef and dairy industries.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The master accession for this WGS sequencing project is AAFC03000000. The individual WGS sequences are AAFC03000001-AAFC03131728, and the scaffold records are CM000177-CM000206 (chromosomes) and DS490632-DS495890 (unplaced scaffolds).

Funded by: the National Human Genome Research Institute (NHGRI U54 HG003273); the U.S. Department of Agriculture's Agricultural Research Service (USDA ARS Agreement No. 59-0790-3-196) and Cooperative State Research, Education, and Extension Service National Research Initiative (Grant No. 2004-35216-14163); the state of Texas; Genome Canada through Genome British Columbia; The Alberta Science and Research Authority; The Commonwealth Scientific and Industrial Research Organization of Australia (CSIRO); Agritech Investments Ltd., Dairy Insight, Inc. and AgResearch Ltd., all of New Zealand; the Research Council of Norway; the Kleberg Foundation; and the National, Texas and South Dakota Beef Check-off Funds.

Appendix

Principal Investigator: Richard A. Gibbs¹

Analysis Project Leadership: Christine G. Elsik^{2,3}, Ross L. Tellam⁴

Sequencing Project Leadership: Richard A. Gibbs¹, Donna M. Muzny¹, George M. Weinstock^{5,1}

Analysis Group Organization: David L. Adelson⁶, Evan E. Eichler^{7,8}, Laura Elnitski⁹, Christine G. Elsik^{2,3}, Roderic Guigó¹⁰, Debora L. Hamernik¹¹, Steve M. Kappes¹², Harris A. Lewin^{13,14}, David J. Lynn¹⁵, Frank W. Nicholas¹⁶, Alexandre Reymond¹⁷, Monique Rijkels¹⁸, Loren C. Skow¹⁹, Ross L. Tellam⁴, Kim C. Worley¹, Evgeny M. Zdobnov^{20,21,22}

Sequencing Project White Paper: Richard A. Gibbs¹, Steve M. Kappes¹², Lawrence Schook¹³, Loren C. Skow¹⁹, George M. Weinstock^{5,1}, James Womack²³

Gene Prediction and Consensus Gene Set: Tyler Alioto¹⁰, Stylianos E. Antonarakis²⁰, Alex Astashyn²⁴, Charles E. Chapple¹⁰, Hsiu-Chuan Chen²⁴, Jacqueline Chrast¹⁷, Francisco Câmara¹⁰, Christine G. Elsik^{2,3} (leader), Olga Ermolaeva²⁴, Roderic Guigó¹⁰, Charlotte N. Henrichsen¹⁷, Wratko Hlavina²⁴, Yuri Kapustin²⁴, Boris Kiryutin²⁴, Paul Kitts²⁴, Felix Kokocinski²⁵, Melissa Landrum²⁴, Donna Maglott²⁴, Kim Pruitt²⁴, Alexandre Reymond¹⁷, Victor Sapojnikov²⁴, Stephen M. Searle²⁵, Victor Solovyev²⁶, Alexandre Souvorov²⁴, Catherine Ucla²⁰, George M. Weinstock^{5,1}, Carine Wyss²⁰

Experimental Validation of Gene Set: Tyler Alioto¹⁰, Stylianos E. Antonarakis²⁰, Charles E. Chapple¹⁰, Jacqueline Chrast¹⁷, Francisco Câmara¹⁰, Roderic Guigó¹⁰ (leader), Charlotte N. Henrichsen¹⁷, Alexandre Reymond¹⁷, Catherine Ucla²⁰, Carine Wyss²⁰

MicroRNA Analysis: Juan M. Anzola³, Daniel Gerlach^{20,21}, Evgeny M. Zdobnov^{20,21,22} (leader)

GC Composition Analysis: Eran Elhaik^{27,28}, Christine G. Elsik^{2,3} (leader), Dan Graur²⁷, Justin T. Reese²

Repeat Analysis: David L. Adelson⁶ (leader), Robert C. Edgar²⁹, John C. McEwan³⁰, Gemma M. Payne³⁰, Joy M. Raison³¹

Protein Ortholog Analysis: Thomas Junier^{19,20}, Evgenia V. Kriventseva³², Evgeny M. Zdobnov^{20,21,22} (leader)

Exon Skipping Analysis: Jacqueline Chrast¹⁷, Eduardo Eyra^{33,34}, Charlotte N. Henrichsen¹⁷, Mireya Plass³⁴, Alexandre Reymond¹⁷ (leader)

Evolutionary Breakpoint Analysis and Oxford Grid: Ravikiran Donthu¹³, Denis M. Larkin^{13,14}, Harris A. Lewin^{13,14} (leader), Frank W. Nicholas¹⁶

Bidirectional Promoter Analysis: Laura Elnitski⁹ (leader), Denis M. Larkin^{13,14}, Harris A. Lewin^{13,14}, James Reecy³⁵, Mary Q. Yang⁹

Segmental Duplication Analysis: David L. Adelson⁶, Lin Chen⁷, Ze Cheng⁷, Carol G. Chitko-McKown³⁶, Evan E. Eichler^{7,8} (leader), Laura Elnitski⁹, Christine G. Elsik^{2,3}, George E. Liu³⁷, Lakshmi K. Matukumalli^{38,37}, Jiuzhou Song³⁹, Bin Zhu³⁹

Analysis of Gene Ontology in Segmental Duplications: Christine G. Elsik^{2,3}, David J. Lynn¹⁵ (leader), Justin T. Reese²

Adaptive Evolution: Daniel G. Bradley⁴⁰, Fiona S.L. Brinkman¹⁵, Lilian P.L. Lau⁴⁰, David J. Lynn¹⁵ (leader), Matthew D. Whiteside¹⁵

Innate Immunity: Ross L. Tellam⁴ (leader), Angela Walker⁴¹, Thomas T. Wheeler⁴²

Lactation: Theresa Casey⁴³, J. Bruce German^{44,45}, Danielle G. Lemay⁴⁵, David J. Lynn¹⁵, Nauman J. Maqbool⁴⁶, Adrian J. Molenaar⁴², Monique Rijnkels¹⁸ (leader)

Metabolism: Harris A. Lewin^{13,14} (leader), Seongwon Seo⁴⁷, Paul Stothard⁴⁸

Adaptive Immunity: Cynthia L. Baldwin⁴⁹, Rebecca Baxter⁵⁰, Candice L. Brinkmeyer-Langford¹⁹, Wendy C. Brown⁵¹, Christopher P. Childers², Timothy Connelley⁵², Shirley A. Ellis⁵³, Krista Fritz¹⁹, Elizabeth J. Glass⁵⁰, Carolyn T.A. Herzig⁴⁹, Antti Iivanainen⁵⁴, Kevin K. Lahmers⁵¹, Loren C. Skow¹⁹ (leader)

Annotation Data Management: Anna K. Bennett², Christopher P. Childers², C. Michael Dickens³, Christine G. Elsik^{2,3} (leader), James G.R. Gilbert²⁵, Darren E. Hagen², Justin T. Reese², Hanni Salih³

Manual Annotation Organization: Jan Aerts⁵⁵, Alexandre R. Caetano⁵⁶, Brian Dalrymple⁴, Christine G. Elsik^{2,3}, Jose Fernando Garcia⁵⁷, Richard A. Gibbs¹, Clare A. Gill^{3,58}, Debora L. Hamernik¹¹, Stefan G. Hiendleder⁵⁹, Erdogan Memili⁶⁰, Frank W. Nicholas¹⁶, James Reecy³⁵, Monique Rijnkels¹⁸, Loren C. Skow¹⁹, Diane Spurlock³⁵, Paul Stothard⁴⁸, Ross L. Tellam⁴, George M. Weinstock^{5,1}, John L. Williams⁶¹, Kim C. Worley¹

cDNA Tissues, Libraries and Sequencing: Lee Alexander⁶², Michael J. Brownstein⁶³, Leluo Guan⁴⁸, Robert A. Holt⁶⁴ (leader), Steven J.M. Jones⁶⁴ (leader), Marco A. Marra⁶⁴ (leader), Richard Moore⁶⁴, Stephen S. Moore⁴⁸ (leader), Andy Roberts⁶², Masaaki Taniguchi^{65,48}, Richard C. Waterman⁶²

Genome Sequence Production: Joseph Chacko¹, Mimi M. Chandrabose¹, Andy Cree¹ (leader), Marvin Diep Dao¹, Huyen H. Dinh¹ (leader), Ramatu Ayiesha Gabisi¹, Sandra Hines¹, Jennifer Hume¹ (leader), Shalini N. Jhangiani¹, Vandita Joshi¹, Christie L. Kovar¹ (leader), Lora R. Lewis¹, Yih-shin Liu¹, John Lopez¹, Margaret B. Morgan¹, Donna M. Muzny¹ (leader), Ngoc Bich Nguyen¹, Geoffrey O. Okwuonu¹, San Juana Ruiz¹, Jireh Santibanez¹, Rita A. Wright¹

Sequence Finishing: Christian Buhay¹ (leader), Yan Ding¹, Shannon Dugan-Rocha¹ (leader), Judith Herdandez¹, Michael Holder¹, Aniko Sabo¹

Automated BAC Assembly: Amy Egan¹, Jason Goodell¹, Katarzyna Wilczek-Boney¹

Sequence Production Informatics: Gerald R. Fowler¹ (leader), Matthew Edward Hitchens¹, Ryan J. Lozado¹, Charles Moen¹, David Steffen^{66,1}, James T. Warren¹, Jingkun Zhang¹

BAC Mapping: Readman Chiu⁶⁴, Steven J.M. Jones⁶⁴, Marco A. Marra⁶⁴ (leader), Jacqueline E. Schein⁶⁴

Genome Assembly: K. James Durbin^{67,1}, Paul Havlak^{68,1}, Huaiyang Jiang¹, Yue Liu¹, Xiang Qin¹, Yanru Ren¹, Yufeng Shen^{1,69}, Henry Song¹, George M. Weinstock^{5,1}, Kim C. Worley¹ (leader)

Sequence Library Production: Stephanie Nicole Bell¹, Clay Davis¹, Angela Jolivet Johnson¹, Sandra Lee¹, Lynne V. Nazareth¹ (leader), Bella Mayurkumar Patel¹, Ling-Ling Pu¹, Selina Vattathil¹, Rex Lee Williams, Jr.¹

BAC Production: Stacey Curry¹, Cerissa Hamilton¹, Erica Sodergren^{5,1} (leader)

Sequence Variation Detection: Lynne V. Nazareth¹, David A. Wheeler¹

Markers and Mapping: David L. Adelson⁶, Jan Aerts⁵⁵, Wes Barris⁴, Gary L. Bennett³⁶, Brian Dalrymple⁴, André Eggen⁷⁰, Clare A. Gill^{3,58}, Ronnie D. Green⁷¹, Gregory P. Harhay³⁶, Matthew Hobbs⁷², Oliver Jann⁵⁰, Steve M. Kappes¹² (leader), John W. Keele³⁶, Matthew P. Kent⁷³, Denis M. Larkin^{13,14}, Harris A. Lewin^{13,14}, Sigbjørn Lien⁷³, John C. McEwan³⁰, Stephanie D. McKay⁷⁴, Sean McWilliam⁴, Stephen S. Moore⁴⁸, Frank W. Nicholas¹⁶, Gemma M. Payne³⁰, Abhirami Ratnakumar^{75,4}, Hanni Salih³, Robert D. Schnabel⁷⁴, Timothy Smith³⁶, Warren M. Snelling³⁶, Tad S. Sonstegard³⁷, Roger T. Stone³⁶, Yoshikazu Sugimoto⁷⁶, Akiko Takasuga⁷⁶, Jeremy F. Taylor⁷⁴, Ross L. Tellam⁴, Curtis P. Van Tassel³⁷, John L. Williams⁶¹

Genomic DNA: Michael D. MacNeil⁶²

Manual Annotation: Antonio R.R. Abatepaulo⁷⁷, Colette A. Abbey³, Jan Aerts⁵⁵, Virpi Ahola⁷⁸, Iassudara G. Almeida⁵⁷, Ariel F. Amadio⁷⁹, Elen Anatriello⁷⁷, Suria M. Bahadue², Cynthia L. Baldwin⁴⁹, Rebecca Baxter⁵⁰, Anna K. Bennett², Fernando H. Biase¹³, Clayton R. Boldt³, Candice L. Brinkmeyer-Langford¹⁹, Wendy C. Brown⁵¹, Alexandre R. Caetano⁵⁶, Jeffery A. Carroll⁸⁰, Wanessa A. Carvalho⁷⁷, Theresa Casey⁴³, Eliane P. Cervelatti⁵⁷, Elsa Chacko⁸¹, Jennifer E. Chapin³, Ye Cheng³⁵, Christopher P. Childers², Jungwoo Choi³, Adam J. Colley⁸², Timothy Connelley⁵², Tatiana A. de Campos⁵⁶, Marcos De Donato⁸³, Isabel K.F. de Miranda Santos^{56,77}, Carlo J.F. de Oliveira⁷⁷, Heather Deobald⁸⁴, Eve Devinoy⁸⁵, C. Michael Dickens³, Kaitlin E. Donohue², Peter Dovc⁸⁶, Annett Eberlein⁸⁷, Shirley A. Ellis⁵³, Carolyn J. Fitzsimmons⁵⁹, Alessandra M. Franzin⁷⁷, Krista Fritz¹⁹, Gustavo R. Garcia⁷⁷, Jose Fernando Garcia⁵⁷, Sem Genini⁶¹, J. Bruce German^{44,45}, James G.R. Gilbert²⁵, Clare A.

Gill^{3,58}, Cody J. Gladney³, Elizabeth J. Glass⁵⁰, Jason R. Grant⁴⁸, Marion L. Greaser⁸⁸, Jonathan A. Green⁷⁴, Darryl L. Hadsell¹⁸, Darren E. Hagen², Hatam A. Hakimov⁸⁹, Rob Halgren⁴³, Jennifer L. Harrow²⁵, Elizabeth A. Hart²⁵, Nicola Hastings^{90,50}, Marta Hernandez⁹¹, Carolyn T.A. Herzig⁴⁹, Stefan G. Hiendleder⁵⁹, Matthew Hobbs⁷², Zhi-Liang Hu³⁵, Antti Iivanainen⁵⁴, Aaron Ingham⁴, Terhi Iso-Touru⁷⁸, Catherine Jamis², Oliver Jann⁵⁰, Kirsty Jensen⁵⁰, Dimos Kapetis⁶¹, Tovah Kerr⁵¹, Sari S. Khalil², Hasan Khatib⁹², Davood Kolbehdari^{48,93}, Charu G. Kumar¹³, Dinesh Kumar^{94,35}, Richard Leach⁵⁰, Justin C-M Lee², Danielle G. Lemay⁴⁵, Changxi Li^{95,48}, George E. Liu³⁷, Krystin M. Logan⁹⁶, Roberto Malinverni⁶¹, Nauman J. Maqbool⁴⁶, Elisa Marques⁴⁸, William F. Martin⁴⁵, Natalia F. Martins⁵⁶, Sandra R. Maruyama⁷⁷, Raffaele Mazza⁹⁷, Kim L. McLean⁸⁴, Juan F. Medrano⁹⁸, Erdogan Memili⁶⁰, Adrian J. Molenaar⁴², Barbara T. Moreno⁵⁷, Daniela D. Moré⁷⁷, Carl T. Muntean³, Hari P. Nandakumar¹⁹, Marcelo F.G. Nogueira⁹⁹, Ingrid Olsaker¹⁰⁰, Sameer D. Pant⁸², Francesca Panzitta⁶¹, Rosemeire C.P. Pastor⁵⁷, Mario A. Poli¹⁰¹, Nathan Poslusny², Satyanarayana Rachagani³⁵, Shoba Ranganathan^{81,102}, Andrej Razpet⁸⁶, James Reecy³⁵, Penny K. Riggs^{3,58}, Monique Rijnkels¹⁸, Gonzalo Rincon⁹⁸, Nelida Rodriguez-Osorio^{60,103}, Sandra L. Rodriguez-Zas¹³, Natasha E. Romero³, Anne Rosenwald², Lillian Sando⁴, Sheila M. Schmutz⁸⁴, Seongwon Seo⁴⁷, Libing Shen², Laura Sherman⁴⁸, Loren C. Skow¹⁹, Bruce R. Southey¹⁰⁴, Diane Spurlock³⁵, Ylva Strandberg Lutzow⁴, Jonathan V. Sweedler¹⁰⁴, Imke Tammen⁷², Masaaki Taniguchi^{65,48}, Ross L. Tellam⁴, Bhanu Prakash V.L. Telugu⁷⁴, Jennifer M. Urbanski², Yuri T. Utsunomiya⁵⁷, Chris P. Verschoor⁸², Ashley J. Waardenberg^{4,105}, Angela Walker⁴¹, Zhiquan Wang⁴⁸, Robert Ward¹⁰⁶, Rosemarie Weikard⁸⁷, Thomas H. Welsh, Jr.^{3,58}, Thomas T. Wheeler⁴², Stephen N. White^{51,107}, John L. Williams⁶¹, Laurens G. Wilming²⁵, Kris R. Wunderlich³, Jianqi Yang¹⁰⁸, Feng-Qi Zhao¹⁰⁹

¹Human Genome Sequencing Center, Department of Molecular and Human Genetics, Baylor College of Medicine, One Baylor Plaza, Houston, TX, 77030, USA

²Department of Biology, 406 Reiss, Georgetown University, 37th & O Streets NW, Washington, DC, 20057, USA

³Department of Animal Science, Texas A&M University, 2471 TAMU, College Station, TX, 77843-2471, USA

⁴Livestock Industries, Commonwealth Scientific and Industrial Research Organization (CSIRO), 306 Carmody Road, St. Lucia, Queensland, 4067, Australia

⁵The Genome Center at Washington University, Washington University School of Medicine, 4444 Forest Park Ave, St. Louis, MO, 63108, USA

⁶School of Molecular and Biomedical Science, School of Agriculture, Food and Wine, The University of Adelaide, Adelaide, SA, 5005, Australia

⁷Department of Genome Sciences, University of Washington, 1705 NE Pacific St, Seattle, WA, 98195-5065, USA

⁸Howard Hughes Medical Institute, Seattle, WA, 98195, USA

⁹National Human Genome Research Institute, National Institutes of Health, 5625 Fishers Lane, Rockville, MD, 20878, USA

¹⁰Center for Genomic Regulation and Grup de Recerca en Informàtica Biomèdica, Institut Municipal d'Investigació Mèdica, Universitat Pompeu Fabra, 08003 Barcelona, Catalonia, Spain

- ¹¹USDA- Cooperative State Research, Education, & Extension Service, 1400 Independence Ave SW, Stop 2220, Washington, DC, 20250-2220, USA
- ¹²National Program Staff, USDA - Agricultural Research Service, 5601 Sunnyside Avenue, Beltsville, MD, USA
- ¹³Department of Animal Sciences, University of Illinois at Urbana-Champaign, 1201 West Gregory Drive, Urbana, IL, 61801, USA
- ¹⁴Institute for Genomic Biology, University of Illinois at Urbana-Champaign, 1201 West Gregory Drive, Urbana, IL, 61801, USA
- ¹⁵Department of Molecular Biology and Biochemistry, Simon Fraser University, 8888 University Drive, Burnaby, BC, V5A 1S6, Canada
- ¹⁶Faculty of Veterinary Science, University of Sydney, Sydney, NSW, 2006, Australia
- ¹⁷Center for Integrative Genomics, University of Lausanne, Lausanne, 1015, Switzerland
- ¹⁸USDA/ARS Children's Nutrition Research Center, Department of Pediatrics-Nutrition, Baylor College of Medicine, 1100 Bates Street, Houston, TX, 77030-2600, USA
- ¹⁹Department of Veterinary Integrative Biosciences, Texas A&M University, College Station, TX, 77843, USA
- ²⁰Department of Genetic Medicine and Development, University of Geneva Medical School, 1 rue Michel-Servet, Geneva, 1211, Switzerland
- ²¹Swiss Institute of Bioinformatics, 1 rue Michel-Servet, Geneva, 1211, Switzerland
- ²²Division of Molecular Biosciences, Imperial College London, South Kensington Campus, London, SW7 2AZ, UK
- ²³Department of Veterinary Pathobiology, Texas A&M University, College Station, TX, 77843, USA
- ²⁴National Center for Biotechnology Information, National Library of Medicine, National Institutes of Health, Bethesda, MD, 20892, USA
- ²⁵Informatics Department, Wellcome Trust Sanger Institute, Hinxton, Cambridge, CB10 1HH, UK
- ²⁶Department of Computer Science, University of London, Royal Holloway, Egham, Surrey, TW20 0EX, UK
- ²⁷Department of Biology and Biochemistry, University of Houston, Houston, TX, 77204, USA
- ²⁸McKusick - Nathans Institute of Genetic Medicine, BRB 579, Johns Hopkins University School of Medicine, 733 N. Broadway, Baltimore, MD, 21205, USA
- ²⁹45 Monterey Drive, Tiburon, CA, 94920, USA
- ³⁰Animal Genomics, AgResearch, Invermay, PB 50034, Mosgiel, 9053, New Zealand
- ³¹eResearch SA, University of Adelaide, North Terrace, Adelaide, South Australia, 5005, Australia

³²Department of Structural Biology and Bioinformatics, University of Geneva Medical School, 1 rue Michel-Servet, Geneva, 1211, Switzerland

³³Catalan Institution for Research and Advanced Studies, 08010 Barcelona, Catalonia, Spain

³⁴Computational Genomics, Universitat Pompeu Fabra, 08003 Barcelona, Catalonia, Spain

³⁵Department of Animal Science, Iowa State University, 2255 Kildee Hall, Ames, IA, 50011-3150, USA

³⁶Meat Animal Research Center, USDA - Agricultural Research Service, Clay Center, NE, 68933, USA

³⁷Bovine Functional Genomics Laboratory, USDA - Agricultural Research Service, BARC-East, Beltsville, MD, 20705, USA

³⁸Department of Bioinformatics and Computational Biology, George Mason University, 10900 University Blvd, Manassas, VA, 20110, USA

³⁹Department of Bioengineering, University of Maryland, College Park, MD, 20742, USA

⁴⁰Smurfit Institute of Genetics, Trinity College Dublin, Dublin 2, Ireland

⁴¹Department of Veterinary Pathobiology, 245 Bond Life Sciences Center, University of Missouri, Columbia, MO, 65211, USA

⁴²Dairy Science and Technology Section, AgResearch, Ruakura Research Centre, East Street, Private Bag 3123, Hamilton, 3240, New Zealand

⁴³Department of Animal Science, Michigan State University, East Lansing, MI, 48824-1225, USA

⁴⁴Nestlé Research Centre, Vers chez les Blanc CH, Lausanne 26, 1000, Switzerland

⁴⁵Department of Food Science and Technology, University of California-Davis, Davis, CA, 95616, USA

⁴⁶Bioinformatics, Mathematics and Statistics, AgResearch, Ruakura Research Centre, East Street, Private Bag 3123, Hamilton, 3240, New Zealand

⁴⁷Division of Animal Science and Resource, Chungnam National University, Daejeon, 305-764, Korea

⁴⁸Department of Agricultural, Food and Nutritional Science, University of Alberta, 410 AgFor Centre, Edmonton, Alberta, T6G 2P5, Canada

⁴⁹Department of Veterinary and Animal Sciences, University of Massachusetts, Amherst, MA, 01003, USA

⁵⁰The Roslin Institute and Royal (Dick) School of Veterinary Studies, University of Edinburgh, Roslin, Midlothian, EH25 9PS, UK

⁵¹Department of Veterinary Microbiology and Pathology, Washington State University, Pullman, WA, 99164, USA

⁵²Division of Infection and Immunity, The Roslin Institute, Royal (Dick) School of Veterinary Science, University of Edinburgh, Roslin, Midlothian, EH25 9RG, UK

⁵³Immunology Division, Institute for Animal Health, Compton, RG20 7NN, UK

⁵⁴Department of Basic Veterinary Sciences, University of Helsinki, POB 66, Helsinki, FIN-00014, Finland

⁵⁵Genome Dynamics and Evolution, Wellcome Trust Sanger Institute, Hinxton, Cambridge, CB10 1SA, UK

⁵⁶Embrapa Recursos Genéticos e Biotecnologia, Final Av. W/5 Norte, Brasília, DF, 70770-900, Brazil

⁵⁷Animal Production and Health Department, UNESP - Sao Paulo State University, Aracatuba, SP, 16050-680, Brazil

⁵⁸Texas AgriLife Research, College Station, TX, 77843, USA

⁵⁹JS Davies Epigenetics and Genetics Group, School of Agriculture, Food & Wine and Research Centre for Reproductive Health, The University of Adelaide, Roseworthy Campus, Roseworthy, SA, 5371, Australia

⁶⁰Department of Animal and Dairy Sciences, Mississippi Agricultural and Forestry Experiment Station, Mississippi State University, Mississippi State, MS, 39762, USA

⁶¹Parco Tecnologico Padano, Via Einstein, Polo Universitario, Lodi, 26900, Italy

⁶²Fort Keogh Livestock and Range Research Laboratory, USDA - Agricultural Research Service, Miles City, MT, 59301, USA

⁶³Building 49, B1EE16, 49 Convent Dr, Bethesda, MD, 20892, USA

⁶⁴Genome Sciences Centre, British Columbia Cancer Agency, 675 West 10th Avenue, Vancouver, British Columbia, V5Z 1L3, Canada

⁶⁵Division of Animal Sciences, National Institute of Agrobiological Sciences, Tsukuba, Ibaraki, 305-8602, Japan

⁶⁶Bioinformatics Research Center, Baylor College of Medicine, One Baylor Plaza, Houston, TX, 77030, USA

⁶⁷Department of Biomolecular Engineering, University of California at Santa Cruz, Santa Cruz, CA, 95064, USA

⁶⁸Department of Computer Science, University of Houston, Houston, TX, 77204-3010, USA

⁶⁹Department of Computer Science and Center for Computational Biology and Bioinformatics, Columbia University, New York, NY, 10032, USA

⁷⁰INRA, Animal Genetics and Integrative Biology, Bovine Genetics and Genomics, 78350 Jouy-en-Josas, France

⁷¹Pfizer Animal Genetics, Pfizer Animal Health, New York, NY, 10017, USA

⁷²Faculty of Veterinary Science, University of Sydney, Camden, NSW, 2570, Australia


- ⁷³Centre for Integrative Genetics and Department of Animal and Aquacultural Sciences, Norwegian University of Life Sciences, Arboretveien 6, Ås, 1432, Norway
- ⁷⁴Division of Animal Sciences, University of Missouri, 920 East Campus Drive, Columbia, MO, 65211, USA
- ⁷⁵Department of Medical Biochemistry and Microbiology, Uppsala University, Uppsala Biomedical Centre Husargatan 3, Uppsala, 75 123, Sweden
- ⁷⁶Shirakawa Institute of Animal Genetics, Nishigo, Fukushima 961-8061, Japan
- ⁷⁷Department of Biochemistry and Immunology, Ribeirão Preto Medical School, University of São Paulo, Av Bandeirantes 3900, Ribeirão Preto, SP, 14049-900, Brazil
- ⁷⁸Biotechnology and Food Research, MTT Agrifood Research Finland, Jokioinen, FI-31600, Finland
- ⁷⁹EEA Rafaela, Instituto Nacional de Tecnología Agropecuaria (INTA), Ruta 34 Km 227, Rafaela, Santa Fe, 2300, Argentina
- ⁸⁰Livestock Issues Research Unit, USDA - Agricultural Research Service, Lubbock, TX, 79403, USA
- ⁸¹Department of Chemistry and Biomolecular Sciences & ARC Centre of Excellence in Bioinformatics, Macquarie University, Sydney, 2109, NSW, Australia
- ⁸²Department of Animal and Poultry Science, University of Guelph, Guelph, ON, N1G2W1, Canada
- ⁸³Instituto de Investigaciones en Biomedicina y Ciencias Aplicadas, Universidad de Oriente, Av. Universidad, Cumana, Sucre, 6101, Venezuela
- ⁸⁴Department of Animal and Poultry Science, University of Saskatchewan, Saskatoon, SK, S7N 5A8, Canada
- ⁸⁵INRA – UR1196 Génomique et Physiologie de la Lactation, F78352 Jouy-en-Josas, France
- ⁸⁶Department of Animal Science, University of Ljubljana, Groblje 3, Domzale, SI-1230, Slovenia
- ⁸⁷Research Unit Molecular Biology, Research Institute for the Biology of Farm Animals (FBN), Dummerstorf, 18196, Germany
- ⁸⁸Department of Animal Sciences, University of Wisconsin-Madison, 1805 Linden Drive, Madison, WI, 53706, USA
- ⁸⁹Department of Molecular and Cellular Biology, University of Guelph, Guelph, ON, N1G 2W1, Canada
- ⁹⁰Cell Biology and Biophysics, EMBL-Heidelberg, Meyerhofstraße 1, Heidelberg, Germany
- ⁹¹Laboratory of Molecular Biology, Instituto Tecnológico Agrario de Castilla y Leon (ITACyL), Ctra. Burgos km 119, Valladolid, 47071, Spain
- ⁹²Department of Dairy Science, University of Wisconsin, Madison, WI, 53706, USA

- ⁹³Monsanto Company, 3302 SE Convenience Blvd, Ankeny, Iowa, 50021, USA
- ⁹⁴Genes & Genetic Resources Molecular Analysis Lab, National Bureau of Animal Genetic Resources, Baldi Bye Pass, Karnal, Haryana, 132001, India
- ⁹⁵Lacombe Research Centre, Agriculture and Agri-Food Canada, Lacombe, Alberta, T4L 1W1, Canada
- ⁹⁶Biomedical Sciences, University of Guelph, Guelph, ON, N1G2W6, Canada
- ⁹⁷Zootechnics Institute, Università Cattolica del Sacro Cuore, via Emilia Parmense 84, Piacenza, 29100, Italy
- ⁹⁸Department of Animal Science, University of California, Davis, Davis, CA, 95616, USA
- ⁹⁹Departamento de Ciências Biológicas, Faculdade de Ciências e Letras, UNESP – São Paulo State University, Av Dom Antônio 2100, Vila Tênis Clube, Assis, SP, 19806-900, Brazil
- ¹⁰⁰Department of Basic Sciences and Aquatic Medicine, Norwegian School of Veterinary Science, P.O. Box 8146 Dep, Oslo, NO-0033, Norway
- ¹⁰¹Instituto de Genética Ewald Favret, Instituto Nacional de Tecnología Agropecuaria (INTA), Las Cabañas y de Los Reseros s/n CC25, Castelar, Buenos Aires, B1712WAA, Argentina
- ¹⁰²Department of Biochemistry, Yong Loo Lin School of Medicine, National University of Singapore, 8 Medical Drive, Singapore, 117597, Singapore
- ¹⁰³Grupo CENTAURO, Universidad de Antioquia, Medellín, Colombia
- ¹⁰⁴Department of Chemistry, University of Illinois, Urbana, IL, 61801, USA
- ¹⁰⁵Eskitis Institute for Cell and Molecular Therapies, Griffith University, Nathan, Queensland, 4111, Australia
- ¹⁰⁶Nutrition and Food Sciences, Utah State University, Logan, UT, 84322, USA
- ¹⁰⁷Animal Disease Research Unit, USDA - Agricultural Research Service, Pullman, WA, 99164, USA
- ¹⁰⁸Department of Pharmacology, 2-344 BSB, University of Iowa, 51 Newton Road, Iowa City, IA, 52242, USA
- ¹⁰⁹Department of Animal Science, 211 Terrill, University of Vermont, 570 Main Street, Burlington, VT, 05405, USA

References

1. Murphy WJ, Pevzner PA, O'Brien SJ. Trends Genet Dec;2004 20:631–639. [PubMed: 15522459]
2. Willham RL. Journal of Animal Science 1986;62:1742–1758.
3. Liu Y, et al. BMC Genomics. In Press.
4. Materials, methods, and additional discussion are available on *Science* online.
5. Nilsen H, et al. Anim Genet Apr;2008 39:97–104. [PubMed: 18307581]
6. Prasad A, et al. BMC Genomics 2007;8:310. [PubMed: 17784962]
7. Malik HS, Eickbush TH. Mol Biol Evol Sep;1998 15:1123–1134. [PubMed: 9729877]
8. Modrek B, Lee CJ. Nat Genet Jun;2003 34:177–180. [PubMed: 12730695]

9. Sorek R, Shamir R, Ast G. *Trends Genet* Feb;2004 20:68–71. [PubMed: 14746986]
10. Everts-van der Wind A, et al. *Proc Natl Acad Sci U S A* Dec 20;2005 102:18526–18531. [PubMed: 16339895]
11. Kordis D, Gubensek F. *Gene* Sep 30;1999 238:171–178. [PubMed: 10570995]
12. Shimamura M, Abe H, Nikaido M, Ohshima K, Okada N. *Mol Biol Evol* Aug;1999 16:1046–1060. [PubMed: 10474901]
13. Bailey JA, Eichler EE. *Nat Rev Genet* Jul;2006 7:552–564. [PubMed: 16770338]
14. Bailey JA, et al. *Science* Aug 9;2002 297:1003–1007. [PubMed: 12169732]
15. Murphy WJ, et al. *Science* Jul 22;2005 309:613–617. [PubMed: 16040707]
16. Hashizume K, et al. *Reprod Fertil Dev* 2007;19:79–90. [PubMed: 17389137]
17. Larson JH, et al. *Physiol Genomics* May 16;2006 25:405–413. [PubMed: 16554549]
18. Zhang SY, et al. *Immunol Rev* Dec;2008 226:29–40. [PubMed: 19161414]
19. Roberts RM, Chen Y, Ezashi T, Walker AM. *Semin Cell Dev Biol* Apr;2008 19:170–177. [PubMed: 18032074]
20. Scocchi M, Wang S, Zanetti M. *FEBS Lett* Nov 17;1997 417:311–315. [PubMed: 9409740]
21. Katze MG, He Y, Gale M Jr. *Nat Rev Immunol* Sep;2002 2:675–687. [PubMed: 12209136]
22. Schmidhauser C, et al. *Mol Biol Cell* Jun;1992 3:699–709. [PubMed: 1498370]
23. Rijnkels M, Elnitski L, Miller W, Rosen JM. *Genomics* Oct;2003 82:417–432. [PubMed: 13679022]
24. Molenaar AJ, et al. *Biomarkers* 2009;14:26–37. [PubMed: 19283521]
25. Mayer B, et al. *J Dairy Res* 2005;72:107–112. Spec No. [PubMed: 16180728]
26. Roopenian DC, Akilesh S. *Nat Rev Immunol* Sep;2007 7:715–725. [PubMed: 17703228]
27. Newby TJ, Stokes CR, Bourne FJ. *Vet Immunol Immunopathol* Jan;1982 3:67–94. [PubMed: 7101722]
28. Clawson ML, et al. *Mamm Genome* Mar;2004 15:227–236. [PubMed: 15014972]
29. Irwin DM. *J Mol Evol* Sep;1995 41:299–312. [PubMed: 7563116]

**Fig. 1.**

Protein orthology comparison among genomes of cattle, dog, human, mouse and rat (*Bos taurus*, *Canis familiaris*, *Homo sapiens*, *Mus musculus*, *Rattus norvegicus*, representing placental mammals), opossum (*Monodelphis domestica*; marsupial), and platypus (*Ornithorhynchus anatinus*; monotreme). (A) The majority of mammalian genes are orthologous, with over half preserved as single-copies (dark blue); a few thousand have species-specific duplications (blue); another few thousand have been lost in specific lineages (orange). We also show those lacking confident orthology assignment (green), and those that are apparently lineage specific [unique (white)]. Placental-specific orthologs are shown in pink. Single- or multiple-copy genes were defined on the basis of representatives in human, bovine or dog, mouse or rat, and opossum or platypus. (B) Venn diagram showing shared orthologous groups (duplicated genes were counted as one) between laurasiatherians (cattle and dog), human, rodents (mouse and rat), and non-placental mammals (opossum and platypus) on the basis of the presence of a representative gene in at least one of the grouped species (as in A). (C) Distribution of ortholog protein identities between human and the other species for a subset of strictly conserved single-copy orthologs. (D) A maximum likelihood phylogenetic tree using all single-copy orthologs supports the accepted phylogeny and quantifies the relative rates of molecular evolution expressed as the branch lengths.

HSA1: 175 – 247 Mbp




Fig. 2. Examples of evolutionary breakpoint regions (EBRs). Ferungulate- artiodactyl- and primate-specific EBRs on HSA1 at 175-247 Mbp (other lineage-specific EBRs not shown). Homologous synteny blocks constructed for the macaque, chimp, cattle, dog, mouse, rat and pig genomes were used for pair-wise comparisons (4). White areas correspond to EBRs. Arrows to the right of the chromosome ideogram indicate positions of representative cattle-specific, artiodactyl-specific (specific to the chromosomes of pigs and cattle), ferungulate-specific (cattle, dog and pig), primate-specific (human, macaque, chimp), and hominoid-specific (human and chimp) rearrangements. Opossum is shown as an outgroup to the eutherian clade, which allows classification of ferungulate-specific EBRs.

Table 1

Changes in the number of genes in innate immune gene families

Gene Family	Bovine	Human	Murine
Cathelicidin	10	1	1
<i>RNase</i>	21	13	25
<i>BPI</i> -like	13	9	11
<i>BPI/LBP</i>	3	2	2
β -Defensin ¹	~106	39	52
Interferon subfamilies ²			
<i>IFNK</i>	1	1	1
<i>IFNE</i>	1	1	1
<i>IFNB</i>	6	1	1
<i>IFNA</i>	13	13	14
<i>IFNW</i>	24	1	0
<i>IFNT</i>	3	0	0
<i>IFNX</i> ⁴	3	0	0
<i>IFNL</i>	0	3	2
<i>IFNZ</i>	0	0	2
C-type Lysozyme	10	1	3
<i>ULBP</i> ³	30	3	1

¹ Many of the β -defensin genes are present in unassigned scaffolds, i.e. they are not yet part of the current assembly. The exact number of genes is uncertain.

² Pseudogenes predicted on the basis of frame-shift mutations or stop codons within the first 100 amino acids of the coding sequence have been excluded from the table.

³ (17)

⁴ The IFNX genes represent a newly discovered subfamily of IFN and are so named for convenience.