



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Yeast 18S rRNA Dimethylase Dim1p: a Quality Control Mechanism in Ribosome Synthesis?

Citation for published version:

Lafontaine, DLJ, Preiss, T & Tollervey, D 1998, 'Yeast 18S rRNA Dimethylase Dim1p: a Quality Control Mechanism in Ribosome Synthesis?' *Molecular and Cellular Biology*, vol 18, no. 4, pp. 2360-2370.

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Molecular and Cellular Biology

Publisher Rights Statement:

RoMEO blue

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Yeast 18S rRNA Dimethylase Dim1p: a Quality Control Mechanism in Ribosome Synthesis?

DENIS L. J. LAFONTAINE,^{1*} THOMAS PREISS,² AND DAVID TOLLERVEY¹

*Institute of Cell and Molecular Biology, The University of Edinburgh, EH9 3JR Edinburgh, Scotland,¹
and European Molecular Biology Laboratory, Gene Expression, Heidelberg, Germany²*

Received 24 October 1997/Returned for modification 5 December 1997/Accepted 21 January 1998

One of the few rRNA modifications conserved between bacteria and eukaryotes is the base dimethylation present at the 3' end of the small subunit rRNA. In the yeast *Saccharomyces cerevisiae*, this modification is carried out by Dim1p. We previously reported that genetic depletion of Dim1p not only blocked this modification but also strongly inhibited the pre-rRNA processing steps that lead to the synthesis of 18S rRNA. This prevented the formation of mature but unmodified 18S rRNA. The processing steps inhibited were nucleolar, and consistent with this, Dim1p was shown to localize mostly to this cellular compartment. *dim1-2* was isolated from a library of conditionally lethal alleles of *DIM1*. In *dim1-2* strains, pre-rRNA processing was not affected at the permissive temperature for growth, but dimethylation was blocked, leading to strong accumulation of nondimethylated 18S rRNA. This demonstrates that the enzymatic function of Dim1p in dimethylation can be separated from its involvement in pre-rRNA processing. The growth rate of *dim1-2* strains was not affected, showing the dimethylation to be dispensable *in vivo*. Extracts of *dim1-2* strains, however, were incompetent for translation *in vitro*. This suggests that dimethylation is required under the suboptimal *in vitro* conditions but only fine-tunes ribosomal function *in vivo*. Unexpectedly, when transcription of pre-rRNA was driven by a polymerase II *PGK* promoter, its processing became insensitive to temperature-sensitive mutations in *DIM1* or to depletion of Dim1p. This observation, which demonstrates that Dim1p is not directly required for pre-rRNA processing reactions, is consistent with the inhibition of pre-rRNA processing by an active repression system in the absence of Dim1p.

In most eukaryotes, three of the four rRNA species are excised from a single large RNA polymerase I (Pol I) transcript (35S pre-rRNA in yeast) which is processed in a complex pathway involving both endonucleolytic cleavages and exonucleolytic digestions (Fig. 1) (reviewed in references 24, 42, and 46). The fourth mature species, 5S rRNA, is transcribed and processed independently. During pre-rRNA processing, many specific nucleotides within the rRNAs are covalently modified. The major types of posttranscriptional modification are the isomerization of uracil to pseudouridine, 2'-O methylation of the ribose moieties, and base methylation. In eukaryotes, interactions between *trans*-acting guide small nucleolar RNAs (snoRNAs) and pre-rRNA identify nucleotides that are to be modified by 2'-O methylation or pseudouridylation (8, 12, 20, 31, 32). Surprisingly, all of the guide snoRNAs tested so far in yeast are nonessential for growth, demonstrating that the corresponding modifications are dispensable.

The only base modifications that have been studied in a eukaryote are the m⁶Am²A doublet present at the 3' end of the small subunit (SSU) rRNA (18S rRNA). In the yeast *Saccharomyces cerevisiae*, these modifications are made by the Dim1p dimethylase. In contrast to 2'-O methylation and pseudouridylation, 18S dimethylation does not appear to involve a guide snoRNA cofactor. Moreover, expression of Dim1p in *Escherichia coli* is able to restore dimethylation to *ksgA* mutants that lack this activity, presumably showing that Dim1p is able to function in the absence of snoRNA cofactors (23).

Dim1p is essential for viability (23), but the requirement for the dimethylated nucleotides was unclear since cells depleted

of Dim1p not only lacked dimethylation but were also strongly defective in the pre-rRNA cleavages at sites A₁ and A₂ that generate 20S pre-rRNA (Fig. 1) (26). 20S pre-rRNA is the immediate precursor to 18S rRNA, so this had the effect of preventing the synthesis of mature but nonmodified rRNA. Pre-rRNA molecules in which the two dimethylated adenosine residues are replaced by guanosine, which cannot be modified, were processed normally, showing that the m⁶Am²A nucleotides are not themselves required for cleavage (26). Expression of 18S rRNA with the double G substitution did not support growth, but it was not clear whether the lack of dimethylation or the substitution of the A residues, which are themselves universally conserved in evolution, was responsible for this defect.

In *E. coli*, *ksgA* strains that lack dimethylation are resistant to kasugamycin, an antibiotic belonging to the aminoglycoside family (14, 15). *ksgA* strains are only marginally affected for growth, but *ksgA* extracts show various defects in translation *in vitro*: (i) initiation requires a higher concentration of IF3 in the absence of IF1 (35), (ii) accuracy is affected (43), and (iii) the affinity between the subunits is decreased (34; reviewed in reference 45). This is consistent with the localization of the m⁶Am²A residues at the interface between the ribosomal subunits at a site where crucial interactions take place during translation (7, 29, 41). However, *in vitro* reconstitution experiments showed that neither dimethylation nor the twin adenosines are crucial for 30S subunit assembly and function (10, 22).

Previous analyses, therefore, had left two outstanding questions. What is the requirement for dimethylated nucleotides in the synthesis and function of eukaryotic ribosomes? What is the basis of the requirement for Dim1p in pre-rRNA processing reactions? Specifically, does a regulatory system monitor the association of Dim1p with pre-rRNA and inhibit process-

* Corresponding author. Mailing address: Institute of Cell and Molecular Biology, University of Edinburgh, Swann Building, King's Buildings, EH9 3JR Edinburgh, Scotland. Phone: 44 131 650 7093. Fax: 44 131 650 7040 or 8650. E-mail: denis.lafontaine@ed.ac.uk.




FIG. 1. Structure of yeast 35S pre-rRNA and the pre-rRNA processing pathway. (A) 35S pre-rRNA. The sequences encoding the mature 18S, 5.8S, and 25S rRNAs are embedded in the 5' and 3' external transcribed spacers (ETS) and in ITS1 and ITS2. Sites of pre-rRNA processing are indicated with uppercase letters (A₀ to E), and the oligonucleotides used for Northern hybridization and primer extension are indicated with lowercase letters (a to h). The site of dimethylation is denoted as m⁶A. (B) Pre-rRNA processing pathway. Processing of the 35S primary transcript starts at site A₀ in the 5' ETS. The resulting 33S pre-rRNA is processed at sites A₁ and A₂, giving rise successively to the 32S pre-rRNA and to the 20S and 27SA₂ precursors. Cleavage at site A₂ separates the pre-rRNAs destined for the small and large ribosomal subunits. The 20S precursor is dimethylated and endonucleolytically cleaved at site D to yield mature 18S rRNA. Cleavage of 27SA₂ at site A₃, by RNase MRP, is rapidly followed by exonucleolytic digestion to site B₁S, generating the 27SB_S precursor. Mature 25S rRNA and 7S pre-rRNA are released from 27SB_S following cleavages at sites C₁ and C₂. 7S pre-rRNA undergoes a rapid 3'→5' exonuclease digestion to site E, generating the mature 3' end of 5.8S rRNA (not represented). For simplicity, only the major processing pathway, from 27SA₂ to 5.8S_S and 25S rRNA, is shown; an alternative pathway generates the minor 5.8S_L rRNA. (C) Pre-rRNA processing in *dim1* TS strains. 35S pre-rRNA is cleaved normally at site A₀. 33S pre-rRNA accumulates and is cleaved at site A₃, providing the 27SA₃ that is normally processed to 5.8S_S and 25S and the aberrant 22S pre-rRNA that is not dimethylated and not processed to 18S rRNA.

ing in its absence, or does Dim1p have an additional function in pre-rRNA processing and/or ribosome assembly that is required for pre-rRNA cleavage? The approach that we adopted was to screen a library of conditionally lethal alleles of *DIMI* in order to isolate mutations that uncouple the enzymatic function of Dim1p in methylation from its involvement in pre-rRNA processing.

MATERIALS AND METHODS

Media and plasmids. Standard *S. cerevisiae* growth and handling techniques were used. 5-Fluoro-orotic acid (5-FOA) was used in minimal medium at 1 g/liter according to the recipe of Boeke et al. (4). The transformation procedure was as described by Gietz et al. (13).

Plasmid pDL31.42 (2 μm *URA3 DIM1*) was isolated from a pFL44L (5)-based yeast genomic library as previously described (23). *DIMI* was recovered from pDL31.42 as a 3,282-bp *SmaI/XhoI* fragment and subcloned in pFL36 (*ARS CEN LEU2*) (5) digested with *PvuII/SmaI* to give plasmid pTL6. A *BamHI HIS3* cassette isolated from plasmid YDp-H (3) was filled and subcloned in the filled *XbaI* site of the terminator region of *DIMI* on plasmids pTL6-dim1-2, pTL6-dim1-9 and pTL6-dim1-1 to give constructs pTL39, pTL42, and pTL43, respec-

tively. Plasmid pTL17 is pJV12 (19) in which the *LEU2* marker of pFL36 (5) was subcloned in *SalI*. In pTL29, the *XhoI/SfiI* ribosomal DNA (rDNA) fragment of pGAL::rDNA (16) was subcloned in pTL17. Plasmids pAT3 and pII2.34Δ33 and plasmid pHT4467 are gifts from T. Preiss (European Molecular Biology Laboratory [EMBL]) and H. Tekotte (EMBL), respectively.

Epitope-tagged versions of Dim1p were constructed in plasmid pTL6 by inserting three copies of the human c-Myc epitope at either the amino- or carboxy-terminal end of the protein (generating plasmids pTL18 and pTL25, respectively). For plasmid pTL18, an *NcoI* 3× Myc cassette was inserted at the ATG codon of *DIMI* at the naturally occurring *NcoI* site. For the carboxy fusion, an in-frame *SacI* site was created at the end of the open reading frame (ORF) by site-directed mutagenesis and a *SacI* 3× Myc cassette was inserted. The 3× Myc cassettes were generated by PCR and have been described previously (25). All constructs were checked by sequencing.

Construction of a library of conditional alleles of *DIMI*. The *DIMI* ORF was mutagenized by PCR under the conditions described by Leung et al. (28). The primers used for amplification with pDL31.42 were 5'-TAAAATTATACCATG GGAAAGGCT-3' and 5'-TGATAAGAGAGCTCATGAAAAATG-3'. The PCR product was subcloned as an *NcoI/SacI* fragment in plasmid pTL6, and the library was amplified in *E. coli*. Random sequencing revealed that the mutation rate was approximately 0.46%. The *DIMI* ORF being ~1 kb, this would give an average of 4 to 5 mutations per gene. Transitions, transversions, and insertions were detected.

TABLE 1. Yeast strains used in this study

Strain	Genotype	Reference or comments
D150	a <i>ura3-52 leu2-3,112 ade1-100 his4-519 GAL⁺</i>	BWG1-7A, L. Guarente
YDL302	As D150 with <i>URA3 GAL10::dim1</i>	26
BMA38	a/α <i>ade2-1/ade2-1 can1-100/can1-100 his3Δ200/his3Δ200 leu2-3,112/leu2-3,112 trp1-1/trp1-1 ura3-1/ura3-1</i>	B. Dujon
YDL303	As BMA38 with <i>dim1Δ::HIS3/DIM1</i>	Full <i>DIM1</i> deletion made by PCR in strain BMA38
YDL304A	a <i>ade2-1 can1-100 his3Δ200 leu2-3,112 trp1-1 ura3-1 dim1Δ::HIS3 + pDL31.42 (URA3 2μm DIM1)</i>	Starting strain, segregant from YDL303 containing a <i>URA3 2μm DIM1</i> plasmid
YDL304B	α <i>ade2-1 can1-100 his3Δ200 leu2-3,112 trp1-1 ura3-1 dim1Δ::HIS3 + pDL31.42 (URA3 2μm DIM1)</i>	YDL304A sister strain
YDL209	a <i>ade2-1 can1-100 his3Δ200 leu2-3,112 trp1-1 ura3-1 dim1Δ::HIS3 + pTL6 (LEU2 ARS/CEN DIM1)</i>	YDL304A in which plasmid pDL31.42 has been shuffled with plasmid pTL6
MBS	a <i>ade2 his3 leu2 trp1 ura3 can1 L-0 M-0</i>	P. Sarnow (17)
YDL321	As MBS with <i>dim1-2</i>	<i>dim1-2</i> integrated as a <i>dim1-2/HIS3</i> cassette isolated from pTL39
YDL324	As MBS with <i>dim1-1</i>	<i>dim1-1</i> integrated as a <i>dim1-1/HIS3</i> cassette isolated from pTL43
CH1462	α <i>ade2 ade3 his3 leu2 ura3 can1</i>	21
YDL314	As CH1462 with <i>dim1-9</i>	<i>dim1-9</i> integrated as a <i>dim1-9/HIS3</i> cassette isolated from pTL42
YDL315	As CH1462 with <i>dim1-1</i>	<i>dim1-1</i> integrated as a <i>dim1-1/HIS3</i> cassette isolated from pTL43
YDL300	a/α <i>ura3-52/ura3-52 lys2-801^{amber}/lys2-801^{amber} ade2-101^{ochre}/ade2-101^{ochre} trp1Δ1/trp1Δ1 his3Δ200/his3Δ200 leu2Δ1/leu2Δ1 dim1Δ::URA3/DIM1</i>	23
YDL100A	a ; as YDL300 + pTL6 (<i>LEU2 ARS/CEN DIM1</i>)	<i>dim1Δ::URA3</i> segregant from YDL300
YDL101A	a ; as YDL300 + pTL18 (<i>LEU2 ARS/CEN 3× Myc-Dim1p</i>)	<i>dim1Δ::URA3</i> segregant from YDL300
YDL102A	a ; as YDL300 + pTL25 (<i>LEU2 ARS/CEN Dim1p-3× Myc</i>)	<i>dim1Δ::URA3</i> segregant from YDL300

Isolation of conditional alleles of *DIM1*. The library was screened according to the cotransformation and plasmid-shuffling procedure described by Sikorski and Boeke (37). The shuffling strain was constructed as follows. A complete deletion of *DIM1* was created in the diploid strain BMA38 by a one-step PCR strategy (2, 25). The oligonucleotides used for amplification with pRS313 (38) were 5'-GGTTATAAGATCGATAAATTAGGAACAGTGCTATTATACAGTCTC TTGGCTCTCTAG-3' and 5'-TTTTTCTTATCTTAGGTAATAGTATA CAAGCACTTACATAATCGTTCAGAAATGACACG-3'. The resulting strain, YDL303, was transformed with plasmid pDL31.42 and sporulated. Haploids containing the deletion rescued by pDL31.42 were identified (strains YDL304A and YDL304B [Table 1]).

The library was transformed into strain YDL304A and plated on minimal medium lacking histidine and leucine (SD -His -Leu) at 30°C. One thousand eight hundred transformants were patched onto SD -His -Leu at 25°C. The resulting master plates were replica plated on 5-FOA at 25°C. After two rounds of selection on 5-FOA, the 5-FOA-resistant clones were tested for growth at various temperatures (18, 25, 30, and 37°C). Approximately 11% of the clones remained 5-FOA sensitive, probably due to nonconditional inactivation. Thirty-two thermosensitive (TS) clones were isolated, some of which were also slightly cryosensitive (CS). No clones showing only a CS phenotype were recovered. Ten conditional alleles, *dim1-1* to *dim1-10*, were selected to be further characterized.

RNA extraction, Northern hybridization, and primer extension. RNA extraction, Northern hybridization, and primer extension were as described previously (26). Oligonucleotides a, b, d, e, f, and g were named d, e, f, g, l, and m, respectively, in reference 26. Oligonucleotides c and h are TCTCTTCAAAGGGTCCG and GCACCGAAGGTACCAG, respectively. RNA analysis of *dim1-1* to *dim1-10* alleles were performed in strain YDL304A cured of plasmid pDL31.42 and bearing the corresponding pTL6-*dim1*-t.s. plasmid. The reference wild-type strain used was YDL209 (Table 1).

For the experiment presented in Fig. 7, *dim1-1 HIS3* and *dim1-9 HIS3* integrative cassettes were recovered from plasmids pTL43 and pTL42 by *EcoRI/XhoI* digestion and transformed in strain CH1462 (21).

To allow *HIS3* selection in strain CH1462 (*ade2 ade3* [Table 1]), an *ADE3* gene was expressed from plasmid pHT4467 (*ADE3 URA3*). The resulting strains, YDL314 (*dim1-9*) and YDL315 (*dim1-1*), were transformed with either pTL29 or pGAL::rDNA and grown at permissive temperature in glucose minimal medium lacking leucine or uracil, respectively, before being transferred to 37°C for 8 h. For *GAL* *Dim1p* depletion, strain YDL302 transformed with pTL29 was grown at 30°C in galactose minimal medium lacking uracil and leucine. Cells were harvested by centrifugation, washed, and resuspended in glucose minimal medium lacking uracil and leucine. During growth, cells were diluted with prewarmed medium and constantly maintained in early exponential phase.

In vitro translation. Cytoplasmic S30 extracts of strains YDL321, YDL324, and the isogenic wild-type control strain MBS were prepared as described previously (17, 40). Strains YDL321 and YDL324 were constructed as follows: integrative cassettes (*dim1* [TS] *HIS3*) were recovered from plasmids pTL39 (*dim1-2*) and pTL43 (*dim1-1*) by *EcoRI/XhoI* digestion and transformed into strain MBS. Poly(A) substrates were prepared as described previously (40) with constructs pAT3 (preprolactin) and p112.34Δ33 (chloramphenicol acetyltransferase [CAT]). Both substrates were capped with the ^{m7}GpppG analog (New England Biolabs). Translation reactions (40) were for 60 min at 24°C with 0, 2, and 10 ng of preprolactin mRNA or 0, 20, and 100 ng of CAT mRNA. Translation products were analyzed by sodium dodecyl sulfate-polyacrylamide gel electrophoresis.

Antibiotic sensitivity assay. As a preliminary experiment, equal amount of cells were plated on yeast extract-peptone-dextrose (YPD), and 5 and 10 μl of various antibiotic stock solutions were spotted on 5-mm sterile filter disks. Stock solutions were as follow: paromomycin (50 mg/ml), G418 (3 mg/ml), hygromycin B (3 mg/ml), cycloheximide (1 mg/ml), and neomycin B (20 mg/ml). Three prokaryote-specific antibiotics, streptomycin (100 mg/ml), erythromycin (50 mg/ml), and kasugamycin (100 mg/ml), were also tested. All antibiotics were purchased from Sigma. *dim1* TS strains were found to be hypersensitive to paromomycin and neomycin B. All strains were also sensitive to cycloheximide but to the same extent as the wild-type isogenic control. The other antibiotics tested showed no effect. For the experiment presented in Fig. 10, paromomycin and neomycin B were used at 0, 250, 375, and 500 μg/μl and 0, 100, 400 and 800 μg/ml, respectively. Plates were incubated at the permissive temperature of 23°C.

Indirect immunofluorescence. Indirect immunofluorescence was performed on strains expressing 3× Myc-*Dim1p* or *Dim1p-3× Myc* fusion proteins (strains YDL101A and YDL102A, respectively [Table 1]). Strain YDL100A, expressing wild-type *Dim1p*, was used as a negative control. Cells were grown in YPD to an optical density at 600 nm of 0.5 to 1.0, fixed in 37% formaldehyde for 1 h, and permeabilized with Zymolyase in sorbitol buffer (1.2 M sorbitol, 20 mM KPI [pH 7.4]). Cells were incubated for 1 h with either anti-Nop1p antibody (monoclonal antibody 66, dilution 1/50; kindly provided by John Aris [1]) or anti-Myc antibody (9E10, dilution 1/50; BAbCO). Both antibodies are mouse monoclonal antibodies; therefore, cells were decorated independently. Detection with Texas red was for 1 h at a dilution of 1/50 (Texas red-conjugated affinipure F(ab')₂ goat anti-mouse immunoglobulin G [H+L]; Dianova). DNA was stained with 4',6-diamidino-2-phenylindole (DAPI). Strains YDL100A, YDL101A, and YDL102A are haploids containing the *dim1Δ::URA3* deletion of strain YDL300 (Table 1) rescued by the wild-type or epitope-tagged proteins expressed from plasmids pTL6 (*Dim1p*), pTL18 (3× Myc-*Dim1p*), and pTL25 (*Dim1p-3× Myc*), respectively. The growth rates of strains YDL100A, YDL101A, and YDL102A are identical, showing the fusion proteins to be fully functional.




FIG. 2. Conditional growth phenotypes of *dim1* TS strains. Dilutions (1× to 10²×) of strains *dim1-1* to *dim1-10*, along with the wild-type isogenic *DIM1* control (strain YDL209) (W.T.), were spotted on minimal plates at 18, 25, 30, and 37°C and incubated for 3 days. Most of the strains are very sensitive to elevated temperatures; some are also slightly CS (*dim1-1* and *dim1-7*).

RESULTS

Isolation of conditional alleles of *DIM1*. A library of conditional alleles of *DIM1* was created by mutagenic PCR and screened by plasmid exchange (37). A diploid strain containing a fully deleted *dim1-Δ* allele was transformed with a multicopy *URA3* plasmid bearing a wild-type copy of *DIM1*. Haploid *dim1-Δ* progeny, rescued by the wild-type plasmid, were used as starting strains (strains YDL304A and YDL304B [Table 1]). The ORF of *DIM1* was amplified under mutagenic PCR conditions and subcloned into an *ARS/CEN LEU2* plasmid (see Materials and Methods). The resulting library was transformed into strain YDL304A. Transformants were replica plated on 5-FOA, which selects for the loss of the wild-type *URA3* plasmid. 5-FOA-resistant clones, which have lost the wild-type *DIM1* plasmid, were screened for growth defects at various temperatures. Among 1,600 5-FOA-resistant clones, 33 showed a conditional phenotype for growth. Most of them were TS, with some being also slightly CS (Fig. 2). No strain showed only a CS phenotype. Ten alleles, *dim1-1* to *dim1-10*, were selected to be further characterized.

The coding sequences of *dim1-1* to *dim1-10* were fully sequenced; all contained from one to six substitution mutations (Fig. 3 and Table 2). It may be significant that mutations at four positions were selected twice (Table 2). Position 218 was mutated in both *dim1-1* and *dim1-4*, and an identical substitu-

tion (S239P) is present in both *dim1-6* and *dim1-10*. Notably, *dim1-4* and *dim1-5* have two identical mutations at identical positions (172 and 197), *dim1-4* having an additional mutation at a position which is also altered in *dim1-1* (position 218). Although mutations were spread over the whole length of the protein, we noticed a degree of clustering centered around positions 170 and 220; in the latter case, these fall in a region which is not conserved in *E. coli* KsgAp.

None of the mutations fell into the predicted binding site for the cofactor *S*-adenosylmethionine (residues 60 to 76) (9, 18). The NXPY residues (positions 128 to 131) are conserved in all known SSU rRNA dimethylases, including human Dim1p (23a). These match the consensus sequence (N/D/S)PP(Y/F), which has been shown to be crucial for catalysis in the case of the *N*-6-adenine methylase *EcoKI* (48). Two mutations (I126V in *dim1-2* and I133V in *dim1-3*) directly flank the NXPY residues, suggesting that they might directly affect catalysis.

***dim1-2* uncouples the two functions of Dim1p.** RNA was extracted from the 10 *dim1* TS strains and the isogenic wild-type strain (strain YDL209 [Table 1]) at the permissive temperature (23°C) and at 2, 8, and 23 h after transfer to 37°C.

Northern hybridization revealed that at the nonpermissive temperature, all *dim1* TS strains are inhibited in cleavage at sites A₁ and A₂, resulting in depletion of the 20S pre-rRNA and 18S rRNA (Fig. 1). These data are presented for five representative alleles (*dim1-1*, *dim1-2*, *dim1-6*, *dim1-7*, and




FIG. 3. Mapping of *dim1* TS mutations. Schematic representation of Dim1p (318 residues) drawn to scale. The *S*-adenosylmethionine binding domain (SAM), the putative catalytic residues (NxPY), and the 32 point mutations (asterisks) identified in the 10 *dim1* TS alleles are represented.

TABLE 2. Nature and positions of mutations in *dim1* TS alleles

Allele	Mutation(s)
<i>dim1-1</i>	E21G; N213D; V218A; D219G; D224G
<i>dim1-2</i>	S25G; I126V; F159S; R174S; Y220N; M273T
<i>dim1-3</i>	I133V
<i>dim1-4</i>	Y172D; R197S; V218E
<i>dim1-5</i>	Y172D; R197S
<i>dim1-6</i>	N80D; N178D; S239P; M265V
<i>dim1-7</i>	C173R; H274R; V313A
<i>dim1-8</i>	N31S; A92E; K278R; L304P
<i>dim1-9</i>	W223R
<i>dim1-10</i>	N41S; I238V; S239P




FIG. 4. Pre-rRNA processing in *dim1* TS strains. (A) Probe against the 5' region of ITS1 (oligonucleotide e [Fig. 1A]). (B) Probes against mature 18S and 25S rRNA (oligonucleotides a and f [Fig. 1A]). RNA was extracted from the *DIM1* (strain YDL209) (W.T.) and *dim1* TS strains following growth at 23°C (0-h lanes) and at intervals following transfer to 37°C (2-, 8-, and 23-h lanes) and separated on 1.2% agarose-formaldehyde gels. The 22S pre-rRNA extends from site A₀ to site A₃ and results from the inhibition of cleavages at sites A₁ and A₂ in the *dim1* TS strains. At the late time point of transfer to 37°C (23 h), the normally minor and barely detectable 23S pre-rRNA that extends from the 5' end of the primary transcript to site A₃ accumulates to the same levels in the wild-type and *dim1* TS strains. In this experiment, all samples, including the wild-type control, showed some accumulation of 35S pre-rRNA and 23S RNA after 23 h at 37°C; this was not observed in other experiments.

dim1-9) in Fig. 4. In the *dim1* mutants, there is seen an aberrant processing pathway in which the 35S pre-rRNA is processed normally at site A₀ but the resulting 33S pre-rRNA is not processed at sites A₁ and A₂. Instead, cleavage at site A₃ in internal transcribed spacer 1 (ITS1) generates the 22S RNA. Concomitant with the appearance of the 22S species, the 20S and 27SA₂ pre-rRNAs are lost (Fig. 4A and data not shown). This is followed by depletion of the mature 18S rRNA (Fig. 4B), indicating that the aberrant 22S RNA cannot be processed to 18S rRNA. No alterations in the level of 27SB pre-rRNA (data not shown) or mature 25S rRNA (Fig. 4B) were observed, indicating that subsequent processing of the 3' region of the pre-rRNA was unaffected.

The pre-rRNA processing defects observed in the *dim1* TS strains closely resemble those seen on depletion of Dim1p in *GAL::dim1* strains (26). The timing of the appearance of the processing defects is strikingly different in the various *dim1* TS

strains. In the *dim1-1*, *dim1-6*, and *dim1-9* strains, processing is strongly inhibited 2 h after transfer to 37°C. In contrast, the 22S RNA is only weakly detected in the *dim1-2* strain 2 h after transfer to 37°C, and the inhibition of processing is much stronger at later times. The 22S RNA is detected in the *dim1-7* strain after 8 h. This may reflect a distinction between mutants in which Dim1p is rapidly inactivated by transfer to 37°C compared to mutations which do not allow synthesis of new, active Dim1p.

Although all strains accumulate 22S pre-rRNA (Fig. 4 and data not shown), there are marked differences in 18S rRNA depletion. *dim1-1*, *dim1-4*, *dim1-6*, *dim1-8*, and *dim1-10* strains all showed strong 18S rRNA depletion, while *dim1-5* and *dim1-9* strains showed only mild depletion and *dim1-2*, *dim1-3*, and *dim1-7* strains were only slightly depleted (Fig. 4 and data not shown). No clear correlation could be established between the locations of the point mutations and the phenotypes, but it




FIG. 5. Overall level of dimethylation in *dim1* TS strains. RNA was extracted from the *DIM1* (strain YDL209) (W.T.) and *dim1* TS strains following growth at 23°C (0-h lanes) and at intervals following transfer to 37°C (2-, 8-, and 23-h lanes) and analyzed by primer extension with oligonucleotide e (Fig. 1A). The positions of primer extension stops due to the presence of the modifications are indicated. A DNA sequence made with the same primer is shown as a size marker.

is notable that *dim1-4*, which contains only one additional mutation (at position 218) compared to *dim1-5*, shows stronger 18S rRNA depletion. A substitution at the same position is present in *dim1-1*, which shows a similar phenotype. *dim1-6* and *dim1-10* strains are also severely affected in 18S rRNA synthesis and share an identical mutation at position 239. No attempt was made to further separate the mutations.

The level of pre-rRNA dimethylation was assessed by primer extension with an oligonucleotide complementary to a sequence in ITS1 located 218 nucleotides downstream of the site of modification (oligonucleotide e [Fig. 1A]). Following transfer to 37°C, the primer extension stop corresponding to dimethylation is strongly reduced in all *dim1* TS strains (Fig. 5). Northern analysis data (Fig. 4) indicate that the overall levels of pre-rRNA species that contain the site of dimethylation are not strongly altered in most *dim1* TS mutants. As with the pre-rRNA processing defects, dimethylation is inhibited to various extents and with different kinetics in the different *dim1* TS alleles (Fig. 5).

The accumulation of nondimethylated 18S rRNA can be monitored by primer extension with an oligonucleotide complementary to the very 3' end of the mature rRNA (oligonucleotide d [Fig. 1A]), since the m²A modification blocks reverse transcription (see the legend to Fig. 6). The level of nondimethylated rRNA and pre-rRNA is indicated by the stop at position c, which represents reverse transcriptase molecules that were able to read through the unmodified site of dimethylation (26). The very faint band visible at this position in the wild-type strain (Fig. 6, lanes 1 to 4) represents primer extension on the unmodified pre-rRNA, which is very much less abundant than the mature rRNA (see reference 26). The *GAL::dim1* strain (Fig. 6, lane 27) does not accumulate nondimethylated 18S rRNA; this is as previously reported (26). In contrast, several *dim1* TS strains accumulate unmodified 18S rRNA. Strong accumulation is seen in the *dim1-2* and *dim1-7* strains (Fig. 6, lanes 5 to 12), even at the permissive temperature, and some accumulation is seen in the *dim1-9* strain at 37°C (Fig. 6, lanes 13 to 16). Little nondimethylated 18S was detected in the *dim1-1* or *dim1-4* strains (Fig. 6, lanes 17 to 24).

Thus, at the nonpermissive temperature, all of the *dim1* TS strains analyzed are impaired both in cleavage of pre-rRNA at sites A₁ and A₂ and in dimethylation of pre-rRNA, although the kinetics and severity of these phenotypes vary between mutants. Some of the *dim1* TS strains accumulate nondimethylated

18S rRNA, showing that the pre-rRNA methylation defect can be uncoupled from the pre-rRNA processing defect. This is particularly true in the *dim1-2* strain, which shows no clear pre-rRNA processing defect at the permissive temperature but has very low levels of pre-rRNA modification.

It is notable that no *dim1* TS strain defective only in pre-rRNA processing was isolated (Fig. 4 and 5). Dim1p must bind its substrate to modify the two adenosines, and this interaction with the pre-rRNA may be both necessary and sufficient to fulfill the requirement for Dim1p in pre-rRNA processing. Since dimethylation is dispensable in vivo (see below), the TS screen would not have isolated strains defective only in methylation without a pre-rRNA processing defect.

Pre-rRNAs transcribed from the *PGK* promoter do not require Dim1p for processing. The studies on pre-rRNA processing described above analyzed pre-rRNA transcribed from the chromosomal rDNA by RNA Pol I. Unexpectedly, the processing of pre-rRNA transcribed by the constitutive RNA Pol II *PGK* promoter is much less sensitive to the *dim1-1* and *dim1-9* mutations than is processing of the Pol I-transcribed pre-rRNA (shown for *dim1-1* in Fig. 7). The Pol II-driven rDNA repeat contains neutral tags within the mature rRNA sequences; these allow the use of hybridization probes that are specific for either the chromosomal or Pol II-transcribed rRNAs. Transfer of the *dim1-1* strain to 37°C for 20 h resulted in underaccumulation of the 18S rRNA synthesized from pre-rRNA transcribed from chromosomal rDNA (Fig. 7, lower panel [compare lane 3 with 4 and lane 5 with 6]) as described above. In contrast, the level of 18S rRNA synthesized from pre-rRNA transcribed from the *PGK* promoter carried on plasmid pTL29 (see Materials and Methods) was reduced to a lesser extent (Fig. 7, upper panel [compare lanes 5 and 6]). Accumulation of 18S rRNA transcribed from a different RNA Pol II promoter, *GAL10* (Fig. 7, upper panel [compare lanes 3 and 4]), was reduced to an extent similar to the rRNA transcribed by Pol I. The same effects were seen in the *dim1-9* strain (data not shown); other *dim1* alleles were not tested.

Similar effects were observed in the *GAL::dim1* strain (YDL302 [Table 1]) following depletion of Dim1p. Following transfer to glucose, the level of 18S rRNA synthesized by Pol I from chromosomal rDNA (Fig. 8B) is strongly depleted. In contrast, little depletion of the 18S rRNA synthesized from the pPGK::rDNA construct is observed (Fig. 8A). The effects of




FIG. 6. Levels of nondimethylated 18S rRNA in *dim1* TS strains. (A) RNA was extracted from the *DIM1* (strain YDL209) (W.T.) and *dim1* TS strains following growth at 23°C (0-h lanes) and at intervals following transfer to 37°C (2-, 8-, and 23-h lanes) and analyzed by primer extension with oligonucleotide d, which is complementary to the very 3' end of 18S rRNA (Fig. 1A). The reactions were performed with dideoxyadenosine nucleotides in place of deoxyadenosine. The site of priming is 3 nucleotides 3' to A₁₇₈₀, and no A residues are incorporated before the site of modification (see panel B). Dimethylation of A₁₇₇₉ A₁₇₈₀ blocks primer extension. Extensions carried out on nondimethylated rRNA extend through the A₁₇₇₉ A₁₇₈₀ site but are blocked 2 nucleotides 5' to A₁₇₇₉ (position U₁₇₇₇). The positions of primer extension stops due to the presence of the modifications are indicated (a, b and c). A DNA sequence made with the same primer is shown as a size marker. Lanes 25 to 27, control lanes (lane 25, no RNA [P denotes primer alone]; lane 26, same as lane 4; lane 27, RNA extracted from the *GAL::dim1* strain (strain YDL302) following transfer to glucose for 60 h). (B) Schematic representation of the 3' end of 18S rRNA. Upper line, rRNA strand (the thick line represents the last 16 nucleotides). Lower line, complementary cDNA strand (the thick broken line represents oligonucleotide d). The three potential extended products are represented by thin lines (a, b, and c). The positions of the primer extension stops due to the presence of the modifications are indicated as a and b; the position of the primer extension stop due to read-through of the dimethylation site is indicated as c.

genetic depletion of two other components required for pre-rRNA processing at sites A₁ and A₂ were also tested. On depletion of snR30 (30) or Rrp5p (47), the processing of the *PGK*-driven pre-rRNAs was inhibited to the same extent as processing of the Pol I-transcribed pre-rRNA (33a).

The processing of the pre-rRNAs transcribed from the *PGK*-driven rDNA cannot readily be assessed in these strains, since Pol I transcription is much stronger than *PGK*-driven transcription. However, we interpret these data as showing that the cleavage of sites A₁ and A₂ is specifically resistant to mutations in *DIM1* or depletion of Dim1p in pre-rRNA molecules that have been transcribed from the *PGK* promoter. The mechanism that relieves the Dim1p dependence may be related to the packaging of the pre-rRNA transcripts with different sets of proteins. Whatever the mechanism, this observation demonstrates that Dim1p is not directly required for pre-rRNA processing. We interpret this as strong evidence for the existence of a system that represses pre-rRNA processing in the absence of Dim1p.

m⁶A₁₇₇₉m⁶A₁₇₈₀ is not essential in vivo but is required for translation in vitro. At the permissive temperature, the *dim1-2* strain showed no pre-rRNA processing defect but predominantly contained nondimethylated 18S rRNA. The *dim1-2* strain showed no clear growth inhibition compared to the oth-

erwise isogenic *DIM1* strain in liquid or solid minimal medium or complete medium at 23°C (data not shown). This demonstrates that 18S rRNA dimethylation is dispensable for translation in vivo.

In order to test whether dimethylation affected ribosome function at a more subtle level, extracts prepared from wild-type and *dim1-2* strains grown at the permissive temperature were tested for the ability to translate exogenous mRNA in vitro. As a control, extracts from a *dim1-1* strain that was not impaired in rRNA methylation at the permissive temperature were also tested.

The *dim1* alleles were integrated into strain MBS (see Materials and Methods and Table 1), which is particularly suitable for such analysis (17). Standardized amounts of extracts prepared from strains YDL321 (*dim1-2*), YDL324 (*dim1-1*), and the isogenic wild-type control strain MBS were incubated with various amounts of either CAT mRNA (Fig. 9B and D) or preprolactin mRNA (Fig. 9A). Figures 9A and B present the analyses of two independently isolated *dim1-2* strains (YDL321-1 and YDL321-2). Translation products were easily detected at the expected lengths when wild-type extracts were incubated with either of the two mRNAs (Fig. 9A, B, and D, lanes 2 and 3). No product was detected for either mRNA substrate when the *dim1-2* extracts were used (Fig. 9A and B, lanes 4 to 9, and




FIG. 7. 18S and 25S accumulation in *dim1* TS strains expressing rDNA from different promoters. Lane 1, RNA extracted from a *DIM1* wild-type strain; lane 2, RNA extracted from a *DIM1* wild-type strain also expressing pre-rRNA from a *GAL* promoter; lanes 3 and 4, RNA extracted from a *dim1-1* strain also expressing pre-rRNA from a *GAL* promoter; lanes 5 and 6, RNA extracted from a *dim1-1* strain also expressing pre-rRNA from a *PGK* promoter. The same Northern filter was hybridized with probes complementary to the 25S and 18S rRNAs. The probes used are specific either for the tagged rRNAs synthesized from the RNA Pol II promoter or for the nontagged rRNAs transcribed from the chromosomal rDNA (RNA Pol I).

Fig. 9D, lanes 4 to 6). Extracts from the *dim1-1* strain were less competent for translation than was the wild-type control (Fig. 9D, lanes 8 and 9) but consistently exhibited substantially greater translation than did the *dim1-2* strain. The reduced translation may be related to the mild processing defect present in the *dim1-1* strain, which accumulated low levels of the 22S pre-rRNA even at the permissive temperature (Fig. 4A, lane 9). Figure 9C shows the analysis of the level of 18S rRNA dimethylation in the cell extracts used for Fig. 9A and B. We concluded that in the *dim1-2* strains, the 40S subunits contain nondimethylated 18S rRNA and are not competent for translation in vitro.

***dim1* TS strains are hypersensitive to aminoglycoside antibiotics.** To identify more subtle effects on translation in vivo, representative *dim1* TS strains were tested for their sensitivities towards a range of antibiotics at the permissive temperature (see Materials and Methods). Strains carrying *dim1-1*,




FIG. 9. In vitro translation analysis of *dim1* TS strains. Cytoplasmic S30 extracts of *dim1-2* strains (YDL321-1 and YDL321-2), a *dim1-1* strain (YDL324), and the wild-type (W.T.) isogenic control (strain MBS) were prepared following growth at 23°C. Standardized amount of extracts were incubated with 0, 2, and 10 ng of preprolactin mRNA (A) or 0, 20, and 100 ng of CAT mRNA (B and D). YDL321-1 and YDL321-2 are two independently isolated integrants of the *dim1-2* allele in the MBS strain (see Materials and Methods and Table 1). The *dim1-2* strain used in panel D is YDL321-2. Translation products were analyzed by sodium dodecyl sulfate-polyacrylamide gel electrophoresis and are indicated at their expected lengths (11 and 25 kDa). (C) 18S rRNA dimethylation in cell extracts, analyzed as described in the legend to Fig. 6. RNA was extracted from strains YDL321-1, YDL321-2, and the wild-type strain (strain MBS) following growth at 23°C. Lanes 1 to 3, control lanes (lane 1, no RNA [P denotes primer alone]; lane 2, RNA extracted from strain MBS following transfer to 37°C for 23 h; lane 3, RNA extracted from the *GAL::dim1* strain [strain YDL302] following transfer to glucose for 60 h).




FIG. 8. pPGK::rDNA transcripts are insensitive to Dim1p depletion. (A) Probes specific to mature 25S and 18S rRNA produced from the pPGK::rDNA construct (oligonucleotides b and g [Fig. 1A]). (B) Probes specific to the mature 25S and 18S rRNA produced from the chromosomal rDNA units (oligonucleotides c and h [Fig. 1A]). RNA was extracted from the *GAL::dim1* strain transformed with the pPGK::rDNA construct following growth in galactose (0-h lanes) and at intervals following transfer to glucose (2- to 60-h lanes) and separated on a 1.2% agarose-formaldehyde gel.

dim1-2, and *dim1-9* were hypersensitive to paromomycin and neomycin B to approximately equal extents (Fig. 10 and data not shown). These two antibiotics belong to the aminoglycoside family and are known to induce misreading and suppression of nonsense mutations in yeast (33, 39). This observation suggested that each of the *dim1* TS strains analyzed had an additional assembly defect which resulted in antibiotic sensitivity. This appears to be unrelated to the degree of rRNA dimethylation.




FIG. 10. Antibiotic sensitivities of *dim1* TS strains. Dilutions (1× and 10×) of *dim1-1* and *dim1-2* strains, along with the isogenic wild-type (W.T.) *DIM1* control strain, were spotted on complete medium supplemented with paromomycin and neomycin B at the concentrations indicated. Plates were incubated at 23°C.

Dim1p is localized to the nucleus with nucleolar enrichment. To determine the subcellular location of Dim1p, the protein was tagged with three copies of the human c-Myc epitope at either the amino- or carboxy-terminal end (see Materials and Methods). Both fusion proteins were expressed in a *dim1-Δ* deleted background and were shown to be fully functional (strains YDL101A and YDL102A [Table 1]).

The fusion proteins were localized by indirect immunofluorescence. Figure 11D presents the results of the carboxy-terminal fusion (strain YDL102A). The amino-terminal fusion gave an identical signal (data not shown). Comparison to DAPI staining of the DNA (Fig. 11C) and the localization of the nucleolar protein Nop1p (Fig. 11B) reveals that the Dim1p fusion protein is localized to the nucleus with enrichment in the nucleolus, which forms a cap-like structure slightly displaced from the DAPI-stained region. Little cytoplasmic staining was detected.

DISCUSSION

We report here that mutations in *DIM1* can uncouple the requirements for Dim1p in pre-rRNA modification and processing. All of the *dim1* alleles that were isolated from a temperature-sensitive library were found to block pre-rRNA processing at the nonpermissive temperature (37°C), although the kinetics of inhibition varied. In addition, the *dim1-2* allele blocks rRNA dimethylation at the permissive temperature (23°C) but shows no pre-rRNA processing defect at this temperature. This resulted in the accumulation of high levels of nondimethylated 18S rRNA. The *dim1-2* strains had no detectable growth defect at 23°C, showing the dimethylation to be dispensable for ribosome function in vivo.

Pre-rRNA molecules in which the two adenosine residues are replaced by guanosines that cannot be modified are processed normally but do not support growth (26). Additionally, these substitutions give rise to a reduced level of 18S rRNA at low temperatures, possibly reflecting an assembly defect, which is often associated with cold sensitivity. From the *dim1-2* analysis, it is clear that the lethality of the $G_{1779}G_{1780}$ *cis* mutation and the cold-sensitive processing phenotype are due to the substitution of the adenosines at these two universally con-

served positions. In *E. coli*, both dimethylation and the twin adenosines were shown to be dispensable for function and assembly in vitro (10, 22), but their importance in vivo has not been assessed.

Strikingly, extracts prepared from *dim1-2* strains grown at 23°C lacked detectable activity for translation in vitro. Extracts prepared from *dim1-1* strains, which have normal levels of rRNA dimethylation, were competent for in vitro translation, although with lower activity than in the wild-type extract. While we cannot exclude the possibility that ribosomes synthesized in the *dim1-2* mutant have some additional defect, it seems likely that dimethylation is required for in vitro translation in yeast extracts. We speculate that the dimethylation fine-tunes the function of the 40S ribosomal subunit in vivo but is essential under the suboptimal in vitro conditions.

Other rRNA modifications, 2'-O methylation and pseudouridine formation, are directed by guide snoRNAs (8, 12, 20, 31, 32). In all cases tested, the guide functions of the snoRNAs were completely dispensable for growth, indicating that these modifications are, like Dim1p dimethylation, dispensable for ribosome function in vivo. The requirements for the 2'-O-methyl and pseudouridine modifications, however, have not been tested in vitro.

Prokaryotic *ksgA* strains lack rRNA dimethylase activity and are resistant to the aminoglycoside antibiotic kasugamycin (14, 15, 44). Hypersensitivity toward the aminoglycoside antibiotics paromomycin and neomycin B was observed in *dim1* TS strains. This could not be correlated, however, with dimethylation; *dim1-1* and *dim1-9* strains, which do not accumulate nondimethylated SSU rRNA, are as sensitive to aminoglycosides as are *dim1-2* strains. We speculate that the *dim1* TS strains analyzed bear an additional assembly defect responsible for antibiotic sensitivity. Hypersensitivity to paromomycin and neomycin B has previously been reported in yeast strains carrying mutations in *NSR1* and *RRP1*, which are also required for normal pre-rRNA processing (11, 27). Neither of these proteins were reported to be involved in rRNA modification, and in both cases an assembly defect is likely to underlie the antibiotic sensitivity observed.

Model for a regulatory mechanism in ribosome synthesis. We reported previously that cells depleted of Dim1p are inhibited in cleavage of the 33S pre-rRNA at sites A_1 and A_2 (26), and this was also the case in all of the *dim1* TS strains at the nonpermissive temperature. However, dimethylation was found to occur on the 20S pre-rRNA, which is the product of cleavage at sites A_1 and A_2 (Fig. 1), consistent with early reports that the formation of $m_2^oAm_2^oA$ is a late event in ribosome synthesis (6, 36). Since the cleavages at sites A_1 and A_2




FIG. 11. Subcellular localization of Dim1p. Indirect immunofluorescence with strain YDL102A (Dim1p-3× Myc). Cells were incubated with either anti-Nop1p (α -Nop1p) or anti-Myc (α -Myc) antibody followed by Texas red (TR) and DAPI staining. Both primary antibodies are mouse monoclonal antibodies; therefore, cells were labelled independently.

occur before dimethylation, they cannot be directly dependent on the presence of the modification. This conclusion was supported by the observation that pre-rRNAs containing the G₁₇₇₉G₁₇₈₀ mutations, which cannot be dimethylated, can be processed at sites A₁ and A₂. We concluded that the Dim1p protein, rather than the modification itself, was required for pre-rRNA processing (26).

Our interpretation was that a quality control system probably inhibited the processing of the pre-rRNA in the absence of Dim1p (26). However, the data left open the alternative possibility that Dim1p had an additional function and was directly required for pre-rRNA processing. During the course of this work, this question was unexpectedly resolved. When transcription of an rDNA unit is driven by an RNA Pol II *PGK* promoter, the pre-rRNAs produced are largely insensitive to *dim1-1* and *dim1-9* mutations that strongly inhibit processing of pre-rRNAs transcribed from the chromosomal rDNA driven by RNA Pol I. This phenomenon is not allele specific: the *PGK*-transcribed pre-rRNAs are also resistant to genetic depletion of Dim1p. Moreover, the otherwise identical pre-rRNA transcribed from a *GAL* promoter was not resistant to the *dim1-1* or *dim1-9* mutations. These observations demonstrate that Dim1p does not have a direct role in pre-rRNA processing.

We speculate that a repression system blocks pre-rRNA processing in the absence of the binding of Dim1p to pre-rRNA. According to this model, Dim1p normally binds to pre-rRNA in the nucleolus at an early stage in ribosome synthesis, and a component of the processing machinery senses this interaction. This could occur through direct interaction with Dim1p or through an interaction with preribosomal particles (a conformational change could be monitored). Cleavages at sites A₁ and A₂ occur in the nucleolus; consistent with this, Dim1p was shown to localize mostly to this cellular compartment. If Dim1p has bound to pre-rRNA, processing at sites A₁ and A₂ proceeds; otherwise, processing is blocked. In mutant strains that lack Dim1p, this leads to the synthesis of a dead-end intermediate, the 22S pre-rRNA, and prevents synthesis of the 18S rRNA. In wild-type strains, pre-rRNA processing is presumably only delayed until Dim1p finds and binds to its target, preventing the formation of mature but nonmodified 18S rRNA. This is desirable since, as shown here, the unmodified ribosomal subunits are impaired in function.

The pre-rRNAs that are transcribed from the *PGK* promoter are very likely to be associated with a different set of hnRNP proteins than are the Pol I transcripts. We speculate that one of these occupies the Dim1p binding site and is detected by the pre-rRNA processing machinery as Dim1p, thus alleviating the need for authentic Dim1p.

The m⁶Am⁶A modification is highly conserved and is present in both bacteria and eukaryotes. In *ksgA* mutant strains of *E. coli* that lack dimethylation, growth is mildly impaired and the nondimethylated ribosomes show defects in translation in vitro (reviewed in reference 45). Expression of Dim1p in *E. coli* restores dimethylation, and *E. coli* KsgAp is highly homologous to Dim1p. However, while bacteria lacking the KsgAp methyltransferase synthesize the unmodified rRNA, eukaryotes have evolved a regulatory system to prevent this. We anticipate that many such quality control mechanisms act to coordinate the numerous steps of eukaryotic pre-rRNA processing, rRNA modification, and ribosome assembly.

ACKNOWLEDGMENTS

We thank E. Petfalski for analysis of the *GAL::RRP5* and *GAL::snr30* strains, Jaap Venema for supplying the *GAL::RRP5* strain, B. Séraphin and M. Hentze (EMBL) for fruitful discussions, G. Berben (Station de Chimie, Gembloux, Belgium) for making the YDp plas-

mids available, H. Tekotte (EMBL) for supplying plasmid pHT4467, and the EMBL sequencing service.

This work was partially supported by the Wellcome Trust. During the course of this work, D. L. J. Lafontaine was the recipient of an EMBO long-term fellowship.

REFERENCES

1. Aris, J. P., and G. Blobel. 1988. Identification and characterization of a yeast nucleolar protein that is similar to a rat liver nucleolar protein. *J. Cell Biol.* **107**:17–31.
2. Baudin, A., O. Ozier-Kalogeropoulos, A. Denouel, F. Lacroute, and C. Cullin. 1993. A simple and efficient method for direct gene deletion in *Saccharomyces cerevisiae*. *Nucleic Acids Res.* **21**:3329–3330.
3. Berben, G., J. Dumont, V. Gilliquet, P.-A. Bolle, and F. Hilger. 1991. The YDp plasmids: a uniform set of vectors bearing versatile gene disruption cassettes for *Saccharomyces cerevisiae*. *Yeast* **7**:475–477.
4. Boeke, J. D., F. Lacroute, and G. R. Fink. 1984. A positive selection for mutants lacking orotidine-5'-phosphate decarboxylase activity in yeast: 5-fluoro-orotic acid resistance. *Mol. Gen. Genet.* **197**:345–346.
5. Bonneaud, N., O. Ozier-Kalogeropoulos, G. Li, M. Labouesse, L. Minvielle-Sebastia, and F. Lacroute. 1991. A family of low and high copy replicative, integrative and single-stranded *S. cerevisiae/E. coli* shuttle vectors. *Yeast* **7**:609–615.
6. Brand, R. C., J. Klootwijk, T. J. M. van Steenberg, A. J. de Kok, and R. J. Planta. 1977. Secondary methylation of yeast ribosomal precursor RNA. *Eur. J. Biochem.* **75**:311–318.
7. Brimacombe, R., P. Mitchell, M. Osswald, K. Stade, and D. Bochkarriov. 1993. Clustering of modified nucleotides at the functional center of bacterial ribosomal RNA. *FASEB J.* **7**:161–167.
8. Cavallé, J., M. Nicoloso, and J.-P. Bachellerie. 1996. Targeted ribose methylation of RNA *in vivo* directed by tailored antisense RNA guides. *Nature* **383**:732–735.
9. Cheng, X., S. Kumar, J. Posfai, J. W. Pflugrath, and R. J. Roberts. 1993. Crystal structure of the *HhaI* DNA methyltransferase complexed with S-adenosyl-L-methionine. *Cell* **74**:299–307.
10. Cunningham, P. R., C. J. Weitzmann, K. Nurse, R. Masurel, P. H. van Knippenberg, and J. Ofengand. 1990. Site-specific mutation of the conserved m⁶Am⁶A residues of *E. coli* 16S ribosomal RNA. Effects on ribosome function and activity of the *ksgA* methyltransferase. *Biochim. Biophys. Acta* **1050**:18–26.
11. Fabian, G. R., and A. K. Hopper. 1987. *RRP1*, a *Saccharomyces cerevisiae* gene affecting rRNA processing and production of mature ribosomal subunits. *J. Bacteriol.* **169**:1571–1578.
12. Ganot, P., M.-L. Bortolin, and T. Kiss. 1997. Site-specific pseudouridine formation in preribosomal RNA is guided by small nucleolar RNAs. *Cell* **89**:799–809.
13. Gietz, D., A. St. Jean, R. A. Woods, and R. H. Schiestl. 1992. Improved method for high efficiency transformation of intact yeast cells. *Nucleic Acids Res.* **20**:1425.
14. Helser, T. L., J. E. Davies, and J. E. Dahlberg. 1971. Change in methylation of 16S ribosomal RNA associated with mutation to kasugamycin resistance in *Escherichia coli*. *Nat. New Biol.* **233**:12–14.
15. Helser, T. L., J. E. Davies, and J. E. Dahlberg. 1972. Mechanism of kasugamycin resistance in *Escherichia coli*. *Nature New Biol.* **235**:6–9.
16. Henry, Y., H. Wood, J. P. Morrissey, E. Petfalski, S. Kearsy, and D. Tollervey. 1994. The 5' end of yeast 5.8S rRNA is generated by exonucleases from an upstream cleavage site. *EMBO J.* **13**:2452–2463.
17. Iizuka, N., L. Najita, A. Franzusoff, and P. Sarnow. 1994. Cap-dependent and cap-independent translation by internal initiation of mRNAs in cell extracts prepared from *Saccharomyces cerevisiae*. *Mol. Cell Biol.* **14**:7322–7330.
18. Ingresso, D., A. V. Fowler, J. Bleibaum, and S. Clarke. 1989. Sequence of the D-aspartyl/L-isoaspartyl protein methyltransferase from human erythrocytes. *J. Biol. Chem.* **264**:20131–20139.
19. Jeeninga, R. E., Y. Van Delft, M. de Graaff-Vincent, A. Dirks-Mulder, J. Venema, and H. A. Raaij. 1997. Variable regions V13 and V3 of *Saccharomyces cerevisiae* contain structural features essential for normal biogenesis and stability of 5.8S and 25S rRNA. *RNA* **3**:476–488.
20. Kiss-László, Z., Y. Henry, J.-P. Bachellerie, M. Caizergues-Ferrer, and T. Kiss. 1996. Site-specific ribose methylation of preribosomal RNA: a novel function for small nucleolar RNAs. *Cell* **85**:1077–1088.
21. Kranz, J. E., and C. Holm. 1990. Cloning by function: an alternative approach for identifying yeast homologs of genes from other organisms. *Proc. Natl. Acad. Sci. USA* **87**:6629–6633.
22. Krzyzosiak, W., R. Denman, K. Nurse, W. Hellmann, M. Boublik, C. W. Gehrke, P. F. Agris, and J. Ofengand. 1987. *In vitro* synthesis of 16S ribosomal RNA containing single base changes and assembly into functional 30S ribosomes. *Biochemistry* **26**:2353–2364.
23. Lafontaine, D., J. Delcour, A.-L. Glasser, J. Desgrès, and J. Vandenhoute. 1994. The *DIM1* gene responsible for the conserved m⁶Am⁶A dimethylation in the 3' terminal loop of 18S rRNA is essential in yeast. *J. Mol. Biol.* **241**:492–497.

- 23a. Lafontaine, D., and D. Tollervey. Unpublished data.
24. Lafontaine, D., and D. Tollervey. 1995. *Trans*-acting factors in yeast pre-rRNA and pre-snoRNA processing. *Biochem. Cell Biol.* **73**:803–812.
25. Lafontaine, D., and D. Tollervey. 1996. One-step PCR mediated strategy for the construction of conditionally expressed and epitope tagged yeast proteins. *Nucleic Acids Res.* **24**:3469–3472.
26. Lafontaine, D., J. Vandenhoute, and D. Tollervey. 1995. The 18S rRNA dimethylase Dim1p is required for pre-ribosomal RNA processing in yeast. *Genes Dev.* **9**:2470–2481.
27. Lee, W. C., D. Zabetakis, and T. Mélése. 1992. *NSR1* is required for pre-rRNA processing and for the proper maintenance of steady-state levels of ribosomal subunits. *Mol. Cell. Biol.* **12**:3865–3871.
28. Leung, D. W., E. Chen, and D. V. Goeddel. 1989. A method for random mutagenesis of a defined DNA segment using a modified polymerase chain reaction. *Technique* **1**:11–15.
29. Maden, B. E. H. 1990. The numerous modified nucleotides in eukaryotic ribosomal RNA. *Prog. Nucleic Acid Res. Mol. Biol.* **39**:241–303.
30. Morrissey, J. P., and D. Tollervey. 1993. Yeast snR30 is a small nucleolar RNA required for 18S rRNA synthesis. *Mol. Cell. Biol.* **13**:2469–2477.
31. Ni, J., A. L. Tien, and M. J. Fournier. 1997. Small nucleolar RNAs direct site-specific synthesis of pseudouridine in ribosomal RNA. *Cell* **89**:565–573.
32. Nicoloso, M., L.-H. Qu, B. Michot, and J.-P. Bachellerie. 1996. Intronic, antisense small nucleolar RNAs: the characterization of nine novel species points to their direct role as guides for the 2'-*O*-ribose methylation of rRNAs. *J. Mol. Biol.* **260**:178–195.
33. Palmer, E., J. M. Wilhelm, and F. Sherman. 1979. Phenotypic suppression of nonsense mutants in yeast by aminoglycoside antibiotics. *Nature* **277**:148–150.
- 33a. Petfalski, E., and D. Tollervey. Unpublished data.
34. Poldermans, B., H. Bakker, and P. H. van Knippenberg. 1980. Studies on the function of two adjacent N^6,N^6 -dimethyladenosines near the 3'-end of 16S ribosomal RNA of *Escherichia coli*. IV. The effect of the methyl groups on ribosomal subunit interaction. *Nucleic Acids Res.* **8**:143–151.
35. Poldermans, B., C. P. J. van Buul, and P. H. van Knippenberg. 1979. Studies on the function of two adjacent N^6,N^6 -dimethyladenosines near the 3'-end of 16S ribosomal RNA of *Escherichia coli*. II. The effect of the absence of the methyl groups on initiation of protein biosynthesis. *J. Biol. Chem.* **254**:9090–9094.
36. Salim, M., and B. E. H. Maden. 1973. Early and late methylations in HeLa cell ribosome maturation. *Nature* **244**:334–336.
37. Sikorski, R. S., and J. D. Boeke. 1991. *In vitro* mutagenesis and plasmid shuffling: from cloned gene to mutant yeast. *Methods Enzymol.* **194**:302–318.
38. Sikorski, R. S., and P. Hieter. 1989. A system of shuttle vectors and yeast host strains designed for efficient manipulation of DNA in *Saccharomyces cerevisiae*. *Genetics* **122**:19–27.
39. Singh, A., D. Ursic, and J. Davies. 1979. Phenotypic suppression and misreading in *Saccharomyces cerevisiae*. *Nature* **277**:146–148.
40. Tarun, S. Z., and A. B. Sachs. 1995. A common function for mRNA 5' and 3' ends in translation initiation in yeast. *Genes Dev.* **9**:2997–3007.
41. Thamana, P., and C. R. Cantor. 1978. Studies on ribosome structure and interactions near the $m_2^6Am_2^6A$ sequence. *Nucleic Acids Res.* **5**:805–823.
42. Tollervey, D. 1996. *Trans*-acting factors in ribosome synthesis. *Exp. Cell Res.* **229**:226–232.
43. van Buul, C. P. J. J., W. Visser, and P. H. van Knippenberg. 1984. Increased translational fidelity caused by the antibiotic kasugamycin and ribosomal ambiguity in mutants harbouring the *ksg4* gene. *FEBS Lett.* **177**:119–124.
44. van Buul, C. P. J. J., J. B. L. Damn, and P. H. van Knippenberg. 1983. Kasugamycin resistant mutants of *Bacillus stearothermophilus* lacking the enzyme for the methylation of two adjacent adenosines in 16S ribosomal RNA. *Mol. Gen. Genet.* **189**:475–478.
45. van Knippenberg, P. H. 1986. Structural and functional aspects of the N^6,N^6 -dimethyladenosines in 16S ribosomal RNA, p. 412–424. *In* B. Hardesty and G. Kramer (ed.), *Structure, function and genetics of ribosomes*. Springer-Verlag Inc., New York, N.Y.
46. Venema, J., and D. Tollervey. 1995. Processing of pre-ribosomal RNA in *Saccharomyces cerevisiae*. *Yeast* **11**:1629–1650.
47. Venema, J., and D. Tollervey. 1996. *RRP5* is required for formation of both 18S and 5.8S rRNA in yeast. *EMBO J.* **15**:5701–5714.
48. Willcock, D. F., D. T. F. Dryden, and N. E. Murray. 1994. A mutational analysis of the two motifs common to adenine methyltransferases. *EMBO J.* **13**:3902–3908.