



# THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### Human artificial chromosome (HAC) vector with a conditional centromere for correction of genetic deficiencies in human cells

**Citation for published version:**

Kim, J-H, Kononenko, A, Erliandri, I, Larionov, V, Kouprina, N, Kim, T-A, Nakano, M, Iida, Y, Masumoto, H, Oshimura, M, Barrett, JC & Earnshaw, WC 2011, 'Human artificial chromosome (HAC) vector with a conditional centromere for correction of genetic deficiencies in human cells' Proceedings of the National Academy of Sciences, vol 108, no. 50, pp. 20048-20053. DOI: 10.1073/pnas.1114483108

**Digital Object Identifier (DOI):**

[10.1073/pnas.1114483108](https://doi.org/10.1073/pnas.1114483108)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Publisher's PDF, also known as Version of record

**Published In:**

Proceedings of the National Academy of Sciences

**Publisher Rights Statement:**

Freely available online through the PNAS open access option

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



# Human artificial chromosome (HAC) vector with a conditional centromere for correction of genetic deficiencies in human cells

Jung-Hyun Kim<sup>a</sup>, Artem Kononenko<sup>a</sup>, Indri Erliandri<sup>a</sup>, Tae-Aug Kim<sup>b</sup>, Megumi Nakano<sup>c</sup>, Yuichi Iida<sup>d</sup>, J. Carl Barrett<sup>e</sup>, Mitsuo Oshimura<sup>d</sup>, Hiroshi Masumoto<sup>c</sup>, William C. Earnshaw<sup>f</sup>, Vladimir Larionov<sup>a</sup>, and Natalay Kouprina<sup>a,1</sup>

Laboratories of <sup>a</sup>Molecular Pharmacology and <sup>b</sup>Tumor and Stem Cell Biology, National Cancer Institute, Bethesda, MD 20892; <sup>c</sup>Kazusa DNA Research Institute, Kisarazu, Chiba 292-0818, Japan; <sup>d</sup>Institute of Regenerative Medicine and Biofunction, Tottori University, Tottori 683-8503, Japan; <sup>e</sup>Translational Sciences for Oncology Innovative Medicine, AstraZeneca, Boston, MA 02451; and <sup>f</sup>Wellcome Trust Centre for Cell Biology, University of Edinburgh, Edinburgh EH14 4AS, Scotland

Edited\* by John Carbon, University of California, Santa Barbara, CA, and approved October 27, 2011 (received for review September 6, 2011)

**Human artificial chromosome (HAC)-based vectors offer a promising system for delivery and expression of full-length human genes of any size. HACs avoid the limited cloning capacity, lack of copy number control, and insertional mutagenesis caused by integration into host chromosomes that plague viral vectors. We previously described a synthetic HAC that can be easily eliminated from cell populations by inactivation of its conditional kinetochore. Here, we demonstrate the utility of this HAC, which has a unique gene acceptor site, for delivery of full-length genes and correction of genetic deficiencies in human cells. A battery of functional tests was performed to demonstrate expression of *NBS1* and *VHL* genes from the HAC at physiological levels. We also show that phenotypes arising from stable gene expression can be reversed when cells are “cured” of the HAC by inactivating its kinetochore in proliferating cell populations, a feature that provides a control for phenotypic changes attributed to expression of HAC-encoded genes. This generation of human artificial chromosomes should be suitable for studies of gene function and therapeutic applications.**

gene delivery vector | gene therapy | transformation-associated recombination cloning

Most advanced current gene therapy systems, such as adenovirus-, lentivirus-, and retrovirus-derived vectors, use cDNA or “minigene” constructs (reviewed in refs. 1–5) that cannot recapitulate the physiological regulation of complex endogenous loci. Viral episomal vectors carrying HSV-1 and EBV amplicons can deliver and express full-length genes as large as approximately 150 kb in size (1, 2). However, several concerns limit the use of HSV-1 and EBV viral vectors: absence of strong copy number control, undesired immunological responses, and occasional integration into the host genome, causing insertional mutagenesis and gene silencing (6).

Human artificial chromosomes (HACs) represent another extrachromosomal gene delivery and gene expression vector system (7–11). Although this technology is less advanced than virus-derived vectors, HACs have several potential advantages over currently used episomal viral vectors for gene therapy applications. First, the presence of a functional centromere provides long-term stable maintenance of HACs as single-copy episomes without integration to the host chromosomes. Second, there is no upper size limit to DNA cloned in a HAC: entire genomic loci with all regulatory elements can be used. Finally, HAC vectors minimize adverse host immunogenic responses and the risk of cellular transformation.

Recent advances have produced a variety of HACs via two different approaches. The “top-down” approach involves modification of human chromosomes in living cells to produce chromosome derivatives (11–18). The “bottom-up” approach involves de novo kinetochore assembly from 50- to 100-kb blocks of centromeric alpha-satellite (i.e., alloid) DNAs. These are multimerized in human cells, forming HACs with sizes of approximately

1 to 3 Mb (7, 8, 10, 19). Several groups have reported successful expression of full-length genes in HACs (9–11, 20, 21). However, in most cases, genes to be expressed were cotransfected with an alloid DNA array into human cells and were incorporated into the forming HAC in vivo. Therefore, gene copy number and location of the gene on the HAC were not predetermined. These factors greatly limited the application of de novo-generated HACs as gene delivery vectors.

Recently, this limitation was overcome by construction of an alloid<sup>tetO</sup>-HAC carrying loxP sites (22, 23). This circular HAC contains approximately 6,000 copies of the tetracycline operator (tetO) sequence embedded in a synthetic alloid DNA array. Although the HAC is mitotically stable, tethering certain tetracycline repressor fusion proteins (tetR-KAP1 or tTS) (22, 24) inactivates the centromere, resulting in HAC loss. Thus, the alloid<sup>tetO</sup>-HAC and any genes it carries can be eliminated from cell populations, allowing for a regulated “hit-and-run” induction of phenotypic changes.

To minimize problems caused by the lack of certain genes from libraries of BAC and YAC clones or the presence of genes fragmented on multiple vectors, we developed a method for direct gene cloning from genomic DNA. Transformation-associated recombination (TAR) cloning allows isolation of any specific allele of a gene of interest as a predetermined DNA fragment (25, 26).

Here, we describe the combination of the TAR gene-cloning technology with the alloid<sup>tetO</sup>-HAC vector for gene delivery. We report a complete cycle starting with selective gene isolation, followed by gene loading into the HAC, and eventually leading to complementation of gene deficiencies in a human cell line (Fig. 1). This approach is useful for studies of gene function and potentially for gene therapy. As a proof of principle, genomic copies of two average-size cancer-associated genes—*VHL* mutated in von Hippel-Lindau syndrome (*VHL*; MIM 193300) and *NBS1* mutated in Nijmegen breakage syndrome (*NBS*; MIM 251260)—were isolated by TAR cloning and loaded into the unique loxP site of the alloid<sup>tetO</sup>-HAC in CHO cells. HAC transfer into human cells deficient in *NBS1* and *VHL* reveals that both genes are expressed normally and complement the defective endogenous alleles.

## Results

**Isolation of Genomic Regions Containing *NBS1* and *VHL* Genes from Human Genome by TAR Cloning.** The TAR cloning scheme for isolating the *NBS1* and *VHL* genes is shown in Fig. 1A. For

Author contributions: J.-H.K., T.-A.K., H.M., V.L., and N.K. designed research; J.-H.K., A.K., I.E., M.N., Y.I., and N.K. performed research; T.-A.K., J.C.B., M.O., H.M., W.C.E., V.L., and N.K. analyzed data; and N.K. wrote the paper.

The authors declare no conflict of interest.

\*This Direct Submission article had a prearranged editor.

Freely available online through the PNAS open access option.

<sup>1</sup>To whom correspondence should be addressed. E-mail: kouprinn@mail.nih.gov.


This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.1114483108/-DCSupplemental](http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1114483108/-DCSupplemental).

cloning purposes, TAR vectors were designed containing short targeting hooks specific to 5' and 3' sequence ends of the genes (*SI Materials and Methods*). Human genomic DNA was transformed into yeast spheroplasts along with linearized TAR vectors. Recombination between the hooks and genomic DNA fragments results in isolation of *NBS1* and *VHL* genes as circular YACs with insert sizes of approximately 55 kb and 25 kb, respectively. Five independent TAR YAC isolates were obtained for each gene. The cloned *NBS1* locus contains approximately 5 kb sequence upstream of the ATG codon and 1.5 kb sequence downstream of the stop codon. The cloned *VHL* locus contains approximately 10 kb sequence upstream of the ATG codon and 7.3 kb sequence downstream of the stop codon. PCR analysis and physical characterization of the TAR isolates for each gene confirmed that the


clones contain all exons (Fig. 1 *F* and *H*) and did not reveal any structural rearrangements. Analysis of the *NBS1* genomic segments is shown in Fig. S1.

**Conversion of *NBS1*- and *VHL*-YACs into BACs with a *LoxP* Cassette for Gene Loading into Alphoid<sup>tetO</sup>-HAC.** A yeast-bacteria-mammalian cell shuttle vector, pJBRV1, was constructed to retrofit YAC gene isolates into YAC/BACs (Figs. S2 and S3). The vector contains a 3' HPRT-*LoxP*-eGFP cassette, allowing gene loading into a unique *LoxP* site of the alphoid<sup>tetO</sup>-HAC in CHO cells. An F' factor origin of replication allows YAC propagation as a BAC molecule. Conversion of the YAC into a BAC is advantageous because purification of circular DNA molecules is much easier from *Escherichia coli* than from yeast cells. The protocol for retrofitting is shown in Fig. 1*B* and Fig. S2. Integrity of *NBS1*- and *VHL*-containing BACs was confirmed by PCR amplification of all exons (Fig. 1 *G* and *I*) and analysis of their inserts by pulse-gel electrophoresis (Fig. S1*D*).

**Loading of *NBS1* and *VHL* Genes into Alphoid<sup>tetO</sup>-HAC by Cre-*LoxP*-Mediated Recombination in CHO Cells.** The alphoid<sup>tetO</sup>-HAC with a unique gene loading site was used for this purpose. This HAC was identified among several HAC clones carrying a *LoxP* cassette (s) (23) by Southern blot hybridization. To insert genomic copies of the *NBS1* and *VHL* genes into the alphoid<sup>tetO</sup>-HAC, the appropriate BAC constructs and a Cre-recombinase expression vector were cotransfected into *hprt*-minus CHO cells carrying the HAC and HPRT-plus colonies were selected on HAT medium (Fig. 1*C*). Insertion of genes into the HAC was confirmed by PCR by using specific primers (Table S1) to detect reconstitution of the *HPRT* gene, which accompanies Cre/*LoxP* targeting. FISH images of alphoid<sup>tetO</sup>-HAC/*NBS1* and alphoid<sup>tetO</sup>-HAC/*VHL* are shown in Fig. 2 *A* and *B*. The HACs are maintained autonomously and the HAC signal colocalizes with *NBS1* and *VHL* gene signals on metaphase chromosome spreads. Immunocytochemistry detected homogeneous pNBS1 expression in the nucleus of all cells (Fig. 2 *C* and *D*). Western blot analysis of the alphoid<sup>tetO</sup>-HAC/*NBS1* clone with human-specific antibodies against pNBS1 revealed that the *NBS1* gene inserted into the



**Fig. 1.** A scheme of consecutive experimental steps from selective gene isolation to its expression in gene-deficient human cells. (A) Direct TAR isolation of the *NBS1* and *VHL* genes from human genomic DNA. TAR vectors contain two gene targeting hooks (yellow and blue boxes). (B) Retrofitting of the circular YAC containing the full-length gene by the pJBRV1 vector containing a 3' HPRT-*LoxP*-eGFP cassette. Recombination of the BamHI-linearized pJBRV1 vector with a YAC in yeast leads to replacement of the ColE1 origin of replication by the F' factor origin of replication, allowing subsequent propagation in BAC form. (C) Gene loading into a unique *LoxP* site of the alphoid<sup>tetO</sup>-HAC by Cre-*LoxP* recombination in CHO cells. (D) MMCT of alphoid<sup>tetO</sup>-HAC/*NBS1* or *VHL* from CHO into gene-deficient cells for complementation analysis. (E) Elimination of the alphoid<sup>tetO</sup>-HAC/*NBS1* or *VHL* from cells by expression of the tTS fusion construct. (F and G) Analysis of TAR clones containing the full-length *NBS1* gene for the presence of exons before (F) and after (G) retrofitting. (H and I) Analysis of TAR clones containing the full-length *VHL* gene for the presence of exons before (H) and after (I) retrofitting.



**Fig. 2.** Expression of *NBS1* and *VHL* genes loaded into the HAC in CHO cells. (A) FISH analysis of the alphoid<sup>tetO</sup>-HAC/*NBS1* in CHO cells with specific probes for the BAC vector (in red) and for gene sequences (green). (B) FISH analysis of alphoid<sup>tetO</sup>-HAC/*VHL* in CHO cells. (C) Immunocytochemical staining of CHO cells and (D) isogenic cells with the alphoid<sup>tetO</sup>-HAC/*NBS1* using Abs against pNBS1 (red) and GFP (green). (E) Western blot analysis of a CHO clone containing alphoid<sup>tetO</sup>-HAC/*NBS1* with human-specific Abs against pNBS1. *NBS1* inserted into the alphoid<sup>tetO</sup>-HAC produces a protein of the predicted size.

alphoid<sup>tetO</sup>-HAC expresses a protein of the predicted size (Fig. 2E). Because antibodies against human pVHL produce a band with the mobility of hVHL on Western blots in naive CHO cell lysates, we used RT-PCR analysis to prove expression of the *VHL* gene. RT-PCR products of the predicted size were obtained (Fig. S4A), and their identity was confirmed by sequencing.


**Functional Complementation of Genetic Deficiency by Alphoid<sup>tetO</sup>-HAC/NBS1.** To prove its use as a genetic complementation vector, the NBS1-containing HAC was transferred to NBS1-deficient GM07166 (657del5) human cells via microcell-mediated chromosome transfer (MMCT; Fig. 1D). FISH analysis confirmed that alphoid<sup>tetO</sup>-HAC/NBS1 propagates autonomously without detectable integration into chromosomes (Fig. 3A). Western blot analysis confirmed that this clone expresses pNBS1 (Fig. 3B). To prove the functionality of the conditional kinetochore and confirm that pNBS1 is expressed from the HAC, we targeted the tTS transcriptional repressor to the HAC centromere in the alphoid<sup>tetO</sup>-HAC/NBS1-containing GM07166 cells (Fig. 1E). This targeting results in a high frequency of HAC loss, as described previously (23, 24) (Fig. S5). Based on FISH and Western blot analysis, cells that lost the HAC no longer expressed pNBS1 (Fig. 3B).

We used physiological tests to prove that alphoid<sup>tetO</sup>-HAC/NBS1 produces functional pNBS1. pNBS1 plays a critical role in damage responses to DNA double-strand breaks (DSBs). Within seconds of generation of DSBs, histone H2A.X molecules are phosphorylated by ATM and ATR on serine residues. The resulting  $\gamma$ -H2A.X spreads along megabase chromatin domains flanking the DSB site (reviewed in refs. 27, 28). pNBS1 then recruits the Mre11/Rad50/NBS1 complex to the break site. Although  $\gamma$ -H2A.X foci formed in NBS1-deficient GM07166 cells

after irradiation, as expected (ref. 29 and references therein), MRE11 protein did not colocalize with  $\gamma$ -H2A.X (Fig. 3F–H). In contrast, in GM07166 cells bearing alphoid<sup>tetO</sup>-HAC/NBS1, localization of NBS1 and MRE11 to damage-induced  $\gamma$ -H2A.X foci was observed (Fig. 3F–H). This colocalization was lost in GM07166 cells that had been “cured” of the HAC (Fig. 3F–H). As a control, we examined colocalization of  $\gamma$ -H2A.X and 53BP1 in the same cells. As shown by others, 53BP1 localization to damage foci is not affected by absence of pNBS1 (ref. 29 and references therein). Indeed,  $\gamma$ -H2A.X and 53BP1 irradiation-induced foci colocalized independently of the presence of functional pNBS1 in GM07166 cells (Fig. 3F–H and Fig. S6).

Analysis of three proteins—ATM, KAP1, and p53—that should be modified in response to irradiation revealed phosphorylation of the predicted amino acid residues in the analyzed cells (Fig. 3E). We then analyzed kinetics of the disassembly of  $\gamma$ -H2A.X foci, which closely parallels the rate of DSB repair. The kinetics slows down in NBS1-deficient cells similar to that in cells deficient for ATM and 53BP1 (30). GM07166 cells bearing alphoid<sup>tetO</sup>-HAC/NBS1 showed a more efficient disassembly of foci compared with cells without HAC after 24 h (Fig. 3C and D). The dynamics of foci disassembly in the cells without and with the HAC was identical to that observed in cultured lymphocytes from patients with NBS and normal individuals, correspondingly (31). Thus, these results indicate a higher rate of rejoining of DSBs in GM07166 cells bearing alphoid<sup>tetO</sup>-HAC/NBS1. Altogether, our observations demonstrate that the alphoid<sup>tetO</sup>-HAC/NBS1 vector expresses a functional protein.

**Functional Complementation of Genetic Deficiency by Alphoid<sup>tetO</sup>-HAC/VHL.** When transferred to 786-0 renal carcinoma cells



**Fig. 3.** pNBS1 expression in NBS1-deficient cells. (A) FISH analysis of the alphoid<sup>tetO</sup>-HAC/NBS1 in GM07166 cells with specific probes for the BAC vector (red) and *NBS1* gene sequences (green). (B) Western blot analysis of NBS1-deficient GM07166 cells, alphoid<sup>tetO</sup>-HAC/NBS1-containing GM07166 cells before and after HAC elimination from the cells by expression of the tTS fusion construct. (C and D) Kinetics of disassembly of  $\gamma$ -H2A.X foci in NBS1-deficient GM07166 cells and the same cells with alphoid<sup>tetO</sup>-HAC/NBS1. (C) Immunostaining of irradiated cells. Cells were stained with anti- $\gamma$ -H2A.X antibodies (red) and with DAPI (blue) after exposition to 1 Gy IR and then collected after the times indicated. (D) Quantitative analysis of disassembly of  $\gamma$ -H2A.X foci. Cells were collected after the time indicated, and the average number of  $\gamma$ -H2A.X foci was quantified over time. (E) Western analysis of GM07166 cells and cells with alphoid<sup>tetO</sup>-HAC/NBS1. As predicted, p53, ATM, and KAP1 are phosphorylated in response to irradiation. (F–H) Analysis of colocalization of  $\gamma$ -H2A.X, MRE11, 53BP1, and NBS1 proteins after irradiation. NBS1-deficient cells (GM07166) and the same cells with alphoid<sup>tetO</sup>-HAC/NBS1 before and after HAC loss were fixed 2 h after irradiation with 3 Gy and were double-stained with anti- $\gamma$ -H2A.X, anti-hMRE11, anti-53BP1, and anti-hNBS1 antibodies. (F) NBS1 is colocalized with  $\gamma$ -H2A.X in cells carrying the alphoid<sup>tetO</sup>-HAC/NBS1. (G)  $\gamma$ -H2A.X and MRE11 are colocalized in foci in a pNBS1-dependent manner. (H) 53BP1 foci formation is not affected by absence of pNBS1.

(RCCs) via MMCT, alphoid<sup>tetO</sup>-HAC/VHL propagated autonomously without integrating into host chromosomes (Fig. 4A). These cells are VHL-negative as a result of a frameshift mutation in the coding region of the gene (311delG). Expression of *VHL* in 786-0 cells bearing alphoid<sup>tetO</sup>-HAC/VHL was confirmed by RT-PCR by using primers that specifically amplify the WT but not the mutant allele of *VHL* (Fig. S4B). To prove that *VHL* is expressed from the HAC, a vector expressing a tTS fusion was introduced into cells to induce HAC loss (23, 24). Several GFP-negative clones were selected. Based on FISH and RT-PCR analyses, these clones lost the *VHL* along with the HAC.

Several specific tests based on the known functions of pVHL were carried out to prove functional stable expression of pVHL from the HAC in recipient cells. A critical role of pVHL in down-regulation of HIF, CyclinD1, and Cdk1/CDC2 is well documented (32–35). pVHL has E3 ubiquitin ligase activity and targets  $\alpha$ -subunits of HIF for rapid degradation by the proteasome under normoxia. Those subunits are stabilized by hypoxia (reviewed in ref. 36). We compared the levels of HIF2 $\alpha$ , CyclinD1, and CDC2 expression in VHL-deficient and isogenic cells containing alphoid<sup>tetO</sup>-HAC/VHL by Western blot analysis. As predicted, levels of HIF2 $\alpha$ , CyclinD1, and CDC2 were significantly lower in cells carrying alphoid<sup>tetO</sup>-HAC/VHL (Fig. 4B), suggesting that pVHL expressed from the HAC restores their normal down-regulation. Indeed, 786-0 cells cured of the alphoid<sup>tetO</sup>-HAC/VHL vector by centromere inactivation expressed the VHL-regulated proteins at their original elevated levels (Fig. 4B). PGK1 and CA9 expression is regulated by hypoxia and more specifically by the HIF1- $\alpha$  isoform (PubMed ID no. 15964822). Because the 786-0 cell line lacks expression of HIF1- $\alpha$ , no reduction of CA9 and PGK1 was observed after induction of pVHL in 786-0 cells (Fig. 4B).


Three additional tests confirmed the functional expression of pVHL in recipient alphoid<sup>tetO</sup>-HAC/VHL cells. In RCCs, hepatocyte growth factor (HGF) and FBS stimulate branching morphogenesis, cell invasiveness, and cell migration via a VHL-dependent pathway (37, 38). We therefore assessed those phenotypes in the VHL-deficient 786-0 RCC parental cell line and its isogenic derivative containing the alphoid<sup>tetO</sup>-HAC/VHL vector (Fig. 4C–E). Addition of exogenous HGF resulted in branching of VHL-deficient 786-0 cells in a growth factor-reduced Matrigel assay. In contrast, stable expression of pVHL from the alphoid<sup>tetO</sup>-HAC/VHL vector abrogated this morpho-

logic response of 786-0 cells to HGF (Fig. 4D). Similarly, the relatively high basal level of invasiveness of VHL-deficient 786-0 cells was markedly stimulated by addition of FBS. This response was dramatically suppressed in the isogenic 786-0 cell line containing the alphoid<sup>tetO</sup>-HAC/VHL vector (Fig. 4C). Finally, migration of 786-0 cells with alphoid<sup>tetO</sup>-HAC/VHL was also reduced relative to the parental cell line (Fig. 4E).

This battery of functional tests shows that the *VHL* gene expressed from the alphoid<sup>tetO</sup>-HAC/VHL produces a functional product that complements the VHL-deficiency in 786-0 cells.


#### Proximity of CENP-A Chromatin Is Compatible with Gene Expression.

During de novo HAC formation, input DNA is multimerized and then assembled into different types of chromatin, including arrays of CENP-A nucleosomes that are interspersed with those carrying different modifications of histone H3 (39). To better understand the chromatin structure of regions flanking the *NBS1* gene in the alphoid<sup>tetO</sup> HAC, alphoid<sup>tetO</sup>-HAC/*NBS1* was transferred back to human HT1080 cells (Fig. S7) via MMCT. The alphoid<sup>tetO</sup>-HAC was generated in this host and its chromatin structure in those cells has been previously characterized (22). After MMCT, two clones (*NBS1* no. 3 and *NBS1* no. 6) were selected (Fig. 5). Immuno-FISH analysis confirmed the assembly of CENP-A centromeric chromatin on the HAC sequences (Fig. 5C). ChIP analysis of these clones showed a chromatin pattern similar to that in the original alphoid<sup>tetO</sup>-HAC clone (AB2.2.18.21) (22) for CENP-A chromatin as well as H3K4me3 (a marker for transcriptionally active chromatin), H3K4me2 (a marker for open chromatin) and H3K9me3 (a marker for heterochromatin) on alphoid<sup>tetO</sup> DNA (Fig. 5A and B). These results confirm that the alphoid<sup>tetO</sup>-HAC/*NBS1* retained its kinetochore chromatin structure after several rounds of MMCT and gene insertion. ChIP analysis of regions flanking the *NBS1* gene revealed a robust level of CENP-A chromatin enrichment at the 3' end of *NBS1* in clone *NBS1* no. 3 and a lower, but still significant, enrichment in clone *NBS1* no. 6 (Fig. 5A). This indicates that the CENP-A chromatin domain extends to beyond the gene insertion site. CENP-A enrichment at the 3' *NBS1* sequence may be explained by the ability of CENP-A chromatin to spread onto noncentromeric DNA (16, 39, 40). The observed stable gene expression in the HAC suggests that close proximity of the CENP-A domain does not prevent expression of Pol II-transcribed genes.



**Fig. 4.** pVHL expression in VHL-deficient cells. (A) FISH analysis of the alphoid<sup>tetO</sup>-HAC/VHL in VHL-deficient 786-0 RCC4 cells. (B) Effect of pVHL on the level of HIF2- $\alpha$ , CyclinD1, and Cdk1/CDC2 proteins. Immunoblots of whole cell extracts isolated from the 786-0 cell line, the same cell line containing alphoid<sup>tetO</sup>-HAC/VHL before and after induction of HAC loss. CA9 and PGK1 proteins, which are not affected by pVHL status in 786-0 cells, were used as controls. (C) The level of invasiveness of VHL-deficient 786-0 cells and the same cell line containing alphoid<sup>tetO</sup>-HAC/VHL in the presence of FBS. (D) pVHL regulation of HGF-mediated (HGF+) 786-0 cell branching morphogenesis in the absence and presence of functional pVHL (with or without alphoid<sup>tetO</sup>-HAC/VHL). (E) pVHL regulation of 786-0 cell migration. pVHL expression in 786-0 cells results in reduction of cellular migration. A clear difference was observed after 7 h when representative fields were photographed. Each experiment was performed in triplicate.

**Fig. 5.** Immuno-FISH and ChIP analyses of the  $\text{alphoid}^{\text{tetO}}$ -HAC/*NBS1*. (A) ChIP analysis of centromeric chromatin in  $\text{alphoid}^{\text{tetO}}$ -HAC/*NBS1* clones 3 and 6. Normal mouse IgG (Top), antibodies against CENP-A, H3K4me2, H3K4me3, and H3K9me3 were used for analysis. The assemblies of these proteins on the  $\text{alphoid}^{\text{tetO}}$ -HAC in AB2.2.18.21 cell line (Left), the  $\text{alphoid}^{\text{tetO}}$ -HAC carrying the human *NBS1* gene in *NBS1* no. 3 and *NBS1* no. 6 cell lines (Right), are shown. The bars show the percentage recovery of the various target DNA loci by immunoprecipitation with each antibody to input DNA. Error bars indicate SD ( $n = 2$  or 3). Analyzed loci are  $\text{alphoid}^{\text{tetO}}$  ( $\text{alphoid}$  DNA with tetO motif on the  $\text{alphoid}^{\text{tetO}}$ -HAC), *bsr* (the marker gene in the HAC vector sequence), and 3' and 5' ends of *NBS1*. rDNA (5S ribosomal DNA),  $\text{alphoid}^{\text{chr.21}}$  (centromeric  $\text{alphoid}$  DNA of chromosome 21), and *sat2* (pericentromeric satellite 2) were used as controls. Immunoprecipitated DNAs were quantified by real-time PCR. (B) Positions of probes for ChIP analysis in the original HAC and in the HAC carrying the *NBS1* gene are shown by colored boxes. (C) Immuno-FISH analysis of metaphase chromosome spreads containing the  $\text{alphoid}^{\text{tetO}}$ -HACs. Cells with the original  $\text{alphoid}^{\text{tetO}}$ -HAC (AB2.2.18.21) and with  $\text{alphoid}^{\text{tetO}}$ -HAC carrying the *NBS1* gene (clones 3 and 6) were used for analysis. Immunolocalization of the centromeric protein CENP-A on metaphases was performed by indirect immunofluorescence with anti-CENP-A antibody and Alexa 488-conjugated secondary antibody (green). HAC-specific DNA sequence (BAC) was used as a FISH probe to detect the HAC (red). CENP-A and BAC signals on the HACs overlap one another.



## Discussion

HACs represent a unique system for delivery and expression of full-length genes that has several advantages over currently used episomal virus-based vectors. Several groups have reported successful expression of full-length genes in HACs (9–11, 20, 21). The most advanced top-down HAC vectors are truncated derivatives of chromosome 21. Determination of the structure of one such derivative, 5-Mb 21 $\Delta$ pqHAC, opened the way for its use in clinical applications (18). Recently, this HAC was successfully used for delivery and expression of several full-length human genes, including the 2-Mb Duchenne Muscular Dystrophy (*DYS*) gene (41). However, a common limitation of all HACs constructed so far is the inability to induce HAC loss to enable transient gene expression or to have a negative control for phenotypic changes attributed to expression of the gene loaded into the HAC.

In our previous work, we developed a synthetic  $\text{alphoid}^{\text{tetO}}$ -HAC with a conditional centromere that can be inactivated (22). This HAC contains a unique loxP site that allows selection for gene loading (23).

In the present study, we analyzed the capacity of the  $\text{alphoid}^{\text{tetO}}$ -HAC to deliver genomic copies of average-size genes into human cells and complement genetic deficiencies. We found that the *VHL* and *NBS1* genes obtained by TAR cloning can be efficiently and accurately introduced into the  $\text{alphoid}^{\text{tetO}}$ -HAC, resulting in artificial chromosomes that can be propagated indefinitely in CHO cells. Because CHO cells exhibit a high efficiency of microcell formation, this enables transfer of the HAC to other cell types, including mouse ES cells, via MMCT (23, 42).

The HACs were transferred into cell lines derived from patients with deficiencies in *VHL* or *NBS1*. Functional expression of pVHL and pNBS1 in recipient cells was demonstrated by a set of specific tests based on the known functions of proteins. Importantly, corresponding controls could be conducted following specific elimination (i.e., curing) of the HAC from the cells following targeted inactivation of the kinetochore.

No significant changes in the level of expression of pNBS1 were detected for more than 3 mo following introduction of the

$\text{alphoid}^{\text{tetO}}$ -HAC into patient-derived cell lines (Fig. S8). Although additional studies are needed to determine whether genes inserted into the  $\text{alphoid}^{\text{tetO}}$ -HAC may eventually be subject to silencing, our results and those obtained with other HACs (21, 41) support the view that proximity to a functional kinetochore does not negatively affect gene expression. It is possible that, in the human genome, centromeric chromatin may be a privileged region for Pol II-transcribed genes, as previously shown for rice centromeres (43) and suggested by the resemblance of the centromeric chromatin profile to the downstream region of transcribed genes in human cells (39, 44). Indirectly, this hypothesis is supported by our data on spreading of CENP-A chromatin to the 3' end of *NBS1* and transcriptional activity of genes in neocentromeres (reviewed in ref. 40).

The  $\text{alphoid}^{\text{tetO}}$ -HAC, with its conditional centromere, has significant advantages over other expression/cloning systems when transient expression of a gene of interest is needed. For example, generation of iPS cells requires only transient expression of specific cellular factors such as OCT4, SOX2, KLF4, cMYC, and LIN28 (45). Therefore, the ability to eliminate the HAC along with any stem cell-inducing genes carried on it could provide a strategy to avoid insertional mutagenesis and cell transformation, complications that are frequently observed during cell reprogramming (46). To adopt the  $\text{alphoid}^{\text{tetO}}$ -HAC vector for such purpose, a HAC inactivation module consisting of the doxycycline-regulated tTS fusion is being incorporated into the HAC allowing induction of its loss by a simple addition of a ligand. For gene therapy, the HAC may be inserted into stem cells in the laboratory and then these stem cells may be introduced into the patient. It is worth noting that efficient formation and stable maintenance of HACs has been demonstrated in stem cells (47).

To summarize, in this work, we demonstrate the utility of an  $\text{alphoid}^{\text{tetO}}$ -HAC vector for delivery of full-length genes and correction of genetic deficiencies in human cells. We also demonstrate the benefit of coupling the TAR gene cloning technology, which provides an effectively unlimited resource of full-length human genes, with the tetO-loxP-HAC gene delivery and expression system. Because the structure of the  $\text{alphoid}^{\text{tetO}}$ -HAC

is determined, this system may be potentially useful for gene therapy.

## Materials and Methods

**Construction of TAR Cloning Vectors.** The TAR circularizing vectors, pVC-NBS1 and pVC-VHL, containing 5' and 3' sequences of the *NBS1* and *VHL* genes, were constructed by using the basic vector pVC604 (26). Details of the construction are described in *SI Materials and Methods*.

**Construction of pJBRV1 Retrofitting Vector and Conversion of NBS1- and VHL-YACs into BACs Containing a loxP Site.** A diagram of the pJBRV1 vector and retrofitting of a circular YAC into a BAC is shown in *Figs. S2* and *S3*. Details are described in *SI Materials and Methods*.

**Loading of Genomic Fragments into Unique loxP Site of Alphoid<sup>tetO</sup>-HAC.** Details of gene loading are described in *SI Materials and Methods*.

**MMCT.** Alphoid<sup>tetO</sup>-HAC/NBS1 and alphoid<sup>tetO</sup>-HAC/VHL were transferred from CHO cells to GM07166 and RCC 786-0 cell lines deficient for *NBS1* and *VHL* genes, respectively, by using a standard MMCT protocol (23, 41, 42). Details of MMCT are described in *SI Materials and Methods*.

**Alphoid<sup>tetO</sup>-HAC Elimination by Its Targeting with Chromatin Modifiers.** Induction of HAC loss by inactivation of its kinetochore was performed as

previously described (23, 24). The kinetics of HAC loss in response to tTS expression are shown in *Fig. S5*.

**Antibodies.** The antibodies used for immunoblotting and immunostaining are described in *SI Materials and Methods*.

**In Vitro Invasion, Cell Migration, and Branching-Morphogenesis Assays.** Cell invasion, cell migration, and branching-morphogenesis assays of 786-0 cells and 786-0 cells containing alphoid<sup>tetO</sup>-HAC/VHL were carried out as described previously (37, 38).

**ChIP Analysis.** ChIP analysis was performed as described previously (22, 24, 44). Details of ChIP are described in *SI Materials and Methods*.

**ACKNOWLEDGMENTS.** We thank Hee-Sheung Lee for technical advice and help with microcell-mediated chromosome transfer experiments. We also thank Olga Aprelikova and Bill Bonner [National Cancer Institute (NCI), National Institutes of Health (NIH)] for helpful discussion. This study was supported by the Intramural Research Program of the NIH NCI Center for Cancer Research (V.L.); grant-in-aid from the Ministry of Education, Science, Sports and Culture of Japan (H.M.); and the 21st Century Center of Excellence program from the Ministry of Education, Culture, Sports, Science and Technology of Japan (M.O.). Work in the laboratory of W.C.E. is funded by the Wellcome Trust, of which W.C.E. is a principal research fellow.

- Lufino MM, Edser PA, Wade-Martins R (2008) Advances in high-capacity extrachromosomal vector technology: Episomal maintenance, vector delivery, and transgene expression. *Mol Ther* 16:1525–1538.
- Epstein AL (2009) Progress and prospects: Biological properties and technological advances of herpes simplex virus type 1-based amplicon vectors. *Gene Ther* 16: 709–715.
- Chailertvanitkul VA, Pouton CW (2010) Adenovirus: A blueprint for non-viral gene delivery. *Curr Opin Biotechnol* 21:627–632.
- Maier P, von Kalle C, Laufs S (2010) Retroviral vectors for gene therapy. *Future Microbiol* 5:1507–1523.
- Mátraí J, Chuah MK, VandenDriessche T (2010) Recent advances in lentiviral vector development and applications. *Mol Ther* 18:477–490.
- Li CM, et al. (2002) Insertional mutagenesis of the mouse acid ceramidase gene leads to early embryonic lethality in homozygotes and progressive lipid storage disease in heterozygotes. *Genomics* 79:218–224.
- Harrington JJ, Van Bokkelen G, Mays RW, Gustashaw K, Willard HF (1997) Formation of *de novo* centromeres and construction of first-generation human artificial microchromosomes. *Nat Genet* 15:345–355.
- Ikeno M, et al. (1998) Construction of YAC-based mammalian artificial chromosomes. *Nat Biotechnol* 16:431–439.
- Grimes BR, Monaco ZL (2005) Artificial and engineered chromosomes: Developments and prospects for gene therapy. *Chromosoma* 114:230–241.
- Basu J, Willard HF (2005) Artificial and engineered chromosomes: Non-integrating vectors for gene therapy. *Trends Mol Med* 11:251–258.
- Monaco ZL, Moralli D (2006) Progress in artificial chromosome technology. *Biochem Soc Trans* 34:324–327.
- Mills W, Critcher R, Lee C, Farr CJ (1999) Generation of an approximately 2.4 Mb human X centromere-based minichromosome by targeted telomere-associated chromosome fragmentation in DT40. *Hum Mol Genet* 8:751–761.
- Shen MH, et al. (2000) A structurally defined mini-chromosome vector for the mouse germ line. *Curr Biol* 10:31–34.
- Yang JW, et al. (2000) Human mini-chromosomes with minimal centromeres. *Hum Mol Genet* 9:1891–1902.
- Spence JM, et al. (2002) Co-localization of centromere activity, proteins and topoisomerase II within a subdomain of the major human X alpha-satellite array. *EMBO J* 21:5269–5280.
- Wong LH, Saffery R, Choo KH (2002) Construction of neocentromere-based human minichromosomes for gene delivery and centromere studies. *Gene Ther* 9:724–726.
- Katoh M, et al. (2004) Construction of a novel human artificial chromosome vector for gene delivery. *Biochem Biophys Res Commun* 321:280–290.
- Kazuki Y, et al. (2010) Refined human artificial chromosome vectors for gene therapy and animal transgenesis. *Gene Ther* 10:1–10.
- Ebersole T, et al. (2005) Rapid generation of long synthetic tandem repeats and its application for analysis in human artificial chromosome formation. *Nucleic Acids Res* 33:e130.
- Moralli D, Simpson KM, Wade-Martins R, Monaco ZL (2006) A novel human artificial chromosome gene expression system using herpes simplex virus type 1 vectors. *EMBO Rep* 7:911–918.
- Breman AM, Steiner CM, Slee RB, Grimes BR (2008) Input DNA ratio determines copy number of the 33 kb Factor IX gene on *de novo* human artificial chromosomes. *Mol Ther* 16:315–323.
- Nakano M, et al. (2008) Inactivation of a human kinetochore by a specific targeting of chromatin modifiers. *Dev Cell* 14:1–16.
- Iida Y, et al. (2010) Human artificial chromosome with a conditional centromere for gene delivery and gene expression. *DNA Res* 17:293–301.
- Cardinale S, et al. (2009) Hierarchical inactivation of a synthetic human kinetochore by a chromatin modifier. *Mol Biol Cell* 20:4194–4204.
- Kouprina N, Larionov V (2006) TAR cloning: Insights into gene function, long-range haplotypes and genome structure and evolution. *Nat Rev Genet* 7:805–812.
- Kouprina N, Larionov V (2008) Selective isolation of genomic loci from complex genomes by transformation-associated recombination cloning in the yeast *Saccharomyces cerevisiae*. *Nat Protoc* 3:371–377.
- Tauchi H, et al. (2002) Nijmegen breakage syndrome gene, NBS1, and molecular links to factors for genome stability. *Oncogene* 21:8967–8980.
- Lamarque BJ, Orazio NI, Weitzman MD (2010) The MRN complex in double-strand break repair and telomere maintenance. *FEBS Lett* 584:3682–3695.
- Nakamura K, et al. (2011) Regulation of homologous recombination by RNF20-dependent H2B ubiquitination. *Mol Cell* 41:515–528.
- Noon AT, et al. (2010) 53BP1-dependent robust localized KAP-1 phosphorylation is essential for heterochromatic DNA double-strand break repair. *Nat Cell Biol* 12: 177–184.
- Porcedda P, et al. (2006) Impaired elimination of DNA double-strand break-containing lymphocytes in ataxia telangiectasia and Nijmegen breakage syndrome. *DNA Repair (Amst)* 5:904–913.
- Maxwell PH, et al. (1999) The tumour suppressor protein VHL targets hypoxia-inducible factors for oxygen-dependent proteolysis. *Nature* 399:271–275.
- Sherr CJ (1995) D-type cyclins. *Trends Biochem Sci* 20:187–190.
- Kim M, et al. (1998) Recombinant adenovirus expressing von Hippel-Lindau-mediated cell cycle arrest is associated with the induction of cyclin-dependent kinase inhibitor p27Kip1. *Biochem Biophys Res Commun* 253:672–677.
- Baba M, et al. (2003) Loss of von Hippel-Lindau protein causes cell density dependent deregulation of CyclinD1 expression through hypoxia-inducible factor. *Oncogene* 22: 2728–2738.
- Baldewijns MM, et al. (2010) VHL and HIF signalling in renal cell carcinogenesis. *J Pathol* 221:125–138.
- Jeffers M, Rong S, Woude GF (1996) Hepatocyte growth factor/scatter factor-Met signaling in tumorigenicity and invasion/metastasis. *J Mol Med (Berl)* 74:505–513.
- Koochekpour S, et al. (1999) The von Hippel-Lindau tumor suppressor gene inhibits hepatocyte growth factor/scatter factor-induced invasion and branching morphogenesis in renal carcinoma cells. *Mol Cell Biol* 19:5902–5912.
- Lam AL, Boivin CD, Bonney CF, Rudd MK, Sullivan BA (2006) Human centromeric chromatin is a dynamic chromosomal domain that can spread over noncentromeric DNA. *Proc Natl Acad Sci USA* 103:4186–4191.
- Alonso A, Hasson D, Cheung F, Warburton PE (2010) A paucity of heterochromatin at functional human neocentromeres. *Epigenetics Chromatin* 3:6.
- Kazuki Y, et al. (2010) Complete genetic correction of ips cells from Duchenne muscular dystrophy. *Mol Ther* 18:386–393.
- Fournier RE, Ruddle FH (1977) Microcell-mediated transfer of murine chromosomes into mouse, Chinese hamster, and human somatic cells. *Proc Natl Acad Sci USA* 74:319–323.
- Nagaki K, et al. (2004) Sequencing of a rice centromere uncovers active genes. *Nat Genet* 36:138–145.
- Bergmann JH, et al. (2011) Epigenetic engineering shows H3K4me2 is required for HJURP targeting and CENP-A assembly on a synthetic human kinetochore. *EMBO J* 30: 328–340.
- Takahashi K, Yamanaka S (2006) Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell* 126:663–676.
- Takahashi K, et al. (2007) Induction of pluripotent stem cells from adult human fibroblasts by defined factors. *Cell* 131:861–872.
- Mandegar MA, et al. (2011) Functional human artificial chromosomes are generated and stably maintained in human embryonic stem cells. *Hum Mol Genet* 20:2905–2913.