



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

The earliest thymic T cell progenitors sustain B cell and myeloid lineage potential

Citation for published version:

Luc, S, Luis, TC, Boukarabila, H, Macaulay, IC, Buza-Vidas, N, Bouriez-Jones, T, Lutteropp, M, Woll, PS, Loughran, SJ, Mead, AJ, Mizukami, T, Matsuoka, S, Ferry, H, Duarte, S, Atkinson, D, Domanski, A, Jacobsen, SEW, Hultquist, A, Anderson, K, Brown, J, Soneji, S, Patient, R, De Bruijn, M, Enver, T, Farley, A, Sanjuan-Pla, A, Nerlov, C, Blackburn, C, Carella, C & Godin, I 2012, 'The earliest thymic T cell progenitors sustain B cell and myeloid lineage potential' Nature Immunology, vol 13, no. 4, pp. 412-419. DOI: 10.1038/ni.2255

Digital Object Identifier (DOI):

[10.1038/ni.2255](https://doi.org/10.1038/ni.2255)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Nature Immunology

Publisher Rights Statement:

RoMEO yellow

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Published in final edited form as:

Nat Immunol. ; 13(4): 412–419. doi:10.1038/ni.2255.

The Earliest Thymic T Cell Progenitors Sustain B Cell and Myeloid Lineage Potentials

Sidinh Luc^{1,2,4}, Tiago C. Luis^{1,9}, Hanane Boukarabila^{1,9}, Iain C. Macaulay¹, Natalija Buza-Vidas^{1,3}, Tiphaine Bouriez-Jones¹, Michael Lutteropp^{1,4}, Petter S. Woll¹, Stephen J. Loughran¹, Adam J. Mead¹, Anne Hultquist², John Brown⁴, Takuo Mizukami¹, Sahoko Matsuoka¹, Helen Ferry^{1,†}, Kristina Anderson², Sara Duarte¹, Deborah Atkinson¹, Shamit Soneji⁴, Aniela Domanski¹, Alison Farley³, Alejandra Sanjuan-Pla³, Cintia Carella⁵, Roger Patient⁴, Marella de Bruijn⁴, Tariq Enver^{4,††}, Claus Nerlov³, Clare Blackburn³, Isabelle Godin^{6,7,8}, and Sten Eirik W. Jacobsen^{1,2,4}

¹Haematopoietic Stem Cell Laboratory, Weatherall Institute of Molecular Medicine, John Radcliffe Hospital, University of Oxford, Headington, Oxford OX3 9DS, United Kingdom

²Hematopoietic Stem Cell Laboratory, Lund Stem Cell Center, Lund University, Klinikgatan 26, 221 84, Lund, Sweden

³Institute for Stem Cell Research, MRC Centre for Regenerative Medicine, University of Edinburgh, EH9 3JQ Edinburgh, UK

⁴MRC Molecular Haematology Unit, Weatherall Institute of Molecular Medicine, University of Oxford, Oxford OX3 9DS, United Kingdom.

⁵EMBL Mouse Biology Unit, Via Ramarini 32, 00015 Monterotondo, Italy

⁶INSERM U1009

⁷Institut Gustave Roussy, 114, rue Edouard Vaillant, Villejuif, F-94805

⁸University of Paris-Sud, Orsay, F-91405

Abstract

The stepwise commitment from hematopoietic stem cells in the bone marrow (BM) to T lymphocyte-restricted progenitors in the thymus represents a paradigm for understanding the requirement for distinct extrinsic cues during different stages of lineage restriction from multipotent to lineage restricted progenitors. However, the commitment stage at which progenitors migrate from the BM to the thymus remains unclear. Here we provide functional and molecular

Correspondence should be addressed to S.E.W.J. (sten.jacobsen@imm.ox.ac.uk).

⁹Equal contribution

[†]Present address: Experimental Medicine Division, Nuffield Department of Clinical Medicine, John Radcliffe Hospital, Oxford OX3 9DU, UK

^{††}Present address: University College London Cancer Institute, London WC1E 6BT, UK.

AUTHOR CONTRIBUTIONS

SEWJ and SL designed and conceptualized the overall research, analyzed the data and wrote the manuscript, which was subsequently reviewed and approved by all authors. JB processed RNA samples and ICM, SS analyzed the microarray data. AM, DA and AH performed quantitative and single cell PCR. AJM, SM, KA performed morphology analyses. HF, SL, ML performed FACS sorting. SL, ML, TBJ, SD, NBV, HB, TCL, AD, SJL contributed in FACS experiments and to in vitro culture experiments. SL, SD, NBV, PSW, TCL, HB performed in vivo transplantations. TE provided expertise in design and analysis of microarray experiments. CB, AF, RP, MdB, IG and TM contributed with expert advice and input on experimental design. CN, ASP and CC generated and provided input on studies of *Vwf*EGFP mice.

Accession codes. All microarray data are deposited at GEO database with project number GSE29382.

Note: Supplementary information is available on the Nature Immunology website.

The authors declare no competing financial interests.

evidence at the single cell level that the earliest progenitors in the neonatal thymus possessed combined granulocyte-monocyte, T and B lymphocyte, but not megakaryocyte-erythroid lineage potential. These potentials were identical to those of thymus-seeding progenitors in the BM, which were closely related at the molecular level. These findings establish the distinct lineage-restriction stage at which the T lineage commitment transits from the BM to the remote thymus.

At the heart of developmental and stem cell biology, as well as regenerative medicine, is the fundamental process of lineage commitment from self-renewing multipotent stem cells to lineage-restricted progenitors. In all species and organ systems this process occurs first during embryonic development, but is recapitulated postnatally and in adult life by adult multipotent stem cells to replenish cell lineages with a limited lifespan. Hematopoiesis represents the mammalian paradigm for how multilineage diversity can be achieved through commitment of multipotent stem cells into lineage-committed progenitors and the establishment of distinct blood cell lineages¹. However, the exact cellular commitment pathways remain unclear^{1,2}.

Whereas lineage-restricted progenitors for all other blood cell lineages can be generated from self-renewing hematopoietic stem cells (HSCs) in the postnatal bone marrow (BM), the final steps of T lymphocyte lineage restriction take place in the thymus³. Because the thymus cannot sustain HSCs, continuous thymopoiesis can only be secured through regular replenishment by BM-resident thymus-seeding progenitors (TSPs)⁴. However, the commitment stage(s) at which these progenitors migrate from the BM to the thymus is unknown. The thymus, as the BM, harbors multiple blood cell lineages^{5,6}, however the identification of multipotent progenitors that match the lineage potential of candidate TSPs in the BM has not been possible so far. The early thymic progenitors (ETPs) have been extensively studied in the adult thymus, but their exact lineage potentials and relationship to candidate TSPs in the BM have remained unclear⁷.

Recent studies that evaluated the lineage potential of ETPs at the single cell level showed that a large fraction of adult ETPs have combined T cell and myeloid (granulocyte-monocyte; GM) potential^{8,9}. B lineage potential was however not detected in the single, highly purified, adult ETPs, suggesting that the most primitive progenitor in the thymus might have a T-GM restricted potential^{8,9}. Similar studies of the fetal thymus have supported the T-GM restricted potential of ETPs, and failed to reveal any B cell potential^{10,11}. However, other studies reported even rarer adult ETPs with combined T and B (but not myeloid) cell potential⁶, and candidate TSPs identified in the BM⁷, such as common lymphoid progenitors (CLPs)¹², lymphoid-primed multipotent progenitors (LMPPs)¹³ and HSCs¹, all possess B cell potential.

The megakaryocyte-erythroid (MkE) potential of ETPs is of particular relevance for the ongoing debate as to whether the first lineage commitment step in hematopoiesis results in a strict separation into common myeloid and lymphoid lineage commitment pathways as depicted in the still prevailing text-book hierarchical model for hematopoiesis^{1,14}, or whether early lymphoid progenitors sustain GM but not MkE potential^{2,13,15}, as recently reported in human studies as well^{16,17}. The MkE potential of purified ETPs has yet to be investigated^{5,6,8,9}.

The fact that no multipotent thymic progenitors with the same lineage potentials as candidate multipotent TSPs in the BM have been identified yet contributes to the considerable gap in understanding the distinct roles of the local BM and thymus environments in promoting distinct pre-thymic and thymic stages of T cell lineage commitment. Here we demonstrate at the single cell level the existence of postnatal ETPs with combined T, GM and B cell but no MkE potential, establishing the exact lineage

commitment step at which the multipotent T-lymphocyte progenitors must migrate to the thymus to allow the final steps of T-lineage restriction to be completed. The data reported here provide further support of a myeloid-based model for hematopoietic lineage commitment.

RESULTS

Neonatal ETPs have combined T, B and GM, but not MkE potential

The current knowledge about candidate TSPs and ETPs can only be reconciled (Supplementary Fig. 1) if either a T-GM restricted progenitor can be identified in the BM, a T-GM progenitor would be generated in the passage from the BM to the thymus and/or a thymic cell population with a combined T, GM and B cell lineage potential could be identified within or outside the ETPs. In the last scenario, the ETP could either be a lymphoid-GM restricted multipotent progenitor or a pluripotent hematopoietic stem- or progenitor cell that also possesses MkE potential. ETPs have mostly been studied in adult mice^{5,6,8,9}. However, thymic involution, the physiological shrinking of the thymus with age that occurs in all vertebrates, implies that thymopoiesis, and therefore thymus seeding, is much more active in the early postnatal thymus¹⁸. The B cell potential of early thymocytes, at a population level, is significantly higher (although still low) in the neonatal thymus compared to the adult thymus¹⁹. In agreement with previous studies, Lin⁻CD4⁻CD8 α ⁻CD25⁻c-Kit^{hi} ETPs represented only 0.01% of adult thymocytes⁵ but as many as 40% of ETPs expressed cell surface Flt3, higher than previously reported²⁰ (Fig. 1a). Also in agreement to previous findings, a low (3.5-4.5%) but highly reproducible frequency of Flt3 expressing ETPs generated B cells^{6,20}, whereas no other adult thymocyte progenitors, including Flt3⁻ ETPs had any detectable B cell potential (Fig. 1b and Supplementary Fig. 2). The frequency of Lin⁻CD4⁻CD8 α ⁻CD25⁻c-Kit^{hi} Flt3^{hi} ETPs increased more than 10-fold in newborn mice compared to adult mice (Fig. 1c), while the frequency of Lin⁻CD4⁻CD8 α ⁻CD25⁻c-Kit^{hi}Flt3^{hi} ETPs with B cell potential was 25% (Fig. 1d and Supplementary Fig. 3). Neither Lin⁻CD4⁻CD8 α ⁻CD25⁻c-Kit^{hi}Flt3⁻ thymocytes or downstream double negative (DN) 2-4 populations in the neonatal thymus had any B cell potential (Fig. 1d). Lin⁻CD4⁻CD8 α ⁻CD25⁻c-Kit^{hi}Flt3^{hi} newborn ETPs also produced B cells *in vivo* when transplanted into irradiated *Rag1*^{-/-} mice, but only very low numbers of short-lived myeloid cells (Supplementary Fig. 4).

Because B cell activity in the thymus might reflect the presence of cells already committed to the B cell lineage^{21,22}, which overlap with the CD25⁻CD44⁺ phenotype of DN1 thymocytes, we performed an *Mb-1* Cre fate mapping experiment, in which all committed B cell progenitors and their progeny are labeled with enhanced yellow fluorescent protein (eYFP)^{23,24}. In agreement with previous studies²³, cells of the B cell lineage, including all CD19⁺B220⁺CD43⁺c-Kit⁺ proB cells, as well as a fraction of Ly-6D⁺ CLPs (Lin⁻CD19⁻B220⁻Sca-1^{low}c-Kit^{low}Flt3⁺IL-7R α ⁺Ly-6D⁺) were labeled in the BM (Supplementary Fig. 5a,b). No eYFP⁺ cells were observed in Lin⁻CD4⁻CD8 α ⁻CD25⁻c-Kit^{hi} ETPs (Fig. 1e) or in Ly-6D⁻CLPs (Lin⁻CD19⁻B220⁻Sca-1^{low}c-Kit^{low}Flt3⁺IL-7R α ⁺Ly-6D⁻) and LMPPs (Lin⁻Sca-1⁺c-Kit^{hi} (LSK)Flt3^{hi}) (Supplementary Fig. 5b,c).

In addition to B cells, newborn Lin⁻CD4⁻CD8 α ⁻CD25⁻c-Kit^{hi}Flt3^{hi} ETPs gave rise efficiently to cells of the T, NK and GM lineages, as previously demonstrated with adult ETPs (Fig. 1f,g and Supplementary Fig. 6a). In contrast, newborn ETPs were completely devoid of MkE potential (Fig. 1f). Adult ETPs lacked Mk potential as well, but, in agreement with previous studies, had GM potential (Supplementary Fig. 6a). Quantitative gene expression analysis showed that purified Lin⁻CD4⁻CD8 α ⁻CD25⁻c-Kit^{hi}Flt3^{hi} newborn ETPs expressed multiple GM and lymphoid, but not Mk and E affiliated genes

(Fig. 1h). Single-cell PCR experiments showed that as much as 65% of newborn $\text{Lin}^{-}\text{CD4}^{-}\text{CD8}\alpha^{-}\text{CD25}^{-}\text{c-Kit}^{\text{hi}}\text{Flt3}^{\text{hi}}$ ETPs co-expressed GM and lymphoid genes, while lacking M κ E gene expression at the single cell level (Fig. 2a).

To establish whether the T, B and GM potential of neonatal ETPs reflects the existence of a multipotent lympo-myeloid progenitor in the thymus, or only a mixture of lineage-restricted progenitors, we assessed the combined lineage potentials of single $\text{Lin}^{-}\text{CD4}^{-}\text{CD8}\alpha^{-}\text{CD25}^{-}\text{c-Kit}^{\text{hi}}\text{Flt3}^{\text{hi}}$ ETPs. Single ETPs were sorted onto OP9 BM stroma to allow cells to expand for 54 hours, after which the expanded cells were split and transferred to OP9 and OP9-DL1 stroma for an additional 1 week to promote B- and combined T- and myeloid cell differentiation, respectively. Although the frequency of ETP-derived clones with GM readout was reduced compared to assays in which only the GM differentiation of ETPs was promoted (Fig. 1f), we could demonstrate the existence of $\text{Lin}^{-}\text{CD4}^{-}\text{CD8}\alpha^{-}\text{CD25}^{-}\text{c-Kit}^{\text{hi}}\text{Flt3}^{\text{hi}}$ single ETPs with combined T, B and GM lineage potentials (9.2% of clones with a lineage readout; Fig. 2b,c and Supplementary Fig. 7a). In fact, all the GM potential from wild-type ETPs was tracked to cells that not only had T cell potential as previously demonstrated^{8,9}, but also possessed B lineage potential (Fig. 2c). Next, we used ETPs purified from *vavP-mcl-1* transgenic mice²⁵ to evaluate whether enhanced cell survival could sustain short-lived myeloid cells to increase the readout for combined myeloid and lymphoid potential. Whereas the B cell potential in *vavP-mcl-1* neonatal thymuses remained restricted to $\text{Lin}^{-}\text{CD4}^{-}\text{CD8}\alpha^{-}\text{CD25}^{-}\text{c-Kit}^{\text{hi}}\text{Flt3}^{\text{hi}}$ ETPs (Supplementary Fig. 7c), the frequency of ETPs with combined T-B-GM readout increased to 20% of all single *vavP-mcl-1* ETPs (as compared to 9.2% of WT ETPs) with a lineage readout (Fig. 2d,e and Supplementary Fig. 7b). These findings in single cell clonal assays establish the existence of thymic ETPs with combined T, B and GM lineage potential.

Lympho-myeloid ETPs are the most multipotent thymic progenitors

We next explored whether the $\text{Lin}^{-}\text{CD4}^{-}\text{CD8}\alpha^{-}\text{CD25}^{-}\text{c-Kit}^{\text{hi}}\text{Flt3}^{\text{hi}}$ ETPs with combined T, B and GM lineage potentials represent the most multipotent progenitors in the neonatal thymus. The lack of detectable M κ and E potential in $\text{Lin}^{-}\text{CD4}^{-}\text{CD8}\alpha^{-}\text{CD25}^{-}\text{c-Kit}^{\text{hi}}\text{Flt3}^{\text{hi}}$ ETPs did not rule out the presence of rare pluripotent stem- or progenitor cells in the neonatal thymus. Thus, we first used highly sensitive flow cytometry analysis to investigate the expression of three antigens, CD150²⁶, CD201²⁷ and Mpl²⁸, each highly expressed on most if not all HSCs as well as multipotent progenitors with sustained M κ E potential. Neither of these antigens was expressed on $\text{Lin}^{-}\text{CD4}^{-}\text{CD8}\alpha^{-}\text{CD25}^{-}\text{c-Kit}^{\text{hi}}$ ETPs (Fig. 3a). Similar to a subfraction of BM LMPPs, all ETPs expressed the RAG1 gene, as assessed using a green fluorescent protein (GFP) reporter under control of the *Rag1* promoter²⁹ and most expressed the chemokine receptor CCR9 (Fig. 3b) in agreement with previous studies of adult ETPs^{30,31}. No BM HSCs expressed either *Rag1*-GFP or CCR9 (Fig. 3b).

Whole thymocytes from neonatal mice transplanted intravenously or intrafemorally (to bypass potential changes in BM homing properties upon thymus entry) reconstituted T cells and low levels of B cells transiently (Fig. 3c,d), but failed to sustain any long-term multilineage reconstitution (Fig. 3e,f), in further support of the idea that the postnatal thymus does not harbor any HSCs. To enhance the detection of HSCs potentially present in the thymus, whole thymocytes were depleted of CD4⁺ and CD8⁺ cells and transplanted intravenously or intrafemorally (Fig. 3g,h). The absence of long-term myeloid reconstitution in all major hematopoietic organs as well as the lack of thymocyte-derived T cell progenitors in the thymus after 13 weeks in all but one transplanted mouse, further corroborated the absence of pluripotent HSCs in the thymus (Fig. 3g,h and Supplementary Fig. 8a-d). Collectively, these results demonstrate the absence of HSCs in the postnatal thymus, compatible with ETPs with combined T, B and GM lineage potential being the most multipotent progenitors in the thymus.

ETPs are closely molecularly related to BM TSPs

Because our findings indicate that $\text{Lin}^{-}\text{CD4}^{-}\text{CD8}\alpha^{-}\text{CD25}^{-}\text{c-Kit}^{\text{hi}}\text{Flt3}^{\text{hi}}$ ETPs in the neonatal thymus have the same lineage potentials as $\text{LSKFlt3}^{\text{hi}}$ LMPPs expressing *Rag1* GFP (which also highly express interleukin 7 receptor α , IL-7R α ; Supplementary Fig. 9a) in the BM^{13,32}, we next investigated the molecular relationship between ETPs and IL-7R α ⁺ LMPPs and HSCs in the BM. We also compared ETPs to the next stages of lineage restriction in the thymus, namely $\text{Lin}^{-}\text{CD44}^{+}\text{CD25}^{+}\text{c-Kit}^{\text{hi}}$ DN2 cells that sustain combined T and GM but no B lineage potential^{8,9}, and $\text{Lin}^{-}\text{CD44}^{-}\text{CD25}^{+}$ DN3 cells, which represent the first T cell restricted progenitors in the thymus³³. Based on global gene expression analysis³⁴, ETPs clustered much closer to LMPPs in the BM than to thymic DN2s and DN3s or BM HSCs. Moreover, LMPPs clustered closer to ETPs than they did to HSCs, and DN2s closer to DN3s than ETPs (Fig. 4a). Because CLPs have been suggested to be candidate TSPs⁷ and have recently been demonstrated to not only have lymphoid potential, but also sustain some myeloid potential similar to LMPPs³⁵, we also compared the molecular profiles of ETPs with the two candidate TSP populations in the BM, LMPPs and $\text{Lin}^{-}\text{CD19}^{-}\text{B220}^{-}\text{Sca-1}^{\text{low}}\text{c-Kit}^{\text{low}}\text{Flt3}^{+}\text{IL-7R}\alpha^{+}\text{Ly-6D}^{-}$ CLPs³⁶ (Fig. 4b). The newborn ETPs clustered closely with CLPs as well as LMPPs and were more distant from HSCs, DN3 and also proB cells. Moreover newborn and adult ETPs clustered closely together along with LMPPs and CLPs (Fig. 4b). These findings establish a close molecular relationship between $\text{Lin}^{-}\text{CD4}^{-}\text{CD8}\alpha^{-}\text{CD25}^{-}\text{c-Kit}^{\text{hi}}\text{Flt3}^{\text{hi}}$ ETPs in the thymus and candidate TSPs with lympho-myeloid potential in the BM (Fig. 4a,b and Supplementary Fig. 9b,c).

To more specifically look at T cell and GM lineage affiliated gene expression, T and GM gene lists were derived from the literature and from previously published data sets^{37,38} (Supplementary Methods). When T and GM lineage affiliated programs were compared, adult and neonatal ETPs clustered closely for both, and closer to LMPPs and CLPs than to DN2 and DN3 in the thymus and HSCs in the BM (Fig. 5a,b and Supplementary Fig. 9d,e). Of particular interest, many T cell affiliated genes that eventually become dramatically upregulated in DN2s and further upon T cell commitment in DN3s, have already been upregulated in LMPPs (as compared to HSCs) and CLPs, with less change from LMPPs and CLPs to ETPs (Fig. 5a). Using quantitative RT-PCR, some of the myeloid, lymphoid, T cell and Notch related genes were investigated in greater detail (Fig. 6). In addition to confirming the combined expression of GM (*Mpo*, *Csf1r*, *Csf2rb*) and lymphoid (sterile IgH, *Ii7r*, *Rag1*) lineage affiliated genes, these data also showed that characteristic early T cell specific genes (*Ptcr*, *Cd3e*, *Cd3g*) were not significantly upregulated in neither multipotent LMPPs nor ETPs. In contrast, *Notch1* was upregulated in LMPPs and further in ETPs, and in agreement with this, the Notch target genes *Ii2ra* (*Cd25*), *Gata3* and in particular *Hes1*³⁹ were upregulated in the transition from LMPPs to ETPs. Finally, whereas HSCs lack expression of *Ccr7* and *Ccr9*^{30,31}, encoding critical chemokine receptors for migration to the thymus, these genes were drastically upregulated in LMPPs, in further support of the idea that LMPPs are TSPs. Collectively, these results demonstrate that ETPs and candidate TSPs such as LMPPs and CLPs, have closely related gene expression profiles, reinforcing that ETPs more likely are derived from lympho-myeloid restricted progenitors than HSCs in the BM.

DISCUSSION

Here we identify ETPs in the neonatal thymus with combined T, B and GM, but no M κ E lineage potential, and we demonstrate a close functional and molecular linkage between ETPs and candidate TSPs in the BM.

The observation that ETPs lack M_kE potential is of key importance towards reconciling the ongoing debate regarding the roadmap for hematopoietic lineage commitment, as the classical model for hematopoietic lineage commitment predicts that the first lineage commitment step of HSCs results in a strict separation of myelo-erythroid and lymphoid commitment pathways^{1,14}. According to this model, any cell with combined lymphoid and GM potential should also possess M_kE potential. However, the current study demonstrates that Lin⁻CD4⁻CD8 α ⁻CD25⁻c-Kit^{hi}Flt3^{hi} ETPs with combined T, B and GM lineage potential were devoid of M_k and E lineage potentials. These cells co-express, at the single cell level, lymphoid and GM, but not M_kE related genes, similar to LMPPs with identical lineage potentials in the BM^{13,34}. Thus, this study provides support for a myeloid-based lineage commitment model^{2,13,15,16,17}, by demonstrating the existence of T-B-GM restricted progenitors in the postnatal thymus. Such cells have been previously identified in the BM and the fetal liver^{13,34}.

It is very likely that the real frequency of neonatal Lin⁻CD4⁻CD8 α ⁻CD25⁻c-Kit^{hi}Flt3^{hi} ETPs with T-B-GM potential is higher than what we were able to demonstrate. Analysis of clones grown from single newborn ETPs demonstrated that most ETPs with T cell potential simultaneously possessed B cell potential, but less than 20% of these also revealed GM potential in WT mice, although under optimized GM conditions more than 50% of ETPs demonstrated GM potential. We speculated that the reduced GM readout in the multilineage clonal assay reflected the short life-span of vulnerable myeloid cells, and in agreement with this MCL-1 enhanced the detection of neonatal ETPs with combined T, B and GM potential, most likely through increased survival of myeloid cells.

Our findings also suggest that the identified T-B-GM restricted progenitor is the most multipotent progenitor in the neonatal thymus, as we failed to detect any M_kE lineage potential or M_kE-specific gene expression within highly purified ETPs. Furthermore, we also demonstrate that there are no phenotypic or *in vivo* reconstituting HSCs or multipotent progenitors in the neonatal thymus.

Previous studies have suggested that the earliest fetal thymic progenitors in the embryo have combined T cell and myeloid, but no B cell lineage potential^{10,11}, raising the possibility that the progenitors seeding the embryonic thymus might be distinct and more committed than in the postnatal thymus. In contrast with the neonatal thymus which we concentrated on in this study, it remains unclear if the adult thymus is also seeded with ETPs with combined T, GM and B lineage potentials. As thymopoiesis is much decreased in the adult compared to the newborn thymus, it can be predicted that the most multipotent ETPs are much more infrequent in adult than in the neonatal thymus. While the low frequency of B lineage potential of adult ETPs^{6,20} does not allow a clear test of the combined lineage potential of these cells using the current clonal lineage potential assay, it is notable that rare Lin⁻CD4⁻CD8 α ⁻CD25⁻c-Kit^{hi}Flt3^{hi} ETPs were the only adult thymocytes with B cell potential^{6,20}. In addition, the current study showed that adult ETPs, like neonatal ETPs, have GM, but not M_k potential, while global gene expression analysis pinpointed a very close molecular relationship between neonatal and adult ETPs. Collectively these datasets suggest that adult thymuses, as the neonatal thymuses, might also be seeded by rare T-B-GM progenitors, which we could unequivocally identify in the neonatal thymus. Nevertheless, there are distinct differences between fetal, neonatal and adult hematopoietic stem and progenitor cells. The regulated migration of TSPs to the thymus might also differ in the newborn and the adult, so it remains possible that the lineage potentials of newborn and adult TSPs might be different.

While these studies establish the extent of ETP multipotentiality (T-B-GM) and the close phenotypic and molecular relationship between ETPs and LMPPs and CLPs with the same

lineage potentials in the BM^{13,32,34,35}, they do not exclude the possibility that other candidate progenitors in the BM might seed the thymus⁴⁰. A recent study suggested that T-GM-restricted progenitors might exist in the BM⁴¹, although such progenitors remain to be purified and characterized in further detail. The GM potential of ETPs is limited, and recent studies have suggested that it has little if any functional significance with regard to these progenitors acting as myeloid progenitors in the thymus^{35,42}. Likewise, it seems unlikely that ETPs have any important physiological role as B cell progenitors. Rather, the significance of these sustained lineage potentials of ETPs is to provide a better understanding of the required lineage restriction steps for lineage commitment from pluripotent HSCs in the BM to a T cell restricted progenitor in the thymus. Specifically, progenitors with combined T-B-GM potential, such as LMPPs and CLPs, are derived in the BM from HSCs that have shut down the M κ E transcriptional programs and lineage potentials. Unlike HSCs, LMPPs and CLPs upregulate CCR9^{30,31}, which enables their transfer to the thymus^{30,31}. Migration to the thymus appears critical for the next T-lineage restriction steps, first to a T-GM^{8,9} and finally to a fully T cell restricted progenitor. In addition to providing novel insights into the normal stepwise process of T lineage commitment in the BM and thymus, the present studies are also of considerable relevance for a recently identified clinically, phenotypically and molecularly distinct group of mixed T-GM acute lymphoblastic leukemias (ALLs) which are predominantly observed in children but also seen in adults, and termed ETP leukemias⁴³. Furthermore, the sustained B cell potential of ETPs might explain why the MLL-AF4 fusion oncogene, highly specific for human B cell malignancies, can give rise to B cell malignancies even if targeted to thymic progenitors⁴⁴.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank the Biomedical Services at Oxford University for expert animal support, S.Clark, T. Furey and B. Wu for technical assistance and E. Zuo and M. Eckart at the Stanford Protein and Nucleic Acid Facility for gene array services. This work was supported by grants from: EU-FP7 STEMEXPAND and a programme grant (H4RPLK0) from the Medical Research Council, UK to SEWJ, as well as the EU-FP7 EuroSyStem Integrated projects to SEWJ, CB and AF. CB and AF were supported by the Leukaemia and Lymphoma Research and AJM holds a Leukaemia and Lymphoma Research Senior Bennett Fellowship. AH is supported by the Crafoord Foundation, The George Danielsson foundation, Swedish Society for Medicine and Swedish Cancer foundation.

Appendix

METHODS

Animals

All animals used were bred and maintained at the Oxford Biomedical Services and all experiments were performed with the approval of the UK Home Office. *Rag1* GFP²⁹ mice were provided by Dr N. Sakaguchi, *Lysozyme M* eGFP45 mice by Dr T. Graf, *vavP-mcl-1* transgenic mice²⁵ by Dr S. Cory, *Mb-1* Cre (Cd79a^{tm1(cre)}Reth)²³ mice by Dr M. Reth, which were crossed with *R26R*-eYFP²⁴ mice from Dr S. Srinivas. *Vwfe*GFP BAC transgenic mice in which all platelets express eGFP were generated by BAC recombineering in bacteria. The eGFP followed by a polyadenylation signal was inserted at the initiation codon within exon 1 of the *Vwf* gene and thereby replacing exon 2. *Vwfe*GFP transgenic mice were generated by intracytoplasmic sperm injection (ICSI) (Nerlov and Jacobsen, manuscript in preparation). Mice were backcrossed multiple generations onto a C57BL/6 background.

Transplantation

Newborn/neonatal thymocytes (total thymocytes or CD4- and CD8 lineage depleted, from one donor to each recipient) or sorted Flt3⁺ ETPs from wild type or *Vwfe*GFP mice were transplanted with 300,000 unfractionated support BM cells into lethally irradiated (2x450cGy) C57BL6 (CD45.1 or CD45.2) or *Rag1*^{-/-} mice (>8 weeks), either intravenously or intraperitoneally. Peripheral blood, spleen, thymus and BM analyses were conducted between 3 and 16-17 weeks post-transplantation. For all lineages, mice showing total donor contribution >0.1% and >0.02% thymocyte-derived lineage chimerism of total cells as well as clear positive events by FACS (>20 events) were considered reconstituted.

Stem- and progenitor cell purification

All cell sorting experiments were performed on a BD FACSAriaIIu cell sorter (BD Biosciences) with a total mean cell sorting purity of 99% ± 2% for Flt3^{hi}ETPs, based on reanalysis for all gating steps and therefore all antigens defining Flt3^{hi}ETPs (Lin⁻CD4⁻CD8α⁻CD25⁺c-Kit^{hi}Flt3^{hi}) (Supplementary Fig. 3). Single cells were seeded by an automated cell deposition unit (ACDU) providing single cells in >99% of the wells, and no wells with more than 1 cell. FACS analyses were performed on a BD LSRII (BD Biosciences). See Supplementary Table 1 for instrument configurations. Subsequent data analyses were performed with the FlowJo analysis software (TreeStar Inc). Cells used in cell sorting experiments were either unenriched or enriched for CD117 with MACS cell separation (Miltenyi Biotec) followed by Fc-block incubation and staining with anti-mouse antibodies (see Supplementary Table 2 for antibody information). Fluorescence-minus-one controls as well as negative populations were used as gate-setting controls.

Microarray analysis

Global gene expression analysis was performed on HSCs (Lin⁻Sca-1⁺c-Kit⁺Flt3⁻CD48⁻CD150⁺), IL-7Rα⁺LMPPs (Lin⁻Sca-1⁺c-Kit⁺Flt3^{hi}IL-7Rα⁺), adult and neonatal ETPs (Lin⁻CD4⁻CD8α⁻CD25⁻c-Kit^{hi}Flt3^{+/hi}), DN2 (Lin⁻CD44⁺CD25⁺c-Kit⁺), DN3 (Lin⁻CD44⁻CD25⁺), CLPs (Lin⁻CD19⁻B220⁻Sca-1^{low}c-Kit^{low}Flt3⁺IL-7Rα⁺Ly-6D⁻ and Pro B cells, CD19⁺B220⁺CD43⁺c-Kit⁺, from one week old mice. For all, except two populations, three individually sorted samples from different pools of mice were prepared. In the remaining two populations, two and four biological replicates were analyzed. Cells (1,600-2,000) were sorted directly into Trizol (Invitrogen) and the RNA extraction carried out as per manufacturer's instructions. Using the same total amount of input RNA, samples were amplified using the NuGEN kit WT-Ovation Pico RNA Amplifications System followed by the WT Ovation cDNA Biotin Module V2 for cDNA labeling (NuGEN) and fragmentation and finally hybridised to Affymetrix Mouse Genome 430 2.0 Arrays using standard protocols (Affymetrix) at the Stanford Protein and Nucleic Acid facility. Data were normalised using the Robust Multi-array Averages (RMA) method in the *affy* Bioconductor/R package. Subsequent data analysis (PCA, cluster analysis) was also carried out in R (see Supplementary Methods).

Single cell and quantitative PCR

Multiplex single-cell RT-PCR was performed as previously described³⁴ (see Supplementary Table 3) on single cells sorted from newborn ETPs. Multiplex quantitative real time PCR analysis was performed using BioMark 48.48 Dynamic Array platform (Fluidigm) and TaqMan Gene Expression Assays (Applied Biosystems) as previously described⁴⁶ on sorted populations. For each cell population, two biological replicates (25 cells/replicate) i.e. individually sorted samples from different mice, or single cells (20 cells/experiment) were prepared. See Supplementary Information and Supplementary Table 4 for TaqMan assays used.

In vitro cultures

GM and Mk lineage potentials were analyzed as previously described³² (see Supplementary Table 5). Mk potential was additionally evaluated using the Megacult collagen-based assay (StemCell Technologies). 200 ETPs or 100,000 unfractionated control BM cells were plated in Megacult-collagen media supplemented with growth factors (see Supplementary Table 5). Mk colonies were evaluated after 7 days with Acetylthiocholiniodide (Sigma) staining according to manufacturer's instructions.

For evaluation of erythroid potential 500-1,000 ETPs and 50 LSKFlt3⁻ control cells were seeded in complete methylcellulose (GF M3434; StemCell Technologies). Cultures were evaluated after 8-11 days using 2,7-diaminofluorene staining (DAF; Sigma) as previously described³⁴.

OP9 and OP9-DL1 stromal cells were kindly provided by Dr A. Cumano. To evaluate the individual GM, B, T and NK cell lineage potentials single ETPs were FACS sorted onto monolayers of OP9 or OP9-DL1 stromal cells as previously described (see Supplementary Table 5). Myeloid lineages formed on OP9 stroma co-cultures were evaluated after 6 days by morphology analysis of May-Grünwald (Sigma) and Giemsa (Fluka) stained slides.

To analyze combined lineage potentials, single cell deposition was performed (ensured as specified above), onto OP9 cells supplemented with hFLT3L, mScf and hIL-7. Approximately 54 hours later, the co-culture was split to a secondary layer of OP9 with the same condition to promote B cells and a layer of OP9-DL1 with cytokines (hFLT3L, mSCF, hIL-7, hIL-6, mIL-3, hG-CSF, mGM-CSF, hCSF-1) to promote T and myeloid cell differentiation. The timing of the first culture period was optimized to give the highest frequency possible for all lineages in the combined lineage potential assay. One week after transfer to the second culture, clones were evaluated for the presence of CD19⁺ B cells, CD4⁺CD8 α ⁺NK1.1⁻ and/or Thy1.2⁺CD25⁺NK1.1⁻ T cells and Mac-1⁺F4/80⁺ myeloid cells. Clones expressing Mac-1 and/or F4/80 were additionally evaluated by morphology. The timing of the readout was optimized to the earliest time-point where T cells could be detected, and at which time myeloid progeny would still be possible to detect. However, as the differentiation time is longer for lymphoid cells, the production of myeloid progeny is most likely underestimated.

All frequencies of single cell assays are expressed as total frequencies, that is relative to the total number of cells plated unless otherwise stated.

REFERENCES

1. Reya T, Morrison SJ, Clarke MF, Weissman IL. Stem cells, cancer, and cancer stem cells. *Nature*. 2001; 414:105–111. [PubMed: 11689955]
2. Katsura Y. Redefinition of lymphoid progenitors. *Nat Rev Immunol*. 2002; 2:127–132. [PubMed: 11910894]
3. Donskoy E, Goldschneider I. Thymocytopoiesis is maintained by blood-borne precursors throughout postnatal life. A study in parabiotic mice. *J Immunol*. 1992; 148:1604–1612. [PubMed: 1347301]
4. Scollay R, Smith J, Stauffer V. Dynamics of early T cells: prothymocyte migration and proliferation in the adult mouse thymus. *Immunol Rev*. 1986; 91:129–157. [PubMed: 3525392]
5. Allman D, et al. Thymopoiesis independent of common lymphoid progenitors. *Nat Immunol*. 2003; 4:168–174. [PubMed: 12514733]
6. Benz C, Bleul CC. A multipotent precursor in the thymus maps to the branching point of the T versus B lineage decision. *J Exp Med*. 2005; 202:21–31. [PubMed: 15983065]

7. Bhandoola A, von Boehmer H, Petrie HT, Zuniga-Pflucker JC. Commitment and developmental potential of extrathymic and intrathymic T cell precursors: plenty to choose from. *Immunity*. 2007; 26:678–689. [PubMed: 17582341]
8. Wada H, et al. Adult T-cell progenitors retain myeloid potential. *Nature*. 2008; 452:768–772. [PubMed: 18401412]
9. Bell JJ, Bhandoola A. The earliest thymic progenitors for T cells possess myeloid lineage potential. *Nature*. 2008; 452:764–767. [PubMed: 18401411]
10. Desanti GE, et al. Clonal analysis reveals uniformity in the molecular profile and lineage potential of CCR9(+) and CCR9(-) thymus-settling progenitors. *Journal of immunology*. 2011; 186:5227–5235. doi:10.4049/jimmunol.1002686. [PubMed: 21421850]
11. Masuda K, et al. Thymic anlage is colonized by progenitors restricted to T, NK, and dendritic cell lineages. *J Immunol*. 2005; 174:2525–2532. [PubMed: 15728458]
12. Kondo M, Weissman IL, Akashi K. Identification of clonogenic common lymphoid progenitors in mouse bone marrow. *Cell*. 1997; 91:661–672. [PubMed: 9393859]
13. Adolfsson J, et al. Identification of Flt3+ lympho-myeloid stem cells lacking erythro megakaryocytic potential a revised road map for adult blood lineage commitment. *Cell*. 2005; 121:295–306. [PubMed: 15851035]
14. Orkin SH, Zon LI. Hematopoiesis: an evolving paradigm for stem cell biology. *Cell*. 2008; 132:631–644. [PubMed: 18295580]
15. Luc S, Buza-Vidas N, Jacobsen SE. Delineating the cellular pathways of hematopoietic lineage commitment. *Semin Immunol*. 2008; 20:213–220. [PubMed: 18752972]
16. Doulatov S, et al. Revised map of the human progenitor hierarchy shows the origin of macrophages and dendritic cells in early lymphoid development. *Nat Immunol*. 2010; 11:585–593. [PubMed: 20543838]
17. Goardon N, et al. Coexistence of LMPP-like and GMP-like leukemia stem cells in acute myeloid leukemia. *Cancer Cell*. 2011; 19:138–152. doi:10.1016/j.ccr.2010.12.012. [PubMed: 21251617]
18. Taub DD, Longo DL. Insights into thymic aging and regeneration. *Immunol Rev*. 2005; 205:72–93. [PubMed: 15882346]
19. Ceredig R, Bosco N, Rolink AG. The B lineage potential of thymus settling progenitors is critically dependent on mouse age. *Eur J Immunol*. 2007; 37:830–837. [PubMed: 17295389]
20. Sambandam A, et al. Notch signaling controls the generation and differentiation of early T lineage progenitors. *Nat Immunol*. 2005; 6:663–670. [PubMed: 15951813]
21. Radtke F, et al. Deficient T cell fate specification in mice with an induced inactivation of Notch1. *Immunity*. 1999; 10:547–558. [PubMed: 10367900]
22. Feyerabend TB, et al. Deletion of Notch1 converts pro-T cells to dendritic cells and promotes thymic B cells by cell-extrinsic and cell-intrinsic mechanisms. *Immunity*. 2009; 30:67–79. [PubMed: 19110448]
23. Hobeika E, et al. Testing gene function early in the B cell lineage in mb1-cre mice. *Proc Natl Acad Sci U S A*. 2006; 103:13789–13794. [PubMed: 16940357]
24. Srinivas S, et al. Cre reporter strains produced by targeted insertion of EYFP and ECFP into the ROSA26 locus. *BMC Dev Biol*. 2001; 1:4. [PubMed: 11299042]
25. Campbell KJ, et al. Elevated Mcl-1 perturbs lymphopoiesis, promotes transformation of hematopoietic stem/progenitor cells, and enhances drug resistance. *Blood*. 2010; 116:3197–3207. [PubMed: 20631380]
26. Kiel MJ, et al. SLAM family receptors distinguish hematopoietic stem and progenitor cells and reveal endothelial niches for stem cells. *Cell*. 2005; 121:1109–1121. [PubMed: 15989959]
27. Balazs AB, Fabian AJ, Esmon CT, Mulligan RC. Endothelial protein C receptor (CD201) explicitly identifies hematopoietic stem cells in murine bone marrow. *Blood*. 2006; 107:2317–2321. [PubMed: 16304059]
28. Solar GP, et al. Role of c-mpl in early hematopoiesis. *Blood*. 1998; 92:4–10. [PubMed: 9639492]
29. Igarashi H, Gregory SC, Yokota T, Sakaguchi N, Kincade PW. Transcription from the RAG1 locus marks the earliest lymphocyte progenitors in bone marrow. *Immunity*. 2002; 17:117–130. [PubMed: 12196284]

30. Zlotoff DA, et al. CCR7 and CCR9 together recruit hematopoietic progenitors to the adult thymus. *Blood*. 2010; 115:1897–1905. [PubMed: 19965655]
31. Krueger A, Willenzon S, Lyszkiewicz M, Kremmer E, Forster R. CC chemokine receptor (CCR) 7 and 9 double-deficient hematopoietic progenitors are severely impaired in seeding the adult thymus. *Blood*. 2010
32. Luc S, et al. Downregulation of Mpl marks the transition to lymphoid-primed multipotent progenitors with gradual loss of granulocyte-monocyte potential. *Blood*. 2008
33. Godfrey DI, Kennedy J, Suda T, Zlotnik A. A developmental pathway involving four phenotypically and functionally distinct subsets of CD3-CD4-CD8-triple-negative adult mouse thymocytes defined by CD44 and CD25 expression. *J Immunol*. 1993; 150:4244–4252. [PubMed: 8387091]
34. Mansson R, et al. Molecular evidence for hierarchical transcriptional lineage priming in fetal and adult stem cells and multipotent progenitors. *Immunity*. 2007; 26:407–419. [PubMed: 17433729]
35. Ehrlich LI, Serwold T, Weissman IL. In vitro assays misrepresent in vivo lineage potentials of murine lymphoid progenitors. *Blood*. 2011; 117:2618–2624. [PubMed: 21163922]
36. Inlay MA, et al. Ly6d marks the earliest stage of B-cell specification and identifies the branchpoint between B-cell and T-cell development. *Genes Dev*. 2009; 23:2376–2381. [PubMed: 19833765]
37. Pronk CJH, et al. Elucidation of the Phenotypic, Functional, and Molecular Topography of a Myeloerythroid Progenitor Cell Hierarchy. *Cell Stem Cell*. 2007; 1:428–442. [PubMed: 18371379]
38. David-Fung ES, et al. Transcription factor expression dynamics of early T-lymphocyte specification and commitment. *Dev Biol*. 2009; 325:444–467. [PubMed: 19013443]
39. Wendorff AA, et al. Hes1 is a critical but context-dependent mediator of canonical Notch signaling in lymphocyte development and transformation. *Immunity*. 2010; 33:671–684. [PubMed: 21093323]
40. Petrie HT, Kincade PW. Many roads, one destination for T cell progenitors. *J Exp Med*. 2005; 202:11–13. [PubMed: 15983068]
41. Chi AW, et al. Identification of Flt3CD150 myeloid progenitors in adult mouse bone marrow that harbor T lymphoid developmental potential. *Blood*. 2011; 118:2723–2732. doi:10.1182/blood-2010-09-309989. [PubMed: 21791413]
42. Schlenner SM, et al. Fate Mapping Reveals Separate Origins of T Cells and Myeloid Lineages in the Thymus. *Immunity*. 2010; 32:426–436. [PubMed: 20303297]
43. Coustan-Smith E, et al. Early T-cell precursor leukaemia: a subtype of very high-risk acute lymphoblastic leukaemia. *Lancet Oncol*. 2009; 10:147–156. [PubMed: 19147408]
44. Metzler M, et al. A conditional model of MLL-AF4 B-cell tumorigenesis using invertebrate technology. *Oncogene*. 2006; 25:3093–3103. doi:10.1038/sj.onc.1209636. [PubMed: 16607274]
45. Faust N, Varas F, Kelly LM, Heck S, Graf T. Insertion of enhanced green fluorescent protein into the lysozyme gene creates mice with green fluorescent granulocytes and macrophages. *Blood*. 2000; 96:719–726. [PubMed: 10887140]
46. Tehrani R, et al. Persistent malignant stem cells in del(5q) myelodysplasia in remission. *N Engl J Med*. 2010; 363:1025–1037. [PubMed: 20825315]





Figure 1.

ETPs are multipotent lympho-myeloid restricted progenitors. (a) Flow cytometry profiles and gating strategies for the detection of $\text{Lin}^- \text{CD4}^- \text{CD8}\alpha^- \text{CD25}^- \text{c-Kit}^{\text{hi}} \text{Flt3}^{\text{hi}}$ ETPs from young adult mice (4–6 weeks). Numbers in plots indicate percent ETPs among total thymocytes. DAPI, DNA-intercalating dye; FSC, forward scatter; SSC, side scatter; -W, width; -H, height. (b) Frequency of B cell potential of cultures seeded with a single $\text{Lin}^- \text{Sca-1}^+ \text{c-Kit}^+ \text{Flt3}^{\text{hi}}$ bone marrow cell (LMPP; $n = 320$); a single Flt3^{hi} ETP ($n = 73$ cells) or ten Flt3^- ETPs ($n = 960$ cells); or other DN thymocyte progenitor populations (DN1–DN4; $n = 2,400$ cells (seeded with 100 cells per well)), all from adult mice. (c) Flow cytometry profiles and gating strategies as in a, for cells from newborn mice (1 d). (d) Frequency of B cell potential as in b, for cultures of cells from newborn mice, seeded as single Flt3^{hi} ETPs ($n = 348$ cells) or single Flt3^- ETPs ($n = 210$ cells), and other DN thymocyte progenitor populations seeded at 100 cells per culture ($n =$

4,200–6,000 cells). (e) Expression of enhanced yellow fluorescent protein (eYFP) in ETPs from neonatal mice ($n = 4$) expressing Cre from the Cd79a promoter. (f) Frequency of cells with B cell potential (B; $n = 348$ cells), T cell potential (T; $n = 204$ cells), natural killer cell potential (NK; $n = 48$ cells), GM potential (grown in liquid (GM(L); $n = 600$ cells) or on stroma (GM(S); $n = 64$ cells)), megakaryocyte potential (grown in liquid (Mk(L); $n = 1,080$ cells) or on semisolid support (Mk(SS); $n = 6$; 200 cells per replicate)) or erythroid potential (E; $n = 8$; 500–1,000 cells per replicate) among Flt3⁺ ETPs from neonatal mice (positive controls, Supplementary Fig. 5). (g) Expression of myeloid markers Mac-1, Gr-1 and lysozyme M (reported as eGFP expression; left and middle), and morphological analysis (right) of sorted granulocytes (top) and monocytes (bottom) from cultured Flt3⁺ ETPs from neonatal mice. Scale bars, 5 μ M. (h) Quantitative analysis of the expression of genes associated with lymphoid cells, myeloid cells and megakaryocytes–erythroid cells by purified Flt3⁺ ETPs from newborn mice ($n = 6$; 25 cells per replicate); results are presented relative to the expression of Hprt (encoding hypoxanthine guanine phosphoribosyl transferase). *, 0.001 (below detection limit). Data are representative of four experiments (a); fourteen experiments (c); seven (b) or sixteen (d) experiments (Flt3^{hi} ETPs); sixteen experiments (bone marrow; b); four experiments (Flt3⁻ ETPs (b) and other DN populations (b,d)); ten experiments (Flt3⁻ ETPs; d); one experiment (e); two to sixteen experiments (f); one experiment (g); or two experiments (h; mean and s.e.m. in b,d,f; average and s.d. of six replicates in h).

**Figure 2.**

ETPs possess combined T, B and GM lineage potentials. **(a)** Single cell gene expression analysis of lymphoid, myeloid and M κ E genes in purified newborn Flt3^{hi} ETPs. Mean (\pm s.d.) frequency of cells expressing specified genes among cells positive for *c-Kit* gene expression (96-98% of total cells) (left; $n=176$ from 2 exp). The frequency (\pm s.d.) of ETPs with combined lymphoid-GM lineage transcriptional priming (right) based on co-expression of one or more genes for the lymphoid program (*Il7r*, sterile IgH), myeloid/GM program (*Csf1r*, *Mpo*) but not the M κ E program (*Gata1*, *Epor*). **(b)** A representative clone from a single newborn WT Flt3⁺ETP cell with combined T, B and myeloid lineage potential determined by FACS and morphology analysis (myeloid, black arrowhead, lymphocyte, white arrowhead). **(c,d)** Cloning frequencies (left), of ETPs generating CD45⁺ cells (white bars) and CD45⁺ cells with a definitive lineage readout (grey bars). Lineage distribution of clones (right) from **(c)** single WT ($n=132$ from 3 exp) and **(d)** single *vavP-mcl-1* ETPs ($n=167$ from 2 exp).




Figure 3.

Absence of pluripotent HSCs in newborn thymus. **(a)** Flow cytometry profiles of CD150, CD201 and Mpl stem cell marker expression in Lin⁻CD4⁻CD8^α⁻CD25⁻c-Kit^{hi} ETPs in newborn mice ($n=2$ exp). **(b)** Flow cytometry profiles of CCR9 and *Rag1* GFP co-expression in ETPs, LMPPs and HSCs ($n=2$ exp). **(c-h)** Peripheral blood analyses showing mean (\pm s.d.) thymocyte contribution to T, B, myeloid and platelet (*Vwf*GFP⁺) lineages at 3-4 weeks (**c**; $n=9$, **d**; $n=7$), 16 weeks (**e**; $n=8$, **f**; $n=7$) and 12 weeks (**g**; $n=8$, **h**; $n=8$) after intravenous or intrafemoral competitive transplantation of total or CD4⁺ and CD8⁺ depleted neonatal thymocytes from *Vwf*GFP or WT mice. The frequency of reconstituted animals is indicated above each lineage.




Figure 4.

ETPs cluster closer to candidate TSPs in the BM than other thymic progenitors. **(a,b)** Two- and three-dimensional principal component analyses of normalized global gene expression profiles of **(a)** purified HSC, LMPP, ETP, DN2 and DN3 from neonatal mice ($n=3$) and **(b)** purified neonatal HSC ($n=3$), LMPP ($n=3$), CLP ($n=4$), ETP ($n=3$), DN3 ($n=3$), ProB ($n=3$) and adult ETP ($n=2$). Each symbol represents a separate biological sample (sorted from different pools of mice). Euclidean distances between average x and y values for each population measured in the first two principal components are shown in panel **(a)**.




Figure 5. ETPs, LMPPs and CLPs have closely related T and myeloid lineage transcriptional profiles. (a,b) Heatmap representation of T (a) and GM (b) lineage-affiliated gene expression represented as normalized median expression values from purified neonatal HSC ($n=6$), LMPP ($n=6$), CLP ($n=4$), ETP ($n=6$), DN2 ($n=3$), DN3 ($n=6$) and adult ETP ($n=2$) populations. Gene lists were established as described in the **Online Methods** section and Supplementary Methods).




Figure 6. Quantitative expression analysis of lymphoid and myeloid genes in neonatal ETPs. Quantitative gene expression analysis showing lymphoid and myeloid genes in purified neonatal HSC ($n=6$), LMPP ($n=4$), ETP ($n=6$), DN2 ($n=6$) and DN3 ($n=6$) populations (25 cells/replicate). Average expression levels (\pm s.d.). (*) 0.001, below detection.