



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Alternative farrowing systems: design criteria for farrowing systems based on the biological needs of sows and piglets

Citation for published version:

Baxter, EM, Lawrence, AB & Edwards, SA 2011, 'Alternative farrowing systems: design criteria for farrowing systems based on the biological needs of sows and piglets' *Animal*, vol 5, no. 4, pp. 580-600., 10.1017/S1751731110002272

Digital Object Identifier (DOI):

[10.1017/S1751731110002272](https://doi.org/10.1017/S1751731110002272)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher final version (usually the publisher pdf)

Published In:

Animal

Publisher Rights Statement:

Copyright 2010 The Animal Consortium

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Alternative farrowing systems: design criteria for farrowing systems based on the biological needs of sows and piglets

E. M. Baxter^{1†}, A. B. Lawrence¹ and S. A. Edwards²

¹Animal Behaviour and Welfare, Sustainable Livestock Systems, Scottish Agricultural College, West Mains Road, Edinburgh, EH9 3JG, UK; ²Newcastle University, School of Agriculture, Food and Rural Development, Agriculture Building, Newcastle upon Tyne NE1 7RU, UK

(Received 16 December 2009; Accepted 30 July 2010; First published online 16 November 2010)

The construction of a suitable farrowing environment is a continuing dilemma: the piglet's needs must be matched with those of the sow and the farmer during the main phases that constitute farrowing: nest building, parturition and lactation. Difficulties exist in resolving the various conflicts of interest between and within these three parties (e.g. sow v. farmer: space needed for nest building v. space needed to maximise the amount of farrowing accommodation, or sow v. sow: ensuring the survival of the current litter v. maintaining condition for future litters). Thus, the challenge is to resolve these conflicts and design a system that maximises sow and piglet welfare while maintaining an economically efficient and sustainable enterprise. In order to successfully design a farrowing and lactation environment, it is necessary to consider the biological needs of both the sow and her litter. The natural behaviour of the sow has been well documented and very little variation exists between reports of peri-parturient behaviour observed in extensively kept domestic sows and their wild counterparts. The failure for domestication to significantly alter these behavioural patterns provides evidence that they are biologically significant and that the commercial farrowing environment should attempt to accommodate this behavioural repertoire. In addition, the behavioural needs of the piglets, as well as the physiological needs of both sows and their offspring should be considered. This article aims to review the considerable body of literature detailing the behavioural repertoire of sows and their offspring during the different phases of farrowing, and the accompanying physiological processes. The focus is on identifying biological needs of the animals involved in order to synthesise the appropriate design criteria for farrowing and lactation systems, which should optimise both welfare and animal production.

Keywords: pig, nest building, farrowing, lactation, welfare

Implications

Designing suitable farrowing and lactation environments that maximise both sow and piglet welfare, while maintaining economically efficient and sustainable enterprises, is a continuing challenge. It is not unreasonable to consider designing housing systems based on the biological specifications of the animals involved. Therefore, this review documents the considerable amount of literature describing biological needs of sows and piglets. It demonstrates the evolutionary significance and thus the function of certain behavioural patterns that continue to be displayed by domestic animals. Identification of biological needs allows synthesis of appropriate design criteria for farrowing and lactation systems that optimise both welfare and animal production.

Introduction

It is not unreasonable to suggest that agricultural practices in livestock farming systems should be based on the biological needs of the animals involved. However, this is rarely the case, with different managerial constraints (e.g. finance, labour and space) often resulting in a compromise with the animals' biological needs. Perhaps the most prevalent illustration of this occurs during the farrowing and lactation period for the domestic sow. During this period the majority of domestic sows (approximately 70% in United Kingdom, BPEX, 2004; 95% in EU and 83% in United States, Johnson and Marchant-Forde, 2009) are housed in farrowing crates, which are behaviourally and physically restrictive and raise serious welfare concerns. However, the farrowing crate was initially developed to reduce building space requirements, provide safe working access to the piglets, reduce labour input and improve piglet survival (Edwards and Fraser, 1997).

† E-mail: Emma.Baxter@sac.ac.uk

Piglet survival is one of the most important outcomes of the farrowing period for the three main stakeholders (sow, piglets and farmer); the piglets obviously need to survive, their survival is the key to profitability for the farmer and the sow can measure her inclusive fitness in terms of reproductive success. To facilitate maximum survival there must be co-ordinated expression between behaviours of the sow and piglets, and their interaction with the environment. These interactions are complex, involving a series of conflicts between parent and offspring who have different needs at different stages of farrowing. The conflicts further increase as the needs of the farmer are taken into account. As a result, designing a suitable farrowing environment is a continuing challenge and one that has been the subject of sustained scientific interest (Edwards, 1996; Arey, 1997; Edwards and Fraser, 1997; Wechsler and Weber, 2007).

Designing a suitable housing system based on the biological specifications of the animals concerned involves asking questions about their needs, both behavioural and physiological. The latter include very obvious requirements including provision of appropriate nutrition and climate to ensure maximum growth, development, health and productivity. The former has received a great deal of attention from ethologists, animal welfare scientists and, increasingly, by a concerned public. This has occurred partly because the consideration of 'behavioural needs' when designing housing systems for livestock is often over-looked within the industry, despite many recommendations by scientists to adjust environments to meet these needs (Brambell, 1965; Farm Animal Welfare Council, 1992; Edwards, 1996; European Food Safety Authority, 2007).

Failure to recognise the importance of behavioural needs may actually be counter-productive in terms of maximising animal productivity, since it has long been accepted that the performance of species-typical behaviour contributes to the biological fitness of an animal (Hamilton, 1964a and 1964b). Agriculture has exploited the animals' natural ability to adapt to their environment (Baxter, 1983). Domestication and intensive breeding strategies have resulted in physiological changes in the pig, which have capitalised on the heritability of traits such as litter size and growth rate, in order to have an 'improved' and efficient farm animal. However, the influence domestication has had on the behavioural needs of the pig is less clear, with certain behaviours apparently robust to the changes brought about by intensive selection (Stolba and Wood-Gush, 1980 and 1981). Nest building behaviour in the peri-parturient sow is an example of a behavioural pattern stubbornly being attempted despite, in the majority of intensive agricultural practices, an environment that precludes its natural performance and appears to render its original function unnecessary. The motivation to perform behaviours that on the surface appear unnecessary in a production environment suggests that they still have some biological significance to the animal.

The concept of 'behavioural needs' has been the subject of much debate and detailed scientific analysis (e.g. Dawkins, 1977; Baxter, 1983; Hughes and Duncan, 1988; Jensen and Toates, 1993). For the purpose of this review, the term 'behavioural need' is generally used to describe the need to

perform a specific behaviour pattern whatever the environment and even if the physiological needs which the behaviour serves are fulfilled in other ways (Baxter, 1983; Jensen and Toates, 1993). Possible behavioural needs can be catalogued by identifying species behaviours with important survival value which occur spontaneously in all environments, and by measuring the preferences of animals (Hughes and Black 1973; Dawkins, 1977). However to demonstrate a true need, it should be shown that failure to meet this need results in a compromise in welfare by demonstrating negative consequences when these actions cannot be performed satisfactorily (e.g. performance of abnormal behaviour, physiological stress response or increased incidence of pathology).

Therefore, this review describes which behavioural patterns are displayed by pigs in natural and semi-natural environments during the various phases of farrowing, and then determines whether they have endured domestication and what their functional significance is. To discuss the motivational framework contributing to these behaviours, the physiological processes that accompany them are explored. Further experimental evidence of true biological needs is reviewed, leading to the synthesis of recommended design criteria to accommodate them in a farrowing environment. The next critical step in an overall design process is to address the challenges of accommodating these criteria while taking account of the needs of the farmer. However, the focus of this review will remain the needs of the animals themselves, with subsequent work considering the issues for the farmer.

Nest building phase: nest-site seeking, isolation and building

Sow behaviour and physiology

Nest building is a well-documented behavioural pattern in pigs (Gundlach, 1968; Frädriich, 1974; Stolba and Wood-Gush, 1984; Jensen, 1986), with a recent review highlighting its continued interest in animal science (Wischner *et al.*, 2009). It is an innate behaviour, unaltered by domestication (Jensen, 2002) and displayed by all members of the family Suidae, with the exception of the Common warthog (*Phacochoerus africanus*) which tends to burrow in order to give birth to the young (MacDonald, 2000). An increase in locomotory activity occurs approximately 2 to 3 days before parturition. The sow seeks isolation from the herd and wanders away in search of a suitable nest site. Jensen (1986) reported that expectant sows travelled 2.5 to 6.5 km during the pre-parturient period in search of suitable sites, often building 'mock-nests' during this search, the functional significance of which is still unknown. This heightened period of activity is also evident in peri-parturient sows kept indoors, with reports of expectant sows in 5 m² pens, 'travelling' on average 30 km (Baxter, 1991). Chosen nest sites are often isolated and at least partially enclosed, affording sows a protected nest site while still being able to maintain vigilance for potential approaching threats (Stolba and Wood-Gush, 1984). In addition, nest-site location is considered with excretory and feeding locations taken into account; observations of wild boar report that

dung is found in concentrated, specific areas 'some' metres away from any nest sites (Wiepkema, 1986). Concentrated areas for dunging may serve some territorial function, but also it is important that these areas are away from nest sites to control the risk of disease. Once chosen, a concave depression is created by hollowing out the ground, or by rooting up soil and vegetation into a pile and then rooting a central depression (Hafez *et al.*, 1962; Graves, 1984). Hollowing out the ground involves digging and rooting behaviours, patterns also observed in domestic sows kept in farrowing crates though re-directed towards floors, bars and drinkers (Lawrence *et al.*, 1994). After hollowing out a nest site, branches are gathered to border the hollow before collecting and arranging grass and leaves to line the nest.

Accompanying nest-building behaviour is a range of endocrine changes that have been the subject of a recent review (Algers and Uvnäs-Moberg, 2007) and therefore will only briefly be outlined here. Widowski *et al.* (1990) demonstrated, by injection of prostaglandin (PGF₂α) into *pre-partum* gilts, which changes in plasma concentrations of prolactin elicited nest-building activities. Further investigations (Castrén *et al.*, 1993a; Damm *et al.*, 2002) have correlated specific nest-building behaviours with specific hormones; time spent carrying and depositing substrate correlated positively with prolactin and progesterone, but negatively with somatostatin concentrations in plasma (Castrén *et al.*, 1993a). Nosing and arranging this substrate correlated negatively with plasma oxytocin (Damm *et al.*, 2002). Therefore, performance and completion of nest building appear to be influenced by both internal and external factors (Jensen and Toates, 1993). There must be a 'switch-off' mechanism because performance of nest building is no longer functional during parturition. Baxter (1983) suggested that udder comfort switches off nesting behaviour, although provision of a comfortable substratum does not abolish the behaviour (Arey *et al.*, 1991). In general, nest-building behaviour abates approximately 4 h before parturition. Oxytocin levels have already started to increase 6 h before parturition (Algers and Uvnäs-Moberg, 2007) and the sow enters a 'quiet phase' before parturition begins.

Functional importance of the nest-building phase

From the detailed descriptions of site choice and construction, as outlined above, the main functions of the nest appear to be protection for piglets from inclement weather conditions and camouflage from potential predators. Thus, its function for wild populations of pigs is obvious, but this does not explain why the action of building a nest should remain so strong in our domestic population of indoor pigs, where the risks from climatic extremes and predation are obsolete. It is possible that the behaviour has persisted simply because of lack of selection against its occurrence, or lack of genetic variation in the control mechanisms within the domesticated population. However, Jensen and Gustafsson (1997) theorized that costly behaviour strategies which are optimal in wild conditions may pay off less in the domesticated situation, and therefore have a negative cost-benefit balance promoting suppression. Nest building is just such a costly

behavioural pattern, involving increased activity and complex construction for as much as 24 h before further energy expenditure during farrowing and lactation. When exploring the importance of behaviours from a life history point of view, a functional aspect of behaviour is based on the contribution that it makes to fitness, as measured by reproductive success (Daan and Tinbergen, 1997). Therefore, we can hypothesise that it is not just the nest that contributes to reproductive success by protecting the offspring, but also the motivation to build and the performance of building that nest. Feedback from building and completing a nest can affect neuro-endocrine regulation of maternal behaviour (Castrén *et al.*, 1993a; Damm *et al.*, 2003; Pedersen *et al.*, 2003; Algers and Uvnäs-Moberg, 2007) during farrowing, with evidence that positive parturient maternal behaviour is influenced by the satisfaction of the nest-building phase (Arey *et al.*, 1991; Jensen, 1993; Damm *et al.*, 2003; Pedersen *et al.*, 2003). For example, several authors have proposed a link between high nest-building activity and reduced risk of crushing (Andersen *et al.*, 2005; Pedersen *et al.*, 2006). The more complete and functional the nest is, the more likely the sow is to end nest building and begin the more somnolent farrowing phase. Sows that continue to display nest-building activity during farrowing are increasing the risk of crushing for their piglets, as well as prolonging farrowing duration with increased risk of piglets being born dead (Baxter *et al.*, 2008). Duration of farrowing has been shown to be longer in sows housed in crates rather than in pens in some (Thodberg *et al.*, 2002; Oliviero *et al.*, 2006), although not all, studies (Cronin *et al.*, 1991; Fraser *et al.*, 1997). It is possible that the inadequacy of nest-building behaviour in crated environments may influence the duration of farrowing and therefore the incidence of stillbirths. The lack of space and substrate when sows are kept in standard farrowing crates precludes feedback from nest-building behaviour and has been shown to constitute a stress for the sow (Lawrence *et al.*, 1994; Jarvis *et al.*, 1997 and 2001; Damm *et al.*, 2003). Further work dissociating the effects of space and substrate demonstrated that space restriction *per se* induced elevated hypothalamic pituitary adrenal (HPA) activity which is deemed indicative of physiological stress (Jarvis *et al.*, 2002). There is also evidence that savaging behaviour is more prevalent in animals housed in farrowing crates (Cronin *et al.*, 1996; Jarvis *et al.*, 1998) and savaging sows are more restless and hyper-responsive to their piglets and environment (Ahlström *et al.*, 2002) than non-savaging sows. It has been proposed that because the restrictive nature of this environment is inadequate to satisfy nest building, the need to nest build may not be 'switched-off'. The sow is thus ill prepared for the somnolent phase of parturition, with potentially fatal consequences for the piglets. Thus despite the absence of offspring at this stage, nest building is arguably a critical phase for piglet survival and consequently remains an evolutionary important behaviour.

Needs-based design criteria for the nest-building phase

From the evidence presented above, it is clear that in order for the species behaviour patterns of the sow to be met

Table 1 Summary of design recommendations to meet the biological needs of sows during farrowing and lactation

| Component of system | Biological specification | Recommendations |
|---------------------|--|--|
| Space* | Increased activity for nest-site seeking | 4.9 m ² (minimum recommendation based on combining two areas the sow can turn around in – see below) ^b |
| | Hygiene – dunging space | Separate area from nest and feed sites ^a . Interaction with air temperature and floor properties |
| | Feeding and foraging | Separate area from nest and dung sites ^a |
| | Turn-around nest space for piglet inspection and gathering behaviour | Floor space = 2.44 m ² , planar space = 3.17 m ² (minimum) ^a . Further research needed for unimpeded turning by the sow |
| | Lateral lying and parturition | 2.79 m ² (minimum) ^a |
| | Thermal comfort via posture changes | 2.44 m ² allows at a minimum getting up and lying down posture changes ^a |
| | Nest-departure | Separate area from nest site ^b |
| | Social contact | Full contact only recommended when mixing familiar groups ^b . Space dependent on group size and needs 'clever' design to accommodate avoidance behaviours ^a . Further research needed on 'personal space' for group suckling behaviour |
| | Gradual separation from piglets and controlled nursing | Separate space unattractive to the piglets ^b . Interaction with air temperature and floor properties |
| Substrate | Nest-building – carrying and manipulating | 2 kg long stemmed straw (minimum) ^a accessible to the sow. Further research needed for alternative substrates with similar properties |
| | Complete nest | 2 kg long stemmed straw and branches (minimum) ^a |
| | Udder comfort | Further research needed, interaction with floor properties |
| | Thermal comfort during nest building | Further research needed to determine sow temperature preferences during nest building |
| | Thermal comfort during parturition | Deep bedding 10 to 12 cm ^a , interactions with floor properties and ambient temperature |
| | Foraging material | Further research needed on suitable materials and required amount |
| Walls | Enclosure/isolation of nest | Three solid-sided walls (cul-de-sac) ^b |
| | Darkness | Dark walls to simulate burrow ^b |
| | Visual and physical contact with non-litter pigs | Vertical barred area outside nest with void wide enough to allow at least nasal contact between pigs for social integration (minimum) ^b |
| | Supported posture changes | Solid sloped or vertical walls ^a |
| | Lack of disturbance | Further research needed to determine benefits of providing a quiet area |
| Flooring | Nest building – digging, rooting and hollowing | Malleable (e.g. earthen) or solid to accommodate deep substrate ^a |
| | Nest building and parturition | Solid to accommodate substrate ^a |
| | Thermal comfort during nest building | Further research needed to determine sow temperature preferences during nest building. Temperature differentials in separate areas allowing choice ^b |
| | Thermal comfort during parturition | High thermal resistance, for example, rubber matting or deep substrate. Temperature differentials in separate areas ^a . Further research needed for recommendation of localised temperature |
| | Thermal comfort during lactation | Low thermal resistance, for example, metal. Temperature differentials in separate areas ^b |
| | Physical comfort – avoiding injury, promoting suckling behaviour | Non-slip surface ^a Minimal abrasiveness, for example, rubber matting or plastic-coated metal ^a Solid to avoid teat injuries ^a |
| | Hygiene | Slatted area ^a Gradation of floor with slope away from lying area ^b . For example, 10% slope for drainage |
| | | High feed intake |
| General | Thermal comfort | Ambient temperature 12°C to 22°C ^a , interactions with substrate and flooring |
| | High feed intake | See space and thermal comfort |

^aRecommendations derived from one or more specific references.

^bRecommendations based on synthesising and interpreting the literature reviewed.

*Space calculations based on an 'improved' sow weighing 350 kg.

during this phase, the following criteria should be considered for design: space, enclosure, substrate and suitable flooring. These criteria need elaboration and attempts are made to quantify and summarise them (Table 1).

Space

It is assumed when determining spatial needs for animals, that they are geometrically similar and that the minimum amount of space required is determined by the animal's size

and body shape (Baxter and Schwaller, 1983). This 'body space' can then be added to, to accommodate behavioural requirements. During the nesting phase, space is required to allow the animal to increase its activity, to 'seek' a nest site, build the nest and to have separate nesting, excretory and feeding sites.

Sows normally circle around when constructing the nest, and this may define the minimum space needed for the nest itself. In an experiment looking at the space preferences in different farrowing crates (Phillips *et al.*, 1992), sows consistently preferred the largest width, which permitted them to turn around. This, together with the previously cited evidence for elevated HPA axis activity when turning is prevented, indicates that the minimum space criterion during this phase should allow this behaviour. Quantification of the required space can be derived from the comprehensive work looking at spatial needs of sows in confinement by Baxter and Schwaller (1983) and Petherick (1983a), who used the principle body dimensions of length, height and width to determine the area occupied by any weight of pig. Using these estimates, Baxter and Schwaller (1983) determined minimum space requirements for the 95th percentile sow weight that they measured (238 kg): 1.90 m² (2334 mm long × 810 mm wide × 870 mm high; see Figure 1). The genetic 'improvement' in mature breeding females has resulted in body weights of many commercial multiparous sows between 300 and 350 kg without undue fatness (Whittemore and Kyriazakis, 2006). Such an observation has also been made by other researchers (Moustsen *et al.*, 2004) and in response to Danish sows steadily increasing in size over the last 20 years without farrowing crates changing to accommodate this increase, these authors took detailed measurements of sows entering and leaving farrowing crates in order to provide an improved biological basis for farrowing crate dimensions. The authors provided their minimum recommendations based on the dimensions of the 95th percentile of the population of sows measured, thus hoping to accommodate the largest of the population. Sows in the 95th percentile weighed 352 kg, were 2.00 m long, 0.47 m wide, 0.95 m tall and measured 0.71 m from the middle of the sow's udder to her back (i.e. breadth). From these data, it is evident that minimum space recommendations should be revised for all forms of accommodation by re-calculation based on the 95th percentile of current average weights. For the biological needs of farrowing sows, the above methods described (Baxter and Schwaller, 1983) were used, and assuming linear relationships, Baxter and Schwaller's graph (Figure 1) was extrapolated and it was estimated that an 'improved' sow of 350 kg needs at least 2.44 m² of floor space to get up and lie down comfortably (2654 mm long × 923 mm wide and 968 mm high; Table 1). By adding to the width of this space, a sow would be able to turn around (Robertson *et al.*, 1972). Curtis *et al.* (1989), using Robertson *et al.*'s (1972) data proposed the allometric equation of $15.4W^{0.33}$ (where W = body weight) to determine width for sows to turn around in gestation crates. However, Robertson *et al.*'s calculations were determined based on designing a stall that a sow could not




Figure 1 Overall, space requirements of sows (dimensions of pen) expressed over weight (W = body weight). Extrapolated from Baxter and Schwaller (1983).

turn around in and therefore involved the animal squeezing against the stall or crate sides to achieve a turning circle. Without conducting detailed experiments, similar to those undertaken by Baxter and Schwaller (1983), the additional width required for turning around unimpeded by a late gestation sow can only be estimated. Lou and Hurnik (1994) designed an Ellipsoid crate that allowed pigs to turn around (average weight 206 kg) with a floor space of 1.80 m² and a planar space (space at sow's shoulder height) of 2.51 m². If we were to over estimate width needed to turn around and assume the sow needed at least its body length in all planes to turn comfortably, then according to Petherick's calculation (1983a), Lou and Hurnik's 206 kg sow needs a crate planar width of 1.74 m to accommodate a comfortable 360° turn. In fact, 1.20 m was provided and sows were able to turn around. Adjusting for the difference between this expected width and actual width of 0.54 m, when applied to a 350 kg sow, results in a planar width of 1.53 m and a planar area of 3.17 m² (sow length × planar width) for turning around, thus giving our current, minimum design criterion for space (Table 1).

In addition to turning around, the sow species behaviour patterns suggest requirement for additional space to increase activity and separate excretion and feeding areas. Overlap of the excretory area with either the nest or feeding area will result in impaired hygiene and is likely to increase the risk of adverse health consequences in this and subsequent lactation stages. In addition, overlap of the nest and feed area will increase activity and divert focus of attention within the nest, thus increasing risk of piglet mortality after farrowing. Evidence regarding preferred excretory and lying sites for sows is rare (Wiepkema, 1986), which is unsurprising given the lack of choice presented to the majority of farrowing sows housed in crates. However, studies looking at gilts (Damm and Pedersen, 2000) and piglets in the farrowing environment showed that in general pigs, when given the opportunity, will keep separate areas for excretion and resting (Watson, 1978; Petherick, 1983b). Without detailed work determining the area needed to satisfy feeding

and dunging needs, we can only estimate this based on the calculations already described. Space required for feeding can be described as 'static', with the requirement that the sow has to be able to stand up and feed. Therefore, as described above, a 350-kg sow would require 2.44 m² of floor space for feeding. In addition, the sow needs to be able to exit the feeding area, however space provided for nesting or excretion could accommodate this requirement. Based on the above calculations, floor space required for a 350-kg sow to have a separate excretion site, in which it can turn around would be the same amount described for nesting (i.e. at least 2.44 m² floor space and 3.17 m² planar space – Table 1). However, it is likely to be the properties of the space, not just the quantity that will determine suitability for meeting needs. For example, pigs prefer to excrete in spaces away from feeding and nesting sites (Baxter, 1982). In addition, excretory areas can be created by providing temperature differentials, whereby intended dunging areas are cooler (Randall *et al.*, 1983) than lying areas (Table 1). With these factors in mind, it is possible that quantity of space could be reduced if substituted by attention to detail and clever design.

Enclosure

In tests looking at the choice of nest site, Hunt and Petchey (1987) demonstrated that sows always chose farrowing locations that were inside, or against a solid wall, never out in the open. This matches the choices of sows under natural and semi-natural conditions (Stolba and Wood-Gush, 1984) where 40% chose total enclosure and 89% chose partial enclosure. However, the amount of enclosure needed to satisfy nest-site choice is still not clear. Providing nest sites that are enclosed on three sides with a side open or partly open would facilitate isolation and vigilance for potential threats (Table 1). Under natural conditions, the nest often includes a roofed area of vegetation, providing protection from inclement weather as well as concealment from potential predators. There are contradictory reports as to the benefits of a roof under indoor farm conditions. Phillips *et al.* (1991) found that providing a 'roof' had no positive influence on site choice between totally enclosed, partially enclosed or open crates and that enclosure was preferred more by younger sows. When a crude form of enclosure was given to animals farrowing in crates by providing a Hessian roof (Cronin and Vanamerongen, 1991), maternal behaviour was influenced positively and it was suggested that the make-shift roof offered a sense of nesting completion for the sow. The crates in this experiment had no further enclosure, and therefore it is possible that the influence was not roof specific.

Substrate

Substrate manipulation, as well as enclosure appears to contribute to perception of nesting completion and, in preference tests with both elements offered, sows made a choice based on the pre-formed nest sites that also offered sufficient quantity of straw to satisfy nest-building behaviour (Arey *et al.*, 1992). Such has been the case in other nest-building species; in farmed mink, for example, when provided

with an artificial nest (nest box), kit survival and vitality was improved and maternal stress *post partum* was reduced (Malmkvist and Palme, 2008). However, there appeared to be greater benefits from providing an artificial nest in conjunction with access to straw, with less variable duration of parturition displayed by these animals with the enriched nests. Hutson (1988) tested a limited number of sows and gilts in an operant conditioning experiment and, based on a low motivation for sows to lift a lever 10 times to obtain 1 kg of straw, it was concluded that straw is unimportant to prepartal sows. However, this claim is unsupported by the majority of other research in this area (Arey *et al.*, 1991 and 1992; Arey, 1992; Damm *et al.*, 2000). Arrangement of nest-building material provides feedback to the expectant sow and, without suitable feedback from this activity to signal that nesting is complete, some sows may continue to be motivated to nest build even during farrowing (Thodberg *et al.*, 1999; Damm *et al.*, 2000). This constitutes a risky situation for the newborns and therefore is counter-productive for reproductive success.

Thus, substrate is important, but the unresolved issue is quantifying the necessary amount and type of substrate. There have been reports of sows making very complex nests outdoors, with 255 kg of substrate in one nest (Zanella and Zanella, 1993). Such extreme quantities are impractical and most likely unnecessary under indoor conditions. The complexity of nesting substrates can influence behaviour; Damm *et al.* (2000) demonstrated that gilts given branches and straw were better able to construct a 'complete' nest than gilts only provided with straw. Arey *et al.* (1991) gave two groups of sows sandy depressions with straw available from a dispenser and absence or presence of pre-formed nests with 23 kg of straw on the floor. The first group removed 23 kg of straw and used it for nest building, while the latter group removed 9.5 kg in addition to the 23 kg already pre-formed. Recent work by Damm and Pedersen (L Pedersen, personal communication) has recorded the amount of substrate carried by sows during the nest-building phase and they have recommended that a minimum of 2 kg of long-stemmed straw be provided to all sows *pre partum* as nesting material (Table 1). Alternative nesting materials have been studied (Widowski and Curtis, 1990), with recommendations that providing a cloth tassel that animals can pull, tear and manipulate may also provide benefits even if the material cannot result in the building of a suitable nest. This is particularly relevant for animals kept in crates, as the majority of the tassel remains at the front of the crate, where sows can continue to access it, whereas straw is often pushed to the back of the crate during nest building, and becomes unavailable (Table 1).

Flooring

Additional preference tests have been performed on the properties of the floor, with 100% of sows choosing to farrow in an earthen pen site that could be hollowed out compared to a concrete floor (Haskell and Hutson, 1994). It is possible that this earthen site offered qualities in addition

to its deformability, including greater udder comfort for the sow or lower thermal conductivity, and clarification of which properties are of importance needs further research. If physical comfort is a key aspect, then quantity of substrate on the floor or area of 'comfortable' flooring should accommodate a sow lying laterally for parturition. Arey *et al.* (1991), in their pre-formed nest experiments already described, noted that 100% of sows farrowed in the sandy depression, potentially because of similar, 'comfortable' qualities. Thodberg *et al.* (1999) suggested that the sand provided in their experiments gave important feedback, in addition to that provided by the substrate available. Thus, flooring with malleable properties to accommodate nest building activities is recommended (Table 1). Thermal comfort is also likely to be a consideration of the nest site and, in choice experiments offering sows a concrete floor or a mat floor with varying amounts of straw, sows appeared to show a preference for flooring that offered the greatest thermal resistance (the mat with the most amount of straw – Hunt and Petchey, 1987), thus limiting the loss of heat via conduction. However, when sows were offered a heated (34°C) or non-heated floor space there was no difference regarding their choice of farrowing area (Pedersen *et al.*, 2007). Thus, further and more detailed investigations are needed on temperature preferences of nesting sows (Table 1).

In addition, the drainage properties of the floor, as well as the provision of space, will facilitate hygiene. Perforated or slatted flooring has been shown to have higher hygiene than solid floors (Rantzer and Svendsen, 2001), however there may be negative consequences from such flooring regarding injury to both piglets and sows (see later section). Defined areas with different floor types and gradation in solid flooring to facilitate pen hygiene by creating a drainage slope, would offer a compromise to satisfy these needs (Table 1).

Parturition, early lactation and nest-occupation

Sow behaviour and physiology

Sow behaviour during this phase is characterised by prolonged lateral lying and udder exposure. Despite only rudimentary maternal care during parturition in this species, sows in semi-natural environments will get up during parturition to inspect their offspring, making nose-to-nose contact before rooting the nest to move piglets out of the way and then resuming lateral lying (Gundlach, 1968; Jensen, 1986). Reactivity patterns of farrowing sows have been described by several authors (Jensen, 1986; Jarvis *et al.*, 1999; Pedersen *et al.*, 2003), with sow's responsiveness being highest in the first 2 h of parturition followed by a prolonged (approximately 6 h) non-responsive phase and then, in conjunction with general activity, responsiveness returns. This pattern is accompanied by endocrine changes (Algers and Uvnäs-Moberg, 2007), but is also influenced by the environment. During farrowing an increase in plasma oxytocin concentrations, promoting uterine contraction for piglet expulsion (Castrén *et al.*, 1993b), also results in continuous let-down of colostrum from the teats. Feedback from the udder, stimulated by massage from piglets, also increases oxytocin concentration and milk let-down

(Algers *et al.*, 1991). Colostrum availability declines exponentially during the first 24 h of secretion, before changing to milk by 30 h *post partum* (Le Dividich *et al.*, 2005). In the early stages of the lactation period, sows generally do not leave the nest site. Observing sows under semi-natural conditions, Jensen (1986) reported that for the first 2 days *post partum* the majority of sows spent 90% of their time in the nest, with the remaining 10% of the time occupied by brief foraging ventures outside. During this nest-occupation phase, sows will spend a large part of the time initiating suckling bouts and nursing their piglets at 30 to 70-min intervals.

Piglet behaviour and physiology

Almost immediately after birth the piglet will start udder and teat seeking behaviours, relying on a mixture of visual, auditory, olfactory and tactile stimuli (Jeppesen, 1981; Morrow-Tesch and McGlone, 1990; Parfet and Gonyou, 1991). Welch and Baxter (1986) suggested that the thermal and tactile properties of the udder are unique in the extrauterine environment, offering salient characteristics similar to those of the intrauterine environment, and therefore highly attractive to the newborn piglets. Colostrum flow is continuous during parturition and piglets have access to relatively plentiful colostrum for approximately 12 h after the birth of the first piglet (Fraser, 1980), before cyclical let-down (approximately every 30 min) of colostrum begins. The diffuse epitheliochorial nature of the porcine placenta means that piglets are born without immune protection, having to acquire maternal antibodies through ingesting colostrum (Gaskins and Kelley, 1995). To acquire passive immunity the small intestine of the newborn piglet undergoes dramatic functional changes immediately *post partum* (for review see Xu *et al.*, 2000). During the first day of life, the neonate's small intestine has the ability to absorb macromolecules, such as intact immunoglobulins (IgGs), across the brush border membrane. The capacity for macromolecule absorption diminishes as early as 24 h (Rooke and Bland, 2002) and is usually complete by 48 h *post partum*, a phenomenon generally referred to as gut closure (Gaskins and Kelley, 1995). Though having an early time window for gut closure limits the time for IgG absorption, it also reduces the chances of pathogens entering the gut and subsequent disease risks. It is noteworthy that the earliest time for sow nest-site departure coincides with gut closure and thus the cessation of passive immunity acquisition by the piglets.

To achieve a high intake of colostrum, gain warmth and establish a teat order, piglets prefer to lie close to the udder during the first 24 h of life. However, there is a trade-off between udder access and the risk of being crushed by the sow. This risk is enhanced when a piglet's energy reserves are low and Weary *et al.* (1996) concluded that crushings are at least partly the result of the nutritional challenge facing piglets. They found that piglets with slow weight gain spent more time in risky areas underneath their sitting or standing mother. Crushing predominantly occurs as the sow makes a posture change from standing to lying or when rolling (Damm *et al.*, 2005; Weary *et al.*, 1998) and the risks associated

with these posture changes are dependent on the environment in which the sow farrows and her maternal behaviour. In a study by Wechsler and Hegglin (1997), it was demonstrated that the quality of the sow's pre-lying behaviour influenced the risk of crushing. Sows that lay down vertically rather than flopping, and those that paid attention to their piglets' location in the nest prior to lying down, had fewer crushed piglets in their litter. Although this study was limited to a small sample size of 11 sows, similar results have been described by other authors (Signoret *et al.*, 1975; Blackshaw and Hagelso, 1990) with sows observed to root through the straw with their noses and group their piglets before lying down considered the more careful mothers.

Newborn piglets are also susceptible to hypothermia as, from the moment they are born and make the transition from the intrauterine and thermoneutral environment to the extrauterine environment, they experience a dramatic reduction in ambient temperature (by approximately 15°C to 20°C – Herpin *et al.*, 2002). Consequently, the lower limit of the thermoneutral zone (the lower critical temperature – approximately 34°C Mount, 1968) is not achieved, and the piglet is at risk of becoming chilled. The physiological immaturity of the newborn piglet contributes to its vulnerability; it is born with very little insulating adipose tissue and no brown fat (Herpin *et al.*, 2002), so can only fuel heat production and increase its core body temperature by mobilising energy reserves, present as glycogen and fat and then catabolism of its skeletal muscle. Glycogen reserves present in muscle are low and depleted quickly (Herpin *et al.*, 1992) and most fat that is present is not available for mobilisation as it is structural fat (Herpin *et al.*, 2002). Catabolism of skeletal muscle is also limited as the process of protein oxidation is poorly developed in the neonatal pig (Schmidt and Herpin, 1998). In addition the domestic, typical 'pink' piglet has lost much of the physical protection from pelage that its wild ancestors (i.e. wild boarlets) have (Foley *et al.*, 1971). The need to preserve homeothermy is therefore closely linked to the need to ingest colostrum, and the post-birth drop in body temperature is only reversed once this is achieved (Noblet and Le Dividich, 1981; Baxter *et al.*, 2008).

After farrowing, as continuous colostrum supply switches to more cyclical milk let-downs every 30 to 70 min (Fraser, 1980; Lewis and Hurnik, 1985), a teat order develops, with strong individual teat fidelity. Suckling not only fulfils basic physiological needs but also assists in establishing bonds between mother and offspring. As with teat-seeking behaviour, recognition appears to involve a mixture of olfactory, auditory and tactile stimuli. Odour cues are important with respect to individual teat recognition by the piglets (Jeppesen, 1982; Morrow-Tesch and McGlone, 1990). Auditory cues are most evident before and during suckling, or in distress situations. The sow commences suckle grunting before let-down, signalling to the piglets to gather at the udder (for review see Fraser, 1980). Further to these olfactory and auditory stimuli, tactile stimuli are evident as the piglets display pre- and post-let-down suckling massage of the udder, and piglets are often observed post-let down going to the sow's head to make nose-to-nose contact.

Functional importance of behaviours during the parturition and nest occupation phase

Once the piglets are born, they face a number of significant challenges arising from an unpredictable mother and environment. In order to survive piglets must get to the udder, show vigour in acquiring and maintaining a functional teat and thus ingest vital colostrum to aid energy balance, preserve homeothermy and promote immune-function (Baxter *et al.*, 2008). This requires effective synchronisation between behaviours of the sow and her piglets. The functional importance of neonatal piglet behaviours is self-evident from the description of physiology. One of the most frequently reported piglet mortality syndromes is hypothermia and starvation leading to piglet lethargy, subsequent crushing and death (English and Morrison, 1984; Weary *et al.*, 1996; Edwards, 2002). However, thermal comfort represents one of the areas where there is potential for parent-offspring conflict. The temperature zone of thermal comfort for a sow is between 12°C and 22°C (Black *et al.*, 1993), and at temperatures above 22°C, adult pigs are increasingly susceptible to heat stress. Heat stress affects lactational output by lowering feed intake, thus contributing to decreased piglet weight gain and increased sow lactational weight loss (Stansbury *et al.*, 1987). In contrast, the small size and immature physiology of the newborn piglet results in a lower critical temperature of 34°C (Herpin *et al.*, 2002). Thus, a compromise must be reached to balance these conflicting needs. Investigations of the thermal properties of nest sites in 'natural' farrowing conditions show that the nests are robust to climatic extremes (Algers and Jensen, 1990), maintaining an average temperature in winter of 20.3°C despite outdoor temperatures averaging –1.5°C. Thermal image data from commercial outdoor huts, taken at the time of farrowing, showed that hut temperature averaged 14.7°C, whereas within the deep straw bedding and at the udder of the sow, temperature averaged 31.5°C (Baxter *et al.*, 2009). This evidence further supports the functional role of nest building behaviour by domestic sows to balance thermal needs and the functional significance of a nest during parturition and nest occupation.

Movement by the sow during farrowing is risky for the piglets, with more passive maternal behaviour thought of as positive to allow safe udder access for the piglets. However, under natural and semi-natural conditions sows will get up, turn around and inspect their piglets (Gundlach, 1968; Jensen, 1986). The functional significance of this was questioned by the authors because of the risks involved. However, in a more natural setting, early interaction with the piglets may enhance the mother–young bonds and litter recognition before returning to the herd. This factor is unlikely to be as important in a conventional indoor situation, where dam and offspring remain together alone until weaning. However, recent work suggests that interaction with piglets is important to allow positive maternal behaviour to develop; Andersen *et al.* (2005) demonstrated that 'non-crushing' mothers were observed to spend more time making nose-to-nose contact with piglets before lying down, as well as reacting to a piglet distress call by making contact with piglets sooner than 'crushers'.

Auditory cues regulate social and nutritional relationships between mother and offspring (Puppe *et al.*, 2003) and not only do the rhythmic vocalisations of the sow during nursing assist with synchronous feeding behaviour, they also provide an auditory 'fingerprint', unique to the sow, that the piglets can use for recognition of their own mother. It appears that, even within the first few hours after birth, piglets are rapidly learning the unique composition of vocalisations by their mother (Walser, 1986). Work by Horrell and Hodgson (1992) showed that, as early as 36-h old, piglets were able to recognise the vocalisations of their own mother over those of an alien sow.

Tactile stimulation performed by the piglets at the udder fulfils several functions; suckling massage releases prolactin that assists with adapting maternal physiology to cope with lactation (Algers *et al.*, 1991) and, dependent on the time spent massaging, increases the teat milk yield for the next suckling – the 'restaurant hypothesis' (Gill and Thomson, 1956; Algers and Jensen, 1985). Nose-to-nose contact between piglet and sow post let-down is thought to signal need and reaffirm bonds and recognition. Recognition of their own mother is an important task for the newborn piglet since pigs are social animals, naturally living in groups where mothers, often themselves related, will synchronously give birth to multiple young (Gundlach, 1968). Frequencies of these nasal contacts under semi-natural conditions were found to decrease substantially with time after farrowing and to stabilise to a low level on day six *post partum* (Stangel and Jensen, 1991), coinciding with the average nest abandonment day also observed by these authors. Under more intensive conditions, it would seem that this recognition is not warranted, as sows and piglets generally do not mix before weaning. Thus, it can be hypothesised that this nose-to-nose contact behaviour has persisted more as a way of signalling need and reaffirming a bond to solicit milk.

Needs-based design criteria for the parturition and nest occupation phase

The needs of the sow during farrowing depend on the satisfaction gained from the nest-building phase. A quiet, enclosed area with a suitable and complete nest should result in a sow that performs increased lateral lying during farrowing (Jarvis *et al.*, 1999). The needs of the piglet during this crucial time are more complex than those of the sow; however, they are invariably linked to the behaviour of the sow. Given that the satisfaction of the nest-building phase directly influences the parturition phase, the criteria for nest building apply here: suitable space, enclosure, substrate and flooring. The evidence presented above suggests that thermal comfort, physical protection of piglets via suitable walls and substrate, as well as lack of disturbance are additional attributes to consider.

Space

Static space required during parturition can be calculated in a very simple manner, with the sow needing to lie laterally and have enough space to facilitate birth and suckling for the

piglets (Table 2). Petherick (1983a) developed an equation for determining space needed for a pig to rest on its sternum or be fully recumbent (Figure 2). Using these equations and assuming that the sow is static while giving birth, a 350-kg sow lying laterally is 2.07-m long and 1.08 m in depth and occupies 2.24 m² of space. In addition, there needs to be dynamic space for the piglets to be born safely and to walk around the sow to access the udder and suckle colostrum.

Using the same equations but applying these to an optimum weight piglet of 1.6 kg (Roehle and Kalm, 2000), whose dimensions would be 0.35-m long and 0.07-m wide, a 350-kg sow giving birth to 1.6 kg piglets would require a minimum floor space of 2.79 m² (Table 2). This is based on a rather crude assumption that in order for the piglet to negotiate access to the udder the existing area allowing lateral recumbency for the sow must now accommodate a piglet being born and then walking around its mother. Therefore, the nest-size space is based on the following equation:

$$(\text{Sow length} + \text{piglet length}) \times (\text{sow lateral width} + \text{piglet ventral width})$$

Suckling space should also be considered, at farrowing and during lactation. Work already described above providing body dimensions of farrowing sows (Moustsen *et al.*, 2004) was added to, to provide body dimensions of suckling piglets (Moustsen and Poulsen, 2004a); these authors recommended that the width between the sow's udder and any solid surface should be a minimum of 0.50 to 0.60 m to allow 4 to 5-week-old piglets to all suckle comfortably in the space envelope between the sow's front and back legs when lying laterally. Thus, the conclusions drawn from these combined data are that the depth would need to be between 1.21 and 1.31 m for lateral lying and comfortable suckling throughout lactation (Table 2).

From the evidence presented above the sow also needs to be able to turn around and inspect her piglets, as well as perform piglet gathering or grouping behaviour before lying down. Therefore, similar calculations for the nest-building phase can be made and a 350 kg sow should have a nest-size that facilitates turning (Table 1). Recent work by Weber *et al.* (2009) compared piglet mortality in loose-housed systems in Switzerland in relation to pen size. They and other groups observed no difference in mortality levels in crateless pens measuring over 5 m² compared with crated accommodation (Cronin *et al.*, 2000; Weber *et al.*, 2007), suggesting that where studies have shown higher mortality in crateless pens than in crates (Blackshaw *et al.*, 1994; Marchant *et al.*, 2000), this was a result of the pens being under 5 m². Consequently, Weber *et al.* (2009) suggest this should be the minimum space requirement for unimpeded piglet gathering behaviour. However, further research is needed to give accurate figures for both comfortable turning and unimpeded gathering behaviour. In addition, during the nest occupation phase, the space needs to facilitate hygiene to avoid detrimental effects on piglet health. Therefore, the

Table 2 Summary of design recommendations to meet the biological needs of piglets during farrowing and lactation

| Component of system | Biological specification | Recommendations |
|---------------------|---|--|
| Space ^c | Parturition | Length of sow and piglet (2.42 m) × width of sow lying laterally and ventral piglet (1.15 m) = 2.79 m ² (minimum) ^a |
| | Udder access for suckling throughout lactation | Depth of sow (0.71 m) + length of 4 to 5 week old piglets (0.50 to 0.60 m) = 1.21 to 1.31 m (minimum) ^a |
| | Protection, safe lying area for parturition and nest occupation | Separate space inaccessible to the sow e.g. 0.80 m ² per 10 to 12 neonates ^a |
| | Protected lying area during lactation | 0.97 m ² (minimum) – 2.32 m ² (maximum) for 14, 4-week-old piglets ^a |
| | Area for feed trough to introduce creep feed Hygiene | Provide in area inaccessible to the sow, interacts with above ^b Separate area for dung site, interacts with flooring ^a |
| Substrate | Foraging, nutritional development Enrichment, social development | Earth-like materials (e.g. peat) ^a Further research needed on quantity. Novelty requires fresh input daily. Complex materials (e.g. branches) preferred ^b |
| | Thermal comfort during parturition | 2.5 cm of straw ^a , interacts with flooring |
| | Physical comfort | Further research needed, interacts with thermal comfort and flooring properties |
| | Protection | Deep bedding – 10 to 12 cm ^a , interacts with flooring |
| Walls | Protection from sow posture changes Social contact (visual and physical) | Escape zones at all pen walls ^b Vertical barred area (minimum) ^b . Further research needed to determine best method to mix pre-weaning. |
| | Hygiene | Solid walls (at least at bottom of penning) separating other litters during first 7 days of life ^b |
| | Thermal comfort | Solid walls in lying area of material to limit heat loss via radiation ^b – interacts with substrate and flooring |
| Flooring | Thermal comfort during parturition and first 24 h of life | High thermal resistance ^a – for example, rubber matting or deep substrate (see above) or under-floor/localised heating (see General) |
| | Thermal comfort during lactation | High thermal resistance ^b – for example, rubber matting or deep substrate (see above) or under-floor/localised heating (see General) |
| | Physical comfort – avoiding injury, promoting suckling behaviour | Solid flooring with minimal abrasiveness and well maintained (e.g. rubber matting or specialised screed with non-slip properties), interacts with substrate |
| | Protection from fatal crushing by the sow Hygiene | Malleable flooring ^b interacts with deep substrate Slatted flooring with void width no more than 10 mm and rounded edges ^a . Interacts with temperature (see General) |
| General | Health – treatment for injuries, vaccines, etc. Promote weaning, reduce nutritional stress and encourage increased feed and water intake | Safe area for handling required to ensure piglet health needs, interacts with space ^b Suitable solid food, inaccessible to the sow – interacts with space and substrate. Provide feed tray with sufficient space to allow social facilitation ^a . Provide water drinker with upward angled bite nipple close to floor ^a |
| | Thermal comfort | Localised heat source set at thermo-neutral temperature (e.g. 34°C at birth) – interacts with substrate ^a |
| | Hygiene | Temperature differentials to encourage dunging outside of nest site – interacts with flooring ^b |

^aRecommendations derived from one or more specific references.

^bRecommendations based on synthesising and interpreting the literature reviewed.

^cSpace recommendations for an 'improved' sow weighing 350 kg and an 'optimum' birth weight piglet of 1.6 kg

space has to provide a defined excretion area away from the nest site, as described in the nest-building phase (Table 1).

Protection

Protection for the newborn piglet is paramount to its survival since crushing by the dam, as either proximate or ultimate cause, is the most frequently recorded reason for mortality. Though, under more natural settings, sows and piglets occupy the same space in the nest, under commercial indoor housing conditions it is generally accepted, and indeed recommended (Defra publications, 2003), that 'a solid and

dry lying area' away from the sow be provided so that the piglets can 'all rest at the same time' and be protected (see also CoE, 2001). This area, often referred to as a creep, also makes it possible to provide different microclimates, as well as separate nutritional provision (see later section) and facilitates easier access for health checks and husbandry procedures. However, it requires additional space (Table 2). Detailed work in Denmark (Moustsen and Poulsen, 2004b) studied piglet dimensions (weight, length, height and depth) in different postures, with the average snout to tail length of 4-week-old piglets measuring 0.56 m, their shoulder width




Figure 2 'Pigtographs' by Petherick (1983a) showing the space requirements for a pig resting fully recumbent or on its sternum (W = body weight in kg; area in m^2).

0.13 m and their depth measuring 0.14 m. Thus, these authors determined that to accommodate 10 to 12 piglets until weaning at 4 weeks of age, a creep should be no less than $0.80 m^2$ (Table 2). Further investigations by these authors have resulted in recommendations that each piglet, when weaning at 4 weeks of age should have a solid, dry area of $0.069 m^2$. Therefore, recommendations for high litter sizes of 14, 4-week-old piglets would require an area of $0.97 m^2$ (Table 2) available to them (V. Moustsen personal communication). However, such minimum recommendations only allow resting in a ventral position. To allow full lateral recumbency, which is a position increasingly adopted as the piglet gets heavier and older (Ekkel *et al.*, 2003), $0.166 m^2$ /piglet would be needed (Table 2). It is possible that such lying areas do not have to be provided solely in the creep; protective space (e.g. underneath sloped walls) may accommodate a number of resting piglets as they get older and heavier. At this time their thermogenic abilities are well developed (Herpin *et al.*, 2002) and therefore the need for all piglets to simultaneously access a heated area may decrease.

Deep bedding (10 to 15 cm depth) also has protective properties (Table 2); it is a malleable, non-uniform mattress cushioning sow posture changes and offering a buffer to piglets from fatal crushing (Baxter *et al.*, 2009). This is most likely more effective when the flooring is also malleable (e.g. earth, sand; Table 2) and under outdoor conditions piglets observed to be crushed by their mothers have appeared up to 4 h later from underneath the sow relatively unharmed and have survived to weaning (E Baxter, unpublished observation).

The care and control with which the sow lies down will influence piglet survival, and providing sows with a supportive surface to lean against during their descent has been shown to reduce crushing (Baxter, 1991; Marchant *et al.*, 2001). In the latter study, 89% of sows preferred to use sloped walls, farrowing rails or creep sides to lie down against rather than descend unsupported. Damm *et al.* (2006) enhanced this work by discriminating which was the most preferred choice of supportive structure. They concurred that the majority preferred support regardless of its form, but found that sows preferred solid sloped or vertical walls over farrowing rails (Table 1). Moreover, a recent survey by Weber *et al.* (2009)

showed no significant influence of the presence of piglet protection bars on piglet losses. The traction and comfort of the floor will also influence lying behaviour, thus protection for the piglets (Damm *et al.*, 2005), as well as the health and welfare of both sow and piglets (Boyle *et al.*, 2000). In the latter study, Boyle *et al.* (2000) demonstrated when rubber mats were given instead of metal slats sow comfort was improved and slipping reduced (Table 1).

Thermal comfort

As stated previously, thermal comfort represents an area of potential parent-offspring conflict and the design challenge involves balancing the dichotomised needs of the sow and piglets. Although sows are at risk from heat stress at temperatures above $22^\circ C$, recent evidence suggests that sows show a preference for warmer areas in the immediate *post-partum* period (Phillips *et al.*, 2000; Pedersen *et al.*, 2007) and certainly will avoid metal flooring, potentially because of the cooling properties of this floor type (Table 1). Both Pedersen *et al.* (2007) and Phillips *et al.* (2000) found the majority of sows chose to lie on floors with temperatures of $34^\circ C$ and $35^\circ C$, respectively, *post partum* and for the following 3 days, after which there was a pronounced change to preference for the cooler floors (Phillips *et al.*, 2000). In both cases, in order to facilitate choice, the sows were loose-housed and thus posture changes to regulate their temperature were possible. Further research is necessary before recommending a floor temperature of $35^\circ C$ as a design criterion in all circumstances, as sows might develop heat stress, particularly if their mobility is restricted. Even if mobility is not restricted, as in loose-housed environments, if sows do not have access to cooler areas they can develop heat stress, as evidenced by Malmkvist *et al.* (2009). These researchers heated the floor fully to $34^\circ C$, limiting behavioural thermoregulation and resulting in increased plasma levels of cortisol and ACTH for sows housed in this treatment.

To meet the thermal needs of the piglet, while safeguarding sow comfort, the provision of facilities to create a microclimate are important. Within the nest area, provision of bedding plays an important role. Deep bedding slows heat loss, having a thermal resistance 11 times greater than that of concrete slats and 22 times greater than solid, wet concrete flooring (Wathes and Whittemore, 2006). Mount (1967) demonstrated that piglets in contact with a concrete floor lost 40% more heat than those in contact with 2.5 cm of straw. Deep bedding will also help reduce evaporative heat loss by speeding the physical removal of placental membranes. In the absence of deep bedding, an artificially heated creep area of suitable size (see above) can provide an effective microclimate provided that piglets can be attracted to use this area at an early age. Creep designs have attempted to exploit the piglet's attraction to the stimuli of warmth and softness (Welch and Baxter, 1986) by providing temperature differentials (Burri *et al.*, 2009; Vasdal *et al.*, 2010), water beds (Ziron and Hoy, 2003) or simulated udders (Lay *et al.*, 1999). However, most piglets prefer to lie at the udder of the sow for at least 24 h after parturition. Nevertheless, a creep area, to provide an effective microclimate

either using heat lamps or heat mats, is recommended (Table 2). If this is not provided then the minimum recommendations to facilitate thermal comfort for piglets at this stage would be provision of straw at a depth of at least 2.5 cm (Table 2), given an ambient room temperature no greater than 22°C to avoid sow heat stress (Table 1).

Lack of disturbance

Isolation of the nest site selected under natural conditions minimises risk of disturbance. There is evidence that disturbance during parturition will negatively influence oxytocin levels (Lawrence *et al.*, 1992). Certainly, the length of the non-responsive phase after the birth of the last piglet may depend on how quiet the nest site is, and may influence subsequent restlessness of the sow and risk of piglet crushing. Analysis of data from both indoor and outdoor farrowing systems indicates that group farrowing in more confined circumstances significantly increases mortality risk (Edwards and Fraser, 1997; Marchant *et al.*, 2000). Sow reactivity is a potential risk factor; though sow responsiveness to the distress calls of a piglet have been considered positive maternal behaviour (Wechsler and Hegglin, 1997; Andersen *et al.*, 2005), when sows farrow in individual pens which are in close proximity, they may respond vigorously to distress calls from outside their own nest, to the potential detriment of their own piglets. Together with disturbance from external activities on the pig unit, this consideration indicates a requirement for further research to evaluate the benefits of a quiet nest site. The degree of enclosure afforded to the nest site will influence disturbance levels; however, clarification of the relative importance of visual and auditory isolation is needed (Table 1).

Nest departure and social integration

Sow and piglet behaviour and physiology

At this stage, in mid lactation, sow and piglet behaviours are reasonably well aligned, with both mother and offspring sharing similar biological goals. Thus, in this section, their behaviours are described together. In a free-range system, piglets will leave the nest site before they are ready for weaning. Gundlach (1968) reported that piglets stayed close to the nest for 7 to 14 days *post partum* and, after venturing out, often returned to suckle and re-establish attachment with the sow via piglet-initiated nose-to-nose contacts. Strategies of newborn ungulates are usually divided into two types; 'hidiers' and 'followers' (Lent, 1974). Hidiers tend to stay in vegetation and await a parental signal to leave this protected site, whereas followers stay close to their parent(s). As a result of building a nest, pigs have generally been described as hidiers (Lent, 1974). However, Jensen and Redbo (1987) described piglets at the time of nest-leaving as entering an intermediate phase in their behavioural development as they change from a 'hider' to a 'follower' strategy. On adventures out of the nest, piglets will follow their mother during foraging bouts. Sows will increase these foraging bouts as lactation progresses and the energy demands become more intense, causing sows to mobilise

their body reserves (Quesnel and Prunier, 1995) and switch from an anabolic to a catabolic state (Valros *et al.*, 2003). Milk production then peaks by the 3rd week in lactation, before gradually declining (Elsley, 1971). When accompanying the sow on ventures out of the nest, piglets mix with other litters and engage in important social behaviours before returning to the nest to suckle. Petersen *et al.* (1989) identified an age-dependent high motivation for piglets to integrate or establish social bonds outwith their litter. Piglets were observed to interact with non-litter mates of the same age between 2 and 6 weeks *post partum* and thereafter they interacted with other group members.

Functional importance of behaviours during this phase

Nest departure and re-integration into the social group serves certain fundamental functions for sows and piglets in natural environments; it allows sows to increase their foraging territory and feed intake during this period of increased metabolic demand for lactation. For the piglets, early adventures out of the nest site allow interaction with a diverse environment, which may have importance for nutritional and long-term behavioural development. Research on the influence of rearing in 'impoverished' conditions (i.e. barren environments such as the farrowing crate) has shown that the complexity of the nursing environment can influence subsequent cognitive abilities of pigs (de Jong *et al.*, 2000), their ability to cope with stressful situations (de Jonge *et al.*, 1996) and may influence future maternal behaviour of any female offspring (Schouten, 1987). Thus, it is possible that piglets reared in a more enriched environment are better adapted for challenges later in life, including social integration. In addition, the foraging activities displayed outside the nest site satisfy the piglets' desire to perform exploratory behaviours, rooting and manipulating substrates and beginning the adaptation to solid food intake. The early socialisation benefits the piglets by reducing the impact of weaning (Pajor *et al.*, 1999; Hessel *et al.*, 2006), particularly reducing the effects of mixing aggression. D'Eath (2005) demonstrated that pre-weaning mixing of piglets reduced aggression levels at weaning and this socialisation appeared to equip piglets with social skills that benefitted them in the long term, enabling them to more rapidly form stable dominance hierarchies during future aggressive encounters with unfamiliar pigs. Social integration at this time is also important for the sow to minimise aggressive interactions associated with re-integration that might impede her fitness. Aggression is a major problem among un-familiar pigs and can lead to severe injury and physiological stress responses which can have detrimental effects on reproductive parameters (Arey and Edwards, 1998). Such overt aggression could also risk injury to the piglets if they share this social space. Under natural or semi-natural conditions pigs maintain established groups where aggression is regulated via 'avoidance order', with specific behavioural patterns reducing risk of attacks by dominant individuals (Jensen, 2002). In studies determining the influence of recognition on aggressive interactions post-mixing (Olsson and Svendsen, 1997)

there was a clear effect of acquaintance with reduced aggression between pigs that recognised each other from their previous groups. Arey (1999) concluded that sows from the same gestational group rarely fought when mixed after 4 weeks separated in farrowing crates, thus recognition is a key factor in post-mixing aggression.

Needs-based design criteria for the nest-departure and social integration phase

Design criteria at this stage must consider the need for enrichment and social contact, and therefore the space and partition design to facilitate the latter. In addition, physical health and nutritional development for the piglet should be considered. For the former, flooring is an important factor.

Space

Space is required in order to leave the nest site. The defined space described in the previous sections to facilitate pen hygiene could accommodate the need to leave the nest site (Tables 1 and 2). Quantifying this criterion is difficult because it is more than likely that once the nest site has been left, qualitative aspects of the space are more important than actual physical area. However, in order to facilitate full social integration, additional space would be required for sows and litters to mix. The extent of this space requirement depends on how many litters and sows would be mixing, and whether the sows were familiar (i.e. came from the same social group or gestation pen). In addition to the sum of each individual's physical space requirements, space for appropriate social interaction must be considered (Baxter, 1985). If animals are unfamiliar space allocation is an important factor to accommodate the agonistic interactions associated with establishing a new hierarchy (Jensen, 1982) and facilitate escape by losers of conflicts. Amongst familiar animals, signalling of submission and avoidance of dominants also requires space (Jensen, 2002). Group size is therefore likely to be a factor influencing requirement for space per sow, with smaller total space allowances resulting in greater aggression; Jensen (1984) showed that in a group of five sows, in order for avoidance behaviours to be displayed, each sow would need 3 m² of space. In these stable groups, aggressive interactions are observed mainly in relation to competition for resources and therefore space requirements are dependent on resource accessibility (Baxter, 1985). Such resources include space for resting, feeding and excretion. However, the work carried out thus far has not included space for nursing, and whether such activity requires additional 'personal space' to that required for simply lying laterally needs additional research (Table 1). In group-housed systems, where sows and litters mix for lactation, space given (per sow and litter) varies considerably (e.g. 27 m² – Kerr *et al.*, 1988 v. 8 m² – Halverson *et al.*, 1997), with group size also varying. Baxter (1985) recommends that instead of extravagant allowances for social space, it is better to consider clever design (Table 1) allowing interaction with the ability to display submission, reduced aggression and specific activities (e.g. nursing).

The social dynamic of the group is crucial when making decisions about pre-weaning mixing and it is recommended that lactational mixing only be attempted when the sows are familiar with each other (i.e. from the same gestational group; Table 1). If this cannot be achieved, some degree of social contact can be afforded by partitions which enable visual, olfactory and even nasal contact between adjacent sows. In pre-weaned pigs, it has been shown that such 'fenceline' contact can reduce aggression at subsequent mixing (Waran and Broom, 1993) and there is anecdotal evidence of similar benefits in sows (Dellmeier and Friend, 1991). Mixing of piglets to facilitate early socialisation prior to weaning could occur without mixing of sows. By providing an additional area to that of the farrowing pen, inaccessible to the sow or providing pop-holes between farrowing pens to be opened during later lactation, unfamiliar piglets could socialise. However, although there are reported benefits (D'Eath, 2005) to pre-weaning mixing, there are also reported negative consequences, including an increase in risk of disease transmission and pre-weaning fighting with the potential to negate the post-weaning benefits (Parratt *et al.*, 2006). There are also reports of the presence of foreign litters increasing teat disputes, disrupting nursing frequency and thus reducing milk intake (Pedersen *et al.*, 1998). Thus, the costs and benefits of pre-weaning mixing make recommendations difficult. Until further definitive evidence is presented, an interim measure, which allows physical contact between neighbouring sows and their litters (e.g. through barred walls) may be beneficial in reducing the negative effects of weaning on both sows and their piglets. Such design would require no extra space, but would require different wall partitions with contact areas (Tables 1 and 2).

Enrichment

Substrate provided regularly could satisfy the need for an enriched and diverse environment. In addition, provision of substrates has been shown to decrease anti-social behaviours, including belly nosing and aggressive interactions with pen mates (Beattie *et al.*, 2000). Any form of enrichment should have properties that accommodate the motivated behaviours for the animals concerned (Van de Weerd *et al.*, 2003) which, in the case of piglets, include foraging and exploratory behaviours. Earth-like materials such as peat, mushroom compost (Beattie *et al.*, 1998) and more complex materials such as branches (Pedersen *et al.*, 2005) have been shown to be preferred as enrichment substrates over straw (Table 2). The amount of substrate necessary to fulfil this enrichment function in young piglets has not, to our knowledge, been quantified. Most enrichment experiments indicate that it is the novel aspect of the enrichment that stimulates exploratory behaviour. In piglets aged between 4 and 8 weeks, Fraser (1985) observed two main peaks in activity that coincided with delivery of fresh straw. These piglets were provided with either 1 or 2.5 kg of fresh straw twice daily in two experimental groups, with activity levels not significantly different. Research in older pigs also suggests that relatively small quantities of straw, if presented

daily, appear to be almost as effective at generating greater straw-directed behaviour and less pig-directed and pen-directed behaviour as large amounts (Kelly *et al.*, 2000). However, Day *et al.* (2002) found that an increasing provision of straw resulted in an increase in the proportional frequency of rooting and ploughing behaviours and an associated decrease in potentially damaging pig-behaviours, such as tail-biting. Further research is needed to determine the quantities of substrate needed to provide pre-weaned piglets with an enriched and diverse environment. Although substrate is likely to encourage foraging behaviours, at this stage, food that aids nutritional development (i.e. creep feed) should also be provided (see later section on nutritional development and weaning; Table 2).

Flooring

If the enrichment provided is to be straw (despite its less preferable rooting and foraging properties than some other substrates) then it is possible that, as substrate is required to fulfil other functions such as physical and thermal comfort, the quantity can first be defined with these needs in mind. The amount of straw needed for thermal comfort will depend on other properties of the environment. Fraser (1985) showed a distinct preference in weaned piglets for bedded areas when the ambient temperature was between 18°C to 21°C and not at 25°C to 27°C. When thermal comfort was not compromised there was no clear preference for lying on bedded or concrete areas. These results suggest that thermal comfort is of greater importance than physical comfort in young pigs. Straw bedding can reduce the prevalence of skin lesions and bursae in older pigs (Moultotou *et al.*, 1999) but pressure from hard flooring is likely to cause less discomfort in younger, lighter piglets. However, hard flooring could cause discomfort and injury for young piglets when they make contact with the floor during the vigorous activity of suckling. Moultotou and Green (1999) found a positive correlation between time spent suckling and incidence of carpal skin abrasions between 4 and 6 days of age. Thus deep straw bedding, at least during the 1st week of life, could have protective benefits for younger piglets (Table 2). In order for this recommendation to be fulfilled, the flooring where the straw is provided needs to be solid to prevent the straw falling through the slats (Tables 1 and 2). It is clear from the literature that the amount of bedding required for physical comfort is poorly defined, especially in relation to the age and weight of pigs, and therefore such quantification requires additional research.

Properties of the floor, such as slip-resistance, abrasiveness, surface profile, hardness, void width (slatted floors) and thermal resistance (already discussed) are important factors that contribute to its injury potential. As with thermal comfort, physical comfort requires balancing sow and piglet needs. For example, slip-resistance, particularly for the sow, needs to be achieved without modifying the floor's characteristics in such a way that predisposes injury through excessive abrasion (McKee and Dumelow, 1995). Abrasion is a common risk for suckling piglets, with mixed reports as to

which flooring material causes the highest incidence of knee damage: Furniss *et al.* (1986) found that the worst knee damage was seen on an old cement screed. In contrast, Clark (1985) found that newly installed concrete screed had the highest abrasive properties, whereas the same floor after 3 months continuous use had the lowest abrasiveness. Therefore, both floor material and its deterioration over time require consideration; repairing solid flooring and using specialised screed offering both grip and low abrasion properties has shown to decrease piglet injuries (Zoric *et al.*, 2009). Any flooring needs to demonstrate these two properties (Tables 1 and 2); straw-bedding (Vellenga *et al.*, 1983; Edwards and Lightfoot, 1986) or solid flooring covered with peat (Zoric *et al.*, 2008) offer the lowest risk of injury. However, hygiene is an important consideration and slatted floors assist with hygiene (Rantzer and Svendsen, 2001), as well as thermal comfort for the sow during lactation in warm conditions. Incorrectly designed slatted flooring can cause injuries to piglet feet (particularly the coronet and footpads, Lewis *et al.*, 2005), as well as damage to sow teats and legs (Edwards and Lightfoot, 1986). If slatted floors are to be used (e.g. in the dunging area), the recommended void width is no greater than 10 mm with rounded edges (Mitchell and Smith, 1978; see Table 2), and plastic-coated metal has been shown to be less injurious than steel (Furniss *et al.*, 1986; Lewis *et al.*, 2005). In addition, void width with respect to use of straw or other substrate that may cause clogging needs to be considered. Many new plastic types of flooring are now available and widely used, but few have been scientifically assessed.

Thus far, the basic needs such as food and water for the sow have been ignored presuming these are already well catered for as fundamental physiological requirements. However, it is important to emphasise that the sow must increase her feed and water intake during this time to accommodate the significant metabolic demands of lactation. Certain aspects of the environment can hamper this, such as increased environmental temperatures (Black *et al.*, 1993), as well as restrictive conditions. It has been reported that sows kept in loose-housed environments have a higher intake of food during lactation than animals kept in farrowing crates (Pajor, 1998). It is known that there is a correlation between food intake and heat stress. Recent work has showed that localised cooling of the sow's shoulders while in the farrowing crate increased sow's feed intake and consequently piglet weight gain (Silva *et al.*, 2006; van Wagenberg *et al.*, 2006). Designing an environment with temperature differentials could facilitate thermal comfort, as well as encourage the animals to use the defined spaces for the functions they are intended for (Table 1).

Nutritional development and weaning

Sow and piglet behaviour and physiology

Studying pigs under semi-natural conditions, Jensen and Recén (1989) observed weaning to occur at 17.2 weeks of age. Prior to weaning, there was gradual separation

between the sow and her litter, with progressive changes in suckling behaviour also evident. For example, sows became less permissive of suckling until, after 4 weeks, sows would terminate 100% of sucklings. This has also been demonstrated in experimental indoor systems in get-away pens (GAPs – Bøe, 1991). In this system, where sows could leave the piglets voluntarily without the piglets being able to follow, suckling frequency decreased gradually from 22.9 to 4.3 bouts every 24 h over a 10-week period. During this experiment, average weaning age was between 11 and 12 weeks and half of the sows were observed to wean their piglets before the piglets were 10 weeks of age (Bøe, 1991). The discrepancy between these weaning ages and those observed by Jensen and Recén (1989), suggests that many sows would wean earlier if given sole responsibility, yet piglets continue to solicit milk for a much greater period of time if allowed to follow the sow. However, under more natural conditions, it is unlikely that the sow would ever be able to totally leave her litter without, at least, the fittest piglets being able to follow. It could therefore be argued that the expectation that the piglets will follow her is part of the innate behaviour of the sow and that results from experiments using GAPs (Bøe, 1991) are inconclusive regarding motivation to wean earlier than reported 'naturally'. Regardless of this, conflict is evident: the sow needs an environment that allows time away from the piglets to gradually decrease milk let-downs and promote the weaning process. In contrast, the piglets need an environment where they can follow the sow to prevent litter desertion and prolong the suckling period to allow gradual transition to solid food.

In natural or semi-natural environments, the piglets' dependency on the sow's milk for nutrition gradually decreases. Piglets have been observed to display foraging behaviours as early as the first few days *post partum* (Petersen, 1994) and after 4 weeks old they gradually increase their time spent foraging. Thus, they can survive without milk at this time provided that alternative food resources are available. This is related to season, with weaning being observed to be later when piglets were born between July and October, with expected weaning during the winter months (Jensen and Recén, 1989).

Functional importance of behaviours during this phase

During the latter stages of lactation, the needs of the sow and her litter become increasingly dichotomised as a power struggle over maternal resources develops because of conflicting evolutionary strategies (Trivers, 1974). The piglets must consider their own survival and soliciting as much milk as possible from the sow will aid in their growth and development. The strategy adopted by the sow is one of balance: she must balance the needs of her current litter with the potential for any future litters, and thus her overall reproductive success. Therefore, in order to maintain body condition while still providing for her current litter, she will reduce the number of milk let-downs per day. This can only be efficiently accomplished by a gradual separation from the litter, thus a control of her lactational output and progressive weaning. Abrupt weaning under more intensive conditions

causes nutritional challenges for the young, can slow growth rate and compromise immune function (Algers, 1984; Pajor *et al.*, 1999). Within 24 h of weaning, there are marked changes in the structure and function of the small intestine resulting from temporary reduction in feed intake (Pluske *et al.*, 1995), with consequential growth check and frequent incidences of diarrhoea. Recently, attention has also been drawn to the possible psychological and long-term consequences of early weaning (Newberry and Swanson, 2008; Weary *et al.*, 2008). Prior to weaning, piglets will be used to regular and synchronised feeding patterns, at this stage controlled by the sow delivering a highly nutritious and palatable milk source (Pluske *et al.*, 1995). Abrupt weaning involves a complete change in the pattern and delivery of food, requiring both behavioural and physiological adaptations by the piglet. If supplementary food is available from at least the 3rd week of lactation, when milk production starts to decline, expression of foraging behaviour has functional consequences for changes in gastric enzyme secretions and gut development (Cranwell, 1995), allowing the piglet to experience a more gradual weaning process. It is possible that, if these foraging needs are met, then the piglet will take a more active role in weaning; Jensen and Recén (1989) proposed the 'fast-food' hypothesis whereby a cost-benefit scenario occurs in which the cost of 'asking' the sow for milk is gradually outweighed by the benefits of foraging for food.

Needs-based design criteria for the nutritional development and weaning phase

From the evidence presented above, the needs of the piglets and sow during this time will be difficult to balance given the trade-offs and parent-offspring conflict described. Assuming the nutritional needs of the sow are met, the design must consider suitable space and flooring to facilitate sow controlled nursing and physical and thermal comfort, respectively. To meet piglet needs, suitable substrate and feed should be provided to promote foraging and encourage nutritional independence. Approaches to ease the transition to weaning should consider minimising both physiological and psychological stressors. A recent review by Weary *et al.* (2008) summarised these approaches as: increasing weaning age, increasing the time the dam and her young spend apart before weaning and making the solid diet more attractive to the pre-weaned young. With respect to the design criteria, the latter two approaches, in particular, require recommendations on how best to accommodate them.

Space

Designing the space required during this time involves a question of balance; the sow needs space to 'escape' her piglets to control lactational output, but the piglets need to avoid total desertion. In GAPs, where the sow could access the feeding and dunging area without the piglets, sows lost less weight and returned to oestrus quicker than sows in confined systems and piglets gained weight and switched to solid food quicker than piglets in confined systems (Pajor, 1998). It has also been shown that in a farrowing environment

which allows 'escape' by the sow, more 'sow-controlled' nursing behaviour developed, which was considered an important aspect of good mothering ability (Thodberg *et al.*, 2002). Though benefits of systems where the sow can leave her piglets freely are evident, there are potential risks of the nursing frequency decreasing to a detrimental level for the piglets or complete abandonment (Bøe, 1993). Environments which encourage voluntary gradual separation are recommended (Weary *et al.*, 2008). Quantifying the space needed to accommodate gradual separation is difficult and, as already mentioned in previous sections, qualitative aspects of the space maybe more important than its actual physical size. Providing defined areas (Table 1), gives the sow options to move away from her piglets. Potentially making certain areas less attractive to the piglets, yet still accessible, would encourage gradual separation; for example, temperature differentials between the defined areas and/or different flooring, enrichment and food provision (Table 1).

Nutritional provision

Design criteria in the previous section to promote social integration and development via appropriate space and substrate will reduce the physical and psychological consequences of early weaning (Newberry and Swanson, 2008) (Tables 1 and 2). In addition, suitable substrate and supplementary feed will encourage foraging behaviours to reduce the impact of nutritional challenges and promote weaning decisions in piglets. It is generally accepted that higher intake of solid feed (or creep feed) prior to weaning results in higher intake post-weaning (Aherne *et al.*, 1982). However, individual differences in pre-weaning creep feed intake are evident; Barnett *et al.* (1989) found total creep feed consumption varied from 13 to 194 g/piglet and from 107 to 1550 g/litter. In addition, some piglets do not eat solid feed at all prior to weaning; using a faecal dyeing technique Callesen *et al.* (2007) identified 78% of piglets as 'eaters' and 22% as 'non-eaters'. Individual piglets appear to vary in their feeding strategies, with some electing to continue suckling and engaging in energy consuming teat stimulation and others prepared to start feeding on solids sooner (Bøe and Jensen, 1995). Since the former strategy results in a more detrimental impact at weaning, encouraging all piglets to sample solid feed prior to weaning should be a design objective in any farrowing system. Appropriate delivery of this supplementary feed to ensure piglet accessibility (Table 2) may influence the success of nutritional development (Appleby *et al.*, 1991 and 1992). Trough designs which increase available feed space and the chance for social facilitation appear to promote feed intake; shallow feeders, rather than traditional hoppers, (Wattanukul *et al.*, 2005) and Playfeeders (Kuller *et al.*, 2010) which stimulate exploratory behaviours have proved most successful (Table 2).

Pre-weaning water intake is an over-looked area, and low feed intake is often compounded with low water intake (Dybkjaer *et al.*, 2006). As with food intake, similar approaches to increasing the attractiveness of water dispensers could influence consumption rates. Larger water dispensers

(e.g. circular bowls) facilitate social drinking and accommodate the development of synchronous behaviour observed in social animals (Phillips and Fraser, 1991). However, in loose-housed environments a water bowl maybe disturbed by the sow and bowls can result in hygiene issues if fouled. After trialling different designs, Phillips and Fraser (2001) showed success via speed of discovery and use with an upward angled bite nipple drinker placed close to the floor (Table 2).

Conclusions

This review has documented the considerable amount of literature describing the biological needs of the sow and her piglets during the different phases of farrowing and lactation. It has highlighted the evolutionary significance, and thus function, of certain behavioural patterns that continue to be displayed under farmed conditions. Identifying and summarising suitable design criteria to accommodate biological needs has highlighted that, despite the abundance of work in this area, there remain out-standing issues requiring additional and detailed research. Quantifying space and substrate need at specific stages, as well as the potential benefits of a quiet nest site are examples of such issues. It is evident that there are a number of trade-offs in operation throughout the phases of farrowing, not only between different animals but also between different goals of the individual, complicating the ability to design a suitable farrowing and lactation environment that meets all biological needs. Though not a focus of this review, further complications will exist when considering the farmer's interests and a successful system should attempt to reconcile the 'triangle of needs' between sow, farmer and piglets, to maximise productivity and welfare.

Acknowledgement

The authors wish to thank the Department for Environment, Food and Rural Affairs (Defra) of the UK for funding.

References

- Aherne FX, Danielsen V and Nielsen HE 1982. The effects of creep feeding on pre- and post-weaning pig performance. *Acta Agriculturae Scandinavica* 32, 155–160.
- Ahlström S, Jarvis S and Lawrence AB 2002. Savaging gilts are more restless and more responsive to piglets during the expulsive phase of parturition. *Applied Animal Behaviour Science* 76, 83–91.
- Algers B 1984. Early weaning and cage rearing of piglets; influence on behaviour. *Zentralblatt für Veterinärmedizin Reihe A* 31, 14–24.
- Algers B and Jensen P 1985. Communication during suckling in the domestic pig. Effects of continuous noise. *Applied Animal Behaviour Science* 14, 49–61.
- Algers B and Jensen P 1990. Thermal microclimate in winter farrowing nests of free-ranging domestic pigs. *Livestock Production Science* 25, 177–181.
- Algers B and Uvnäs-Moberg K 2007. Maternal behavior in pigs. *Hormones and Behavior* 52, 78–85.
- Algers B, Madej A, Rojanasthien S and Uvnäs-Moberg K 1991. Quantitative relationships between suckling-induced teat stimulation and the release of prolactin, gastrin, somatostatin, insulin, glucagon and vasoactive intestinal polypeptide in sows. *Veterinary Research Communications* 15, 395–407.

- Andersen IL, Berg S and Bøe KE 2005. Crushing of piglets by the mother sow (Sus scrofa) – purely accidental or a poor mother? *Applied Animal Behaviour Science* 93, 229–243.
- Appleby MC, Pajor EA and Fraser D 1991. Effects of management options on creep feeding by piglets. *Animal Production* 53, 361–366.
- Appleby MC, Pajor EA and Fraser D 1992. Individual variation in feeding and growth of piglets – effects of increased access to creep food. *Animal Production* 55, 147–152.
- Arey DS 1992. Straw and food as reinforcers for prepartal sows. *Applied Animal Behaviour Science* 33, 217–226.
- Arey DS 1997. Behavioural observations of peri-parturient sows and the development of alternative farrowing accommodation: a review. *Animal Welfare* 6, 217–229.
- Arey DS 1999. Time course for the formation and disruption of social organisation in group-housed sows. *Applied Animal Behaviour Science* 62, 199–207.
- Arey DS and Edwards SA 1998. Factors influencing aggression between sows after mixing and the consequences for welfare and production. *Livestock Production Science* 56, 61–70.
- Arey DS, Petchey AM and Fowler VR 1991. The preparturient behavior of sows in enriched pens and the effect of preformed nests. *Applied Animal Behaviour Science* 31, 61–68.
- Arey DS, Petchey AM and Fowler VR 1992. The effect of straw on farrowing site choice and nest building behavior in sows. *Animal Production* 54, 129–133.
- Barnett KL, Kornegay ET, Risley CR, Lindemann MD and Schurig GG 1989. Characterization of creep feed consumption and its subsequent effects on immune response, scouring index and performance of weanling pigs. *Journal of Animal Science* 67, 2698–2708.
- Baxter MR 1982. Environmental determinants of excretory and lying areas in domestic pigs. *Applied Animal Ethology* 9, 195.
- Baxter MR 1983. Ethology in environmental design for animal production. *Applied Animal Ethology* 9, 207–220.
- Baxter MR 1985. Social space requirements for pigs. In *Social space requirements for domestic animals* (ed. R Zayan), pp. 116–127. Martinus Nijhoff Publishers, Dordrecht, The Netherlands.
- Baxter MR 1991. The freedom farrowing system. *Farm Building Progress* 104, 9–15.
- Baxter MR and Schwaller C 1983. Space requirements for sows in confinement. In *Farm animal housing and welfare* (ed. SH Baxter, MR Baxter and JAD MacCormack), pp. 181–195. Martinus Nijhoff Publishers, The Hague, The Netherlands.
- Baxter EM, Jarvis S, D'Eath RB, Ross DW, Robson SK, Farish M, Nevison IM, Lawrence AB and Edwards SA 2008. Investigating the behavioural and physiological indicators of neonatal survival in pigs. *Theriogenology* 69, 773–783.
- Baxter EM, Jarvis S, Sherwood L, Robson SK, Ormandy E, Farish M, Smurthwaite KM, Roehe R, Lawrence AB and Edwards SA 2009. Indicators of piglet survival in an outdoor farrowing system. *Livestock Science* 124, 266–276.
- Beattie VE, Walker N and Sneddon IA 1998. Preference testing of substrates by growing pigs. *Animal Welfare* 7, 27–34.
- Beattie VE, O'Connell NE and Moss BW 2000. Influence of environmental enrichment on the behaviour, performance and meat quality of domestic pigs. *Livestock Production Science* 65, 71–79.
- Black JL, Mullan BP, Lorschy ML and Giles LR 1993. Lactation in the sow during heat stress. *Livestock Production Science* 35, 153–170.
- Blackshaw JK and Hagelsø AM 1990. Getting-up and lying-down behaviors of loose-housed sows and social contacts between sows and piglets during day-1 and day-8 after parturition. *Applied Animal Behaviour Science* 25, 61–70.
- Blackshaw JK, Blackshaw AW, Thomas FJ and Newman FW 1994. Comparison of behavior patterns of sows and litters in a farrowing crate and a farrowing pen. *Applied Animal Behaviour Science* 39, 281–295.
- Bøe K 1991. The process of weaning in pigs: when the sow decides. *Applied Animal Behaviour Science* 30, 47–59.
- Bøe K 1993. The effect of age at weaning and post-weaning environment on the behaviour of pigs. *Acta Agriculturae Scandinavica, Section A – Animal Science* 43, 173–180.
- Bøe K and Jensen P 1995. Individual differences in suckling and solid food intake by piglets. *Applied Animal Behaviour Science* 42, 183–192.
- Boyle LA, Regan D, Leonard FC, Lynch PB and Brophy P 2000. The effect of mats on the welfare of sows and piglets in the farrowing house. *Animal Welfare* 9, 39–48.
- BPEX 2004. An industry update on farrowing systems. BPEX/MLC, Milton Keynes, UK.
- Brambell FWR 1965. Report of the technical committee to enquire into the welfare of animals kept under intensive livestock husbandry systems. HMSO, London, UK.
- Burri M, Wechsler B, Gygas L and Weber R 2009. Influence of straw length, sow behaviour and room temperature on the incidence of dangerous situations for piglets in a loose farrowing system. *Applied Animal Behaviour Science* 117, 181–189.
- Callesen J, Halas D, Thorup F, Bach Knudsen KE, Kim JC, Mullan BP, Wilson RH and Pluske JR 2007. The influence of nutritional and management factors on piglet weight gain to weaning in a commercial herd in Denmark. *Livestock Science* 108, 117–119.
- Castrén H, Algers B, de Passillé A-M, Rushen J and Uvnäs-Moberg K 1993a. Preparturient variation in progesterone, prolactin, oxytocin and somatostatin in relation to nest building in sows. *Applied Animal Behaviour Science* 38, 91–102.
- Castrén H, Algers B, de Passillé AM, Rushen J and Uvnäs-Moberg K 1993b. Early milk ejection, prolonged parturition and periparturient oxytocin release in the pig. *Animal Production* 57, 465–471.
- Clark M 1985. Farrowing pen floor abrasiveness measured using a rubber-block drag test. *Farm Building Progress* 80, 29–32.
- Council of Europe 2001. Commission Directive 2001/93/EC of 9 November 2001 amending Directive 91/630/EEC laying down minimum standards for the protection of pigs. *Official Journal of the European Communities* L316, 1–3. http://eur-lex.europa.eu/pri/en/oj/dat/2001/l_316/l_31620011201en00010004.pdf
- Cranwell PD 1995. Development of the Neonatal Gut. In *The neonatal pig development and survival* (ed. MA Varley), pp. 99–154. CABI Publishing, Wallingford, UK.
- Cronin GM and Vanamerongen G 1991. The effects of modifying the farrowing environment on sow behavior and survival and growth of piglets. *Applied Animal Behaviour Science* 30, 287–298.
- Cronin GM, Simpson GJ and Hemsworth PH 1996. The effects of the gestation and farrowing environments on sow and piglet behaviour and piglet survival and growth in early lactation. *Applied Animal Behaviour Science* 46, 175–192.
- Cronin GM, Lefebvre B and McClintock S 2000. A comparison of piglet production and survival in the Werribee farrowing pen and conventional farrowing crates at a commercial farm. *Australian Journal of Experimental Agriculture* 40, 17–23.
- Cronin GM, Barnett JL, Hodge FM, Smith JA and McCallum TH 1991. The welfare of pigs in 2 farrowing lactation environments – cortisol responses of sows. *Applied Animal Behaviour Science* 32, 117–127.
- Curtis SE, Hurst RJ, Gonyou HW, Jensen AH and Muehling AJ 1989. The physical space requirement of the sow. *Journal of Animal Science* 67, 1242–1248.
- D'Eath RB 2005. Socialising piglets before weaning improves social hierarchy formation when pigs are mixed post-weaning. *Applied Animal Behaviour Science* 93, 199–211.
- Daan S and Tinbergen JM 1997. Adaptation of life histories. In *Behavioural ecology: an evolutionary approach* (ed. JR Krebs and NB Davis), pp. 311–333. Wiley-Blackwell, UK.
- Damm BI and Pedersen LJ 2000. Eliminative behaviour in preparturient gilts previously kept in pens or stalls. *Acta Agriculturae Scandinavica Section A-Animal Science* 50, 316–320.
- Damm BI, Forkman B and Pedersen LJ 2005. Lying down and rolling behaviour in sows in relation to piglet crushing. *Applied Animal Behaviour Science* 90, 3–20.
- Damm BI, Vestergaard KS, Schröder-Petersen DL and Ladewig J 2000. The effects of branches on prepartum nest building in gilts with access to straw. *Applied Animal Behaviour Science* 69, 113–124.
- Damm BI, Pedersen LJ, Marchant-Forde JN and Gilbert CL 2003. Does feed-back from a nest affect preparturient behaviour, heart rate and circulatory cortisol and oxytocin in gilts? *Applied Animal Behaviour Science* 83, 55–76.
- Damm BI, Bildsøe M, Gilbert C, Ladewig J and Vestergaard KS 2002. The effects of confinement on periparturient behaviour and circulating prolactin, prostaglandin F₂[alpha] and oxytocin in gilts with access to a variety of nest materials. *Applied Animal Behaviour Science* 76, 135–156.

Farrowing pen design on the basis of biological needs

- Damm BI, Moustsen V, Jørgensen E, Pedersen LJ, Heiskanen T and Forkman BR 2006. Sow preferences for walls to lean against when lying down. *Applied Animal Behaviour Science* 99, 53–63.
- Dawkins M 1977. Do hens suffer in battery cages? Environmental preferences and welfare. *Animal Behaviour* 25, 1034–1046.
- Day JEL, Burfoot A, Docking CM, Whittaker X, Spoolder HAM and Edwards SA 2002. The effects of prior experience of straw and the level of straw provision on the behaviour of growing pigs. *Applied Animal Behaviour Science* 76, 189–202.
- Defra publications 2003. Code of recommendations for the welfare of livestock pigs. Defra publications, Admail 6000, London SW1A 2XX.
- de Jong IC, Prelle IT, van de Burgwal JA, Lambouij E, Korte SM, Blokhuis HJ and Koolhaas JM 2000. Effects of environmental enrichment on behavioral responses to novelty, learning, and memory, and the circadian rhythm in cortisol in growing pigs. *Physiology & Behavior* 68, 571–578.
- de Jonge FH, Bokkers EAM, Schouten WGP and Helmond FA 1996. Rearing piglets in a poor environment: developmental aspects of social stress in pigs. *Physiology & Behavior* 60, 389–396.
- Dellmeier GR and Friend TH 1991. Behavior and extensive management of domestic sows (*Sus-Scrofa*) and litters. *Applied Animal Behaviour Science* 29, 327–341.
- Dybkaer L, Jacobsen AP, Togersen FA and Poulsen HD 2006. Eating and drinking activity of newly weaned piglets: effects of individual characteristics, social mixing, and addition of extra zinc to the feed. *Journal of Animal Science* 84, 702–711.
- Edwards SA 1996. Designing systems to meet behavioural needs: the family pen system for pigs. In *Animal behavior and the design of livestock and poultry systems*, pp. 115–125. Northeast Regional Agricultural Engineering Service, Ithaca, USA.
- Edwards SA 2002. Perinatal mortality in the pig: environmental or physiological solutions? *Livestock Production Science* 78, 3–12.
- Edwards SA and Lightfoot AL 1986. The effect of floor type in farrowing pens on pig injury. II. Leg and teat damage of sows. *British Veterinary Journal* 142, 441–445.
- Edwards SA and Fraser D 1997. Housing systems for farrowing and lactation. *The Pig Journal* 39, 77–89.
- Ekkel ED, Spoolder HAM, Hulsegge I and Hopster H 2003. Lying characteristics as determinants for space requirements in pigs. *Applied Animal Behaviour Science* 80, 19–30.
- Elsley FWH 1971. Nutrition and lactation in the sow. In *Lactation* (ed. ED Falconer), pp. 393–411. Butterworths, London.
- English PR and Morrison V 1984. Causes and prevention of piglet mortality. *Pig News Information* 5, 369–375.
- European Food Safety Authority 2007. Animal health and welfare aspects of different housing and husbandry systems for adult breeding boars, pregnant, farrowing sows and unweaned piglets, pp. 1–13. Scientific Opinion of the Panel on Animal Health and Welfare, EFSA.
- Farm Animal Welfare Council 1992. FAWC updates the five freedoms. *Veterinary Record* 131, 357.
- Foley CW, Seerley RW, Hansen WJ and Curtis SE 1971. Thermoregulatory responses to cold environment by neonatal wild and domestic piglets. *Journal of Animal Science* 32, 926–929.
- Frädrich H 1974. A comparison of behaviour in the Suidae. In *The Behavior of Ungulates and its relation to management* (ed. V Geist), pp. 133–143. IUCN Publication New Series, University of Calgary, Alberta, Canada.
- Fraser D 1980. A review of the behavioural mechanism of milk ejection of the domestic pig. *Applied Animal Ethology* 6, 247–255.
- Fraser D 1985. Selection of bedded and unbedded areas by pigs in relation to environmental-temperature and behavior. *Applied Animal Behaviour Science* 14, 117–126.
- Fraser D, Phillips PA and Thompson BK 1997. Farrowing behaviour and stillbirth in two environments: an evaluation of the restraint-stillbirth hypothesis. *Applied Animal Behaviour Science* 55, 51–66.
- Furniss SJ, Edwards SA, Lightfoot AL and Spechter HH 1986. The effect of floor type in farrowing pens on pig injury. 1. Leg and teat damage of suckling piglets. *British Veterinary Journal* 142, 434–440.
- Gaskins HR and Kelley KW 1995. Immunology and neonatal mortality. In *The neonatal pig: development and survival* (ed. MA Varley), pp. 39–56. CABI Publishing, Wallingford, UK.
- Gill JC and Thomson W 1956. Observations on the behaviour of suckling pigs. *The British Journal of Animal Behaviour* 4, 46–51.
- Graves HB 1984. Behavior and ecology of wild and feral swine (*Sus Scrofa*). *Journal of Animal Science* 58, 482–492.
- Gundlach H 1968. Brutfürsorge, Brutpflege, Verhaltensontogenese und Tagesperiodik beim Europäischen Wildschwein *Sus Scrofa L* (translation: maternal care, ontogeny of behaviour and daily periodicity in European wild boars, *Sus Scrofa L*). *Zeitschrift für Tierpsychologie* 25, 955–995.
- Hafez ESE, Sumption LJ and Jakway JS 1962. The behaviour of swine. In *The behavior of domestic animals* (ed. ESE Hafez), pp. 334–369. The Williams and Wilkins Co., Baltimore, USA.
- Halverson M, Honeyman MS and Adams M 1997. Swedish deep-bedded group nursing systems for feeder pig production. Swine system options for Iowa. Extension Publication p. 12 Iowa State University, Ames.
- Hamilton WD 1964a. The genetical evolution of social behaviour. I. *Journal of Theoretical Biology* 7, 1–16.
- Hamilton WD 1964b. The genetical evolution of social behaviour. II. *Journal of Theoretical Biology* 7, 17–52.
- Haskell MJ and Hutson GD 1994. Factors affecting the choice of farrowing site in sows. *Applied Animal Behaviour Science* 39, 259–268.
- Herpin P, Le Dividich J and Vanos M 1992. Contribution of colostrum fat to thermogenesis and glucose-homeostasis in the newborn pig. *Journal of Developmental Physiology* 17, 133–141.
- Herpin P, Damon M and Le Dividich J 2002. Development of thermoregulation and neonatal survival in pigs. *Livestock Production Science* 78, 25–45.
- Hessel EF, Reiners K and Van den Weghe HFA 2006. Socializing piglets before weaning: effects on behavior of lactating sows, pre- and postweaning behavior, and performance of piglets. *Journal of Animal Science* 84, 2847–2855.
- Horrell I and Hodgson J 1992. The bases of sow-piglet identification. 1. The identification by sows of their own piglets and the presence of intruders. *Applied Animal Behaviour Science* 33, 319–327.
- Hughes BO and Black AJ 1973. The preference of domestic hens for different types of battery cage floor. *British Poultry Science* 14, 615–619.
- Hughes BO and Duncan IJH 1988. Behavioural needs: can they be explained in terms of motivational models? *Applied Animal Behaviour Science* 19, 352–355.
- Hunt K and Petchey AM 1987. A study of the environmental preferences of sows around farrowing. *Farm Building Progress* 89, 11–14.
- Hutson GD 1988. Do sows need straw for nest-building? *Australian Journal of Experimental Agriculture* 28, 187–194.
- Jarvis S, Calvert SK, Stevenson J, van Leeuwen N and Lawrence AB 2002. Pituitary-adrenal activation in pre-parturient pigs (*Sus scrofa*) is associated with behavioural restriction due to lack of space rather than nesting substrate. *Animal Welfare* 11, 371–384.
- Jarvis S, Lawrence AB, Mclean KA, Deans LA, Chirside J and Calvert SK 1997. The effect of environment on behavioural activity, ACTH, beta-endorphin and cortisol in pre-farrowing gilts. *Animal Science* 65, 465–472.
- Jarvis S, Lawrence AB, Mclean KA, Chirside J, Deans LA and Calvert SK 1998. The effect of environment on plasma cortisol and beta-endorphin in the parturient pig and the involvement of endogenous opioids. *Animal Reproduction Science* 52, 139–151.
- Jarvis S, Mclean KA, Calvert SK, Deans LA, Chirside J and Lawrence AB 1999. The responsiveness of sows to their piglets in relation to the length of parturition and the involvement of endogenous opioids. *Applied Animal Behaviour Science* 63, 195–207.
- Jarvis S, van der Vegt BJ, Lawrence AB, Mclean KA, Deans LA, Chirside J and Calvert SK 2001. The effect of parity and environmental restriction on behavioural and physiological responses of pre-parturient pigs. *Applied Animal Behaviour Science* 71, 203–216.
- Jensen P 1982. An analysis of agonistic interaction patterns in group-housed dry sows – aggression regulation through an ‘avoidance order’. *Applied Animal Ethology* 9, 47–61.
- Jensen P 1984. Effects of confinement on social interaction patterns in dry sows. *Applied Animal Behaviour Science* 12, 93–101.
- Jensen P 1986. Observations on the maternal behaviour of free-ranging domestic pigs. *Applied Animal Behaviour Science* 16, 131–142.
- Jensen P 1993. Nest building in domestic sows: the role of external stimuli. *Animal Behaviour* 45, 351–358.

- Jensen P 2002. Behaviour of pigs. In *The ethology of domestic animals* (ed. P Jensen), pp. 159–172. CABI Publishing, Wallingford, UK.
- Jensen P and Redbo I 1987. Behaviour during nest leaving in free-ranging domestic pigs. *Applied Animal Behaviour Science* 18, 355–362.
- Jensen P and Recén B 1989. When to wean – observations from free-ranging domestic pigs. *Applied Animal Behaviour Science* 23, 49–60.
- Jensen P and Toates FM 1993. Who needs 'behavioural needs'? Motivational aspects of the needs of animals. *Applied Animal Behaviour Science* 37, 161–181.
- Jensen P and Gustafsson M 1997. Towards a functional view on domestication. In *Proceedings of 31st International Congress of the International Society for Applied Ethology*, pp. 43–44. Prague, Polygrafia SAV, Czech Republic.
- Jeppesen LE 1981. An artificial sow to investigate the behaviour of sucking piglets. *Applied Animal Ethology* 7, 359–367.
- Jeppesen LE 1982. Teat-order in groups of piglets reared on an artificial sow. II. Maintenance of teat-order with some evidence for the use of odour cues. *Applied Animal Ethology* 8, 347–355.
- Johnson AK and Marchant-Forde JN 2009. Welfare of pigs in the farrowing environment. In *The welfare of pigs* (ed. JN Marchant-Forde), pp. 141–188. Springer, The Netherlands.
- Kelly HRC, Bruce JM, English PR, Fowler VR and Edwards SA 2000. Behaviour of 3-week weaned pigs in straw-flow, deep straw and flatdeck housing systems. *Applied Animal Behaviour Science* 68, 269–280.
- Kerr SGC, Wood-Gush DGM, Moser H and Whittemore CT 1988. Enrichment of the production environment and the enhancement of welfare through the use of the Edinburgh family pen system of pig production. *Research and Development in Agriculture* 5, 171–186.
- Kuller WI, Tobias TJ and van Nes A 2010. Creep feed intake in unweaned piglets is increased by exploration stimulating feeder. *Livestock Science* 129, 228–231.
- Lawrence AB, Petherick JC, Mclean K, Gilbert CL, Chapman C and Russell JA 1992. Naloxone prevents interruption of parturition and increases plasma oxytocin following environmental disturbance in parturient sows. *Physiology & Behavior* 52, 917–923.
- Lawrence AB, Petherick JC, Mclean KA, Deans LA, Chirside J, Vaughan A, Clutton E and Terlouw EMC 1994. The effect of environment on behavior, plasma-cortisol and prolactin in parturient sows. *Applied Animal Behaviour Science* 39, 313–330.
- Lay DCJ, Haussmann MF, Buchanan HS and Daniels MJ 1999. Danger to pigs due to crushing can be reduced by the use of a simulated udder. *Journal of Animal Science* 77, 2060–2064.
- Le Dividich J, Rooke JA and Herpin P 2005. Nutritional and immunological importance of colostrum for the new-born pig. *The Journal of Agricultural Science* 143, 469–485.
- Lent PC 1974. Mother-infant relationships in ungulates. In *The behaviour of ungulates and its relationship to management* (ed. V Geist and F Walther), pp. 14–55. International Union for the Conservation of Nature and Natural Resources, Morges, Switzerland.
- Lewis NJ and Hurnik JF 1985. The development of nursing behaviour in swine. *Applied Animal Behaviour Science* 14, 225–232.
- Lewis E, Boyle LA, O'Doherty JV, Brophy P and Lynch PB 2005. The effect of floor type in farrowing crates on piglet welfare. *Irish Journal of Agricultural and Food Research* 44, 69–81.
- Lou ZS and Hurnik JF 1994. An ellipsoid farrowing crate – its ergonomic design and effects on pig productivity. *Journal of Animal Science* 72, 2610–2616.
- MacDonald AA 2000. Comparative anatomy, physiology and ecology of pregnancy and lactation in wild pigs: a review. In *Zoo animal nutrition* (ed. J Nijboer, JM Hatt, W Kaumans and U Ganslober), pp. 213–236. Filander, Furth.
- Malmkvist J and Palme R 2008. Periparturient nest building: Implications for parturition, kit survival, maternal stress and behaviour in farmed mink (*Mustela vison*). *Applied Animal Behaviour Science* 114, 270–283.
- Malmkvist J, Damgaard BM, Pedersen LJ, Jørgensen E, Thodberg K, Chaloupková H and Bruckmaier RM 2009. Effects of thermal environment on hypothalamic-pituitary-adrenal axis hormones, oxytocin, and behavioral activity in periparturient sows. *Journal of Animal Science* 87, 2796–2805.
- Marchant JN, Broom DM and Corning S 2001. The influence of sow behaviour on piglet mortality due to crushing in an open farrowing system. *Animal Science* 72, 19–28.
- Marchant JN, Rudd AR, Mendl MT, Broom DM, Meredith MJ, Corning S and Simmins PH 2000. Timing and causes of piglet mortality in alternative and conventional farrowing systems. *Veterinary Record* 147, 209–214.
- Mckee CI and Dumelow J 1995. A review of the factors involved in developing effective non-slip floors for pigs. *Journal of Agricultural Engineering Research* 60, 35–42.
- Mitchell CD and Smith WJ 1978. Piglet foot dimensions for design of slotted floors. *Farm Buildings Progress* 51, 7–9.
- Morrow-Tesch J and McGlone JJ 1990. Sources of maternal odors and the development of odor preferences in baby pigs. *Journal of Animal Science* 68, 3563–3571.
- Mount LE 1967. The heat loss from new-born pigs to the floor. *Research in Veterinary Science* 8, 175–186.
- Mount LE 1968. *The climatic physiology of the pig*. Edward Arnold, London, UK.
- Moustsen VA and Poulsen HL 2004a. Anbefalinger vedr. dimensioner på fareboks og kassesti. Landsudvalget for Svin, Danske Slagterier, Faglig Publikation 0414.
- Moustsen VA and Poulsen HL 2004b. Pattegrise dimensioner. Landsudvalget for Svin, Danske Slagterier, Faglig Publikation 0432.
- Moustsen VA, Poulsen HD and Nielsen MBF 2004. Krydsningssoer dimensioner. Landsudvalget for Svin, Danske Slagterier, Faglig Publikation 649.
- Mouttoto N and Green LE 1999. Incidence of foot and skin lesions in nursing piglets and their association with behavioural activities. *Veterinary Record* 145, 160–165.
- Mouttoto N, Hatchell FM and Green LE 1999. The prevalence and risk factors associated with forelimb skin abrasions and sole bruising in preweaning piglets. *Preventive Veterinary Medicine* 39, 231–245.
- Newberry RC and Swanson JC 2008. Implications of breaking mother-young social bonds. *Applied Animal Behaviour Science* 110, 3–23.
- Noblet J and Le Dividich J 1981. Energy-metabolism of the newborn pig during the 1st 24-h of life. *Biology of the Neonate* 40, 175–182.
- Oliviero C, Heinonen M, Valros A, Halli O and Peltoniemi OAT 2006. Duration of farrowing is longer in sows housed in farrowing crates than in pens. *Reproduction in Domestic Animals* 41, 367–367.
- Olsson ACh and Svendsen J 1997. The importance of familiarity when grouping gilts, and the effect of frequent grouping during gestation. *Swedish Journal of Agricultural Research* 27, 33–43.
- Pajor EA 1998. Parent-offspring conflict and its implications for maternal housing systems in domestic pigs. PhD, McGill University.
- Pajor EA, Weary DM, Fraser D and Kramer DL 1999. Alternative housing for sows and litters: 1. Effects of sow-controlled housing on responses to weaning. *Applied Animal Behaviour Science* 65, 105–121.
- Parfet KA and Gonyou HW 1991. Attraction of newborn piglets to auditory, visual, olfactory and tactile stimuli. *Journal of Animal Science* 69, 125–133.
- Parratt CA, Chapman KJ, Turner C, Jones PH, Mendl MT and Miller BG 2006. The fighting behaviour of piglets mixed before and after weaning in the presence or absence of a sow. *Applied Animal Behaviour Science* 101, 54–67.
- Pedersen LJ, Malmkvist J and Jørgensen E 2007. The use of a heated floor area by sows and piglets in farrowing pens. *Applied Animal Behaviour Science* 103, 1–11.
- Pedersen LJ, Studnitz M, Jensen KH and Giersing AM 1998. Suckling behaviour of piglets in relation to accessibility to the sow and the presence of foreign litters. *Applied Animal Behaviour Science* 58, 267–279.
- Pedersen LJ, Damm BI, Marchant-Forde JN and Jensen KH 2003. Effects of feedback from the nest on maternal responsiveness and postural changes in primiparous sows during the first 24 h after farrowing onset. *Applied Animal Behaviour Science* 83, 109–124.
- Pedersen LJ, Holm L, Jensen MB and Jørgensen E 2005. The strength of pigs' preferences for different rooting materials measured using concurrent schedules of reinforcement. *Applied Animal Behaviour Science* 94, 31–48.
- Pedersen LJ, Jørgensen E, Heiskanen T and Damm BI 2006. Early piglet mortality in loose-housed sows related to sow and piglet behaviour and to the progress of parturition. *Applied Animal Behaviour Science* 96, 215–232.
- Petersen V 1994. The development of feeding and investigatory behaviour in free-ranging domestic pigs during their first 18 weeks of life. *Applied Animal Behaviour Science* 42, 87–98.

- Petersen HV, Vestergaard K and Jensen P 1989. Integration of piglets into social groups of free-ranging domestic pigs. *Applied Animal Behaviour Science* 23, 223–236.
- Petherick JC 1983a. A biological basis for the design of space in livestock housing. In *Farm animal housing and welfare* (ed. SH Baxter, MR Baxter and JAD MacCormack), pp. 103–121. Martinus Nijhoff Publishers, The Hague, The Netherlands.
- Petherick JC 1983b. A note on the space use for excretory behaviour of suckling piglets. *Applied Animal Ethology* 9, 367–371.
- Phillips PA and Fraser D 1991. Discovery of selected water dispensers by newborn piglets. *Canadian Journal of Animal Science* 71, 233–236.
- Phillips PA and Fraser D 2001. Modifying water nipples for newborn piglets. *Canadian Biosystems Engineering* 43, 5.1–5.4.
- Phillips PA, Fraser D and Thompson BK 1991. Preference by sows for a partially enclosed farrowing crate. *Applied Animal Behaviour Science* 32, 35–43.
- Phillips PA, Fraser D and Thompson BK 1992. Sow preference for farrowing-crate width. *Canadian Journal of Animal Science* 72, 745–750.
- Phillips PA, Fraser D and Pawluczuk B 2000. Floor temperature preference of sows at farrowing. *Applied Animal Behaviour Science* 67, 59–65.
- Pluske JR, Williams IH and Aherne FX 1995. Nutrition of the neonatal pig. In *The neonatal pig development and survival* (ed. MA Varley), pp. 187–235. CABI Publishing, Wallingford, UK.
- Puppe B, Schön PC, Tuchscherer A and Manteuffel G 2003. The influence of domestic piglets' (*Sus scrofa*) age and test experience on the preference for the replayed maternal nursing vocalisation in a modified open-field test. *Acta Ethologica* 5, 123–129.
- Quesnel H and Prunier A 1995. Endocrine bases of lactational anoestrus in the sow. *Reproduction Nutrition Development* 35, 395–414.
- Randall JM, Armsby AW and Sharp JR 1983. Cooling gradients across pens in a finishing piggery. 2. Effects on excretory behavior. *Journal of Agricultural Engineering Research* 28, 247–259.
- Rantzer D and Svendsen J 2001. Slatted versus solid floors in the dung area of farrowing pens: effects on hygiene and pig performance, birth to weaning. *Acta Agriculturae Scandinavica Section A-Animal Science* 51, 167–174.
- Robertson AM, Alliston JC and Bruce AI 1972. Dimensions of individual pens for dry and in-pig sows. *Farm Building Progress* 29, 29–36.
- Roehle R and Kalm E 2000. Estimation of genetic and environmental risk factors associated with pre-weaning mortality in piglets using generalized linear mixed models. *Animal Science* 70, 227–240.
- Rooke JA and Bland IM 2002. The acquisition of passive immunity in the new-born piglet. *Livestock Production Science* 78, 13–23.
- Schmidt I and Herpin P 1998. Carnitine palmitoyltransferase I (CPT I) activity and its regulation by malonyl-CoA are modulated by age and cold exposure in skeletal muscle mitochondria from newborn pigs. *Journal of Nutrition* 128, 886–893.
- Schouten WGP 1987. Effects of rearing conditions on the behavior of gilts around farrowing. *Applied Animal Behaviour Science* 17, 367–367.
- Signoret JP, Baldwin BA, Fraser D and Hafez ESE 1975. The behaviour of swine. In *The behaviour of domestic animals* (ed. ESE Hafez), pp. 349–389. Baillière Tindall and Cassell, London, UK.
- Silva BAN, Oliveira RFM, Donzele JL, Fernandes HC, Abreu MLT, Noblet J and Nunes CGV 2006. Effect of floor cooling on performance of lactating sows during summer. *Livestock Science* 105, 176–184.
- Stangel G and Jensen P 1991. Behaviour of semi-naturally kept sows and piglets (except suckling) during 10 days postpartum. *Applied Animal Behaviour Science* 31, 211–227.
- Stansbury WF, McGlone JJ and Tribble LF 1987. Effects of season, floor type, air temperature and snout coolers on sow and litter performance. *Journal of Animal Science* 65, 1507–1513.
- Stolba A and Wood-Gush DGM 1980. Arousal and exploration in growing pigs in different environments. *Proceedings Conference SVE, Applied Animal Ethology* 6, 382–383.
- Stolba A and Wood-Gush DGM 1981. The assessment of behavioural needs of pigs under free-range and confined conditions. *Applied Animal Ethology* 7, 388–389.
- Stolba A and Wood-Gush DGM 1984. The identification of behavioural key features and their incorporation into a housing design for pigs. *Annales De Recherches Veterinaires* 15, 287–299.
- Thodberg K, Jensen KH and Herskin MS 2002. Nest building and farrowing in sows: relation to the reaction pattern during stress, farrowing environment and experience. *Applied Animal Behaviour Science* 77, 21–42.
- Thodberg K, Jensen KH, Herskin MS and Jørgensen E 1999. Influence of environmental stimuli on nest building and farrowing behaviour in domestic sows. *Applied Animal Behaviour Science* 63, 131–144.
- Trivers RL 1974. Parent-offspring conflict. *Integrative and Comparative Biology* 14, 249–264.
- Valros A, Rundgren M, Špinko M, Saloniemi H, Rydhmer L, Hultén F, Uvnäs-Moberg K, Tománek M, Krejčí P and Algers B 2003. Metabolic state of the sow, nursing behaviour and milk production. *Livestock Production Science* 79, 155–167.
- Van de Weerd HA, Docking CM, Day JEL, Avery PJ and Edwards SA 2003. A systematic approach towards developing environmental enrichment for pigs. *Applied Animal Behaviour Science* 84, 101–118.
- van Wagenberg AV, van der Peet-Schwering C, Binnendijk GP and Claessen PJPW 2006. Effect of floor cooling on farrowing sow and litter performance: field experiment under Dutch conditions. *Transactions of the ASABE* 49, 1521–1527.
- Vasdal G, Møgedal I, Bøe KE, Kirkden R and Andersen IL 2010. Piglet preference for infrared temperature and flooring. *Applied Animal Behaviour Science* 122, 92–97.
- Vellenga L, Vanveen HM and Hoogerbrugge A 1983. Mortality, morbidity, and external injuries in piglets housed in 2 different housing systems. 1. Farrowing house. *Veterinary Quarterly* 5, 101–106.
- Walser EES 1986. Recognition of the sow's voice by neonatal piglets. *Behaviour* 99, 177–188.
- Waran NK and Broom DM 1993. The influence of a barrier on the behavior and growth of early-weaned piglets. *Animal Production* 56, 115–119.
- Wathes C and Whittemore CT 2006. Environmental management of pigs. In *Whittemore's science and practice of pig production* (ed. I Kyriazakis and CT Whittemore), pp. 533–592. Blackwell Publishing, Oxford, UK.
- Wattanukul W, Bulman CA, Edge HL and Edwards SA 2005. The effect of creep feed presentation method on feeding behaviour, intake and performance of suckling piglets. *Applied Animal Behaviour Science* 92, 27–36.
- Weary DM, Jasper J and Hotzel MJ 2008. Understanding weaning distress. *Applied Animal Behaviour Science* 110, 24–41.
- Weary DM, Pajor EA, Thompson BK and Fraser D 1996. Risky behaviour by piglets: a trade off between feeding and risk of mortality by maternal crushing? *Animal Behaviour* 51, 619–624.
- Weary DM, Phillips PA, Pajor EA, Fraser D and Thompson BK 1998. Crushing of piglets by sows: effects of litter features, pen features and sow behaviour. *Applied Animal Behaviour Science* 61, 103–111.
- Weber R, Keli N, Fehr M and Horat R 2007. Piglet mortality on farms using farrowing systems with or without crates. *Animal Welfare* 16, 277–279.
- Weber R, Keil NM, Fehr M and Horat R 2009. Factors affecting piglet mortality in loose farrowing systems on commercial farms. *Livestock Science* 124, 216–222.
- Wechsler B and Hegglin D 1997. Individual differences in the behaviour of sows at the nest-site and the crushing of piglets. *Applied Animal Behaviour Science* 51, 39–49.
- Wechsler B and Weber R 2007. Loose farrowing systems: challenges and solutions. *Animal Welfare* 16, 295–307.
- Welch AR and Baxter MR 1986. Responses of newborn piglets to thermal and tactile properties of their environment. *Applied Animal Behaviour Science* 15, 203–215.
- Watson TS 1978. The development of dunging preferences in piglets. *Applied Animal Ethology* 4, 293.
- Whittemore CT and Kyriazakis I 2006. Growth and body composition changes in pigs. In *Whittemore's science and practice of pig production* (ed. I Kyriazakis and CT Whittemore), pp. 65–103. Blackwell Publishing, Oxford.
- Widowski TM and Curtis SE 1990. The influence of straw, cloth tassel or both on the prepartum behavior of sows. *Applied Animal Behaviour Science* 27, 53–71.
- Widowski TM, Curtis SE, Dziuk PJ, Wagner WC and Sherwood OD 1990. Behavioral and endocrine responses of sows to prostaglandin F₂ alpha and cloprostenol. *Biology of Reproduction* 43, 290–297.
- Wiepkema PR 1986. Remarks on the behaviour of wild boar. *Applied Animal Behaviour Science* 15, 179–180.

Baxter, Lawrence and Edwards

Wischner D, Kemper N and Krieter J 2009. Nest-building behaviour in sows and consequences for pig husbandry. *Livestock Science* 124, 1–8.

Xu RJ, Wang F and Zhang SH 2000. Postnatal adaptation of the gastrointestinal tract in neonatal pigs: a possible role of milk-borne growth factors. *Livestock Production Science* 66, 95–107.

Zanella AJ and Zanella EL 1993. Nesting material used by free-range sows in Brazil. In *Proceedings of the 3rd Joint Meeting of the International Congress on Applied Ethology* (ed. M Nichelmann, HK Wierenga and S Braun), p. 411. KTBL, Berlin, Darmstadt, Germany.

Zoric M, Nilsson E, Lundeheim N and Wallgren P 2009. Incidence of lameness and abrasions in piglets in identical farrowing pens with four different types of floor. *Acta Veterinaria Scandinavica* 51, 23.

Zoric M, Nilsson E, Mattsson S, Lundeheim N and Wallgren P 2008. Abrasions and lameness in piglets born in different farrowing systems with different types of floor. *Acta Veterinaria Scandinavica* 50, 37.

Ziron M and Hoy S 2003. Effect of a warm and flexible piglet nest heating system – the warm water bed – on piglet behaviour, live weight management and skin lesions. *Applied Animal Behaviour Science* 80, 9–18.