



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Volatile and semi-volatile organic compounds in smoke exposure of firefighters during prescribed burning in the Mediterranean region

Citation for published version:

Barboni, T, Cannac, M, Pasqualini, V, Simeoni, A, Leoni, E & Chiaramonti, N 2010, 'Volatile and semi-volatile organic compounds in smoke exposure of firefighters during prescribed burning in the Mediterranean region' *International Journal of Wildland Fire*, vol 19, no. 5, pp. 606-612., 10.1071/WF08121

Digital Object Identifier (DOI):

[10.1071/WF08121](https://doi.org/10.1071/WF08121)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Author final version (often known as postprint)

Published In:

International Journal of Wildland Fire

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



VOLATILE AND SEMI-VOLATILE ORGANIC COMPOUNDS IN SMOKE EXPOSURE OF FIREFIGHTERS DURING PRESCRIBED BURNING IN THE MEDITERRANEAN REGION

Toussaint Barboni*, Magali Cannac, Vanina Pasqualini, Albert Simeoni, Eric Leoni
and Nathalie Chiaramonti

Université de Corse, UMR CNRS 6134, Sciences Pour l'Environnement, BP 52,
20250 Corte, France.

* Corresponding author

Abstract

Prescribed fires can be used as a forest management tool to reduce the severity of wildfires. Thus, over prolonged and repeated periods, firefighters are exposed to toxic air contaminants. This work consisted in collecting and analysing smoke released by typical Mediterranean vegetation during prescribed burning. Sampling was performed at five active zones on the island of Corsica. Seventy-nine compounds were identified: volatile organic compounds and semivolatile organic compounds, including polycyclic aromatic hydrocarbons. Depending on exposure levels, the toxins present in smoke may cause short-term or long-term damage to firefighters' health. The dangerous compounds emitted, benzene, toluene, ethylbenzene and xylenes, were quantified. Their concentrations varied as a function of the study site. These variations were due to the intrinsic and extrinsic characteristics of the fire site (e.g. plant species, fire intensity and wind). Our results show that benzene concentration is high during prescribed burning, close to the exposure limit value or short-term exposure limit. Benzene can be considered as a toxicity tracer for prescribed burning because its concentration was above the exposure limit value at all the study sites. The authors suggest that respirators should be used to protect staff during prescribed burning operations.

Additional keywords: air quality, BTEXs, firefighter exposure, forest fire smoke, SVOC, VOC.

Introduction

Every year, thousands of hectares of forest are destroyed by wildfires in the Mediterranean basin. In France, 4 million hectares of scrubland and 1 million hectares of pine forest are affected by fire during the summer. To prevent wildfires, prescribed burning is used as a forest management tool (Rigolot 2003; Fernandes and Botelho 2004). Prescribed burning: (i) reduces fuel load and fire risks; (ii) recreates natural disturbance dynamics; and (iii) restores the ecosystem (Weber and Taylor 1992; Graham et al. 2004; Fernandes and Rigolot 2007). During prescribed burning operations, personnel are repeatedly exposed to smoke and volatile organic compound (VOC) emissions for prolonged periods. In France, operating staff have no respiratory protection, so there is a real health risk (Reinhardt and Ottmar 2004; Stefanidou and Athanaselis 2004), because pollutants and particulate matter can penetrate the upper respiratory tract, and then down into the lungs. Smoke is considered as one of the damaging effects of prescribed burning. Many studies have been carried out on the

analysis of smoke during forest fires and prescribed burning in the USA (Kelly 1992; Reinhardt et al. 2000), Greece (Statheropoulos and Karma 2007), SE Asia (Muraleedharan et al. 2000), Australia (Reisen et al. 2006; Reisen and Brown 2009) and Russia (Lee et al. 2005). Determining the composition of smoke is complicated by the numerous compounds emitted from the external environment (e.g. plastics, fertilisers, pesticides), leading to secondary compounds (Statheropoulos and Karma 2007). The reaction of partial combustion releases a large number of molecular pollutants and particulates (Wade and Ward 1973; Ward 1989; Brown and Bradshaw 1994; Korontzi et al. 2003; Miranda et al. 2005; Statheropoulos and Karma 2007; Wise 2008). The compounds emitted in smoke have an impact on air quality and human health. Smoke consists of permanent gases, VOCs, semi-volatile organic compounds (SVOCs) and particulates. The permanent gases in the smoke from forest fires are CO, which is the major toxic component, CO₂ and NO_x (NO+NO₂) (Muraleedharan et al. 2000). The VOCs emitted include ethane (Ward 1989; Miranda 2004) and aldehydes (Schauer et al. 2001; Reinhardt and Ottmar 2004; Statheropoulos and Karma 2007), together with benzene, toluene, ethylbenzene and xylene (Reh and Deitchman 1992; Hoerning et al. 1996; Friedli et al. 2001; Statheropoulos and Karma 2007), and the SVOCs include polycyclic aromatic hydrocarbons (PAHs; Kelly 1992; Muraleedharan et al. 2000) and organic acids (Ward 1989; Dost 1991). Exposure to high levels of carbon monoxide (CO), respirable particulates, aldehydes, VOCs and SVOCs can cause short-term adverse health effects to firefighters, such as coughing, eye irritation, shortness of breath, headaches, dizziness and nausea (Dost 1991; Reisen and Brown 2009).

No epidemiological studies have been carried out on the firefighters involved. However, epidemiological studies of indoor and community exposure to biomass smoke indicate a consistent relationship between exposure and increased symptoms of respiratory troubles, increased risk of respiratory illness and decreased lung function (Johnson et al. 2002; Chen et al. 2006). The health effects of long-term exposure to smoke from forest fires can include nose and eye irritation, acute respiratory infections and lung function problems (Malilay 1999; Reinhardt et al. 2000). Short-term exposure has been associated with lung function decrease and increases in the airway sensitivity of firefighters (Fowler 2003; Slaughter et al. 2004). Bushfires and prescribed burning episodically expose humans to high levels of smoke pollutants for short periods of time (Morawska and Zhang 2002). Concentrations exceeding exposure limit values of PM₁₀ (particulate matter 10 µm in size) (Reh and Deitchman 1992; Mott et al. 2005), PM_{2.5} (particulate matter 2.5 µm in size) (Statheropoulos and Karma 2007), CO and formaldehyde (Reinhardt et al. 2000) have been reported. The VOCs determined were the products of several metabolic pathways (endogenous compounds) on inhaled contaminants (exogenous compounds). Fires are known to produce a variety of these compounds, but little characterisation work has been done. If inhaled, benzene causes headaches, dizziness, nausea, confusion and respiratory tract irritation. In the long term, it is a human carcinogen (Margossian 2002). Long-term exposure to toluene can cause organic psychosyndromes and cardiomyopathy. No detrimental effects of metabolites due to ethylbenzene have been reported in humans. Xylenes can cause vomiting and a risk of pulmonary oedema in the short term (Margossian 2002). The possible synergistic effects of exposure to a mixture of toxic substances in forest fire smoke are unknown. To our knowledge, no study has been carried out on the toxicity of smoke from the prescribed burning of typical Mediterranean vegetation in Europe.

The type and the concentration of compounds emitted into the atmosphere

from burning plants depend on the type of fuel, fireline intensity, fuel moisture, meteorology and geographical location (Miranda et al. 2005, 2007). The forest-firefighting community has expressed its concern about the health risks due to short- or long-term exposure to this smoke.

The general aim of the present work is to contribute to the identification of this source of atmospheric pollution, obtain data regarding smoke emissions in Mediterranean forests and inform firefighters of the risks faced owing to smoke inhalation.

This specific aims of this study are: (i) to identify the VOCs and SVOCs emitted during prescribed burning in order to evaluate the impact of smoke on the firefighters; (ii) to quantify the BTEXs and (iii) to compare the benzene, toluene, ethylbenzene and xylenes (BTEX) toxicity with the short-term exposure limit (STEL) and measure the health risks for firefighters, at five sites in Corsica with typical Mediterranean vegetation. The next section presents the materials and methods used for the present work, followed by a presentation of the results with a discussion of the health risks encountered by prescribed burning operating personnel.

Materials and methods

Experimental plots and sampling

The sampling was conducted on the island of Corsica (France–western Mediterranean, Fig. 1) at five different sites. Table 1 lists the sites and their characteristics. The sites were selected by the ONF (Office National des Forêts) and forest-fire prevention operating staff. Sampling was performed by the same operating personnel who performed the prescribed burning at the five sites. VOC and SVOC emissions were measured on mixed samples (i.e. without distinguishing between the flaming and smouldering phases). The distance between the sampling area and the fire front was between 1 and 10 m, corresponding to the position of the operating staff. The staff consisted of four persons who controlled the fire by means of flaming torches. Each person was equipped with a portable sampling pump. The operating staff performed the sampling with Tenax TA[®] tubes (Supelco[®], Saint-Quentin Fallavier, France) at three different times during the prescribed burn, in order to provide triplicate measurements. Sampling was carried out for a period of 15 min at a flow-rate of 150 mL min⁻¹. Twelve samples were collected at each site and analysed. In total, 60 samples were studied. The Tenax TA sorbent tubes used were multibed glass tubes, with the dimensions 11.5 x 6 mm outer diameter (o.d.) x 4 mm inner diameter (i.d.), in order to capture low - and medium - volatility organic compounds. The sorbent tubes were transferred to a freezing box in the laboratory and analysed the next day using automatic thermal desorption–gas chromatography–mass spectrometry (ATD-GC/MS). It should be noted that sorbent tubes have previously been used in other studies for sampling VOCs and SVOCs (Reinhardt et al. 2000; Reisen et al. 2006) before GC analysis.

Exposure limit

All over the world, national health and safety commissions or agencies set concentrations of substances in the air at or below which exposure is considered to be acceptable. In the USA, the National Institute for Occupational Safety and Health (NIOSH) advises the Occupational Safety and Health Administration (OSHA) on health hazards in the workplace and establishes recommended exposure limits that are based on thorough reviews of available scientific information. In France, CIRC

(Centre International de la Recherche sur le Cancer) and other agencies (Institut National de la Recherche et de la Santé, Agence Française de Sécurité Sanitaire de l'Environnement et du Travail) publish lists of carcinogenic substances or those suspected of being so. An important parameter is the STEL, a maximal concentration to which workers can be continuously exposed for up to 15 min without adverse effect. In the present work, different products involved in burning biomass were identified and quantified and their values are compared with the STEL.

ATD-GC/MS analysis of VOCs

The analyses were carried out 1 day after each experiment in the laboratory using an automatic thermal desorber Perkin Elmer_ ATD Turbomatrix. For the thermal desorption of VOCs, helium (He) flow was set at 30 mL min⁻¹ with a column head pressure at 3626 x 10⁻⁶ Pa (25 psi). The sorbent tube was brought to 280°C over 10 min and a carrier gas flushed the sample towards a cold trap at 5°C. In a second step, the cold trap (22 cm, 0.53mm i.d.; Supelco) was programmed to increase temperature from 5 to 280°C at 40°C s⁻¹, then held isothermally at 280°C for 3 min. The compounds were then desorbed onto the chromatograph under helium as carrier via a heated transfer line maintained at 280°C. The injector temperature was set to 280°C. The ionization energy for mass detection was set to 70 eV and electron ionisation mass spectra were acquired over the mass range 35–350 Da. The chromatograph and the mass spectrometer were Perkin Elmer Clarus 500[®] apparatus (Waltham, MA, USA). The chromatograph was equipped with a non-polar column (Rtx[®]-1 fused silica column, dimethylsiloxane, Resteck, Lisses, France), length 60 m and i.d. 0.22 µm. This column was coupled with the mass detector. Detection was carried out by a quadripolar analyser made up of an assembly of four parallel electrodes of cylindrical section. The oven temperature of the chromatograph was programmed from 50 to 260°C at 2°C min⁻¹ and then held isothermally at 260°C for 10 min.

Identification and quantification

The methodology used to identify individual components was based on the comparison of their GC retention indices (RI) on non-polar and polar columns with those of authentic compounds or literature data (König et al. 2001; NIST WebBook, see <http://webbook.nist.gov/chemistry>, accessed 2006). The RIs on nonpolar and polar columns were determined relative to the retention time of a series of n-alkanes with linear interpolation (van Den Dool and Kratz 1963). The results were computer-matched with commercial mass spectrum libraries (Adams 2001; König et al. 2001; <http://webbook.nist.gov/chemistry>) and spectra were compared with those in our own library of authentic compounds.

The majority of these compounds were commercial standard components. The relative amounts of the individual components were calculated on the basis of their GC peak areas, obtained on a capillary Rtx-1 column, without flame ionisation detector (FID) response factor correction.

Quantification was performed for benzene, toluene, ethylbenzene and xylenes (o-xylene, m-xylene and p-xylene). The method of external calibration was used with a commercial mixture of BTEX in methanol from Restek[®] (Lisses, France) in the range 5–250 mg mL⁻¹. Triplicate injections of standards were made each set of concentrations for the curves of external calibration standards. The correlation coefficient (R²) for the linear regression of the curves of external calibration standards varied between 0.996 and 0.999, indicating a good correlation between the detector's

response and the concentration of injected products.

Statistical analysis

Analyses of variance (ANOVA test) were used when the conditions of application were satisfied (variance homogeneity and normality of the populations). Pearson's test was carried out to find correlations between BTEXs concentrations and many parameters such as abiotic factors and fire data. The Statgraphics for Windows program of Statistical Graphics Corporation_ (Waltham, MA, USA) was used for this test.

Results and discussion

Identification of VOCs during prescribed burning

ATD-GC/MS analysis of the five prescribed burning sites led to the identification of 79 compounds (Table 2), including benzene and benzene derivatives, phenol and phenol derivatives, organic acids, terpenic compounds, alkanes, alkenes and aldehyde compounds. Other ring systems included furans, 1H-indene, pyrroles, and naphthalene and their derivatives. These compounds were reported at sites A–E (Table 2). Many other compounds (VOCs and SVOCs) could not be identified because their mass spectra or retention indices were not included in our databank or in commercial databanks. Among the 79 individual compounds, 65 were identified by comparing their electronimpact (EI) mass spectra and retention indices with those of our own library. Compounds 12, 16, 27, 36, 38, 45, 47, 49, 63, 65, 66, 68, 69 and 71 were identified by comparing their EI-mass spectra and retention indices with those of commercial libraries or literature data (Ward 1989; Reinhardt et al. 2000; Andreae and Merlet 2001; Reisen et al. 2006; Statheropoulos and Karma 2007). The same compounds (reported in Table 2) had previously been found in forest fires (Ward 1989; Andreae and Merlet 2001; Statheropoulos and Karma 2007) and prescribed burning operations (Reinhardt et al. 2000; Reisen et al. 2006). The smoke composition (VOCs and SVOCs) was approximately the same at the five sampling sites and in the literature. Indeed, many studies (Austin et al. 2001; Austin and Wang 2002) show that the composition of the smoke produced by fires of various kinds is similar.

Only at site E were five monoterpenes (33, 44, 46, 54 and 64; Table 2) and five sesquiterpenes (75–79; Table 2) observed. This burning operation was conducted under *Pinus laricio*. The needles of this species emit these compounds by vapour-transpiration (biogenic volatile organic compounds; Rezzi et al. 2001).

Quantification of BTEX during prescribed burning

The concentrations of BTEX identified for all sites are given in Table 3. Total BTEX concentrations varied significantly between the different sites (ANOVA, $P < 0.05$). Benzene concentrations were not statistically different at sites A and B (ANOVA, $P < 0.05$). Toluene concentrations differed significantly between sites A and E, B and D, and site C (ANOVA, $P < 0.05$). There is a statistically significant difference between sites A, C and E regarding ethylbenzene concentrations, and between all the sites regarding xylene concentrations (ANOVA, $P < 0.05$).

The highest benzene concentrations were found at sites C (54 mg m^{-3}) and D (40 mg m^{-3}). The benzene levels at sites A, B, C and D were above the exposure limit value (ELV) and STEL (benzene: ELV, 30 mg m^{-3} ; STEL, 16 mg m^{-3}). For sites A

and B, the benzene levels were close to this limit value. The highest BTEX contents were measured at sites C and D, but their concentrations were much lower than the ELV. These higher concentrations can be explained by the high fuel load and the proportion of cellulose. The vegetation was dense at both sites (Table 1), resulting in much less complete combustion. The vegetation density of sites A, B and E was lower, resulting in lower BTEX concentrations. The correlation test between data plots (Table 1) and quantities of BTEX showed no correlation.

In comparison with literature data (Reinhardt et al. 2000; Reisen et al. 2006; Statheropoulos and Karma 2007), we found that BTEX concentrations were a 1000 times higher a few metres from prescribed burning operations than near forest fires. This can be explained by the distance between the sampling position and the flame front, which was between 1 and 10 m. In forest-fire studies, that distance is between 70 and 150m (Statheropoulos and Karma 2007).

In the present study, prescribed burning was carried out on different vegetation types, with a wide variety of parameters such as meteorological factors (pre-ignition temperature, fuel moisture hydrometry, wind, etc.), plant species, fuel load and fire intensity. All these parameters can influence the quantitative concentrations of BTEXs. Dokas et al. (2007) explain that the concentration of smoke from forest fires depends on meteorological data such as wind speed and direction, temperature and relative humidity. Ward (1989) show that PAH emissions increase with vegetation density. Faix et al. (1991) studied the relationship between the degradation of cellulose and the production of benzene derivatives. Low-diameter fuel loads burn almost completely at the combustion stage, which is not the case with broader-diameter fuels. Alén et al. (1996) show that the proportion of different compounds varies with degradation temperature. Butt (2006) explains that the concentration of phenols increases with oxygen concentration.

All these changes may be responsible for the variation in BTEXs between the study sites. The Pearson test was performed to investigate the correlation between the concentrations of BTEXs and the various parameters. No particular predominant factor was identified with regard to the variations in these compounds. The fire intensity of prescribed burning is considerably lower than that of a forest fire, resulting in a slower combustion and more often incomplete combustion process.

Health risks

Table 3 shows that the concentration of total BTEX was well above the ELV at sites A–D. It was only below the ELV at site E. Concentrations of benzene, which is considered a potential carcinogen, exceeded STEL (and ELV) limits at all the sites. Exposure to benzene at 160 mg m^{-3} (for 60 min) leads to symptoms of illness, and exposure to between 16 and 50 mg m^{-3} (for 5 hours) results in headaches, lassitude and weakness (Statheropoulos and Karma 2007). In our experiments, benzene concentrations were between 12 and 54 mg m^{-3} , implying shortterm intoxication.

Timothy et al. (2000) established toxic potential P(t) by summing the concentrations of different toxic compounds. Applying this P(t) to TEX (toluene, ethylbenzene and xylenes) levels at all the study sites gave a toxic potential of TEXs less than 1. TEXs are molecules with relatively low toxicity and high ELVs. In the literature, the concentrations of VOCs measured were not greater than limit values (Reinhardt et al. 2000; Miranda et al. 2005; Reisen et al. 2006; Statheropoulos and Karma 2007).

There are other compounds of importance with regard to human health risk. Phenolic compounds (phenol and derivatives identified in the smoke from the sites;

Table 2) are important because they are serious irritants, produced in abundance by the partial oxidation of cellulose fuels. Naphthalene (59) and naphthalene derivatives (67, 70) are carcinogens. Some aldehydes are extremely irritating to human mucous membranes (12, 24, 32, 34, 45 and 61).

The long-term health effects have not been studied, making it difficult to correlate exposure to toxic smoke with inherent risks. On the basis of our results, the authors suggest the use of respirators to protect staff involved in prescribed burning operations.

Conclusions

The present work consisted in the chemical analysis of the smoke released by prescribed burning operations in Corsica. ATD-GC/MS analysis of smoke from burning biomass led to the identification of 79 compounds. These compounds were derived from the distillation and decomposition of biopolymers. Some of them are toxic or harmful to humans, such as BTEXs. Although the toxicity of the smoke from prescribed burning was not clearly established, we showed that there was a potential risk for the health of firefighters. Benzene concentrations were close to the ELV or STEL, implying that this compound can be considered as a tracer of toxicity for prescribed burning.

At the international scale, the wearing of a respirator is not compulsory. In France, firefighters are only protected from exposure to large particles by wearing a hood. As a precaution, the authors consider that firefighters involved in prescribed burning operations should wear a protective respirator, which would keep out fine particles and decrease the quantities of dangerous compounds inhaled.

Acknowledgments

We thank the ONF and the forest firefighters for carrying out prescribed burning on our study sites. The authors acknowledge the European Community for partial financial support (Programme Interêt Communautaire). We are pleased to acknowledge financial support from the Ministry of Agriculture and Fishing (Groupement d'Intérêt Scientifique Incendie program) and the National Research Agency under contract no. NT05-2_44411.

References

- Adams RP (2001) 'Identification of Essential Oils by Capillary Gas Chromatography/Mass Spectroscopy.' (Allured: Carol Stream, IL)
- Alén R, Kuoppala E, Oesch P (1996) Formation of the main degradation compound groups from wood and its components during pyrolysis. *Journal of Analytical and Applied Pyrolysis* 36, 137–148.
- Andreae MO, Merlet P (2001) Emission of trace gases and aerosols from biomass burning. *Global Biogeochemical Cycles* 15, 955–966.
- Austin CC, Wang D (2002) VOC chemical fingerprinting: combustion sources and environmental monitoring. In 'EnviroAnalysis: the 4th Biennial International Conference on Chemical Measurement and Monitoring of the Environment', 27–30 May, Toronto, ON, Canada.
- Austin CC, Wang D, Ecobichon DJ, Dussault G (2001) Characterization of volatile organic compounds in smoke at municipal structural fires. *Journal of Toxicology and Environmental Health. Part A* 63, 437–458.

Brown JK, Bradshaw LS (1994) Comparisons of particulate-emissions and smoke impacts from presettlement, full suppression, and prescribed natural fire period in the Selway-Bitterroot Wilderness. *International Journal of Wildland Fire* 4(3), 143–155.

Butt D (2006) Formation of phenols from the low-temperature fast pyrolysis of Radiata pine (*Pinus radiata*) – Part II. Interaction of molecular oxygen and substrate water. *Journal of Analytical and Applied Pyrolysis* 76, 48–54.

Chen L, Verrall K, Tong S (2006) Air particulate pollution due to bushfires and respiratory hospital admissions in Brisbane, Australia. *International Journal of Environmental Health Research* 16, 181–191.

Dokas I, Statheropoulos M, Karma S (2007) Integration of field chemical data in initial risk assessment of forest fire smoke. *The Science of the Total Environment* 376, 72–85.

Dost FN (1991) Acute toxicology of components of vegetation smoke. *Reviews of Environmental Contamination and Toxicology* 119, 1–46.

Faix O, Fortman I, Bremer J, Meier D (1991) Thermal degradation products of wood, gas chromatographic separation and mass spectrometric characterization of polysaccharide-derived products. *Holz als Roh und Werkstoff* 49, 213–219.

Fernandes P, Botelho H (2004) Analysis of prescribed burning practice in the pine forest of north-western Portugal. *Journal of Environmental Management* 70, 15–26.

Fernandes P, Rigolot E (2007) The fire ecology and management of maritime pine (*Pinus pinaster* Ait.). *Forest Ecology and Management* 241, 1–13.

Fowler CT (2003) Human health impacts of forest fires in the southern United States: a literature review. *Journal of Ecological Anthropology* 7, 39–59.

Friedli HR, Atlas E, Stroud VR, Giovani L, Campas T, Radke LF (2001) Volatile organic trace gases emitted from North American wildfires. *Global Biochemistry Cycles* 15, 435–452.

Graham RT, McCaffrey S, Jain TB (2004) Science basis for changing forest structure to modify wildfire behavior and severity. USDA Forest Service, Rocky Mountain Research Station, General Technical Report RMRS-GTR-120. (Fort Collins, CO)

Hoerning JM, Evans MA, Aerts DJ, Ragland KW (1996) Organic emissions from combustion of pine, plywood, and particleboard. *Energy & Fuels* 10, 299–304.

Jennings W, Shibamoto T (1980) ‘Qualitative Analysis of Flavor and Fragrance Volatiles by Glass Capillary Gas Chromatography.’ (Academic Press: New York)

Johnson FH, Kavanagh AM, Bowman DM, Scott RS (2002) Exposure to bushfire smoke and asthma: an ecological study. *The Medical Journal of Australia* 176, 535–538.

Joulain D, König WA (1998) ‘The Atlas of Spectral Data of Sesquiterpene Hydrocarbons.’ (EB-Verlag: Hamburg, Germany)

Kelly J (1992) Health Hazard Evaluation Report. USDI, National Park Service, Gallatin National Forest, Montana. HETA 91-312-2185. (US Department of Health and Human Services, Public Health Service, Centers for Disease Control and National Institute for Occupational Safety and Health: Cincinnati, OH)

König WA, Hochmuth DH, Joulain D (2001) ‘Terpenoids and Related Constituents of Essential Oils.’ Library of Massfinder 2.1. (University of Hamburg, Institute of Organic Chemistry: Hamburg)

Korontzi S, Ward DE, Susott RA, Yokelson RJ, Justice CO, Hobbs PV,

Smithwick EAH, Hao WM (2003) Seasonal variation and ecosystem dependence of emission factors for selected trace gases and PM_{2.5} for southern African savanna fires. *Journal of Geophysical Research* 108, 4758–4762.

Lee KH, Kim JE, Kim YJ, Kim J, Hoyningen-Huene W (2005) Impact of the smoke aerosol from Russian forest fires on the atmospheric environment over Korea during May 2003. *Atmospheric Environment* 39, 85–99.

Malilay J (1999) A review of factors affecting the human health impacts of air pollutants from forest fires. In 'Health Guidelines for Vegetation Fire Events: Background Papers'. pp. 255–270. (World Health Organization, Geneva)

Margossian N (2002) 'Risque Chimique.' (Dunod: Paris)

Miranda AI (2004) An integrated numerical system to estimate air quality effects of forest fires. *International Journal of Wildland Fire* 13, 217–226.

Miranda AI, Ferreira J, Valente J, Santos P, Amorim JH, Borrego C (2005) Smoke measurements during Gestosa-2002 experimental field fires. *International Journal of Wildland Fire* 14, 107–116.

Miranda AI, Monteiro A, Martins V, Carvalho A, Schaap M, Builjes P, Borrego C (2007) Forest Fires Impact on Air Quality over Portugal. In '29th International Technical Meeting on Air Pollution Modelling and its Application', 24–28 September, Aveiro, Portugal. (Ed. A Carvalho) pp. 185–192.

Morawska LH, Zhang J (2002) Combustion sources of particles. 1. Health relevance and source signatures. *Chemosphere* 49, 1045–1058.

Mott JA, Mannino DM, Alverson CJ, Kiyub A, Hashimb J, Lee T, Falther K, Redd SC (2005) Cardiorespiratory hospitalizations associated with smoke exposure during the 1997 South-east Asian forest fires. *International Journal of Hygiene and Environmental Health* 208, 75–85.

Muraleedharan TR, Radojevic M, Waugh A, Caruana A (2000) Chemical characterisation of the haze in Brunei Darussalam during the 1998 episode. *Atmospheric Environment* 34, 2725–2731.

RehCM, Deitchman SD (1992) Health hazard evaluation report, USDI National Park Service, HETA 88-320-2176. (Yellowstone National Park, WY)

Reinhardt TE, Ottmar RD (2004) Baseline measurements of smoke exposure among wildland firefighters. *Journal of Occupational and Environmental Hygiene* 1, 593–606.

Reinhardt TE, Ottmar RD, Hanneman A (2000) Smoke exposure among firefighters at prescribed burns in the Pacific Northwest. USDA Forest Service, Pacific Northwest Research Station, Research Paper PNW-RP-526. (Seattle, WA)

Reisen F, Brown SK (2009) Australian firefighters' exposure to air toxics during bushfire burns of autumn 2005 and 2006. *Environment International* 35, 342–352.

Reisen F, Brown S, Cheng M (2006) Air toxics in bushfire smoke – firefighters exposure during prescribed burns. *Forest Ecology and Management* 234(Suppl. 1), S144.

Rezzi S, Bighelli A, Mouillot D, Casanova J (2001) Composition and chemical variability of the needle essential oil of *Pinus nigra* subsp. *laricio* from Corsica. *Flavour and Fragrance Journal* 16, 379–383.

Rigolot E (2003) Le feu domestiqué: outil de gestion des espaces méditerranéens. *Forêt méditerranéenne* 1, 37–44.

Schauer JJ, Kleeman MJ, Cass G, Simoneit BT (2001) Measurement of emissions from air pollution sources. 3.C1–C29 organic compounds from fireplace combustion of wood. *Environmental Engineering Science* 35, 1716–1728.

Slaughter JC, Koenig JQ, Reinhardt TE (2004) Association between lung function and exposure to smoke among firefighters at prescribed burns. *Journal of Occupational and Environmental Hygiene* 1, 45–49.

Statheropoulos M, Karma S (2007) Complexity and origin of the smoke components as measured near the flame-front of a real forest fire incident: a case study. *Journal of Analytical and Applied Pyrolysis* 78, 430–437.

Stefanidou M, Athanaselis S (2004) Toxicological aspects of fire. *Veterinary and Human Toxicology* 46, 196–199.

van Den Dool H, Kratz PDA (1963) Generalization of the retention index system including linear temperature programmed gas-liquid partition chromatography. *Journal of Chromatography. A* 11, 463–471.

Wade DD, Ward DE (1973) An analysis of the Air Force Bomb Range Fire. USDA Forest Service, Southeastern Forest Experiment Station, Research Paper SE-105. (Asheville, NC)

Ward DE (1989) Air toxics and fireline exposure. In ‘Tenth Conference on Fire and Forest Meteorology’, 17–21 April 1989, Ottawa, Canada. (Eds DC McIver, H Auld, R Whitewood) pp. 184–193. (Forestry Canada: Ottawa, ON)

Weber MG, Taylor SW (1992) The use of prescribed fire in the management of Canada’s forested lands. *Forestry Chronicle* 68, 324–334.

Wise EK (2008) Meteorologically influenced wildfire impacts on urban particulate matter and visibility in Tucson, Arizona, USA. *International Journal of Wildland Fire* 17, 214–223.

Table 1. Main characteristics of the study sites and main parameters concerning prescribed fire

	Site A	Site B	Site C	Site D	Site E
Locality	Sainte Lucie de Porto-Vecchio	Cunoli	Corte	Quezra	Ospédale
Coordinates	41°42'N 09°20'E	41°59'N 08°54'E	42°18'N 09°09'E	41°46'N 09°08'E	41°41'N 09°12'E
Elevation (m)	0	650	400	805	950
Main species	<i>Quercus ilex</i> L., <i>Olea europaea</i> L., <i>Arbutus unedo</i> L., <i>Cistus monspeliensis</i> L., <i>Cytisus triflorus</i> L., <i>Erica arborea</i> L.	<i>Arbutus unedo</i> L., <i>Erica arborea</i> L., <i>Cistus monspeliensis</i> L.	Leaf of <i>Quercus ilex</i> L.	<i>Jasiperus nana</i> L.	Scrub burning under <i>Pinus laricio</i>
Date of prescribed fire	12 July 2004	10 June 2005	14 June 2005	16 June 2005	10 April 2007
Slope (%)	0–30	0–10	0	0–20	5–10
Vegetation height (cm)	80–250	80–150	1–10	10–20	20–35
Fuel load (g m ⁻²)	1160	1200	1370	1450	960
Vegetation cover (%)	50–60	50–60	60–70	70–80	30–40
Burning area (ha)	0.06	2	0.4	2	1
Relative humidity, RH (%)	20–23	22–25	24–25	30–35	21–23
Temperature (°C)	30–32	22–24	22–25	14–16	24–26
Wind velocity (km h ⁻¹)	40	20	<5	<5	<5

Table 2. Volatile organic compounds (VOCs) and semi-volatile organic compounds (SVOCs) identified during prescribed fires

Compounds are listed in order of elution from the apolar (Rtx-1) column. Compounds listed were present at 80% to the five sites (A–E) and collected in 12 samples. RII, literature retention indices on the apolar column (Jennings and Shibamoto 1980; Joulain and König 1998; NIST WebBook, see <http://webbook.nist.gov/chemistry>, accessed 2006). RIa, retention indices on the apolar Rtx-1 column. A correct isomer was not identified for data marked with an asterisk (*). N, the total number of samples for the all sites. Superscript letters represent the sites sampled

N	Compounds (n = 60)	RII	RIa	N	Compounds (n = 60)	RII	RIa
1	1-Propene ^{A,B,C,D,E}	290	<500	41	Phenol ^{A,B,C,D,E}	1002	998
2	Methyl chloride ^{A,B,C,D,E}	325	<500	42	Benzofuran ^{A,B,C,D}	1005	1002
3	2-Methylpropene ^{A,B,C,D,E}	388	<500	43	1,2,3-Trimethylbenzene ^{A,B,C,D,E}	–	1007
4	1-Pentene ^{A,B,C,D,E}	487	<500	44	<i>p</i> -Cymene ^E	1015	1010
5	Furan ^{A,B,C,D,E}	500	<500	45	4-Hydroxybenzaldehyde ^{A,B,C,D}	–	1013
6	Acetone ^{A,B,C,D}	530	515	46	Limonene ^{D,E}	1021	1020
7	2-Methylpropanal ^{A,B,C,D}	540	518	47	*Trimethylbenzene ^{A,B,C,D}	–	1025
8	3-Buten-2-one ^{A,B,C,D}	561	535	48	1H-Indene ^{A,B,C,D,E}	1071	1031
9	2-Methylpentane ^{A,B,C,D}	562	540	49	*Trimethylbenzene ^{A,B,C}	–	1037
10	2,3-Butanedione ^{A,B,C,D}	575	565	50	<i>o</i> -Cresol ^{A,B,C,D,E}	1048	1044
11	Hexane ^{A,B,C,D}	600	600	51	<i>p</i> -Cresol ^{A,B,C,D,E}	1051	1055
12	3-Methyl-butanal ^{A,B,C,D}	630	615	52	<i>m</i> -Cresol ^{A,B,C,D,E}	1048	1064
13	Benzene ^{A,B,C,D,E}	651	644	53	2-Methoxyphenol ^{A,B,C,D,E}	1074	1067
14	Acetic acid ^{A,B,C,D,E}	674	651	54	α -Terpinolene ^E	1080	1078
15	1-Heptene ^{A,B,C,D,E}	689	685	55	Nonanal ^{A,B,C,D,E}	1087	1083
16	2,5-Dimethylfuran ^{A,C,D}	697	690	56	Undecene ^{A,B,C,D,E}	1088	1086
17	Heptane ^{A,B,C,D,E}	700	700	57	Undecane ^{A,B,C,D,E}	1100	1100
18	1-Methylpyrrole ^{A,B,D,E}	715	711	58	1-Methyl-1H-indene ^{A,B,C,D,E}	1148	1130
19	Pyrrole ^{A,B,C,D,E}	768	721	59	Naphthalene ^{A,B,C,D,E}	1185	1161
20	Toluene ^{A,B,C,D,E}	756	749	60	4-Methyl-2-methoxy-phenol ^{A,B,C,D,E}	1190	1178
21	Cyclopentanone ^{A,B,E}	775	758	61	Decanal ^{A,B,C,D,E}	1188	1185
22	1-Octene ^{A,D,E}	797	764	62	Dodecane ^{A,B,C,D,E}	1200	1200
23	Octane ^{A,B,C,D,E}	800	800	63	4-Hydroxystyrene ^{A,B,C}	1215	1205
24	Furfural ^{A,B,C,D,E}	815	836	64	Linalylacetate ^E	1239	1240
25	Ethylbenzene ^{A,B,C,D,E}	854	845	65	1,2-Benzenediol ^{A,B,C}	–	1205
26	(<i>m</i> + <i>p</i>)-Xylene ^{A,B,C,D,E}	860	853	66	2,3-Dihydro-1H-inden-1-one ^{A,B,C,D,E}	1292	1275
27	Cyclohexanone ^{B,C,E}	875	870	67	2-Methylnaphthalene ^{A,B,C,D,E}	–	1276
28	Styrene ^{A,B,C,D,E}	880	873	68	4-Ethyl-2-methoxyphenol ^{A,B,C,D,E}	1270	1280
29	<i>o</i> -Xylene ^{A,B,C,D,E}	884	880	69	Vinyl guaiaicol ^{A,C}	–	1284
30	1-Nonene ^{A,B,C,D,E}	886	883	70	1-Methylnaphthalene ^{A,B,C,D,E}	1276	1291
31	Nonane ^{A,B,C,D,E}	900	900	71	2-Methoxy-4-vinylphenol ^{A,B,C,D,E}	1306	1301
32	Benzaldehyde ^{A,B,C,D}	947	929	72	2,6-Dimethoxyphenol ^{A,B,C,D,E}	1349	1319
33	α -Pinene ^{A,B,C,D,E}	942	931	73	Eugenol ^{C,E}	1351	1337
34	5-Methyl-2-furancarboxaldehyde ^{A,B,C,D}	944	940	74	Biphenyl ^{A,B}	1366	1359
35	Propylbenzene ^{A,B,C,D}	949	944	75	Longifolene ^E	1411	1413
36	*Ethyltoluene ^{A,B,C,D,E}	–	952	76	<i>trans</i> - β -Caryophyllene ^E	1421	1424
37	Benzonitrile ^{A,B,C,D,E}	965	962	77	β -Copaene ^E	1430	1431
38	*Ethyltoluene ^A	–	971	78	Germaacrene ^{D,E}	1480	1480
39	3-Methylstyrene ^{A,B,C,D}	988	984	79	δ -Cadinene ^E	1524	1515
40	Decene ^{A,B,C,D}	991	987				

Table 3. Concentrations of BTEXs (benzene, toluene, ethylbenzene and xylenes; mg m⁻³ ± standard deviation) identified at all sites in prescribed burning (in 15 min during the prescribed burning) and exposure limit value (ELV) and short-term exposure limit (STEL) values Data marked with a superscript letter (e.g. A, B, C, D, E) are significantly different among stations: ANOVA P < 0.05

Compounds	Site A (n = 12)	Site B (n = 12)	Site C (n = 12)	Site D (n = 12)	Site E (n = 12)	ELV	STEL
Benzene	30 ^A ± 2.6	27 ^A ± 2.9	54 ^B ± 3.0	40 ^C ± 4.3	12 ^D ± 1.8	30	16
Toluene	6 ^A ± 0.9	13 ^B ± 2.1	78 ^C ± 6.1	15 ^B ± 3.1	6 ^A ± 0.9	550	574
Ethylbenzene	5 ^A ± 0.8	23 ^B ± 1.9	62 ^C ± 6.9	23 ^B ± 2.8	2 ^D ± 0.3	650	662
Xylene	8 ^A ± 0.8	26 ^B ± 1.8	54 ^C ± 3.1	32 ^D ± 2.8	1 ^E ± 0.5	650	541
Total BTEX	49 ^A ± 5.1	89 ^B ± 8.7	224 ^C ± 28.0	110 ^D ± 12.0	21 ^E ± 3.5		




Fig. 1. Location of the study sites on Corsica (French) island.