



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Draft Genome Sequences of Strains of *Pasteurella multocida* Isolated from the United Kingdom and the United States

Citation for published version:

Lainson, FA, Dagleish, MP, Fontaine, M, Bayne, C & Hodgson, JC 2013, 'Draft Genome Sequences of Strains of *Pasteurella multocida* Isolated from the United Kingdom and the United States' *Genome announcements*, vol 1, no. 5, e00773-13., 10.1128/genomeA.00773-13

Digital Object Identifier (DOI):

[10.1128/genomeA.00773-13](https://doi.org/10.1128/genomeA.00773-13)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher final version (usually the publisher pdf)

Published In:

Genome announcements

Publisher Rights Statement:

Copyright © 2013 Lainson et al.

This is an open-access article distributed under the terms of the Creative Commons Attribution 3.0 Unported license.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Draft Genome Sequences of Strains of *Pasteurella multocida* Isolated from the United Kingdom and the United States

Frederick A. Lainson, Mark P. Dagleish, Michael Fontaine, Colin Bayne,* J. Christopher Hodgson

Moredun Research Institute, Edinburgh, United Kingdom

* Present address: Colin Bayne, Neogen Europe Ltd., the Dairy School, Auchincruive, Ayr, Scotland.

***Pasteurella multocida* is a major pathogen of farm animals and has worldwide distribution. Here we report the draft genome sequences of four strains that were isolated from animals in the United Kingdom and the United States and represent pathogenic and commensal presentation of the bacterium.**

Received 3 September 2013 Accepted 10 September 2013 Published 10 October 2013

Citation Lainson FA, Dagleish MP, Fontaine M, Bayne C, Hodgson JC. 2013. Draft genome sequences of strains of *Pasteurella multocida* isolated from the United Kingdom and the United States. *Genome Announc.* 1(5):e00773-13. doi:10.1128/genomeA.00773-13.

Copyright © 2013 Lainson et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to Frederick A. Lainson, alex.lainson@moredun.ac.uk.

The Gram-negative facultatively anaerobic bacterium *Pasteurella multocida* is a pathogen that affects cattle, sheep, goats, pigs, and poultry worldwide and causes severe respiratory or septicemic infection. It is also found commonly as a commensal colonizing the upper respiratory tract in apparently healthy animals. Hotchkiss et al. (1) used multilocus sequence typing to characterize variation in isolates and relate this to host range, geographical range, and virulence. This approach delineated broad groups such as bovine, ovine, and avian isolates; however, in order to develop effective control or diagnostic strategies it is necessary to have a more detailed understanding of the genetic variability within and between groups. Here we present draft genome sequences for three ovine or bovine isolates from the United Kingdom and one bovine isolate from United States.

Genome sequencing was performed using an Illumina Solexa Genome Analyzer at the GenePool sequencing facility at the University of Edinburgh, <http://genepool.bio.ed.ac.uk/>. De novo assembly was carried out using Velvet version 0.7 (2), and the resulting sequence data were submitted to NCBI analysis using the Prokaryotic Genomes Automatic Annotation Pipeline (PGAAP). Details of the origins of these strains are shown in Table 1. The cumulative sequence lengths and the numbers of predicted coding sequences are consistent with those of related *P. multocida*

strains sequenced previously (accession numbers NC_017764, NC_016808, NC_17027, and NC_002663). This indicates that coverage is relatively complete for each of these sequenced genomes.

Assembled contig sequences were mapped to the reference genome PM70 (3) (accession number NC_002663) using Nucmer (4). This showed extensive sequence similarity, with a high proportion of each genome mapping to PM70 with an identity of 98% or greater, as shown in Table 1. Each of these Nucmer alignments was also inspected using Mapview (4) and this showed extensive synteny, with each genome being colinear to PM70 along the majority of their length.

Despite the variation in host range and the diverse virulence characteristics among this collection of strains, their sequence variability appears to be quite localized and this may provide an insight into genetic adaptation to particular niches.

Nucleotide sequence accession numbers. The genome sequences for these four *P. multocida* strains have been deposited in the NCBI and assigned the following accession numbers: strain 2000, [ARNW000000000](https://www.ncbi.nlm.nih.gov/nuccore/ARNW000000000); P1933, [ARNY000000000](https://www.ncbi.nlm.nih.gov/nuccore/ARNY000000000); R11F, [ARNZ000000000](https://www.ncbi.nlm.nih.gov/nuccore/ARNZ000000000); and 1500E, [AQTL000000000](https://www.ncbi.nlm.nih.gov/nuccore/AQTL000000000). In each case the version described here is NNNN000000000.1.

TABLE 1 Details of the four sequenced *Pasteurella multocida* strains reported here

Strain name	Yr of isolation	Geographical location of isolation	Host species and tissue site of isolation	No. of contigs	Cumulative contig length (bp)	No. of proteins predicted ^d	% of genome with ≥98% identity to PM70 ^b
2000	2000	United Kingdom, Scotland	Bovine, lung	261	2,210,534	2,141	93.34
P1933	2004	United States	Bovine, lung	269	2,441,686	2,441	92.65
R11F	1999	United Kingdom	Bovine, vaginal	243	2,195,634	2,129	98.58
1500E	2000	United Kingdom, Scotland	Bovine, nasal	265	2,210,576	2,148	93.37

^a Number of proteins predicted in the PGAAP (Prokaryotic Genomes Automatic Annotation Pipeline) annotation process.

^b Percentage length of each genome that has an identity of 98% or greater to the reference genome PM70.

ACKNOWLEDGMENT

We thank the Rural and Environment Science and Analytical Services Division of the Scottish Government (RESAS) for financial support.

REFERENCES

1. Hotchkiss EJ, Dagleish MP, Willoughby K, McKendrick IJ, Finlayson J, Zadoks RN, Newsome E, Brulisauer F, Gunn GJ, Hodgson JC. 2010. Prevalence of *Pasteurella multocida* and other respiratory pathogens in the nasal tract of Scottish calves. *Vet. Rec.* 167:555–560.
2. Zerbino DR, Birney E. 2008. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res.* 18:821–829.
3. May BJ, Zhang Q, Li LL, Paustian ML, Whittam TS, Kapur V. 2001. Complete genomic sequence of *Pasteurella multocida*, Pm70. *Proc. Natl. Acad. Sci. U. S. A.* 98:3460–3465.
4. Delcher AL, Salzberg SL, Phillippy AM. 2003. Using MUMmer to identify similar regions in large sequence sets. *Curr. Protoc. Bioinformatics* 10:10.3. doi:10.1002/0471250953.bi1003s00.