



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Draft Genome Sequence of the Wolbachia Endosymbiont of *Drosophila suzukii*

Citation for published version:

Siozios, S, Cestaro, A, Kaur, R, Pertot, I, Rota-Stabelli, O & Anfora, G 2013, 'Draft Genome Sequence of the Wolbachia Endosymbiont of *Drosophila suzukii*' *Genome announcements*, vol 1, no. 1., 10.1128/genomeA.00032-13

Digital Object Identifier (DOI):

[10.1128/genomeA.00032-13](https://doi.org/10.1128/genomeA.00032-13)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher final version (usually the publisher pdf)

Published In:

Genome announcements

Publisher Rights Statement:

Freely available in PMC.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Draft Genome Sequence of the *Wolbachia* Endosymbiont of *Drosophila suzukii*

Stefanos Siozios, Alessandro Cestaro, Rupinder Kaur, Ilaria Pertot, Omar Rota-Stabelli, Gianfranco Anfora

Research and Innovation Centre, Fondazione Edmund Mach, San Michele all'Adige (TN), Italy

***Wolbachia* is one of the most successful and abundant symbiotic bacteria in nature, infecting more than 40% of the terrestrial arthropod species. Here we report the draft genome sequence of a novel *Wolbachia* strain named “wSuzi” that was retrieved from the genome sequencing of its host, the invasive pest *Drosophila suzukii*.**

Received 14 January 2013 Accepted 17 January 2013 Published 28 February 2013

Citation Siozios S, Cestaro A, Kaur R, Pertot I, Rota-Stabelli O, Anfora G. 2013. Draft genome sequence of the *Wolbachia* endosymbiont of *Drosophila suzukii*. *Genome Announc.* 1(1):e00032-13. doi:10.1128/genomeA.00032-13.

Copyright © 2013 Siozios et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to Stefanos Siozios, stefanos.siozios@fmach.it.

Drosophila suzukii (Matsumura) (Diptera *Drosophilidae*) is an invasive and destructive crop pest native to Southeast Asia that recently invaded Western countries, threatening both European and American fruit production (1, 2). The exploitation of symbiotic microorganisms for insect pest control has received considerable interest over the last few years (3). Owing to a pandemic distribution and unique ability to manipulate host reproduction, *Wolbachia* symbiotic bacteria are considered to be great candidates for their biocontrol potential (4, 5). Advances in genomics provide new opportunities for improving our understanding of *Wolbachia* biology.

As a serendipitous byproduct of *D. suzukii* genome sequencing (L. Ometto, A. Cestaro, S. Ramasamy, A. Grassi, S. Revadi, S. Siozios, M. Moretto, P. Fontana, C. Varotto, D. Pisani, T. Dekker, N. Wrobel, R. Viola, I. Pertot, D. Cavalieri, M. Blaxter, G. Anfora, and O. Rota-Stabelli, submitted for publication) the genome of its *Wolbachia* endosymbiont (“wSuzi”) was sequenced. By using the genome sequences of the wMel, wRi, wAna, wWil, and wSim *Wolbachia* strains as probes, we searched the raw sequences from *D. suzukii* sequencing and retrieved 1,082,694 *Wolbachia* matched reads (IlluminaHiSeqII 100-bp reads). Among the 1,082,694 reads, 1,054,920 were paired, corresponding to two pair-end libraries with average insert sizes of 180 and 300 bp, while the remaining 27,774 reads were singletons. *De novo* assembly was performed using the packages MIRA (6) under default parameters and Velvet (7) using a kmer of 65; we further assisted assembly using the AMOS_cmp assembler (8) and the genome of the wRi strain infecting *D. simulans* as a reference (9). To identify ambiguously assembled contigs, the three assemblies were simultaneously mapped against the wRi genome using the Geneious mapping algorithm (10). The final assembly yielded approximately 1.35 Mbp in 110 contigs, with a maximum length of 89,713 bp, a mean of 12,272 bp, and an average depth of coverage of 60×. The genome has an average G+C content of 35.2% and contains approximately 1,262 open reading frames (ORFs), as predicted by Glimmer v3.02 (11), and one copy of the 16S, 23S, and 5S rRNA genes and 34 tRNA genes, as predicted by tRNAscan_SE (12).

The draft genome of wSuzi displays great similarity with the

wRi genome and covers approximately 98% of its length. Indeed, all of the markers commonly used to discriminate between different *Wolbachia* strains (13–17) revealed no substitutions between wRi and wSuzi. The only exception was *dnaA*, with only one non-synonymous substitution separating the two strains. However, we were able to identify and validate with PCR several structural variations, such as indels and genomic rearrangements. The most relevant deletion is of approximately 21 kbp and is upstream of the WORIB prophage; this region includes mainly ankyrin repeat genes. There are also two large-scale rearrangements distinguishing the two genomes. The largest one corresponds to an inversion of a segment at least 80 kbp long. Finally, we found that most of the polymorphism between the two strains involves transposable elements: we could detect approximately 34 deletions of insertion sequence (IS) elements in wSuzi compared to the wRi genome. Our data indicate that *D. suzukii* carries a novel strain of *Wolbachia* whose close relationship with wRi hampers its discrimination with the classical MLST approach (16, 17). Further genetic and comparative studies will provide new insights into the biology of this *Drosophila*-*Wolbachia* association.

Nucleotide sequence accession numbers. The genome sequence of the *Wolbachia* endosymbiont of *D. suzukii* has been deposited at EMBL under accession numbers CAOU02000001 to CAOU02000110. The sequences are also available from the website of Fondazione Edmund Mach (<http://genomics.research.iasma.it/ds/twiki/bin/view/Main/ScaricaGenomi>).

ACKNOWLEDGMENT

The genome of *D. suzukii* was sequenced by the GenePool Genomics Facility of the University of Edinburgh.

We did not receive any specific funding for this work.

REFERENCES

1. Cini A, Ioriatti C, Anfora G. 2012. A review of the invasion of *Drosophila suzukii* in Europe and a draft research agenda for integrated pest management. *Bull. Insectol.* 65:149–160.
2. Rota-Stabelli O, Blaxter M, Anfora G. 2013. Quick guide: *Drosophila suzukii*. *Curr. Biol.* 23:R8–R9.
3. Douglas AE. 2007. Symbiotic microorganisms: untapped resources for insect pest control. *Trends Biotechnol.* 25:338–342.

4. Bourtzis K. 2008. *Wolbachia*-based technologies for insect pest population control, p 104–113. In Aksoy S (ed), *Transgenesis and the management of vector-borne disease*, vol 627. Springer Verlag, New York, NY.
5. Zabalou S, Riegler M, Theodorakopoulou M, Stauffer C, Savakis C, Bourtzis K. 2004. *Wolbachia*-induced cytoplasmic incompatibility as a means for insect pest population control. *Proc. Natl. Acad. Sci. U. S. A.* 101:15042–15045.
6. Chevreux B, Pfisterer T, Drescher B, Driesel AJ, Müller WEG, Wetter T, Suhai S. 2004. Using the miraEST assembler for reliable and automated mRNA transcript assembly and SNP detection in sequenced ESTs. *Genome Res.* 14:1147–1159.
7. Zerbino DR, Birney E. 2008. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res.* 18:821–829.
8. Pop M, Phillippy A, Delcher AL, Salzberg SL. 2004. Comparative genome assembly. *Brief. Bioinform.* 5:237–248.
9. Klasson L, Westberg J, Sapountzis P, Naslund K, Lutnaes Y, Darby AC, Veneti Z, Chen L, Braig HR, Garrett R, Bourtzis K, Andersson SG, Naslund K, Lutnaes Y, Darby AC, Veneti Z, Chen L, Braig HR, Garrett R, Bourtzis K, Andersson SGE. 2009. The mosaic genome structure of the *Wolbachia* wRi strain infecting *Drosophila simulans*. *Proc. Natl. Acad. Sci. U. S. A.* 106:5725–5730.
10. Drummond AJ, Ashton B, Cheung M, Heled J, Kearse M, Moir R, Stones-Havas S, Thierer T, Wilson A. 2008. Geneious, V 4.0. <http://www.geneious.com/>.
11. Delcher AL, Bratke KA, Powers EC, Salzberg SL. 2007. Identifying bacterial genes and endosymbiont DNA with glimmer. *Bioinformatics* 23:673–679.
12. Lowe TM, Eddy SR. 1997. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res.* 25:955–964.
13. O’Neill SL, Giordano R, Colbert AM, Karr TL, Robertson HM. 1992. 16S rRNA phylogenetic analysis of the bacterial endosymbionts associated with cytoplasmic incompatibility in insects. *Proc. Natl. Acad. Sci. U. S. A.* 89:2699–2702.
14. Werren JH, Zhang W, Guo LR. 1995. Evolution and phylogeny of *Wolbachia*: reproductive parasites of arthropods. *Proc. R. Soc. Lond. B Biol. Sci.* 261:55–63.
15. Zhou WG, Rousset F, O’Neill S. 1998. Phylogeny and PCR-based classification of *Wolbachia* strains using *wsp* gene sequences. *Proc. R. Soc. Lond. B Biol. Sci.* 265:509–515.
16. Baldo L, Hotopp JCD, Jolley KA, Bordenstein SR, Biber SA, Choudhury RR, Hayashi C, Maiden MCJ, Tettelin H, Werren JH. 2006. Multilocus sequence typing system for the endosymbiont *Wolbachia pipientis*. *Appl. Environ. Microbiol.* 72:7098–7110.
17. Paraskevopoulos C, Bordenstein SR, Wernegreen JJ, Werren JH, Bourtzis K. 2006. Toward a *Wolbachia* multilocus sequence typing system: discrimination of *Wolbachia* strains present in *Drosophila* species. *Curr. Microbiol.* 53:388–395.