



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

P21-PARP-1 pathway is involved in cigarette smoke-induced lung DNA damage and cellular senescence

Citation for published version:

Yao, H, Sundar, IK, Gorbunova, V & Rahman, I 2013, 'P21-PARP-1 pathway is involved in cigarette smoke-induced lung DNA damage and cellular senescence' PLoS One, vol 8, no. 11, e80007., 10.1371/journal.pone.0080007

Digital Object Identifier (DOI):

[10.1371/journal.pone.0080007](https://doi.org/10.1371/journal.pone.0080007)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher final version (usually the publisher pdf)

Published In:

PLoS One

Publisher Rights Statement:

Copyright: 2013 Yao et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



P21-PARP-1 Pathway Is Involved in Cigarette Smoke-Induced Lung DNA Damage and Cellular Senescence

Hongwei Yao^{1*}, Isaac K. Sundar¹, Vera Gorbunova², Irfan Rahman^{1*}

1 Department of Environmental Medicine, Lung Biology and Disease Program, Rochester, New York, United States of America, **2** Department of Biology, University of Rochester Medical Center, Rochester, New York, United States of America

Abstract

Persistent DNA damage triggers cellular senescence, which may play an important role in the pathogenesis of cigarette smoke (CS)-induced lung diseases. Both p21^{CDKN1A} (p21) and poly(ADP-ribose) polymerase-1 (PARP-1) are involved in DNA damage and repair. However, the role of p21-PARP-1 axis in regulating CS-induced lung DNA damage and cellular senescence remains unknown. We hypothesized that CS causes DNA damage and cellular senescence through a p21-PARP-1 axis. To test this hypothesis, we determined the levels of γ H2AX (a marker for DNA double-strand breaks) as well as non-homologous end joining proteins (Ku70 and Ku80) in lungs of mice exposed to CS. We found that the level of γ H2AX was increased, whereas the level of Ku70 was reduced in lungs of CS-exposed mice. Furthermore, p21 deletion reduced the level of γ H2AX, but augmented the levels of Ku70, Ku80, and PAR in lungs by CS. Administration of PARP-1 inhibitor 3-aminobenzamide increased CS-induced DNA damage, but lowered the levels of Ku70 and Ku80, in lungs of p21 knockout mice. Moreover, 3-aminobenzamide increased senescence-associated β -galactosidase activity, but decreased the expression of proliferating cell nuclear antigen in mouse lungs in response to CS. Interestingly, 3-aminobenzamide treatment had no effect on neutrophil influx into bronchoalveolar lavage fluid by CS. These results demonstrate that the p21-PARP-1 pathway is involved in CS-induced DNA damage and cellular senescence.

Citation: Yao H, Sundar IK, Gorbunova V, Rahman I (2013) P21-PARP-1 Pathway Is Involved in Cigarette Smoke-Induced Lung DNA Damage and Cellular Senescence. PLoS ONE 8(11): e80007. doi:10.1371/journal.pone.0080007

Editor: Amit Gaggar, University of Alabama-Birmingham, United States of America

Received: August 7, 2013; **Accepted:** October 7, 2013; **Published:** November 11, 2013

Copyright: © 2013 Yao et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This study was supported by the National Institutes of Health (NIH) 1R01HL092842, 2R01HL085613, 1R01HL097751 (to IR), American Lung Association RG-266456-N (to HY), and NIEHS Environmental Health Science Center grant P30-ES01247. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: Hongwei_Yao@urmc.rochester.edu (HY); irfan_rahman@urmc.rochester.edu (IR)

Introduction

Cigarette smoke (CS) is an important risk factor in chronic inflammatory pulmonary diseases including chronic obstructive pulmonary disease (COPD), due to increased inflammation, oxidative stress, apoptosis/proliferation, and premature senescence/aging [1,2]. Recently, we have shown that CS exposure causes premature senescence in lung cells, leading to airspace enlargement and lung function decline [3]. Persistent or severe DNA damage is able to trigger cellular senescence [4–6]. Indeed, increased DNA damage is observed in lungs of patients with COPD [7–12]. However, the molecular mechanism of CS-induced DNA damage and subsequent lung cellular senescence is unknown.

Accumulating evidence has shown that p21^{CDKN1A} (p21), the first identified inhibitor of cyclin/cyclin-dependent kinase complex, participates in the DNA damage response [13]. We, and others, have shown that CS increased the level of p21 protein in lung macrophages and epithelial cells *in vitro* and in mouse lungs *in vivo* [3,14–16]. Poly(ADP-ribose) polymerase 1 (PARP-1), an NAD⁺-dependent ADP-ribosyltransferase 1, is known as a damage sensor, which binds to damaged DNA and thereby promotes the cellular response to DNA single-strand breaks and double-strand breaks (DSB) [17,18]. For example, PARP-1-mediated PARylation promotes non-homologous end joining (NHEJ) through recruitment or retention of repair factors at DSB sites [19].

Recent studies have shown a physical association between p21 and PARP-1 [20], implicating an involvement of p21-PARP-1 pathway in DNA damage and repair. We, therefore, hypothesized that CS causes DNA damage and premature senescence through p21-PARP-1 signals. To test this hypothesis, we exposed the p21 knockout (p21^{-/-}) and C57BL/6J wild-type (WT) mice to CS for 3 days and determined the role of p21 in DNA damage and repair. Furthermore, a selective PARP-1 inhibitor 3-aminobenzamide (3-AB) was administered in p21^{-/-} and WT mice to study the role of p21-PARP-1 in CS-induced DNA damage and cellular senescence.

Materials and Methods

Ethics statement

All experimental protocols were performed in accordance with the standards established by the United States Animal Welfare Act, as set forth by the National Institutes of Health guidelines. The research protocol for these studies was approved by the University of Rochester Committee on Animal Research.

Mice and 3-AB administration

The p21^{-/-} mice was described previously [3,16], which were obtained from Dr. Michael O'Reilly (University of Rochester) who backcrossed them 10 generations to C57BL/6J after receiving from Dr. Philip Leder at Harvard Medical School (Boston, MA).




Figure 1. CS exposure induces DNA damage and cellular senescence associated with increased p21 level in mouse lungs. CS increased γ H2AX level (A), but reduced the levels of Ku70 in mouse lungs (B). Both SA- β -gal activity (C) and p21 level (D) were increased in lungs of CS-exposed mice. Gel pictures shown are representative of at least 3 separate mice. Relative density ratio is indicative of results after normalizing to corresponding GAPDH. Data are shown as mean \pm SEM (n = 3 to 6 per group). Original magnification, $\times 200$. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ vs air-exposed mice.

doi:10.1371/journal.pone.0080007.g001

3-AB (10 mg/kg, Calbiochem) was administered through intraperitoneal injection at 2 h prior to CS exposure daily for 3 d [21]. Control mice were intraperitoneally injected with saline.

CS exposure

Eight to ten weeks old mice were used for CS exposure as described previously [16,22]. The research grade cigarettes (3R4F, University of Kentucky) were used to generate smoke, and mice were exposed to CS according to the Federal Trade Commission protocol (1 puff/min of 2-s duration and 35-ml volume) with a Baumgartner-Jaeger CSM2072i automatic CS generating machine (CH Technologies). The smoke concentration was set at a value of ~ 300 mg/m³ total particulate matter by adjusting the flow rate of the diluted medical air, and the level of carbon monoxide in the chamber was ~ 350 ppm [22]. Mice received two 1 hour exposures (one hour apart) daily for 3 consecutive days, and were sacrificed at 24 h post-last exposure. Control mice were exposed to filtered air in an identical chamber according to the same protocol described for CS exposure.

Bronchoalveolar lavage (BAL)

The mouse lungs were lavaged three times with 0.6 ml of 0.9% NaCl after intraperitoneal injection of pentobarbiturate (100 mg/kg body weight) [22]. The BAL cell pellet was resuspended in 1 ml of 0.9% NaCl, and the total cell number was determined by counting on a hemocytometer. Differential cell counts (minimum of 500 cells per slide) were performed on cytopsin-prepared slides (Thermo Shandon) stained with Diff-Quik (Dade Behring).

Immunohistochemical staining of proliferating cell nuclear antigen (PCNA)

The PCNA staining was performed using a PCNA kit (Invitrogen, Camarillo, CA). The deparaffinized and rehydrated

lung sections were exposed to 3% H₂O₂ in methanol for 30 min to quench endogenous peroxidase activity after antigen retrieval using the citrate buffer (0.01 M, pH 6.0). Nonspecific binding of antibodies on the tissue sections were blocked by incubating the sections with 5% normal goat serum in PBS for 30 min. Lung tissue sections were incubated with a primary PCNA antibody at a titer of 1:100 overnight at 4°C. After being washed, sections were incubated with secondary antibody biotinylated anti-rabbit Ig (DAKO Corp.) for 1 h, and DAB (DAKO) was used as peroxidase substrate. The counterstaining with hematoxylin was performed before examination under a light microscope.

Preparation of whole cell lysate

The preparation of whole cell lysate from lung tissue was described previously [22]. Briefly, lung tissue (100 mg) was mechanically homogenized with 0.5 ml of radioimmunoprecipitation assay buffer, and the tissue homogenates were kept on ice for 45 min to allow complete cell lysis. Following centrifugation at 13,000 *g* in an eppendorf tube for 5 min, the supernatant was collected as whole cell lysate. Protein level in samples was measured with a BCA kit (Pierce).

Immunoblot

Protein samples were separated on a 6%–10% sodium dodecyl sulfate- polyacrylamide gel electrophoresis, and separated proteins were electroblotted onto nitrocellulose membranes (Amersham). The membranes were blocked for 1 h at room temperature with 5% BSA, and then probed with a 1:400–1:1000 diluted antibodies of anti-p21, anti-poly(ADP-ribose) (pADPr, PAR), anti-PCNA (Santa Cruz), anti-Ku70, anti-Ku80 (Abcam), and anti- γ H2AX (Millipore), anti-PARP-1, and anti-GAPDH (Cell Signaling). After three washing steps (10 min each), the levels of protein were detected using secondary antibody (1:5,000 dilution in 2.5% BSA in PBS containing 0.1% Tween (v/v) 20 for one hour) linked to




Figure 2. p21 deletion decreases the level of γ H2AX but augments the levels of Ku70 and Ku80 in mouse lung, which was affected by PARP-1 inhibitor. γ H2AX level was decreased, whereas the levels of Ku70 and Ku80 were augmented in lungs of p21^{-/-} mice compared to WT mice exposed to CS. 3-AB treatment further increased γ H2AX level, but reduced the levels of Ku70 in WT mice. Gel pictures shown are representative of at least 3 separate mice. Relative density ratio is indicative of results after normalizing to corresponding GAPDH. Data are shown as mean \pm SEM (n = 3 per group). * P < 0.05, *** P < 0.001 vs air group; †† P < 0.01, ††† P < 0.001, vs Veh group; † P < 0.05, †† P < 0.01, ††† P < 0.001 vs WT mice. doi:10.1371/journal.pone.0080007.g002

horseradish peroxidase (Dako), and bound complexes were detected using ECL method (Perkin Elmer). Equal loading of the samples was determined by quantitation of proteins and by reprobing membranes for GAPDH.

Senescence-associated β -galactosidase (SA- β -gal) activity assay

The SA- β -gal staining was performed in OCT-embedded frozen lung tissues using a commercial kit (Cell Signaling) [3]. Briefly, the lungs were fixed in 2% formaldehyde containing 0.2% glutaraldehyde for 15 min. After washed with PBS, the samples were incubated at 37°C for 24 h in the staining solution (pH 6.0) containing 1 mg/ml of X-gal. The cells with blue color are SA- β -gal positive. The SA- β -gal activity in lung homogenate was also measured by the rate of conversion of 4-methylumbelliferyl- β -D-galactopyranoside to the fluorescent hydrolysis product 4-methylumbelliferone at pH 6.0 [3]. Briefly, the lung tissues were homogenized in the lysis buffer (pH 6.0), and kept on ice for one hr. The lysates were centrifuged for 5 min at 12,000 g, and the supernatant was mixed with 2 \times reaction buffer containing 1.7 mM of 4-methylumbelliferyl- β -D-galactopyranoside, which was placed into a 37°C water bath for 3 h. Finally, 50 μ l of the reaction mix was added to 500 μ l of 400 mM sodium carbonate stop solution (pH 11.0), which was read at 150 μ l per well in a 96-well plate using a SpectrumMax M5 plate reader (Molecular Devices) with excitation at 360 nm, emission at 465 nm, 40 μ s integration, and gain held constant at 46. Normalized SA- β -gal activity is expressed as observed fluorescence divided by micrograms of total protein.

Statistical analysis

Statistical analysis of significance was calculated using one-way Analysis of Variance (ANOVA) for multigroup comparisons using STATVIEW. The results are shown as the mean \pm SEM. The P < 0.05 is considered as statistically significant.

Results

CS exposure causes DNA damage and cellular senescence along with increased p21 level

An early cellular response to DSB is the rapid phosphorylation of H2AX at Ser139 (γ H2AX), which is a sensitive molecular marker for DNA damage [23,24]. To determine whether CS causes DNA damage, we performed the immunoblotting for γ H2AX in mouse lungs. As shown in **Figure 1A**, the level of γ H2AX was significantly increased in mouse lungs exposed to CS for 3 days. We also observed that the levels of NHEJ proteins Ku70, but not Ku80, were reduced in CS-exposed mouse lungs (**Figure 1B**). Persistent DNA damage has been shown to cause cellular senescence [4,5,25]. Therefore, we determined cellular senescence via measuring SA- β -gal activity and p21 levels; and found that CS caused an increase in SA- β -gal activity and p21 level in mouse lungs (**Figure 1C and 1D**). These findings suggest that CS-induced DNA damage and cellular senescence are associated with increased p21 level.

Deletion of p21 attenuates CS-induced DNA damage and NHEJ repair impairment

To further determine the causal role of p21 in CS-mediated DNA damage, the p21^{-/-} and WT mice were exposed to CS for 3




Figure 3. p21 deletion increased the levels of PAR, which was reduced by 3-AB treatment. The levels of PAR and PARP-1 were determined by Western blot in lungs of C57BL/6J (A), and p21^{-/-} mice as well as WT littermates (B) in response to CS. PAR levels were increased in lungs of p21^{-/-} mice as compared to WT mice. 3-AB treatment reduced PAR level in lungs of both p21^{-/-} and WT mice. Intact PARP-1 level was not altered by either p21 deficiency or 3-AB treatment. Gel pictures shown are representative of at least 3 separate mice. Fold change is indicative of the alteration of PAR and PARP-1 compared with air-exposed and vehicle (Veh)-treated WT mice after normalizing to corresponding GAPDH or β -actin. Data are shown as mean \pm SEM (n=3-13 per group). * P <0.05 vs air group; + P <0.05, ++ P <0.01, +++ P <0.001 vs Veh group; +++ P <0.001 vs WT mice. doi:10.1371/journal.pone.0080007.g003

days in order to compare the lung DNA damage response. We found that p21 deletion decreased the level of γ H2AX in CS, but not in air-exposed mouse lungs (Figure 2). Further, the levels of Ku70 and Ku80 were increased in lungs of p21^{-/-} mice as compared to WT mice regardless of air or CS exposure (Figure 2). We then focused on whether p21-mediated DNA damage is associated with the regulation of PARP-1 since a physical interaction between p21 and PARP-1 is reported recently [20]. As expected, p21 deletion increased the level of PAR (reflecting PARP-1 activity) in both air- and CS-exposed mouse lungs, although CS exposure did not have any effect on the level of PAR in WT or p21^{-/-} mice (Figure 3A and 3B). The level of full length PARP-1 in lungs was increased by CS exposure in WT but not in p21^{-/-} mice (Figure 3B). These results suggest that p21 deletion attenuates CS-induced DNA damage, which is associated with PARP-1 activation.

PARP-1 inhibitor increases DNA damage and cellular senescence, but has no effect on lung inflammatory response to CS

As shown in Figures 3, PARP-1 activation upon p21 deletion was associated with protection against CS-induced DNA damage in mouse lungs. Therefore, we determined the role of PARP-1 in p21-mediated DNA damage by administering a selective PARP-1

inhibitor 3-AB to p21^{-/-} and WT mice prior to CS exposure. As expected, lung PARP-1 activity, which is reflected by PAR level, was inhibited after 3-AB treatment (Figure 3). Administration of 3-AB increased the levels of γ H2AX in lungs of WT, but did not increase in p21^{-/-} mice exposed to CS (Figure 2). 3-AB treatment significantly reduced the levels of Ku70 in lungs of both WT and p21^{-/-} mice exposed to CS (Figure 2). Furthermore, the level of Ku80 was decreased by 3-AB treatment in p21^{-/-}, but not in WT mice by CS exposure (Figure 2).

It is known that persistent DNA damage is able to induce cellular senescence [4-6,26]. Therefore, we determined the cellular senescence in response to PARP-1 inhibition by 3-AB; and found that 3-AB treatment increased SA- β -gal activity in lungs of WT, but not in p21^{-/-} mice exposed to CS (Figure 4A). Further, p21 deletion increased the expression of PCNA (a proliferation biomarker) in mouse lungs exposed to CS, which was reduced by 3-AB treatment (Figure 4B). However, the number of neutrophils in BAL fluid was not altered by 3-AB administration either in p21^{-/-} or WT mice in response to CS exposure, although p21 deletion reduced CS-induced neutrophil influx in BAL fluid (Figure 4C). Altogether, our data show that p21 deletion protects against CS-induced DNA damage and cellular senescence via PARP-1 activation.




Figure 4. 3-AB increases CS-induced cellular senescence, but does not affect neutrophil influx in mouse lungs. 3-AB augmented CS-induced increase in SA-β-gal activity in WT, but not p21^{-/-} mice (A). p21 deletion increased the expression of PCNA in lungs as compared to WT mice exposed to CS, which was reduced by 3-AB treatment (B). p21 deletion attenuated CS-induced neutrophil influx in BAL fluid, which was not affected by 3-AB (C). Original magnification, ×200. Gel pictures shown are representative of at least 3 separate mice. Relative density ratio is indicative of results after normalizing to corresponding GAPDH. Data are shown as mean ± SEM (n=3-4 per group). *P<0.05, **P<0.01 vs air group; †P<0.05, ††P<0.001 vs Veh group; †††P<0.001, ††††P<0.0001 vs WT mice. doi:10.1371/journal.pone.0080007.g004

Discussion

We have shown that cellular senescence and premature aging play important roles in the development of COPD/emphysema [2,3]. Persistent DNA damage has been shown to trigger cellular




Figure 5. A schematic model showing the role of p21-PARP-1 in acute cigarette smoke (CS)-induced DNA damage and cellular senescence. CS exposure causes DNA damage including double-strand break (DSB). CS exposure also increases the level of p21, and p21 gene deletion augments PARP-1 activity and the levels of non-homologous end joining (NHEJ) proteins to repair damaged DNA. Consequently, CS-induced cellular senescence is attenuated by p21 deletion. doi:10.1371/journal.pone.0080007.g005

senescence [4,5]. However, the mechanisms of CS-induced DNA damage and cellular senescence remain unknown. In this study, we found that DNA damage and NHEJ impairment occurred in lungs of CS-exposed mice. Furthermore, p21 deletion reduced DNA damage, but increased NHEJ repair proteins in mouse lungs in response to acute CS exposure. These findings suggest that p21 is an important mediator in CS-induced DNA damage and DNA repair impairment. However, the link between p21 and DNA damage/repair as well cellular senescence is not known.

It has been shown that p21 accumulates rapidly at DNA damage sites [27–29], which may be a marker for DNA damage. We found that the levels of both γH2AX (recognition of DNA damage) and p21 were increased in lungs of CS-exposed mice. This is in agreement with the findings that elevation of p21 level in senescent cells correlates with the accumulation of γH2AX [30]. The DSB is the most dramatic form of DNA damage, which is repaired predominantly by NHEJ [31–33]. Except for NHEJ pathway of DNA repair, Ku proteins (Ku70 and Ku80) also function as a molecular scaffold, to which other proteins involved in NHEJ can bind [34]. The levels of NHEJ proteins Ku70 were decreased by CS exposure. Hence, CS exposure not only causes DNA damage but also impairs NHEJ repair. It is possible that the regulation of other NHEJ factors (e.g., DNA ligase IV and DNA-PKcs) are also affected in response to CS exposure, which requires further studies. Interestingly, p21 deletion reduced the level of γH2AX, but increased the levels of Ku70 and Ku80 in mouse lungs, suggesting an important role of p21 in CS-induced DNA damage and repair impairment. This is consistent with the findings that p21 causes DNA damage, and inhibits nucleotide excision repair as well as NHEJ via a redox dependent mechanism [35–42]. However, recent studies have also demonstrated that p21 protein helps DNA repair and homologous recombination (HR)

[41,43,44]. The discrepancy among these studies may be due to the different experimental conditions (e.g., low and high levels of DNA damage imposed by oxidative stress) and DNA repair pathways (HR vs NHEJ) as well as the use of different cell (*in vitro*) and rodent (*in vivo*) model systems [13,45].

PARP-1, the most abundant nuclear enzyme of PARP family, catalyzes the formation of the polymer pADPr (PAR) using NAD⁺ as substrate. The level of PAR represents the relative activity of PARP-1. PARP-1 is activated in response to DNA damage, which participates in DNA repair and genomic integrity [17,18,46–49]. This is in agreement with ours and others' findings that PARP-1 level is increased during CS-induced DNA damage [47–51], whereas conflicting results were obtained in other studies [52,53]. Nevertheless, the role of PARP-1 in DNA damage and repair during the pathogenesis of COPD is implicated [54]. A recent study has shown a physical association between p21 and PARP-1 [20], implicating the regulation of p21-PARP-1 axis in CS-mediated DNA damage. We found that p21 knockout mice showed augmented PARP-1 activity in lungs when compared to WT mice regardless of air or CS exposure. This is corroborated by the findings that p21 negatively regulates PARP-1 activity [20]. Administration of PARP-1 inhibitor 3-AB increased the level of γ H2AX, but reduced Ku70 level in mouse lungs exposed to CS. This suggests a positive regulation of PARP-1 in NHEJ DNA repair [19,55]. We and others have shown that PARP-1 also regulates HR, which is used by cells to accurately repair breaks on both strands of DNA [46,56,57]. Hence, PARP-1 may control HR (e.g., RAD51) in response to CS exposure, which needs to be investigated. Overall, PARP-1 activation upon p21 deletion may mediate the protection against CS-induced DNA damage via NHEJ repair. PARP-1 can localize on the promoters, and impact chromatin integrity via PARylation [19,58], which may regulate the expression of genes and their encoding proteins such as Ku70. This may explain the alteration of Ku70 after PARP-1 inhibitor treatment in mouse lungs.

Persistent or severe DNA damage has been shown to trigger cellular senescence [4,5]. We found increased activity of SA- β -gal and p21 level along with augmented DNA damage in mouse lungs exposed to CS. Deletion of p21 reduced CS-induced DNA damage and cellular senescence [3]. Interestingly, PARP-1 inhibitor further increased CS-induced cellular senescence (i.e.,

increased SA- β -gal activity) in WT mouse lungs. This is in line with a study showing induction of cellular senescence via inhibition of PARP by veliparib [59]. We noticed that treatment with a PARP-1 inhibitor (3-AB) did not increase lung γ H2AX level or SA- β -gal activity as compared to vehicle control in p21^{-/-} mice in response to CS exposure. This may be due to the sufficient endogenous NHEJ or compensatory PARP-1 activity in these mice for repairing damaged DNA. Altogether, p21 deletion augments PARP-1 activity and NHEJ repair, thereby reduces CS-mediated DNA damage and subsequent cellular senescence (**Figure 5**). However, further studies are required to investigate the role of p21-PARP-1 in regulating lung DNA damage and repair as well as cellular senescence during the development of COPD/emphysema.

Senescent cells develop a PARP-1 and NF- κ B-associated secretome, and these cells are prone to secrete inflammatory mediators, such as IL-6 and IL-8 [60]. Interestingly, inhibition of PARP-1 by 3-AB did not exhibit any effect on CS-induced neutrophil influx. In fact, the effect of PARP-1 on inflammatory response is controversial. Most of the studies showed the pro-inflammatory effect of PARP-1 in lung inflammation [61–63], whereas increased lung inflammation occurred in PARP-1 knockout mice in response to hyperoxia [64]. The discrepancies among these studies are unresolved, which need further investigation.

In conclusion, our findings reveal the important role of p21-PARP-1 axis in regulating CS-induced lung DNA damage and NHEJ impairment as well as cellular senescence. Therefore, both PARP-1 and p21 would be potential targets in intervening CS-induced DNA damage and premature aging as well as subsequent chronic pulmonary diseases, such as COPD.

Acknowledgments

We thank Suzanne E Cook and Stephanie Uhrinek for their technical assistance.

Author Contributions

Conceived and designed the experiments: HY IKS VG IR. Performed the experiments: HY IKS. Analyzed the data: HY IKS. Contributed reagents/materials/analysis tools: VG. Wrote the paper: HY IKS IR.

References

1. Yao H, Rahman I (2011) Current concepts on oxidative/carbonyl stress, inflammation and epigenetics in pathogenesis of chronic obstructive pulmonary disease. *Toxicol Appl Pharmacol* 254: 72–85.
2. Hwang JW, Yao H, Caito S, Sundar IK, Rahman I (2013) Redox regulation of SIRT1 in inflammation and cellular senescence. *Free Radic Biol Med* 61C: 95–110.
3. Yao H, Chung S, Hwang JW, Rajendrasozhan S, Sundar IK, et al. (2012) SIRT1 protects against emphysema via FOXO3-mediated reduction of premature senescence in mice. *J Clin Invest* 122: 2032–2045.
4. Rodier F, Coppe JP, Patil CK, Hoeijmakers WA, Munoz DP, et al. (2009) Persistent DNA damage signalling triggers senescence-associated inflammatory cytokine secretion. *Nat Cell Biol* 11: 973–979.
5. Rodier F, Munoz DP, Teachenor R, Chu V, Le O, et al. (2011) DNA-SCARS: distinct nuclear structures that sustain damage-induced senescence growth arrest and inflammatory cytokine secretion. *J Cell Sci* 124: 68–81.
6. Campisi J (2013) Aging, cellular senescence, and cancer. *Annu Rev Physiol* 75: 685–705.
7. Caramori G, Adcock IM, Casolari P, Ito K, Jazrawi E, et al. (2011) Unbalanced oxidant-induced DNA damage and repair in COPD: a link towards lung cancer. *Thorax* 66: 521–527.
8. Pastukh VM, Zhang L, Ruchko MV, Gorodnya O, Bardwell GC, et al. (2011) Oxidative DNA damage in lung tissue from patients with COPD is clustered in functionally significant sequences. *Int J Chron Obstruct Pulmon Dis* 6: 209–217.
9. Ceylan E, Kocyigit A, Gencer M, Aksoy N, Selek S (2006) Increased DNA damage in patients with chronic obstructive pulmonary disease who had once smoked or been exposed to biomass. *Respir Med* 100: 1270–1276.
10. Deslee G, Woods JC, Moore C, Conradi SH, Gierada DS, et al. (2009) Oxidative damage to nucleic acids in severe emphysema. *Chest* 135: 965–974.
11. Neofytou E, Tzortzaki EG, Chatziantoniou A, Siafakas NM (2012) DNA Damage Due to Oxidative Stress in Chronic Obstructive Pulmonary Disease (COPD). *Int J Mol Sci* 13: 16853–16864.
12. Shi Y, Cao J, Gao J, Zheng L, Goodwin A, et al. (2012) Retinoic acid-related orphan receptor-alpha is induced in the setting of DNA damage and promotes pulmonary emphysema. *Am J Respir Crit Care Med* 186: 412–419.
13. Cazzalini O, Scovassi AI, Savio M, Stivala LA, Prospero E (2010) Multiple roles of the cell cycle inhibitor p21(CDKN1A) in the DNA damage response. *Mutat Res* 704: 12–20.
14. Marwick JA, Kirkham P, Gilmour PS, Donaldson K, Mac NW, et al. (2002) Cigarette smoke-induced oxidative stress and TGF-beta1 increase p21waf1/cip1 expression in alveolar epithelial cells. *Ann N Y Acad Sci* 973: 278–283.
15. Tomita K, Caramori G, Lim S, Ito K, Hanazawa T, et al. (2002) Increased p21(CIP1/WAF1) and B cell lymphoma leukemia-x(L) expression and reduced apoptosis in alveolar macrophages from smokers. *Am J Respir Crit Care Med* 166: 724–731.
16. Yao H, Yang SR, Edirisinghe I, Rajendrasozhan S, Caito S, et al. (2008) Disruption of p21 attenuates lung inflammation induced by cigarette smoke, LPS, and fMLP in mice. *Am J Respir Cell Mol Biol* 39: 7–18.
17. Woodhouse BC, Dianov GL (2008) Poly ADP-ribose polymerase-1: an international molecule of mystery. *DNA Repair (Amst)* 7: 1077–1086.
18. Robu M, Shah RG, Petitclerc N, Brind'Amour J, Kandan-Kulangara F, et al. (2013) Role of poly(ADP-ribose) polymerase-1 in the removal of UV-induced DNA lesions by nucleotide excision repair. *Proc Natl Acad Sci U S A* 110: 1658–1663.

19. Couto CA, Wang HY, Green JC, Kiely R, Siddaway R, et al. (2011) PARP regulates nonhomologous end joining through retention of Ku at double-strand breaks. *J Cell Biol* 194: 367–375.
20. Cazzalini O, Dona F, Savio M, Tillhon M, Maccario C, et al. (2010) p21CDKN1A participates in base excision repair by regulating the activity of poly(ADP-ribose) polymerase-1. *DNA Repair (Amst)* 9: 627–635.
21. Genovese T, Mazzon E, Di Paola R, Muia C, Threadgill MD, et al. (2005) Inhibitors of poly(ADP-ribose) polymerase modulate signal transduction pathways and the development of bleomycin-induced lung injury. *J Pharmacol Exp Ther* 313: 529–538.
22. Yao H, Edirisinghe I, Rajendrasozhan S, Yang SR, Caito S, et al. (2008) Cigarette smoke-mediated inflammatory and oxidative responses are strain-dependent in mice. *Am J Physiol Lung Cell Mol Physiol* 294: L1174–1186.
23. Mah LJ, El-Osta A, Karagiannis TC (2010) gammaH2AX: a sensitive molecular marker of DNA damage and repair. *Leukemia* 24: 679–686.
24. Redon C, Pilch D, Rogakou E, Sedelnikova O, Newrock K, et al. (2002) Histone H2A variants H2AX and H2AZ. *Curr Opin Genet Dev* 12: 162–169.
25. Yao H, Rahman I (2012) Role of histone deacetylase 2 in epigenetics and cellular senescence: implications in lung inflammation and COPD. *Am J Physiol Lung Cell Mol Physiol* 303: L557–566.
26. te Poelc RH, Okorokov AL, Jardine L, Cummings J, Joel SP (2002) DNA damage is able to induce senescence in tumor cells in vitro and in vivo. *Cancer Res* 62: 1876–1883.
27. Koike M, Yutoku Y, Koike A (2011) Accumulation of p21 proteins at DNA damage sites independent of p53 and core NHEJ factors following irradiation. *Biochem Biophys Res Commun* 412: 39–43.
28. Jakob B, Scholz M, Taucher-Scholz G (2002) Characterization of CDKN1A (p21) binding to sites of heavy-ion-induced damage: colocalization with proteins involved in DNA repair. *Int J Radiat Biol* 78: 75–88.
29. Perucca P, Cazzalini O, Mortusewicz O, Necchi D, Savio M, et al. (2006) Spatiotemporal dynamics of p21CDKN1A protein recruitment to DNA-damage sites and interaction with proliferating cell nuclear antigen. *J Cell Sci* 119: 1517–1527.
30. Herbig U, Wei W, Dutriaux A, Jobling WA, Sedivy JM (2003) Real-time imaging of transcriptional activation in live cells reveals rapid up-regulation of the cyclin-dependent kinase inhibitor gene CDKN1A in replicative cellular senescence. *Aging Cell* 2: 295–304.
31. Seluanov A, Mittelman D, Pereira-Smith OM, Wilson JH, Gorbunova V (2004) DNA end joining becomes less efficient and more error-prone during cellular senescence. *Proc Natl Acad Sci U S A* 101: 7624–7629.
32. Mao Z, Bozzella M, Seluanov A, Gorbunova V (2008) DNA repair by nonhomologous end joining and homologous recombination during cell cycle in human cells. *Cell Cycle* 7: 2902–2906.
33. Mao Z, Bozzella M, Seluanov A, Gorbunova V (2008) Comparison of nonhomologous end joining and homologous recombination in human cells. *DNA Repair (Amst)* 7: 1765–1771.
34. Mandal PK, Blanpain C, Rossi DJ (2011) DNA damage response in adult stem cells: pathways and consequences. *Nat Rev Mol Cell Biol* 12: 198–202.
35. Pan ZQ, Reardon JT, Li L, Flores-Rozas H, Legerski R, et al. (1995) Inhibition of nucleotide excision repair by the cyclin-dependent kinase inhibitor p21. *J Biol Chem* 270: 22008–22016.
36. Passos JF, Nelson G, Wang C, Richter T, Simillion C, et al. (2010) Feedback between p21 and reactive oxygen production is necessary for cell senescence. *Mol Syst Biol* 6: 347.
37. Bendjennat M, Boulaire J, Jascur T, Brickner H, Barbier V, et al. (2003) UV irradiation triggers ubiquitin-dependent degradation of p21(WAF1) to promote DNA repair. *Cell* 114: 599–610.
38. Cooper MP, Balajee AS, Bohr VA (1999) The C-terminal domain of p21 inhibits nucleotide excision repair in vitro and in vivo. *Mol Biol Cell* 10: 2119–2129.
39. Podust VN, Podust LM, Goubin F, Ducommun B, Hubscher U (1995) Mechanism of inhibition of proliferating cell nuclear antigen-dependent DNA synthesis by the cyclin-dependent kinase inhibitor p21. *Biochemistry* 34: 8869–8875.
40. Jurk D, Wang C, Miwa S, Maddick M, Korolchuk V, et al. (2012) Postmitotic neurons develop a p21-dependent senescence-like phenotype driven by a DNA damage response. *Aging Cell* 11: 996–1004.
41. Mauro M, Rego MA, Boisvert RA, Esashi F, Cavallo F, et al. (2012) p21 promotes error-free replication-coupled DNA double-strand break repair. *Nucleic Acids Res* 40: 8348–8360.
42. Dolan D, Nelson G, Zupanic A, Smith G, Shanley D (2013) Systems modelling of NHEJ reveals the importance of redox regulation of Ku70/80 in the dynamics of dna damage foci. *PLoS One* 8: e55190.
43. Lee JY, Kim HS, Kim JY, Sohn J (2009) Nuclear translocation of p21(WAF1/CIP1) protein prior to its cytosolic degradation by UV enhances DNA repair and survival. *Biochem Biophys Res Commun* 390: 1361–1366.
44. Stivala LA, Riva F, Cazzalini O, Savio M, Prospero E (2001) p21(waf1/cip1)-null human fibroblasts are deficient in nucleotide excision repair downstream the recruitment of PCNA to DNA repair sites. *Oncogene* 20: 563–570.
45. Romanov VS, Pospelov VA, Pospelova TV (2012) Cyclin-Dependent Kinase Inhibitor p21(Waf1): Contemporary View on Its Role in Senescence and Oncogenesis. *Biochemistry (Mosc)* 77: 575–584.
46. Mao Z, Hinc C, Tian X, Van Meter M, Au M, et al. (2011) SIRT6 promotes DNA repair under stress by activating PARP1. *Science* 332: 1443–1446.
47. Liu X, Conner H, Kobayashi T, Kim H, Wen F, et al. (2005) Cigarette smoke extract induces DNA damage but not apoptosis in human bronchial epithelial cells. *Am J Respir Cell Mol Biol* 33: 121–129.
48. Kovacs K, Erdelyi K, Hegedus C, Lakatos P, Regdon Z, et al. (2012) Poly(ADP-ribose)ylation is a survival mechanism in cigarette smoke-induced and hydrogen peroxide-mediated cell death. *Free Radic Biol Med* 53: 1680–1688.
49. Caito S, Hwang JW, Chung S, Yao H, Sundar IK, et al. (2010) PARP-1 inhibition does not restore oxidant-mediated reduction in SIRT1 activity. *Biochem Biophys Res Commun* 392: 264–270.
50. Kamp DW, Srinivasan M, Weitzman SA (2001) Cigarette smoke and asbestos activate poly-ADP-ribose polymerase in alveolar epithelial cells. *J Investig Med* 49: 68–76.
51. Hwang JW, Chung S, Sundar IK, Yao H, Arunachalam G, et al. (2010) Cigarette smoke-induced autophagy is regulated by SIRT1-PARP-1-dependent mechanism: implication in pathogenesis of COPD. *Arch Biochem Biophys* 500: 203–209.
52. Rico de Souza A, Zago M, Pollock SJ, Sime PJ, Phipps RP, et al. (2011) Genetic ablation of the aryl hydrocarbon receptor causes cigarette smoke-induced mitochondrial dysfunction and apoptosis. *J Biol Chem* 286: 43214–43228.
53. Banerjee S, Chattopadhyay R, Ghosh A, Koley H, Panda K, et al. (2008) Cellular and molecular mechanisms of cigarette smoke-induced lung damage and prevention by vitamin C. *J Inflamm (Lond)* 5: 21.
54. Oit-Wiscombe I, Virag L, Soomets U, Altraja A (2013) Increased DNA damage in progression of COPD: a response by poly(ADP-ribose) polymerase-1. *PLoS One* 8: e70333.
55. Wang M, Wu W, Rosidi B, Zhang L, Wang H, et al. (2006) PARP-1 and Ku compete for repair of DNA double strand breaks by distinct NHEJ pathways. *Nucleic Acids Res* 34: 6170–6182.
56. Claybon A, Karia B, Bruce C, Bishop AJ (2010) PARP1 suppresses homologous recombination events in mice in vivo. *Nucleic Acids Res* 38: 7538–7545.
57. Schultz N, Lopez E, Saleh-Gohari N, Helleday T (2003) Poly(ADP-ribose) polymerase (PARP-1) has a controlling role in homologous recombination. *Nucleic Acids Res* 31: 4959–4964.
58. Lapucci A, Pittelli M, Rapizzi E, Felici R, Moroni F, et al. (2011) Poly(ADP-ribose) polymerase-1 is a nuclear epigenetic regulator of mitochondrial DNA repair and transcription. *Mol Pharmacol* 79: 932–940.
59. Efimova EV, Mauceri HJ, Golden DW, Labay E, Bindokas VP, et al. (2010) Poly(ADP-ribose) polymerase inhibitor induces accelerated senescence in irradiated breast cancer cells and tumors. *Cancer Res* 70: 6277–6282.
60. Ohanna M, Giuliano S, Bonet C, Imbert V, Hofman V, et al. (2011) Senescent cells develop a PARP-1 and nuclear factor- κ B-associated secretome (PNAS). *Genes Dev* 25: 1245–1261.
61. Hageman GJ, Larik I, Pennings HJ, Haenen GR, Wouters EF, et al. (2003) Systemic poly(ADP-ribose) polymerase-1 activation, chronic inflammation, and oxidative stress in COPD patients. *Free Radic Biol Med* 35: 140–148.
62. Boulares AH, Zoltoski AJ, Sherif ZA, Jolly P, Massaro D, et al. (2003) Gene knockout or pharmacological inhibition of poly(ADP-ribose) polymerase-1 prevents lung inflammation in a murine model of asthma. *Am J Respir Cell Mol Biol* 28: 322–329.
63. Liaudet L, Pacher P, Mabley JG, Virag L, Soriano FG, et al. (2002) Activation of poly(ADP-Ribose) polymerase-1 is a central mechanism of lipopolysaccharide-induced acute lung inflammation. *Am J Respir Crit Care Med* 165: 372–377.
64. Pagano A, Metrailler-Ruchonnet I, Aurrand-Lions M, Lucattelli M, Donati Y, et al. (2007) Poly(ADP-ribose) polymerase-1 (PARP-1) controls lung cell proliferation and repair after hyperoxia-induced lung damage. *Am J Physiol Lung Cell Mol Physiol* 293: L619–629.