



# THE UNIVERSITY of EDINBURGH

## Edinburgh Research Explorer

### Runx1 is required for progression of CD41+ embryonic precursors into HSCs but not prior to this

**Citation for published version:**

Liakhovitskaia, A, Rybtsov, S, Smith, T, Batsivari, A, Rybtsova, N, Rode, C, de Brujin, M, Buchholz, F, Gordon-Keylock, S, Zhao, S & Medvinsky, A 2014, 'Runx1 is required for progression of CD41+ embryonic precursors into HSCs but not prior to this' *Development*, vol. 141, no. 17, pp. 3319-3323. DOI: 10.1242/dev.110841

**Digital Object Identifier (DOI):**

[10.1242/dev.110841](https://doi.org/10.1242/dev.110841)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Publisher's PDF, also known as Version of record

**Published In:**

Development

**Publisher Rights Statement:**

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0/>), which permits unrestricted use, distribution and reproduction in any medium provided that the original work is properly attributed.

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



## RESEARCH REPORT

## STEM CELLS AND REGENERATION

# Runx1 is required for progression of CD41<sup>+</sup> embryonic precursors into HSCs but not prior to this

Anna Liakhovitskaia<sup>1</sup>, Stanislav Rybtsov<sup>1</sup>, Tom Smith<sup>1</sup>, Antoniana Batsivari<sup>1</sup>, Natalia Rybtsova<sup>1</sup>, Christina Rode<sup>2</sup>, Marella de Bruijn<sup>2</sup>, Frank Buchholz<sup>3</sup>, Sabrina Gordon-Keylock<sup>1</sup>, Suling Zhao<sup>1</sup> and Alexander Medvinsky<sup>1,\*</sup>

**ABSTRACT**

Haematopoiesis in adult animals is maintained by haematopoietic stem cells (HSCs), which self-renew and can give rise to all blood cell lineages. The AGM region is an important intra-embryonic site of HSC development and a wealth of evidence indicates that HSCs emerge from the endothelium of the embryonic dorsal aorta and extra-embryonic large arteries. This, however, is a stepwise process that occurs through sequential upregulation of CD41 and CD45 followed by emergence of fully functional definitive HSCs. Although largely dispensable at later stages, the Runx1 transcription factor is crucially important during developmental maturation of HSCs; however, exact points of crucial involvement of Runx1 in this multi-step developmental maturation process remain unclear. Here, we have investigated requirements for Runx1 using a conditional reversible knockout strategy. We report that Runx1 deficiency does not preclude formation of VE-cad+CD45–CD41<sup>+</sup> cells, which are phenotypically equivalent to precursors of definitive HSCs (pre-HSC Type I) but blocks transition to the subsequent CD45<sup>+</sup> stage (pre-HSC Type II). These data emphasise that developmental progression of HSCs during a very short period of time is regulated by precise stage-specific molecular mechanisms.

**KEY WORDS:** AGM region, CD41, HSC, Runx1, Mouse

**INTRODUCTION**

Embryonic development of the haematopoietic stem cell lineage occurs through sequential maturation stages (Cumano and Godin, 2007; Dzierzak and Speck, 2008; Medvinsky et al., 2011). By mid-gestation, definitive HSCs (dHSCs) emerge in the aorta-gonad-mesonephros (AGM) region, as well as in placenta, large extra-embryonic vessels, yolk sac and perhaps head (de Bruijn et al., 2000; Dzierzak and Robin, 2010; Gekas et al., 2005; Gordon-Keylock et al., 2013; Li et al., 2012; Medvinsky and Dzierzak, 1996). The current prevailing view that HSCs originate in the dorsal aorta is supported by strong evidence in lower vertebrates, mouse and human (Bertrand et al., 2010, 2005; Ciau-Uitz et al., 2000; Dieterlen-Lievre, 1975; Ivanovs et al., 2011; Kissa and Herbomel, 2010; Medvinsky and Dzierzak, 1996; Medvinsky et al., 1993;

Swiers et al., 2013; Taoudi and Medvinsky, 2007). Definitive HSCs (dHSC) originate from the mesoderm that generates the VE-cadherin<sup>+</sup> endothelium, part of which becomes haematogenic and in turn generates the haematopoietic compartment marked by CD41 and subsequently by CD45 (Cumano and Godin, 2007; Dzierzak and Speck, 2008; Medvinsky et al., 2011; Mikkola et al., 2003; Rybtsov et al., 2011; Taoudi et al., 2008). Runx1 is a transcription factor playing a key role in development of the haematopoietic system; however, it is largely dispensable for the maintenance of adult bone marrow HSCs (Ichikawa et al., 2004; North et al., 1999; Okuda et al., 1996; Putz et al., 2006; Wang et al., 1996). Germline *Runx1* homozygous deletion blocks both erythro-myeloid haematopoietic progenitors (CFU-C) and HSC formation (Cai et al., 2000), which leads to severe anaemia and embryonic death by E12.5. Conditional genetic and cell fate analysis using VE-cadherin-Cre deleter mice indicates that Runx1 is crucial for the endothelial-haematopoietic transition during HSC formation (Chen et al., 2009). However, this transition involves at least three sequential stages of maturation marked by continuous expression of VE-cadherin and sequential upregulation of haematopoietic markers, first CD41 (pre-HSC Type I: VE-cad+CD41<sup>+</sup>CD45–) and subsequently CD45 (pre-HSC Type II: VE-cad+CD45<sup>+</sup>), before they become fully functional definitive HSCs (Rybtsov et al., 2011; Taoudi et al., 2008). Time specific-induced inactivation shows that Runx1 is critically important for HSC development even at E11.5 (Tober et al., 2013). This raises the question of exactly when HSC development in *Runx1* null embryos is blocked. Here, using Runx1 conditional reversible knockouts (Liakhovitskaia et al., 2009; Samokhvalov et al., 2006), we show that contrary to previously held opinion, the HSC lineage in *Runx1* knockout embryos develops up to the point when it expresses CD41, considered to be a haematopoietic commitment marker in development (Ferkowicz et al., 2003; Mikkola et al., 2003; Mitjavila-Garcia et al., 2002). Although the CD41<sup>+</sup> cell population is smaller in knockout embryos, it is clearly detectable and is less apoptotic than haematopoietic cells in wild-type embryos. Accordingly, conditional restoration of the *Runx1* locus using CD41-Cre deleter mice rescues definitive HSCs (for experimental design, see supplementary material Fig. S1A). In summary, we show that in the complete absence of Runx1, haematopoietic specification of the HSC lineage in the embryo is initiated towards the CD41<sup>+</sup> stage, but cannot progress.

**RESULTS AND DISCUSSION**


We investigated whether Runx1-deficient embryos show any haematopoietic commitment beyond primitive erythropoiesis (Okuda et al., 1996) and detected by RT-PCR an early haematopoietic marker, CD41, in E11.5 *Runx1* knockout embryos (Fig. 1A). Flow cytometry analysis confirmed the presence of a low

<sup>1</sup>MRC Centre for Regenerative Medicine, University of Edinburgh, Edinburgh EH16 4UU, UK. <sup>2</sup>MRC Molecular Haematology Unit, Weatherall Institute of Molecular Medicine, University of Oxford, Oxford OX3 9DS, UK. <sup>3</sup>Max Planck Institute of Molecular Cell Biology and Genetics, 01307 Dresden, Germany.

\*Author for correspondence (a.medvinsky@ed.ac.uk)

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0/>), which permits unrestricted use, distribution and reproduction in any medium provided that the original work is properly attributed.

Received 1 April 2014; Accepted 9 July 2014




**Fig. 1.** *Runx1* knockout embryos develop CD41<sup>+</sup> cells. (A) CD41 (left panel) and CD45 (right panel) mRNA are detected in wild-type and *Runx1* knockout embryos by RT-PCR. E9.5 *Runx1* knockout body and yolk sac (YS) (lanes 1 and 2, respectively); E9.5 wild-type body and YS (lanes 3 and 4, respectively); E11.5 *Runx1* knockout and wild-type YS (lanes 5 and 6, respectively); H<sub>2</sub>O control (lane 7). (B) Flow cytometry analysis of SSC vs CD41 and VEcad<sup>+</sup>CD45<sup>-</sup>CD41<sup>lo</sup> cells bearing the pre-HSCs Type I phenotype (7AAD+Ter119<sup>+</sup>VE-cad<sup>-</sup> cells are excluded). (Bottom row) CD45<sup>+</sup> cells are absent in *Runx1* knockout embryos but are rescued in [CD41-Cre::Runx1<sup>LacZ/Δ</sup>] embryos (7AAD+Ter119<sup>+</sup> cells are excluded). (C) CD41<sup>+</sup> and CD31<sup>+</sup> cells in the E10.5 dorsal aorta of wild-type and *Runx1*<sup>LacZ/Δ</sup> knockout embryos (confocal microscopy). Scale bars: 50 µm.

CD41-expressing (CD41<sup>lo</sup>) cell population and very few bright CD41-expressing (CD41<sup>hi</sup>) cells in *Runx1* knockout embryos compared with heterozygous littermate controls (Fig. 1B; supplementary material Fig. S2A). CD41<sup>lo</sup> cells in *Runx1* knockout embryos could be identified in the area of the dorsal

aorta using immunofluorescence (Fig. 1C). By contrast, the CD45<sup>+</sup> population is practically non-detectable (Fig. 1A,B). We found that in *Runx1* heterozygous and wild-type embryos, a large proportion of both CD41<sup>+</sup> and CD45<sup>+</sup> haematopoietic cells were apoptotic, as evidenced by annexin V staining (Fig. 2A; supplementary material Fig. S2A; data not shown) and active caspase 3 staining of many cells in intra-aortic clusters (Fig. 2B). In individual embryos, 25–55% of intra-aortic clusters contained at least one active caspase 3<sup>+</sup> cell, and some clusters were entirely apoptotic (supplementary material Fig. S2B). In layers surrounding the dorsal aorta, 31–47% of CD45<sup>+</sup> cells were apoptotic (supplementary material Fig. S2B). However, the CD41<sup>lo</sup> population in *Runx1* knockout embryos was less apoptotic than in littermate controls (Fig. 2A). A similar tendency was observed in CD41<sup>hi</sup> cells, which were produced in considerably smaller numbers in *Runx1* knockout embryos (Fig. 2A). Flow cytometry analysis has shown that phenotypic equivalents of pre-HSC Type I (VE-cad<sup>+</sup>CD45<sup>-</sup>CD41<sup>lo</sup>) can be detected in *Runx1* mutants (Fig. 1B, middle panel). We therefore investigated whether the block in HSC development occurs in the CD41 compartment and can be overcome by restoration of *Runx1* expression in CD41<sup>+</sup> cells. To this end, [CD41-Cre :: Runx1<sup>LacZ/Δ</sup>] embryos were generated in which both *Runx1* alleles are non-functional, of which one is stably deleted (*Runx1*<sup>Δ</sup>) and the other (*Runx1*<sup>LacZ</sup>) can be reactivated through Cre-mediated recombination, hereafter referred to as *Runx1*<sup>Re</sup> (see Materials and Methods; supplementary material Fig. S1) (Samokhvalov et al., 2006). In contrast to *Runx1*<sup>LacZ/Δ</sup> knockout embryos, *Runx1*<sup>Re/Δ</sup> embryos showed clear signs of rescued haematopoiesis. Both the E10.5 AGM region and the yolk sac developed CD41<sup>hi</sup> cells similar to *Runx1* heterozygous littermates (Fig. 1B and data not shown). CD45<sup>+</sup> populations were also observed in the AGM region and yolk sac of rescued embryos (Fig. 1B and data not shown). In contrast to *Runx1* mutants, *Runx1*<sup>Re/Δ</sup> embryos were no longer dying by E12.5 and survived until birth, but as expected were not found alive after that due to other non-haematopoietic defects (Liakhovitskaia et al., 2010).

To test whether development of HSCs was rescued, foetal liver cells from E14.5 *Runx1*<sup>Re/Δ</sup> embryos were transplanted into irradiated recipients. This led to successful long-term multi-lineage donor-derived engraftment, with only one exception (Fig. 3B). All donor-derived lymphoid and myeloid lineages were represented similar to control *Runx1* heterozygous transplants (Fig. 3E). Transplantations into secondary recipients also gave multi-lineage donor-derived haematopoietic engraftment (data not shown). However, when we tested whether HSCs are rescued in the AGM region, we found that, in contrast to *Runx1* heterozygous AGM regions, transplantation of E11.5 *Runx1*<sup>Re/Δ</sup> AGM regions did not produce haematopoietic repopulation (Fig. 3C). One out of five yolk sacs and one of six placentas were able to repopulate irradiated recipients (not shown). To test the possibility of delayed HSC development in rescued embryos, AGM region explants were cultured for 4 days in conditions supporting HSC development followed by transplantation into irradiated recipients (Fig. 3D). All four recipients transplanted showed high levels of donor-derived multi-lineage haematopoietic engraftment, thus demonstrating the presence of rescued pre-HSCs in the AGM region of *Runx1*<sup>Re/Δ</sup> embryos (Fig. 3D). None of the five *Runx1*<sup>LacZ/Δ</sup> AGM explants, which did not harbour the Cre transgene, were able to repopulate recipient mice.

In previous reports, inactivation of *Runx1* in the VE-cad<sup>+</sup> population suggested that *Runx1* is essential for endothelial-haematopoietic transition but not subsequently, when CFU-Cs and HSCs start expressing Vav (Chen et al., 2009). However, continuous expression of VE-cadherin over several HSC developmental stages within the



**Fig. 2. Development of hematopoietic cells in *Runx1* knockout and rescued embryos.** (A) Representative plots of annexin V staining in CD41<sup>+</sup> Ter119<sup>+</sup> cells in E10.5 AGM regions (7AAD+Ter119+CD41<sup>-</sup> cells were gated out). (Top) The CD41<sup>lo</sup> population in Runx1<sup>LacZ/Δ</sup> embryos contains a smaller proportion of annexin V<sup>+</sup> cells than in heterozygous Runx1<sup>Δ/wt</sup> embryos (red line separates CD41<sup>lo</sup> and CD41<sup>hi</sup> subsets); the same tendency but to a smaller degree is observed in the CD41<sup>hi</sup> population. (Bottom) Proportion of annexin V<sup>+</sup> cells in CD41<sup>lo</sup> and CD41<sup>hi</sup> fractions in knockout Runx1<sup>LacZ/Δ</sup> and control Runx1<sup>Δ/wt</sup> E10.5 AGM regions. Each circle represents an individual embryo. Data were obtained from five independent experiments. (B) Active caspase 3 expression in intra-aortic hematopoietic clusters in the wild-type E11.5 AGM region (confocal microscopy). Dotted lines show the endothelial lining of the dorsal aorta. D-V and A-P indicate the dorsoventral and anteroposterior axes, respectively.

VE-cad/Vav expression time window obscures the exact initial point at which Runx1 deficiency blocks this process. Our data demonstrate that, in *Runx1* knockout embryos, initial hematopoietic specification does occur. Indeed, while Runx1<sup>LacZ/Δ</sup> knockout embryos develop VE-cad<sup>+</sup>CD41<sup>low</sup>CD45<sup>-</sup> cells, low numbers of CD45<sup>+</sup> cells are generated at E9.5 but disappear by E11.5. However, CD41<sup>low</sup> cells in *Runx1* null embryos are stably present and are less apoptotic than in control *Runx1* heterozygous embryos. This explains why successful restoration of the *Runx1* functional allele in the CD41<sup>+</sup> population of mutant embryos rescued both CFU-Cs and definitive HSCs. Therefore, *in vivo* *Runx1* is required for transition of CD41<sup>+</sup> cells into the CD45<sup>+</sup> cells but not prior to that. This result contradicts previous reports indicating that Runx1 deficiency blocks transition from CD41-negative endothelial into CD41<sup>+</sup> hematopoietic cells (Bertrand et al., 2008; Lancrin et al., 2009). This discrepancy could be due to the use of an ES cell system as a model system in which hematopoietic differentiation may deviate from the *in vivo* development, or due to differences in sensitivity of methods of CD41 detection. However, an early study reported that Runx1-deficient

ES cells can generate CD41<sup>+</sup> cells lacking Kit expression (Mikkola et al., 2003). Of note, some Runx1-deficient zebrafish do recover from a larval ‘bloodless’ phase and develop to fertile adults with multilineage hematopoiesis (Sood et al., 2010), which might be explained at least partly by initiation of the hematopoietic programme in the absence of Runx1. It would be interesting to investigate whether *Runx1*-deficient cells in zebrafish, which die attempting to undergo endothelial-hematopoietic transition in the dorsal aorta, acquire the CD41<sup>+</sup> phenotype prior to that (Kissa and Herbomel, 2010). Apoptosis observed during normal early hematopoietic development is an interesting phenomenon. Significant reduction of apoptosis in hematopoietic cells of Runx1-deficient embryos concurrent with blockade of hematopoietic differentiation suggests that apoptosis is an attribute of hematopoietic differentiation and not of the most immature CD41 fraction.

In summary, we demonstrate that, in the absence of Runx1, the HSC lineage progresses to the CD41<sup>+</sup> stage but ceases further development. Therefore, transition from the CD41<sup>-</sup> endothelium into the hematopoietically committed CD41<sup>+</sup> stage is Runx1 independent. This study provides a better understanding of Runx1-dependent checkpoints during HSC development, which may be required for generating definitive HSCs from pluripotent ES/iPS cells *in vitro*.

## MATERIALS AND METHODS

### Mice

All mice used to generate embryos were bred to the C57BL/6 (CD45.2/2) background. Transgenic mice used in this study have been described previously: CD41-Cre deleter mice (Emambokus and Frampton, 2003; Rybtsov et al., 2011); activatable Runx1<sup>LacZ/wt</sup> mice (Samokhvalov et al., 2006); and conditional Runx1<sup>f/f</sup> knockout mice (Putz et al., 2006). Runx1<sup>f/wt</sup> mice were used to generate Runx1<sup>Δ/wt</sup> by Cre-mediated excision. For experimental crossings, we always used [CD41-Cre :: Runx1<sup>Δ/wt</sup>] males and Runx1<sup>LacZ/wt</sup> females (supplementary material Fig. S1). The morning of discovery of the vaginal plug was designated embryonic day 0.5. [CD41-Cre :: Runx1<sup>Δ/wt</sup>] embryos in the figures and figure legends are presented as Runx1<sup>Re/Δ</sup> for brevity. Mice were bred and used in experiments under UK Home Office regulations with approval of the University of Edinburgh Ethical Review Committee.

### Long-term repopulation assay


Cell suspensions from embryos at different stages were injected into irradiated adult recipients (CD45.1/1) either directly (suspensions from AGM region or E14.5 foetal livers) or after culture (E11.5 AGM region explants), along with 80,000 CD45.2/1 bone marrow carrier cells. Recipients were irradiated by a split dose (600+550 rad with 3 h interval) of  $\gamma$  irradiation. Donor-derived chimerism was monitored in blood at different time points after transplantation using LSRFortessa (BD). The peripheral blood was collected by bleeding the lateral tail vein into 500  $\mu$ l of 5 mM EDTA/PBS, and erythrocytes were depleted using Pharm Lyse (BD). Cells were stained with anti-CD16/32 (Fc-block), CD45.1-APC (clone A20) and anti-CD45.2-PE (clone 104) monoclonal antibodies (eBioscience). Appropriate isotype controls were used. Dead cells were excluded using 7AAD (eBioscience).

### Flow cytometry analysis

Donor-derived contribution into different hematopoietic lineages in blood or organs was determined by exclusion of recipient and carrier CD45.1<sup>+</sup> cells and staining with lineage-specific monoclonal antibodies to Mac1, CD3e, Gr1, B220 and Ter119 conjugated with PE, FITC, APC or biotin. Biotinylated antibodies were detected by incubation with streptavidin APC or PE (BD). All analyses were performed using FlowJo software (Tree Star). Statistical analyses were performed in GraphPad Prism6 software.

### AGM region explant culture

E11.5 AGM regions were dissected and cultured for 5 days on floating 0.8  $\mu$ m Millipore membranes at the liquid-gas interface with IMDM<sup>+</sup>



media consisting of 20% FCS, L-Gln, P/S IMDM and growth factors (100 ng/ml IL-3, 100 ng/ml SCF, and 100 ng/ml Flt3 ligand; all from PeproTech) as previously described (Taoudi et al., 2008). After culture, explants were dissociated enzymatically as previously described and long-term repopulation assays were performed.

#### Genotyping of embryos and assessment of Cre-mediated recombination

Genotyping was performed by Southern blotting as described previously (Liakhovitskaia et al., 2009). Specific recombination in [CD41-Cre :: Runx1<sup>ΔLacZ</sup>] embryos was always controlled by analysing the recombination in blood and separately in the tail of the embryo.

#### Confocal microscopy

Whole-mount immunostaining was performed as previously described (Yokomizo and Dzierzak, 2010). Briefly, embryos were dissected from the yolk sac, fixed with 2% paraformaldehyde (PFA), dehydrated in ascending concentration of methanol, and the head, limbs and one lateral body wall removed. Samples were then rehydrated by 50% methanol, washed with PBS and blocked in 50% FCS/0.5% Triton X-100. Embryos were incubated overnight with primary antibodies: unconjugated rabbit anti-mouse active caspase 3 (C92-605, BD Pharmingen, 1:100) and goat anti-mouse CD45 (AF114, R&D Systems, 1:100). Secondary antibodies used were anti-rabbit

NL557 (NL004, R&D Systems, 1:100) and anti-rat Alexa 488 (A-21208, Invitrogen, 1:100). Then the embryos were washed, dehydrated in methanol and cleared with BABB solution. Images were acquired using an inverted confocal microscope (Leica SP8) and processed using Volocity software.

For immunostaining on sections, embryos were fixed in 4% PFA, washed with PBS, incubated in 15% sucrose, embedded in OCT compound and snap-frozen on dry-ice/ethanol. Frozen sections (10 µm) mounted on slides were washed in 10% FCS/PBS (with penicillin/streptomycin), incubated in PBS containing 10% FCS, penicillin/streptomycin and 0.05% Tween 20, and peroxidase quenched with 3% hydrogen peroxide to be used with Tyramide Amplification kit (Molecular Probes, #T30955). After applying blocking buffer and washing in PBS, slides were incubated with anti-mouse CD31 antibody (BD #553370, 1:30) for 1 h and washed with PBS. After incubation with secondary Alexa Fluor 488 goat anti-rat antibody for 1–2 h and washing with PBS, sections were blocked with PBS containing 10% FCS, 0.05% Tween 20 and 5% normal rat serum. Endogenous avidin and biotin were blocked for 30 min each (Abcam, #ab3387). Following 1 h staining with biotinylated CD41 antibody (eBioscience, #13-0411-82, 1:100), sections were incubated with SAV-HRP and Tyramide Alexa Fluor 555 according to the Tyramide Amplification kit instructions. Sections were mounted with HardSet Vectashield containing DAPI. Images were taken on Zeiss 510 confocal microscope using the 63× oil objective and processed with ZEN2011, Adobe Photoshop and Illustrator.

## RT-PCR analysis

RNA was isolated using the RNeasy mini kit (Qiagen) and treated with DNase I (Ambion). DNA-free RNA (1 µg) was used as a template for the random primed reverse transcription reaction using the Retroscript first-strand synthesis kit for RT-PCR (Ambion). Ten percent of the RT reaction were used for PCR with CD41 (5'-GTTGGGAAGAAGGAAGATGGC-3' and 5'-ATTCCACCGCTCCAAGG-3'), CD45 (5'-GGCAACACC-TACACCCAGTGA-3' and 5'-CCATGGGTTAGATGCAG-3') and actin (5'-CCAGAGCAAGAGAGGTATC-3' and 5'-TGGAAGGTGGAC-AGTGAG-3') primers.

## Acknowledgements

We thank Evan Stamatidis and Michael Stockton for technical assistance; C. Manson and J. Verth for animal maintenance and recipient irradiation; and S. Monard and O. Rodriguez for assistance with flow cytometry. We thank Dr Darren Shaw (Roslin Institute) and Professor S. Foss (Heriot-Watt University) for initial conversations regarding experimental design.

## Competing interests

The authors declare no competing financial interests.

## Author contributions

A.L., S.R., T.S., A.B., N.R., C.R. and S.Z. carried out experiments. A.L., S.R., S.G.-K., M.deB. and A.M. analysed and assembled data. F.B. provided essential experimental materials. A.M. designed the research. A.L., S.G.-K. and A.M. wrote the paper.

## Funding

This research was supported by funding from the Medical Research Council, Leukaemia and Lymphoma Research, the Biotechnology and Biological Sciences Research Council and the Wellcome Trust. Deposited in PMC for immediate release.

## Supplementary material

Supplementary material available online at  
<http://dev.biologists.org/lookup/suppl/doi:10.1242/dev.110841/-DC1>

## References

- Bertrand, J. Y., Giroux, S., Golub, R., Klaine, M., Jalil, A., Boucontet, L., Godin, I. and Cumano, A. (2005). Characterization of purified intraembryonic hematopoietic stem cells as a tool to define their site of origin. *Proc. Natl. Acad. Sci. USA* **102**, 134–139.
- Bertrand, J. Y., Kim, A. D., Teng, S. and Traver, D. (2008). CD41(+) cmyb(+) precursors colonize the zebrafish pronephros by a novel migration route to initiate adult hematopoiesis. *Development* **135**, 1853–1862.
- Bertrand, J. Y., Chi, N. C., Santoso, B., Teng, S., Stainier, D. Y. R. and Traver, D. (2010). Haematopoietic stem cells derive directly from aortic endothelium during development. *Nature* **464**, 108–111.
- Cai, Z., de Bruijn, M., Ma, X., Dortland, B., Luteijn, T., Downing, R. J. and Dzierzak, E. (2000). Haploinsufficiency of AML1 affects the temporal and spatial generation of hematopoietic stem cells in the mouse embryo. *Immunity* **13**, 423–431.
- Chen, M. J., Yokomizo, T., Zeigler, B. M., Dzierzak, E. and Speck, N. A. (2009). Runx1 is required for the endothelial to haematopoietic cell transition but not thereafter. *Nature* **457**, 887–891.
- Ciau-Uitz, A., Walmsley, M. and Patient, R. (2000). Distinct origins of adult and embryonic blood in Xenopus. *Cell* **102**, 787–796.
- Cumano, A. and Godin, I. (2007). Ontogeny of the hematopoietic system. *Annu. Rev. Immunol.* **25**, 745–785.
- de Bruijn, M. F. T. R., Speck, N. A., Peeters, M. C. E. and Dzierzak, E. (2000). Definitive hematopoietic stem cells first develop within the major arterial regions of the mouse embryo. *EMBO J.* **19**, 2465–2474.
- Dieterlen-Lievre, F. (1975). On the origin of haemopoietic stem cells in the avian embryo: an experimental approach. *J. Embryol. Exp. Morphol.* **33**, 607–619.
- Dzierzak, E. and Robin, C. (2010). Placenta as a source of hematopoietic stem cells. *Trends Mol. Med.* **16**, 361–367.
- Dzierzak, E. and Speck, N. A. (2008). Of lineage and legacy: the development of mammalian hematopoietic stem cells. *Nat. Immunol.* **9**, 129–136.
- Emambokus, N. R. and Frampton, J. (2003). The glycoprotein IIb molecule is expressed on early murine hematopoietic progenitors and regulates their numbers in sites of hematopoiesis. *Immunity* **19**, 33–45.
- Ferkowicz, M. J., Starr, M., Xie, X., Li, W., Johnson, S. A., Shelley, W. C., Morrison, P. R. and Yoder, M. C. (2003). CD41 expression defines the onset of primitive and definitive hematopoiesis in the murine embryo. *Development* **130**, 4393–4403.
- Gekas, C., Dieterlen-Lievre, F., Orkin, S. H. and Mikkola, H. K. (2005). The placenta is a niche for hematopoietic stem cells. *Dev. Cell* **8**, 365–375.
- Gordon-Keylock, S., Sobiesiak, M., Rybtsov, S., Moore, K. and Medvinsky, A. (2013). Mouse extra-embryonic arterial vessels harbor precursors capable of maturing into definitive HSCs. *Blood* **122**, 2338–2345.
- Ichikawa, M., Asai, T., Chiba, S., Kurokawa, M. and Ogawa, S. (2004). Runx1/AML-1 ranks as a master regulator of adult hematopoiesis. *Cell Cycle* **3**, 720–722.
- Ivanovs, A., Rybtsov, S., Welch, L., Anderson, R. A., Turner, M. L. and Medvinsky, A. (2011). Highly potent human hematopoietic stem cells first emerge in the intraembryonic aorta-gonad-mesonephros region. *J. Exp. Med.* **208**, 2417–2427.
- Kissa, K. and Herberman, P. (2010). Blood stem cells emerge from aortic endothelium by a novel type of cell transition. *Nature* **464**, 112–115.
- Lancrin, C., Sroczynska, P., Stephenson, C., Allen, T., Kouskoff, V. and Lacaud, G. (2009). The haemangioblast generates haematopoietic cells through a haemogenic endothelium stage. *Nature* **457**, 892–895.
- Li, Z., Lan, Y., He, W., Chen, D., Wang, J., Zhou, F., Wang, Y., Sun, H., Chen, X., Xu, C. et al. (2012). Mouse embryonic head as a site for hematopoietic stem cell development. *Cell Stem Cell* **11**, 663–675.
- Liakhovitskaia, A., Gribi, R., Stamatidis, E., Villain, G., Jaffredo, T., Wilkie, R., Gilchrist, D., Yang, J., Ure, J. and Medvinsky, A. (2009). Restoration of Runx1 expression in the Tie2 cell compartment rescues definitive hematopoietic stem cells and extends life of Runx1 knockout animals until birth. *Stem Cells* **27**, 1616–1624.
- Liakhovitskaia, A., Lana-Eiola, E., Stamatidis, E., Rice, D. P., van't Hof, R. J. and Medvinsky, A. (2010). The essential requirement for Runx1 in the development of the sternum. *Dev. Biol.* **340**, 539–546.
- Medvinsky, A. and Dzierzak, E. (1996). Definitive hematopoiesis is autonomously initiated by the AGM region. *Cell* **86**, 897–906.
- Medvinsky, A. L., Samoylina, N. L., Müller, A. M. and Dzierzak, E. A. (1993). An early pre-liver intraembryonic source of CFU-S in the developing mouse. *Nature* **364**, 64–67.
- Medvinsky, A., Rybtsov, S. and Taoudi, S. (2011). Embryonic origin of the adult hematopoietic system: advances and questions. *Development* **138**, 1017–1031.
- Mikkola, H. K. A., Fujiwara, Y., Schlaeger, T. M., Traver, D. and Orkin, S. H. (2003). Expression of CD41 marks the initiation of definitive hematopoiesis in the mouse embryo. *Blood* **101**, 508–516.
- Mitjavila-Garcia, M. T., Cailleret, M., Godin, I., Nogueira, M. M., Cohen-Solal, K., Schiavon, V., Lecluse, Y., Le Pesteur, F., Lagrue, A. H. and Vainchenker, W. (2002). Expression of CD41 on hematopoietic progenitors derived from embryonic hematopoietic cells. *Development* **129**, 2003–2013.
- North, T., Gu, T. L., Stacy, T., Wang, Q., Howard, L., Binder, M., Marin-Padilla, M. and Speck, N. A. (1999). Cbfa2 is required for the formation of intra-aortic hematopoietic clusters. *Development* **126**, 2563–2575.
- Okuda, T., van Deursen, J., Hieber, S. W., Grosfeld, G. and Downing, J. R. (1996). AML1, the target of multiple chromosomal translocations in human leukemia, is essential for normal fetal liver hematopoiesis. *Cell* **84**, 321–330.
- Putz, G., Rosner, A., Nuesslein, I., Schmitz, N. and Buchholz, F. (2006). AML1 deletion in adult mice causes splenomegaly and lymphomas. *Oncogene* **25**, 929–939.
- Rybtsov, S., Sobiesiak, M., Taoudi, S., Souilhol, C., Senserrick, J., Liakhovitskaia, A., Ivanovs, A., Frampton, J., Zhao, S. and Medvinsky, A. (2011). Hierarchical organization and early hematopoietic specification of the developing HSC lineage in the AGM region. *J. Exp. Med.* **208**, 1305–1315.
- Samokhalov, I. M., Thomson, A. M., Lalancette, C., Liakhovitskaia, A., Ure, J. and Medvinsky, A. (2006). Multifunctional reversible knockout/reporter system enabling fully functional reconstitution of the AML1/Runx1 locus and rescue of hematopoiesis. *Genesis* **44**, 115–121.
- Sood, R., English, M. A., Belele, C. L., Jin, H., Bishop, K., Haskins, R., McKinney, M. C., Chahal, J., Weinstein, B. M., Wen, Z. et al. (2010). Development of multilineage adult hematopoiesis in the zebrafish with a runx1 truncation mutation. *Blood* **115**, 2806–2809.
- Swiers, G., Baumann, C., O'Rourke, J., Giannoulatou, E., Taylor, S., Joshi, A., Moignard, V., Pina, C., Bee, T., Kokkaliaris, K. D. et al. (2013). Early dynamic fate changes in haemogenic endothelium characterized at the single-cell level. *Nat. Commun.* **4**, 2924.
- Taoudi, S. and Medvinsky, A. (2007). Functional identification of the hematopoietic stem cell niche in the ventral domain of the embryonic dorsal aorta. *Proc. Natl. Acad. Sci. USA* **104**, 9399–9403.
- Taoudi, S., Gonneau, C., Moore, K., Sheridan, J. M., Blackburn, C. C., Taylor, E. and Medvinsky, A. (2008). Extensive hematopoietic stem cell generation in the AGM region via maturation of VE-cadherin+CD45+ pre-definitive HSCs. *Cell Stem Cell* **3**, 99–108.
- Tober, J., Yzaguirre, A. D., Piwarzky, E. and Speck, N. A. (2013). Distinct temporal requirements for Runx1 in hematopoietic progenitors and stem cells. *Development* **140**, 3765–3776.
- Wang, Q., Stacy, T., Binder, M., Marin-Padilla, M., Sharpe, A. H. and Speck, N. A. (1996). Disruption of the Cbfa2 gene causes necrosis and hemorrhaging in the central nervous system and blocks definitive hematopoiesis. *Proc. Natl. Acad. Sci. USA* **93**, 3444–3449.
- Yokomizo, T. and Dzierzak, E. (2010). Three-dimensional cartography of hematopoietic clusters in the vasculature of whole mouse embryos. *Development* **137**, 3651–3661.