



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Macrophage proliferation, provenance, and plasticity in macroparasite infection

Citation for published version:

Rückerl, D & Allen, JE 2014, 'Macrophage proliferation, provenance, and plasticity in macroparasite infection' *Immunological reviews*, vol 262, no. 1, pp. 113-33., 10.1111/imr.12221

Digital Object Identifier (DOI):

[10.1111/imr.12221](https://doi.org/10.1111/imr.12221)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher final version (usually the publisher pdf)

Published In:

Immunological reviews

Publisher Rights Statement:

© 2014 The Authors. *Immunological Reviews* published by John Wiley & Sons Ltd. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Dominik Rückerl
Judith E. Allen

Macrophage proliferation, provenance, and plasticity in macroparasite infection

Authors' address

Dominik Rückerl¹, Judith E. Allen¹

¹Institute for Immunology and Infection Research, School of Biological Sciences, University of Edinburgh, Edinburgh, UK.

Correspondence to:

Judith E. Allen

Institute for Immunology and Infection Research
School of Biological Sciences
University of Edinburgh
Edinburgh EH9 3JT, UK
Tel.: +44 (0)131 650 7014
Fax: +44 (0)131 650 6564
e-mail: j.allen@ed.ac.uk

Acknowledgements

This work was funded by the Medical Research Council UK (G0600818 and MR/K01207X/1 to J. E. Allen), with additional support from the Wellcome Trust (strategic award for Centre for Immunity, Infection, and Evolution). The authors declare no financial conflicts of interests.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

This article is part of a series of reviews covering Monocytes and Macrophages appearing in Volume 262 of *Immunological Reviews*.

Immunological Reviews 2014

Vol. 262: 113–133

Printed in Singapore. All rights reserved

© 2014 The Authors. *Immunological Reviews* published by John Wiley & Sons Ltd.
Immunological Reviews
0105-2896

© 2014 The Authors. *Immunological Reviews* published by John Wiley & Sons Ltd.
Immunological Reviews 262/2014

Summary: Macrophages have long been center stage in the host response to microbial infection, but only in the past 10–15 years has there been a growing appreciation for their role in helminth infection and the associated type 2 response. Through the actions of the IL-4 receptor α (IL-4R α), type 2 cytokines result in the accumulation of macrophages with a distinctive activation phenotype. Although our knowledge of IL-4R α -induced genes is growing rapidly, the specific functions of these macrophages have yet to be established in most disease settings. Understanding the interplay between IL-4R α -activated macrophages and the other cellular players is confounded by the enormous transcriptional heterogeneity within the macrophage population and by their highly plastic nature. Another level of complexity is added by the new knowledge that tissue macrophages can be derived either from a resident prenatal population or from blood monocyte recruitment and that IL-4 can increase macrophage numbers through proliferative expansion. Here, we review current knowledge on the contribution of macrophages to helminth killing and wound repair, with specific attention paid to distinct cellular origins and plasticity potential.

Keywords: IL-4 receptor, monocyte, macrophage, wound repair, inflammation, helminth infection

Introduction

Infection with multicellular parasites represents one of the most significant challenges to both veterinary and public health. Medically relevant multicellular parasites are called 'helminths', a term that does not accurately reflect the vast phylogenetic diversity of this group, which includes the platyhelminths (flukes and tapeworms) and nematodes (roundworms). These animal phyla diverged over 1 billion years ago, predating the split between vertebrates and invertebrates (1). Despite the phylogenetic distance between them, helminths as well as many insects induce a characteristic immune response that at its core, features the production of interleukin-4 (IL-4) and IL-13 by T-helper 2 (Th2) cells and a range of innate cells. Thus, we are apparently hardwired to mount a type 2 response on exposure to multicellular parasites (2).

In humans, helminths predominantly cause disease in developing countries but are a major veterinary problem throughout the world and often establish long-lasting infections that can last years and even decades. As a consequence, these infections are rarely lethal but can have devastating effects on life quality (3, 4). Often, however, infections are asymptomatic or associated with less obvious repercussions (e.g. growth retardation, blunted mental development, etc.) caused by nutritional restrictions imposed on the host (5). To maintain fitness, the infected individual must cope with large motile foreign organisms that cannot easily be digested or removed, and for which an aggressive host response would be self-damaging. Further, these parasites often directly damage host tissue during migration or feeding. The host must thus strike a balance between immune suppression to minimize self-harm, the repair of damaged tissue and a sufficiently robust anti-worm response to keep parasite numbers below a level that compromises fitness. Type 2 immunity appears to be the response mammals have evolved to cope with this challenge. Th2 cells that produce IL-4, IL-5, IL-9, IL-10, and IL-13 and B cells producing immunoglobulin E (IgE) are the central players in the adaptive type 2 response and the innate cells associated with type 2 cytokines include eosinophils, mast cells, basophils, macrophages, and type 2 innate lymphoid cells (ILC2). Together, these pathways promote parasite killing and expulsion but are also key components of tissue repair pathways that are fundamentally non-inflammatory (2, 6, 7).

In this review, we focus on the role of macrophages in keeping the host fit in the context of a type 2 response induced by helminth infection. Macrophages have long been center stage as the target of Th1-type responses with a clear role in containment of microbial pathogens, particularly intracellular bacteria and protozoa. The function of macrophages activated by Th2 cytokines has been far more difficult to establish, despite their abundance in almost every helminth infection as well as other type 2 conditions such as allergy and asthma. While IL-4R α ligation by either IL-4 or IL-13 has a dramatic effect on the macrophage transcriptional landscape (8, 9), the specific functions of macrophages activated via the IL-4R α have yet to be fully defined. Here, we use the term alternatively activated macrophage (AAM Φ) to define macrophage phenotypes that rely on the IL-4R α (10). However, even when considering activation via a single receptor there is a broad heterogeneity of context-dependent macrophage phenotypes observable. This makes it difficult to identify unifying functional properties of AAM Φ . With an emphasis on our own research, we

highlight here the knowledge that has been gained from studying macrophage activation in helminth models, which has provided broad insight into macrophage biology.

Macrophage proliferation

We have been using a murine model of filariasis, *Litomosoides sigmodontis*, to understand the mechanisms of resistance and susceptibility to filarial nematode infection. Infective larvae (L3) invade via the skin, migrate through the lymphatics, and enter the pleural cavity, where they mature to adulthood. In susceptible BALB/c mice, the adult worms produce blood-circulating offspring (microfilariae) by day 50 postinfection. In resistant C57BL/6 mice the parasites are killed in the pleural cavity prior to reaching sexual maturity. This has proved a powerful model for the study of filariasis (11) and highlighted a critical role for T-regulatory cells in the susceptible phenotype of BALB/c mice (12). To investigate the contribution of macrophages to parasite killing in the resistant C57BL/6 strain, we depleted blood monocytes to prevent their recruitment to the infection site. To our surprise, we saw no reduction in the high macrophage numbers accumulating in the pleural cavity of infected mice. Time course assessment over the first 10 days of infection demonstrated that although macrophages were steadily increasing in number, their flow cytometry profile remained essentially the same as in naive mice, with no evidence of monocyte or neutrophil infiltration (13). This suggested that the resident macrophage population was expanding by other means than recruitment from the blood. Indeed, using both Ki67 staining and a 3 h BrdU pulse, extensive proliferation of the resident F4/80^{hi} population was observed (13).

Macrophage expansion and survival had previously been associated with CSF1R signaling and differentiation of monocyte derived macrophages (14–17). However, because helminth-driven macrophage proliferation was independent from blood monocyte recruitment and the numbers of cells accumulating at the site of infection were significantly reduced in IL-4^{-/-} mice (13), we tested the possibility that IL-4 was directly responsible. An injection of IL-4 complex (IL-4C) [a mixture of IL-4 with anti-IL-4 for increased bioactivity (18)] into the peritoneal cavity led to dramatic macrophage proliferation (13). Remarkably, proliferative expansion was not restricted to the serous cavities but was observed in the liver (13), spleen, and lungs (authors' unpublished observation). This was consistent with previously published data showing that IL-4 delivery, either in complex form or by mini-osmotic pump, led to

macrophage hyperplasia in the liver, spleen and bone marrow (18). In both the Milner et al. study and ours (13, 18–20), macrophage expansion following IL-4C delivery was independent of the adaptive immune system, occurring in RAG^{-/-} mice but not in Stat-6^{-/-} or IL-4R α ^{-/-} animals.

IL-4C injection allowed us to take a reductionist approach to address the factors that contribute to helminth-driven macrophage proliferation. Using macrophage-intrinsic competitive bone marrow (BM) chimeras in which Ly5.1⁺ WT mice are reconstituted with a 1:1 mix of Ly5.1⁺ WT and IL-4R α ^{-/-} Ly5.2⁺ BM, we established that IL-4 acts directly on the macrophages and IL-4R α expression is essential to initiate a program of proliferation (19). However, important differences between the IL-4C model and *L. sigmodontis* infection were observed using these 50:50 WT:IL-4R α ^{-/-} chimeras. No BrdU incorporation into IL-4R α ^{neg} cells was observed following IL-4C injection, but during the early stages of *L. sigmodontis* infection, IL-4R α ^{neg} cells did incorporate BrdU, albeit at significantly lower level than the WT cells. As discussed below, we subsequently identified CSF1-dependent mechanisms to be responsible for the residual proliferative expansion of these cells during infection. Critically, IL-4R α positive macrophages have a competitive advantage, proliferating to a greater extent and eventually outnumbering their IL-4R α negative counterparts (19). Further, both macrophage proliferation and numbers are reduced in IL-4^{-/-} and IL-4R α ^{-/-} mice in all infection models tested, emphasizing the central role of IL-4 in driving macrophage accumulation during infection. Because IL-13 can also signal through the IL-4R α we addressed whether IL-13 was also capable of inducing macrophage proliferation. We generated an IL-13 complex and demonstrated that it was equivalently able to drive macrophage proliferation when delivered intraperitoneally (19). However, a role for IL-13-mediated proliferation in a physiological setting has yet to be established.

The source of IL-4 (or IL-13) that drives macrophage proliferation during helminth infection is likely to vary with tissue and stage of infection (6) and may include mast cells, eosinophils, or innate lymphoid cells among others. However, our data suggest that during infection, CD4⁺ lymphocytes are required to generate significant macrophage numbers. In the absence of either RAG genes or class II, there is no expansion or alternative activation of macrophages in a peritoneal implant model of filarial infection (21). In the *L. sigmodontis* infection model, proliferation does not occur until the onset of the adaptive immune response and does not occur in infected RAG-deficient mice (13, 19).

Along with the need for high concentrations of IL-4 (19), these data strongly suggest a requirement for cognate T-helper interactions to initiate macrophage proliferation in the infection context. Although macrophage accumulation in both IL-4C and infection models is the direct result of IL-4 driven proliferation, anti-apoptotic properties of IL-4 may also make a significant contribution to the final numbers.

Macrophage provenance

Our discovery that IL-4 could drive expansion of resident macrophages occurred as a new paradigm in macrophage biology was unfolding. The established dogma had been that tissue-resident macrophages were derived from circulating bone marrow-derived monocytes. With new fate mapping technologies it became apparent that in most tissues including the serous cavities (22, 23) the resident macrophages are established prenatally and are retained throughout the life of the animal through proliferative self-renewal (reviewed in 24, 25). The notable exceptions to this rule are macrophages in the skin and GI tract (26, 27).

To verify the source of proliferating macrophages in our *L. sigmodontis* infection model, we used tissue-protected BM chimeras (13) in which resident macrophages in the body cavities are protected from radiation damage and thus not replaced by BM-derived cells. Using the recipient/donor ratio found in the blood versus the body cavity of naive animals, it is possible to determine whether the macrophages in infected tissues are of BM origin or not. Using this method, we were able to show that the expanding macrophage population following both IL-4C injection and *L. sigmodontis* infection were derived from tissue-resident cells (13). The tissue-protected BM chimeras thus allowed us to formally demonstrate that the expanding cells were not derived from circulating blood monocytes (13). However, AAM Φ in the GI tract of nematode infected mice are largely monocyte-derived (28), and we have observed proliferation of macrophages in the lamina propria of mice infected with the GI nematode, *Heligmosomoides polygyrus* (Fig. 1). We thus investigated whether IL-4-driven proliferative expansion was restricted to tissue-resident macrophages or if IL-4 could act similarly on monocyte-derived cells. Using the tissue-protected chimeras, we demonstrated that injection of thioglycollate together with IL-4C stimulates the proliferation of recruited bone marrow-derived monocytes, while IL-4C alone stimulates resident cell expansion. Thus the ability of IL-4 to stimulate proliferation is not restricted to resident cells (13, 20).




Fig. 1. Proliferative expansion of macrophages (MΦ) is induced in various helminth infection models. Representative dotplots of one naive (N) and one infected (I) animal shown. Graphs indicate percent of F4/80⁺ macrophages that are Ki67⁺. Data points depict individual animals. Detailed description of materials and methods used in similar experiments can be found in Jenkins et al. (19). Data from *N. brasiliensis* infected alveolar macrophages isolated from whole tissue digests (CD11b^{low} SiglecF⁺ F4/80⁺ CD11c^{high}) kindly provided by Dr. Tara E. Sutherland.

Macrophage self-renewal in most tissues relies on the CSF-1 receptor (CSF1R) (24), and CSF-1 has been shown to be required for both the steady state maintenance of resident peritoneal macrophages as well as recovery of the resident population by proliferative expansion following an inflammatory response (29). In our experiments with 50:50 WT:IL-4Rα^{-/-} bone marrow chimeras, steady-state proliferation in PBS-treated animals was not influenced by IL-4Rα signaling (19), confirming that IL-4 is not needed for local self-renewal of peritoneal macrophages (30). We hypothesized that instead, IL-4 functioned by enhancing steady state proliferation, perhaps by CSF1R upregulation or increased ligand production. To test this, the CSF1R was blocked during IL-4C delivery. IL-4-induced proliferation was undiminished in the presence of the CSF1R antibody demonstrating complete independence from the CSF1R (19). However, CSF1R blockade during *L. sigmodontis* infection suggested that CSF-1 does contribute to macrophage proliferation early in infection but is superseded by IL-4 presumably upon entry of Th2 cells. Supporting this, CSF1R blockade at day 8 post-infection inhibited macrophage proliferation only in the animals that did not yet exhibit evidence of high IL-4 exposure (19). Thus, IL-4 rather than acting via the CSF1R appears to unleash macrophages from their dependence on the CSF1R for proliferation.

IL-4 has the capacity to induce alternative activation as well as proliferation in both resident and recruited settings (13). This raised the fundamental question of whether IL-4Rα-activated macrophages of distinct origin are

functionally similar. To address this P'ng Loke and colleagues (20) compared the transcriptome of resident versus recruited macrophages exposed to IL-4 using delivery of IL-4 into naive or thioglycollate-injected mice. The results dramatically illustrated that cellular origins impart a far larger stamp on the transcriptome than IL-4 exposure. They further demonstrated that many of the markers traditionally associated with alternative activation were only present on the blood monocyte derived population, including the mannose receptor, Raldh2 and PD-L2. The difference in Raldh2 expression translated into the ability of only monocyte derived cells to promote the differentiation of FoxP3 cells from naive CD4⁺ cells via retinoic acid production. In contrast, some IL-4-induced gene products, such as Uncoupling protein 1, were seen only in the resident derived cells. These phenotypic differences between cells of embryonic origin versus bone marrow origin were verified in *L. sigmodontis* versus *Schistosoma mansoni* infections, respectively (20). Thus, macrophage provenance is of critical importance in understanding the outcome of a type 2 immune response.

In addition to the expansion of resident pleural macrophages in the *L. sigmodontis* model, we also observed that infection with the purely gastrointestinal nematode, *H. polygyrus*, caused expansion of peritoneal resident cells. This was somewhat surprising, as the parasite does not enter the peritoneal space. However, previous studies have shown the presence of Th2 cells in the peritoneal cavity of infected mice (31), consistent with our prediction that T-helper cells are instrumental in inducing proliferation. In both the *L. sigmodontis* and

H. polygyrus models, shielded bone marrow chimeras were used to verify the resident origin of the pleural and peritoneal macrophages, respectively. However, these infections may prove to be exceptions to the rule and in many, if not most, Th2 infections, there is likely to be recruitment of monocytes from the blood. Because the importance of macrophage origins has only recently become recognized, little information is available in different model systems, and thus we are relatively ignorant of what determines when a recruited population is dominant over resident expansion. Even in the *L. sigmodontis* model, over time there is increasing numbers of recruited monocytes (unpublished observation). We presume that ‘non-inflammatory’ entry of the infective larvae combined with local induction of IL-4 by innate cells means blood cell recruitment is initially avoided. It may be that only subsequent parasite death is sufficient to trigger monocyte recruitment. In settings in which bacteria are present, such as infection of the gastro-intestinal tract, monocyte recruitment will likely dominate (28) (Fig. 2).

Two important papers have recently addressed macrophage origins in the liver of *S. mansoni* infected mice (32, 33). During patent *S. mansoni* infection, eggs become trapped in the liver and these induce a host-protective Th2-mediated granuloma, in which AAM Φ are a major component. Using intra-vital imaging and cell transfer experiments with Cx3Cr1^{GPE/+} mice, Girgis et al. (33) demonstrated that

AAM Φ in the liver granulomas arise predominantly from blood monocytes. Similarly, Nascimento et al. (32) using careful phenotyping of the developing granuloma and pulse chase experiments demonstrate that the majority of macrophages are bone marrow derived. Importantly, Nascimento et al. (32) show that there is robust infection-induced proliferation of the resident macrophages, but this does not translate into significant cellular expansion. The dominance of the recruited monocyte population may reflect an inflammatory environment caused by leakage of intestinal bacteria into the portal bloodstream that may occur as eggs successfully cross into the gut lumen (32). Indeed, the authors show that the macrophage phenotype is highly heterogeneous, with clear evidence of alternative activation but also the production of iNOS and TNF. We have observed macrophage proliferation in the serous cavities, lamina propria, and the lung in different Th2-inducing models (Fig. 1). Nascimento et al. (32) extend these observations to the liver, but illustrate that the dynamics by which macrophages accumulate are highly complex. Importantly, even when proliferation is not observed, this may be due to the narrow time window in which proliferation can be seen (19). Thus, even when it is straightforward to assess the tissue-resident versus BM origin of macrophages in inflamed tissue, determining whether the final numbers are the result of proliferation, recruitment, or enhanced survival will still be challenging.




Fig. 2. In vivo recruitment/expansion of macrophages (M Φ) is dependent on the infectious agent. Macrophage numbers can expand in the tissues either due to recruitment, proliferation, or a combination of both. This figure illustrates different infection models or experimental manipulations that can lead to these distinct outcomes.

With the new understanding of macrophage origins and proliferation capacity, the functional potential of macrophages activated in type 2 immunity gains even more complexity. However, at a more fundamental level we are still remarkably ignorant of the specific contribution of IL-4R α activated macrophages to disease outcome in most type 2 disease settings. In the following sections, we discuss what is currently known about the specific functions of AAM Φ but also new leads from transcriptional analysis comparing WT versus IL-4R α ^{-/-} macrophages (9).

Helminth killing

Immune-mediated killing of parasitic worms has been repeatedly demonstrated in animal models, and, with rare exceptions, type 2 immunity is required. In the case of several gastrointestinal models, delivery of IL-4 and/or IL-13 alone triggers mucus production, smooth muscle contraction and epithelial turnover, which is sufficient to mediate expulsion from the gut (34). More broadly IL-4 or IL-4 receptor deficiency confers susceptibility on resistant murine strains for a range of helminth infections (6). Helminth infected humans are reported to display a 'modified Th2' response that results from sophisticated adaptations employed by the parasites to divert the host immune reaction as well as host adaptations to avoid immune-mediated damage (35–37). Together these allow chronic infections to establish and can make deciphering the appropriate protective response difficult. This is further complicated by the dynamics of helminth life cycles and life histories such that the killing mechanisms depend on multiple factors that include the developmental stage of the parasite as well as the tissue through which the parasite migrates and eventually resides.

This complexity is exemplified by the model filarial nematode, *L. sigmodontis*. In primary infection, over half of the infective-stage L3 larvae are killed in the skin immediately after infection or during their migration to the pleural cavity (38). Once the remaining larvae make it to the pleural cavity and start to undergo the molting process into L4 stage larvae and adult worms, further killing is relatively slow. Furthermore, the mechanisms by which the larvae and adults are killed by the immune system differ (39). In susceptible but immunocompetent mouse strains (e.g. BALB/c) this process can take several months (>90 days) and even in resistant strains (e.g. C57BL/6) the parasites survive for several weeks (approximately 40–50 days) (38). Similarly, on secondary infection with the gastrointestinal parasite *H. polygyrus* larvae are killed, but adult parasites implanted directly

into the gut lumen of the immune recipient can establish patent infection (40). In general, early developmental stages are more vulnerable to immune attack than later stages, perhaps because of the superior ability of parasites that survive to adulthood to modulate the host response (41). Some parasites create their own 'immunoprivileged' sites by forming cysts (e.g. *Echinococcus granulosus*) which show a high degree of immunomodulatory activity diverting potentially protective immune responses (42). Thus, one aim of the immune response is to trap parasites during their migration/development before a successful immune evasion strategy is in place. This appears to be the basis of most successful model vaccines (43). For example, *L. sigmodontis* larvae exhibit limited migratory capacity in vaccinated mice (44), and *H. polygyrus* larvae become trapped inside the gut wall in resistant or immune mice (45, 46). A range of type 2 cytokines, cell types, and antibodies have been implicated in protective immunity in many different helminth models, but only recently have macrophages been receiving due attention (47).

The specific contribution of macrophages and, moreover, IL-4R α -dependent activation in the killing and expulsion of helminths is still unclear. Macrophages make up a large proportion of the cells present at the site of various helminth infections (48) and are a major constituent of granulomas forming around parasites (49–51). Macrophage depletion using clodronate liposomes has demonstrated their importance for expulsion of intestinal parasites like *Nippostrongylus brasiliensis* or *H. polygyrus* (28, 52). Similarly interference with macrophage function through injection of carrageenan or carbon particles enhances the survival of *Brugia malayi* and *B. pahangi* larvae in the peritoneal cavity of infected animals (53, 54). Of particular interest is recent data suggesting that while macrophages are important in larval killing, they don't act alone. Bonne-Année et al. (55, 56) demonstrated that *in vitro* macrophages and neutrophils work together to kill *Strongyloides stercoralis* larvae.

The evidence that macrophages are involved in worm killing is strong, but whether this requires alternative activation or is possibly hampered by it is still contentious. In a very recent study, expression of alternative activation markers positively correlated with resistance to infection with *H. polygyrus* and depletion of macrophages using clodronate liposomes resulted in reduced resistance (51). However, the study did not directly assess the role of the IL-4R α , and it is possible that although macrophages themselves might be important for efficient killing of invading parasites the

IL-4R α -mediated activation might not. Most of the studies attempting to resolve the specific role of AAM Φ have used mice with LysM-Cre mediated deletion of the floxed IL-4R α gene, which is relatively inefficient particularly on peripheral blood monocytes (57). Additionally, even if IL-4R α -induced macrophage genes contribute to worm killing, these may be countered by proteins that inhibit Th2-induced effector pathways. For example, Arginase and RELM α , products strongly associated with alternative macrophage activation, can act as negative feedback regulators directly limiting the Th2 response (58, 59). Thus, it may not be surprising that efficient elimination of the gastrointestinal nematodes *N. brasiliensis* or *Trichinella spiralis* was unimpaired in LysMCre IL-4R α^{floxed} mice relative to WT (60–62). Critically, a role (or lack thereof) for AAM Φ in one infection system may not apply to all nematode infections as the Th2 effector mechanisms involved can be quite distinct (34, 63).

Data from our laboratory using the filarial *L. sigmodontis* model suggest that there is a role for IL-4R α -mediated macrophage activation in the control of this tissue nematode infection but also revealed a key difficulty with the LysMCre

mice in the context of IL-4R α expression. Infected LysMCre IL-4R α^{floxed} mice showed increased number of circulating microfilaria at late stages of infection (Fig. 3) indicating a role for AAM Φ in either killing the circulating microfilaria or impairing female fertility. To our surprise, standard markers of alternative activation (i.e. RELM α and YM1) in the macrophage specific KO while absent as expected at day 10 were no different to wildtype animals at the late stage of infection (d60). Because LysMCre-mediated deletion of the floxed IL-4R α gene is relatively inefficient (57) we predicted that chronic IL-4 exposure was causing an outgrowth of IL-4R α sufficient macrophages, such that by day 60 post infection all macrophages were wildtype. This was consistent with our data utilizing 50:50 WT:IL-4R $\alpha^{-/-}$ chimeras injected with IL-4, in which the WT macrophages outcompete their gene deficient counterparts (19). The competitive advantage during nematode infection was confirmed with *H. polygyrus*, in which the proportion of RELM α positive peritoneal macrophages in LysMCre IL-4R α^{floxed} mice increased from 4% in naive mice to 25% at day 14 and 70% by day 28, by which point >50% of the macrophages expressed the IL-4R α (19).




Fig. 3. Expression of IL-4R α by myeloid cells is required for optimal immune responses to filarial infection. BALB/c (blue squares; WT), IL-4R α -deficient (black circles; KO), IL-4R α heterozygous (green upward triangles; het), or LysMCre IL-4R α^{floxed} (red downward triangles; Cre) were infected by subcutaneous injection of 25 *L. sigmodontis* L3 stage larvae. (A) At the indicated time points, 10 μ l venous blood were collected from each mouse and the number of microfilaria present assessed by microscopic analysis. Data points indicate mean and SEM of 4–5 mice per group. (B) Number of adult parasites present in the pleural cavity of the mice analyzed in (A), 60 days after infection. Data points indicate individual animals and lines indicate mean. (C) Percent Relm- α expressing macrophages 10 days postinfection in naive (N) or infected (I) BALB/c (WT) or LysMCre IL-4R α^{floxed} (Cre) mice. (D) Percent Relm- α expressing macrophages in the mice analyzed in A, 60 days postinfection. Detailed description of materials and methods used in similar experiments can be found in van der Werf et al. (36) (A, B) and in Jenkins et al. (13) (C, D).

Because actual IL-4R α deletion occurred only early in infection, the data suggest AAM Φ act at early stages of infection to influence later resistance. That early events can determine the later outcome of infection is consistent with data in which T-regulatory cell depletion just prior to *L. sigmodontis* infection dramatically decreases microfilaria numbers at day 60 (64). This hypothesis would also be in line with data showing that host immune status at the time of infection influences parasite developmental rate and subsequent fecundity (65). The failure of macrophages to efficiently alternatively activate early in infection could therefore lead to an altered parasite developmental program which allows the parasite to produce more offspring. One caveat is that LysMCre will also mediate IL-4R α deletion in granulocytes, and thus we cannot be certain that the impaired microfilarial killing is specifically due to macrophages.

A recent report by Esser-von Bieren *et al.* (46) sheds new light on the potential contribution of macrophages to nematode killing. Using mice that lack antibodies (JH^{-/-}) or activating Fc receptors (FcR γ ^{-/-}), they demonstrate in a model of secondary *H. polygyrus* infection that antibodies activate macrophages to trap and immobilize infective larvae. Arginase was important for killing as discussed below. In this setting, the ability of antibodies to induce arginase was independent of the IL-4R α , but in its absence macrophages fail to accumulate at the infection site (46). Thus, even if the IL-4R α is not always needed to induce macrophage effector molecules, it may be essential to expand macrophages in sufficient numbers to perform non-IL-4R α -dependent functions.

It is therefore still early in our understanding of the mechanisms by which AAM Φ may contribute to parasite control. In the sections below, we discuss some of the findings about well-known markers of alternative activation and their role in resistance to helminth infection as well as more recent data from our laboratory identifying potential new effector molecules not previously associated with AAM Φ . Due to the fact that parasites can survive for very long periods of time in their respective hosts, it is reasonable to expect that a range and combination of different effector mechanisms need to work in concert to achieve effective parasite expulsion or killing. Of note, several AAM Φ products have been associated with an immune modulatory function rather than direct anti-parasitic effector responses. Thus, in many circumstances AAM Φ may not have a primary role in parasite killing but act to contain excessive inflammation and maintain tissue homeostasis.

Effector molecules and pathways

Chitinase-like proteins

YM1 [chitinase-3-like-3 (Chi3l3)] is one of the most highly expressed genes in alternatively activated macrophages (8, 9, 66). Upon activation with IL-4, macrophages accumulate large amounts of intracellular YM1 that can be readily detected by flow cytometry even without the use of Brefeldin A or Monensin and is simultaneously released in high quantities into the surrounding tissue (13, 67). YM1 is part of the glycoside hydrolase family 18, which includes enzymatically active, chitin-degrading enzymes (i.e. chitotriosidase and AMCase) as well as enzymatically inactive, chitinase-like proteins (CLPs), e.g. YM1, BRP39, YM2 (reviewed in 68). Whereas the chitin-degrading enzymes are highly conserved across the animal kingdom, the CLPs represent relatively recent gene duplication events in mammals and are highly diverse between species (69). Because of the absence of direct homology between mouse YM1 and human CLPs, the importance of studying murine CLPs is sometimes questioned. However, CLP family members are present in almost all mammalian species and are undergoing apparently rapid evolution. This high diversity and upregulation by IL-4 strongly suggests CLPs are involved in protection against infection, and the fundamental mechanism of action is likely to be shared across species.

AMCase and chitotriosidase, which actively degrade chitin, are thought to be important in the defense against fungal infections as well as in the degradation of inhaled chitin (68, 70–72). Nance *et al.* (73) have elegantly demonstrated that AMCase activity is essential to control chitin-containing cysts in the brains of mice chronically infected with *Toxoplasma gondii*. AMCase in humans and mice is strongly associated with Th2 responses (68, 70), and blockade of AMCase activity has profound consequences for Th2 mediated, allergic airway inflammation (74). It is therefore very likely that these molecules are involved in protection against Th2-inducing chitin-containing nematodes. They may also contribute to degradation of chitin-containing parasite-debris that is shed during molting. Additionally, because chitin itself has profound effects on the immune system (75), the ability to break down chitin or even just bind to it, means these proteins will be involved in shaping the immune response triggered by chitin-containing parasites.

In contrast to the active chitinases, the function of YM1 and the other CLPs is still something of a mystery. This is

despite a remarkable depth of knowledge on the structural biology of these proteins, due to the propensity for YM1 to form crystals (76–78). Although crystal formation has so far only been described in the context of lung pathology, it is feasible to speculate that the large quantities of YM1 produced in the direct environment of parasites are taken up during feeding. Crystal formation could lead to internal damage in the gastrointestinal system of the parasite. This is unlikely to be sufficient for parasite killing, as both C57BL/6 and BALB/c mice produce relatively large quantities of YM1 in response to *L. sigmodontis* infection but only C57BL/6 mice are able to kill the parasites before patency (19, 79). Unlike many other CLPs, YM1 does not actually bind chitin but has been shown to bind components of the extracellular matrix such as heparan sulfate (80). This would be consistent with a tissue remodeling or repair function rather than active parasite killing. YM1 has also been implicated in the immune regulation of T-cell responses by IL-4 activated dendritic cells *in vitro* and *in vivo* (81–83) and has direct effects on CD8⁺ T-cell proliferation (84). Taken together, YM1 is likely to have a role in the defense against parasitic infections but this may be through regulatory and tissue remodeling functions rather than direct anti-parasite action.

RELM α / β

Another key molecule strongly associated with alternative macrophage activation and thus potentially important for the resistance against parasite infections is RELM α (*Retnla*) (also called FIZZ-1). Indeed, more than arginase or YM1, which can be induced by classical stimuli such as LPS (85, 86), upregulation of RELM α is typically very reliant on the IL-4R α for enhanced expression (13). It is important to note, however, that macrophages are not the only RELM α source. Eosinophils, neutrophils, and dendritic cells as well as non-hematopoietic cells (e.g. epithelium) express RELM α under certain circumstances (87–90). One argument for a function in host defense is that RELM β , a closely related family member produced by epithelial cells (91), plays a direct role in the attrition and expulsion of intestinal, lumen dwelling parasites (92). This effect seems to mainly be mediated by binding of RELM β to chemosensory organs of parasites (93) limiting worm chemotaxis and the location of their food source eventually leading to starvation and death (92). In contrast no direct interaction of RELM α with parasites or parasite products has been described so far and somewhat surprisingly RELM α has instead been associated with suppression of Th2 immune responses and reduced

resistance to gastrointestinal parasite infection (59, 87). Furthermore RELM α from ‘alternatively activated’ dendritic cells during the priming of a Th2 response *in vitro* leads to enhanced IL-10 production from T cells (82) and RELM α has been implicated in the induction of Th17-responses during bacterial infection (94). Thus, one of the main effects of RELM α seems to be the modulation of the adaptive immune response limiting Th2 responses and enhancing Th17- or regulatory responses. Nonetheless, RELM α may still promote parasite resistance. For example, RELM α can act as a chemoattractant for eosinophils (95, 96), which are strongly connected with resistance to certain parasite infections (97).

A critical function for RELM α may prove to be in the regulation of host energy metabolism and glucose homeostasis during infection. RELM α is related to resistin, a hormone associated with insulin resistance (98) and has been linked to glucose tolerance and the regulation of serum leptin levels during colonic inflammation (90). RELM α itself has a cholesterol-lowering effect and protects against atherosclerosis in murine models (99). Although the impact of this metabolic control on helminth infection has yet to be determined, it is increasingly recognized that alternative macrophage activation not only induces a very distinct metabolic phenotype within the macrophage but also affects the metabolism of the whole organism (100–102). Thus, RELM α seems to adopt very diverse roles dependent on the context and the infection site, making its role in resistance to parasite infection difficult to dissect. Moreover, whether these differences in functionality are associated with different cellular sources or specific modifications of the RELM α protein (e.g. proteolytic cleavage, glycosylation etc.) remain to be elucidated. Overall, unlike its sibling, RELM β , RELM α may have no direct anti-parasite effects but instead may modulate the host immune responses that affect worm survival and promote host tolerance to infection.

Arginase

The last of the ‘Big Three’ molecules associated with alternative activation of macrophages is arginase 1 (103). Arginase 1 is the enzyme that most typifies the immunomodulatory and cross-regulatory nature of macrophage activation. Arginase 1 competes with inducible nitric oxide synthase (iNOS/NOS2) for their mutual substrate L-arginine. Because iNOS is important in the production of reactive nitrogen species and microbial killing, preferential expression of arginase-1 in alternatively activated macrophages serves the dual

purpose of creating important anti-nematode mediators (see below) as well as inhibiting concomitant NOS2 activity. Importantly this inhibitory effect is not restricted to just NO-production but also limits the release of pro-inflammatory cytokines by macrophages and restricts inflammatory cell recruitment during endotoxemia (104). Because arginase activity can cause systemic changes in arginine availability, the immunosuppressive effects extend beyond macrophage intrinsic effector functions and lead to impaired T cell responses (104–107). Thus like YM1 and RELM α , arginase 1 functions as a modulator of immune responsiveness. However, unlike YM1 and RELM α , arginase-1 has been directly linked to resistance to helminth infections.

Depletion of macrophages or inhibition of arginase during secondary infection with *H. polygyrus* results in loss of a protective memory response and failure to expel the parasites (28, 51). Thus, although arginase expression can limit immune responses including T cell activation, it also negatively impacts on the invading parasite. Whether this is due to a reduction in the availability of L-arginine to the growing parasite remains to be tested. More recent work suggests an alternate mechanism of arginase 1-mediated parasite control. L-ornithine and polyamines, products of arginase 1 mediated arginine catabolism, reduced the migratory capacity of *H. polygyrus* larvae *in vitro* (46) and inhibition of arginase 1 prevented retention and trapping of *N. brasiliensis* larvae in the skin of infected animals (108). Proline and hydroxyproline, further downstream products of arginase activity, are central constituents of collagen (109), a major component of extracellular matrix, and are found at high levels in *S. mansoni*-induced liver granulomas and therefore potentially important in restricting larval migration. Thus, the effects of arginase on parasite survival and resistance to infection may in part be mediated through its well-known role in tissue remodeling rather than immune modulation or arginine removal. Of note, the Wynn laboratory (58) found the consequences of arginase 1 expression depend on the cellular source. In their model of liver fibrosis caused by *S. mansoni* egg deposition, macrophage-derived arginase expression was central to immunosuppressive and regulatory but not pro-fibrotic functions. They propose that fibroblast derived arginase is more important for extracellular matrix deposition (58). Taken together, these studies illustrate that both arginase expression and macrophages can be of central importance in the resistance to helminth infections, but the specific contribution of macrophage-derived arginase will need to be assessed in each model.

INOS/ROS

As mentioned above, helminth infections are most commonly associated with a protective Th2 immune response and alternative activation of macrophages. However, anti-bacterial effector molecules such as reactive nitrogen or oxygen species due to enhanced activity of the corresponding enzymes (i.e. INOS, NOX) have been shown to effectively kill helminth parasites *in vitro* (110, 111). Because this protection would come at the cost of enhanced pathology and tissue destruction (7), such responses are usually tightly controlled and suppressed. Nonetheless, several findings indicate that components of type 1 immunity do contribute to parasite killing and expulsion of helminth infections. For example, a lack of IFN γ and consequently reduced neutrophil accumulation has been associated with prolonged survival/persistence of adult *L. sigmodontis* worms in the pleural cavity of infected mice (112). Similarly formation of extracellular traps (NET), normally associated with the trapping of bacteria and fungi, have been shown to effectively kill *S. stercoralis* larvae in a collaborative effort of alternatively activated macrophages and neutrophils (55). In the case of the tapeworm *Taenia crassiceps*, type 1 responses and iNOS are essential for resistance at the early stages of infection (113). Thus, although pro-inflammatory responses are typically suppressed during helminth infections, localized expression (possibly in the context of granulomas as discussed below) with tightly managed release of cytotoxic mediators might provide additional means by which the immune system can control helminth infections. It may also be that when the type 2 effectors alone cannot effectively control infection, the contribution of more inflammatory type 1 pathways are a necessary risk, especially in the context of more lethal infections, such as those caused by some tapeworm species.

Multinucleated giant cell formation

One characteristic of many helminth infections is the formation of dense granulomas around the invading parasite (50, 114). These usually consist of highly organized layers of immune cells with macrophages forming a central part. Some of these macrophages then fuse together to form multinucleated giant cells (MNGCs). MNGCs and granulomas were originally described in the context of Th1-biased *Mycobacterium tuberculosis* infection and were thought to help prevent the spread of infectious mycobacteria (115). It is becoming more apparent, however, that granulomatous reactions are also a central part of many Th2-mediated

responses, and indeed seem to be an integral part of the alternative activation program of macrophages. Many of the molecules thought to be involved in the formation of MNGC (e.g. E-cadherin, CD36, Mannose receptor) are also markers of AAM Φ , and incubation of bone marrow derived macrophages with IL-4 or IL-13 *in vitro* will lead to fusion and MNGC-formation (116–119). The consequences of macrophage fusion and granuloma formation during helminth infections are not yet clear, but the encapsulation of parasites may prevent them from causing excessive tissue damage or, as discussed above, help to trap them in tissues more easily accessible to immune cells (120). In this context, Rajan *et al.* (49) showed that macrophages attached to *B. malayi* larvae lead to deformations/changes in the underlying cuticle of the parasite which allowed eosinophils to transmigrate underneath the cuticle and potentially damage the parasite. Partial encapsulation of parasites or simple attachment of immune cells alone, however, does not seem to be sufficient to efficiently kill adult parasites as we often observe healthy looking motile filarial nematodes in which a section of the parasite is encased in a granulomatous structure. Thus entrapment might just be the first step allowing the immune system enough time to deploy additional defense mechanisms.

Attachment of macrophages to introduced foreign material (e.g. implants) creates a privileged space between the cell-membrane and the surface of the material into which degradative enzymes, acid, and reactive oxygen or nitrogen intermediates can safely be released without risk of deactivation or damage to the surrounding tissue (116, 121). Macrophage fusion increases the surface area covered by the privileged space (122). MNGC-formation during nematode infection might thus constitute a form of foreign body reaction where the inability of individual macrophages to engulf a particular parasite leads to their fusion in a process called 'frustrated phagocytosis' (121). These fused macrophages help create a 'restricted' environment surrounding the parasite in which secretion of actively 'killing' molecules or removal of essential nutrients to starve the parasite, could limit worm survival. A major chicken and egg question remains: do parasites get encapsulated because they are dying/damaged or does the encapsulation cause their death?

Complement

To develop both a broader and more detailed picture of the function of the IL-4R α on macrophages during nematode infection, we recently performed an in-depth RNAseq analysis of IL-4R α -deficient and -sufficient macrophages elicited

by implantation of the peritoneal cavity with the nematode *B. malayi* or exposure to thioglycollate (9). Surprisingly, one of the most differentially regulated pathways in AAM Φ as compared to thioglycollate elicited macrophages was the complement and coagulation cascade (9). Although the upregulation of the associated transcripts was not entirely IL-4R α dependent the sheer number of transcripts associated with this pathway exceeded all expectations. C3, the central component of all complement activation pathways, was among the most abundant gene transcripts found in nematode elicited AAM Φ . Under normal circumstances hepatocytes are the major source of C3 in serum. However, many more cell types including macrophages and monocytes can express complement factors locally (123). The functional relevance of local versus systemic complement production, however, is not well understood.

In the context of filarial nematode infection, the absence of cellular recruitment might prevent the leakage of fluids and serum components from the vasculature that would normally occur during inflammatory processes (13). Thus, local sources of these components may be required to allow efficient accumulation of complement components at the infection site, perhaps binding to the parasite and facilitating the adherence of immune cells. The contribution of complement to nematode control has been demonstrated in recent studies (46, 56), and the importance of local C3-production is highlighted by the fact that parasitic nematodes have evolved counter-measures to inactivate C3 (124). Of note the induction of these counter-mechanisms seems to be stage specific with early stage larvae being more susceptible than adult parasites (125). Thus, local production of complement factors by macrophages might allow rapid attack of invading parasites and avoid the induction of inflammatory processes that might lead to tissue damage. This hypothesis is also supported by an earlier finding that activated C3 acts as a chemokine attracting eosinophils but not neutrophils and thus circumventing normal inflammatory recruitment pathways (126).

It may also be that complement components are contributing to the anti-inflammatory environment of helminth infection. C1q and C3, which are both highly upregulated in AAM Φ (9), have been found to exhibit profound anti-inflammatory properties. Pretreatment with C3 containing serum enhances phagocytosis and uptake of *Francisella tularensis* by human macrophages but simultaneously reduces NF- κ B signaling and pro-inflammatory cytokine secretion (127). Similarly, uptake of C1q-bound apoptotic cells by human macrophages alters LPS-induced cytokine production to a

more immunoregulatory phenotype with increased release of IL-10, IL-27, and IL-33 and reduced inflammasome activation (128). Thus, similar to the many AAM Φ effector molecules, release of complement components during helminth infection might serve a dual or even multiple purposes; complement factors may allow efficient attachment of cells to invading larvae and help to prevent patent infection, while at the same time limiting excessive pro-inflammatory responses and hence pathological sequelae. Dissecting the contribution of complement to both host resistance to nematode infection, and host tolerance through suppression of inflammatory responses is an exciting area of future research.

Metabolic re-programming and nutrient sequestration

Another very prominent characteristic evident from our RNAseq data is the dramatic change in metabolic pathways utilized by macrophages following parasite infection (9). Changes in energy metabolism have been described for IL-4 activated macrophages before (100, 129) and indeed have been described as a cardinal feature of alternative activation (130). However, the extent to which these changes dominate the alternative activation phenotype have not been fully appreciated. Twenty-three of the 27 most highly regulated pathways in nematode elicited macrophages were connected to metabolic processes (9). Interestingly many of these pathways were also associated with amino acid degradation or processing (alanine, aspartate and glutamate metabolism; valine, leucine and isoleucine degradation; valine, leucine and isoleucine biosynthesis; arginine and proline metabolism; glycine serine and threonine metabolism). Given the well documented role of nutritional immunity, i.e. the withholding of essential nutrients from invading pathogens, in resistance to bacterial and intracellular infections (131, 132), it is feasible that these pathways perform a similar role in helminth infection.

IL-4R α has a major impact on the expression of specific amino acid transporters in AAM Φ . In particular, there is increased expression of transporters associated with arginine uptake (Slc7a2, Slc7a4, and Slc36a2) and downregulation of a transporter involved in arginine efflux (Slc7a8) (9). However, we do not know whether the differential regulation of these pathways provides a means of removing essential nutrients from the direct environment of the parasite and eventually leading to its starvation, or whether they provide building blocks for host metabolic processes. In this context, it is important to note that transcripts associated with uptake

of metal ions (e.g. transferrin, S100a1) are also expressed at enhanced levels by AAM Φ . Thus, AAM Φ seem to increase the uptake of several essential nutrients. The metabolic changes in nematode elicited macrophages provide an exciting new avenue of investigation. Further research is required to ascertain the functional consequence for the resistance to helminth infection or indeed, tissue homeostasis during infection, as discussed below.

Tissue repair

Macrophages are always present following tissue injury and regulate all aspects of repair, from the initial inflammatory phase, to cellular proliferation, angiogenesis, wound contraction, matrix deposition, and tissue remodeling and/or scar formation (133, 134). Although the initial inflammatory phase is typically associated with M1-type macrophages, there is a large and growing literature documenting that a reprogramming of macrophages away from an M1 phenotype promotes tissue repair and regeneration (reviewed in 135). These 'repair' macrophages are typically called M2, but this encompasses an enormous range of potential phenotypes, and the specific contribution of IL-4R α signaling to repair and regeneration processes still needs to be elucidated.

The evidence that IL-4R α signaling on macrophages contributes to repair is mainly circumstantial although increasingly strong. The IL-4R α dependent production of arginase is one of the earliest and best examples. Arginase contributes to tissue remodeling and repair because ornithine generated by arginase activity can be converted to polyamines and proline, supporting cell proliferation and collagen synthesis, respectively (109). These properties also explain the frequent association of arginase with fibrosis and in particular asthma, where it is believed to contribute to pathological tissue remodeling (136). Although it was the production of arginase by 'M2' macrophages that was largely responsible for them being associated with tissue repair, it seems that the pro-fibrotic activity of arginase may not be macrophage-derived. When arginase deficiency is restricted to macrophages, the T-cell suppressive but not the pro-repair/fibrosis functions are mediated by macrophages (58). Nonetheless, the ability to control inflammation is a critical component of wound repair (137) and thus macrophage-derived arginase is still likely to be an important player in many injury contexts.

In addition to arginase, RELM α and the CLP YM1 are highly but transiently upregulated in response to incisional

wounding (21). Despite the reported failure to detect IL-4 or IL-13 in wounds (138), we found that the expression of these markers was entirely IL-4R α dependent during wounding (21). There is considerable evidence for a pro-repair role for RELM α . Several papers have documented the angiogenic as well as mitogenic properties of RELM α *in vitro* (139, 140). Indeed, instillation of recombinant RELM α into the lungs of mice induced a similar phenotype as seen after pneumonectomy with enhanced lung hyperplasia and epithelial cell proliferation, typical signs of wound healing (141). However, the contribution of RELM α to repair may be highly complicated by its ability to suppress Th2 responses (58, 59). So although RELM α has direct repair functions, it also acts in a negative feedback loop to control fibrosis. Thus, its role in injury and wound healing is likely highly context dependent. CLPs, and in particular YM1, are frequently identified at sites of injury. However, until recently, direct evidence for involvement of CLPs in repair, was lacking. In a bleomycin model of lung injury, Zhou *et al.* (142) have now shown that Chi311 (BRP39) promotes repair through augmented alternative macrophage activation, fibroblast formation, and matrix deposition.

Many other repair proteins are regulated by IL-4 or IL-13 in macrophages contributing to the evidence that tissue protection is a key function for AAM Φ . For example, transcriptionally, we observed that extracellular matrix degrading matrix metalloproteases (MMPs) are actively downregulated by IL-4R α -mediated signaling in macrophages, while their inhibitors (TIMP1 and TIMP2) are upregulated (9). Similarly, we observe IL-4R α dependent induction of insulin-like growth factor 1 (IGF-1) during helminth infection (9) as previously demonstrated in response to Th2 cytokines *in vitro* (143). IGF-1 has a long established role in repair in part through its ability to stimulate the proliferation and survival of fibroblasts and myofibroblasts and to promote matrix production and wound closure (144, 145). The importance of IGF-1 in repair during helminth infection was illustrated in a recent study by Chen *et al.* (146), in which IGF-1-producing AAM Φ were needed to prevent excessive tissue destruction and hemorrhage caused by lung migrating larvae.

Although more direct evidence is needed, there is little question that macrophages activated via the IL-4R α contribute to repair. Wound repair, especially in adults, is often associated with scar tissue and when excessive leads to fibrosis. It has been proposed that wound healing is evolutionarily optimized for speed of healing under dirty

conditions and thus scar tissue may be the price mammals must pay to close wounds sufficiently rapidly to avoid infection (147). Because it seems likely that a fundamental function of type 2 immunity is to accelerate repair, it is not surprising that AAM Φ and their products are associated with fibrosis (2). The future challenge will be to identify the specific functions of AAM Φ -derived repair molecules in specific settings and, if at all possible, to identify pro-repair versus pro-fibrotic pathways. This will require testing the quality and rate of repair in macrophage specific deletions of the IL-4R α as well as individual effector molecules.

Because macrophages are rapidly recruited to the site of tissue injury, the predominant source of 'wound' macrophages are likely to be blood-circulating monocytes. The currently held view is that the early Ly6Chi monocytes are replaced by Ly6Clo 'patrolling' monocytes as repair progresses. Indeed, the inflammatory monocytes may only be needed to control potential infection, with the later stage macrophages more essential for full repair (133). However, with our growing understanding of the importance and longevity of tissue-resident macrophages, it seems likely that in many contexts tissue-resident macrophages will contribute, particularly at later stages of repair, when the inflammatory monocytes have gone. If so, they will need to expand in number, potentially by IL-4, or perhaps CSF-1 driven proliferation.

Suppressing inflammation

IL-4 has long been considered an anti-inflammatory cytokine (148), and macrophages activated by IL-4 have important functions in suppressing immune responses (reviewed in 48). Inflammation needs to be controlled to allow effective wound repair (137), and thus, the anti-inflammatory nature of AAM Φ is intimately linked to their wound repair phenotype. This makes evolutionary sense in the context of the adaptive type 2 response: a host infected with macroparasites would want to repair any damage caused by a large tissue migrating parasite but also avoid the serious collateral consequences of mounting an inflammatory response to the pathogen.

The data supporting an anti-inflammatory role for AAM Φ have come from many laboratories and has been mostly based on the evidence that AAM Φ are important sources of immunosuppressive cytokines. These include TGF- β (149, 150), PGE2 (151), and the IL-1 receptor antagonist (149, 152). The chemokine expression profile is also strongly associated with a non-inflammatory role (153) indicated by

specific downregulation of key pro-inflammatory chemokines via IL-4 (8, 9, 152).

The ability of a single protein to be both anti-inflammatory and a central mediator of repair is demonstrated by TGF- β , which is known for both immune suppression and fibrosis (154). This functional duality is typical of many Th2-activated macrophage products. As discussed earlier, Arg-1 promotes collagen deposition on one hand but inhibits nitric oxide production and T-cell activation on the other. 12/15 lipoxygenase, a protein strongly associated with IL-4 activation of macrophages in both mice and people, is needed for effective wound repair (155) but attenuates pro-inflammatory macrophage activation (156). RELM α exhibits angiogenic properties (139) and YM1/2 bind extracellular matrix (157), but these are also implicated in regulating inflammation (59, 81, 87). Similarly, Chi311 contributed to repair in a model of bleomycin-induced lung injury both through direct repair functions and suppression of inflammation (142). Thus, modulation of inflammation is intrinsically linked to wound repair and an important characteristic of AAM Φ .

Although IL-10 is very strongly associated with a M2 phenotype and is often listed as a prototypic cytokine associated with alternative activation (47, 149), new sequencing technologies and tools for assessing cell-specific expression *in vivo*, have made the story more complicated. Despite abundant expression of the IL-10 receptor, YM1, RELM α , arginase and many other AA markers, RNAseq analysis of nematode-elicited macrophages had sufficient depth of coverage to be able to say that IL-10 is not produced by F4/80hi macrophages in this context (9). The important source of IL-10 following both hookworm migration through the lung and filarial nematode infection appears to be T cells (146, 158). This is consistent with recent data showing anti-inflammatory functions of gut macrophages are mediated not by their ability to produce IL-10 but by IL-10 receptor expression (159). The lack of IL-10 production by AAM Φ may be surprising considering the known ability of macrophages to produce IL-10 in response to antibody cross-linking (160), and indeed we would expect antibodies to be present in the helminth models discussed above. However, IL-10 production by macrophages may require additional signals, such as TLR ligands and type 1 interferons (161, 162), that may not be present in these nematode infection sites. The absence of macrophage derived IL-10 despite high levels of parasite specific antibody and antigen, may be further explained by Fc γ receptor usage on AAM Φ during helminth infection. We

found that IL-4R α signaling drives a switch from the activating Fc receptors involved in IL-10 production (160) to the inhibitory Fc γ RIIb (9, 163).

The combined anti-inflammatory/wound healing function of AAM Φ is strongly supported by a study of *S. mansoni* infection in mice that lack the IL-4R α specifically on macrophages and neutrophils but have otherwise intact Th2 responses (60). Following *S. mansoni* infection, these mice died from overwhelming inflammatory responses in the intestine and leakage of bacteria into the blood. Although not conclusive evidence, the data strongly suggest that in the absence of AAM Φ , these mice were unable to repair the damage caused by egg migration through the intestinal wall and the subsequent septic inflammation (60). The specific roles Th2-induced proteins play in the complex orchestra of tissue repair and remodeling are still to be established, but a rapid shutdown of the inflammatory response to injury is likely an important contribution.

Immune regulation

A paradox that still needs to be resolved is that AAM Φ , which actively proliferate *in vivo*, are profoundly antiproliferative *ex vivo*. AAM Φ suppress proliferation of cells with which they are co-cultured *ex vivo* in an IL-4-dependent manner (150, 164). This property was our first insight into the distinct function of macrophages under IL-4R α control and has been reproduced in many laboratories where suppression has been attributed to arginase depletion (58), TGF- β (165), PD-L pathway (166, 167), and 12/15 lipoxygenase (168). Of note, rapid division of immune cells (i.e. effector T cells) has previously been shown to be highly reliant on glucose and the Warburg effect (169). In contrast, alternative activation and also IL-4-driven proliferation of macrophages are dependent on fatty acid oxidation and oxidative phosphorylation (OXPHOS) (100) (Fig. 4). The reasons for these metabolic differences are not yet completely clear. Studies in T cells suggest that aerobic glycolysis allows for quick accumulation of biomass needed for cellular expansion whereas OXPHOS is associated with the enhanced lifespan of memory T cells (170, 171). In an analogous manner, the utilization of OXPHOS in AAM Φ might impart slightly slower proliferative expansion but allow the expanded population to survive for a longer period of time. Another hypothesis is that metabolic differences are linked to the anatomical location in which cellular expansion is likely to occur. T cells normally expand within lymphoid structures which are surrounded by adipose tissue and well vascularized (172, 173). Thus, it is feasible to assume that




Fig. 4. Inhibition of fatty acid oxidation blocks macrophage alternative activation and proliferation in vivo. C57BL/6 mice were injected i.p. with 0.5 mg Etomoxir (Eto, Sigma Aldrich) to inhibit fatty acid oxidation 1 h prior to treatment with IL-4C. Detailed description of similar experiments can be found in Ruckerl et al. (67). (A) 24 h later, the proliferative expansion of F4/80^{high} cells was determined by 3hr-BrdU pulse and flow cytometry. Data points indicate individual animals, and lines indicate mean. (B) Intracellular RELM α expression in the cells analyzed in (A). (C) Number of F4/80^{high} cells present in the pleural cavity of the mice analyzed in (A). (D) Concentration of RELM α in the lavage fluid isolated from the pleural cavity of the mice analyzed in (A).

these structures provide ample nutrients for optimal, rapid cellular expansion. In contrast AAM Φ expand in the tissues in response to IL-4 (13). In the presence of a large, tissue-destructive macroparasite and the accompanying wound healing process, these might suffer from acute shortage of essential nutrients, especially glucose. Thus, the adoption of OXPHOS as a means of generating energy might be necessary to allow the expansion of tissue-resident macrophages during helminth infection. In this context, it is interesting to note that certain tumor cell lines, normally well known for their high glucose consumption and glycolytic energy metabolism, adopt OXPHOS to maintain cellular proliferation under low glucose conditions (174). Furthermore, we observe a highly significant increase in expression of the arginine transporter CAT2 (Slc7a2), as well as other amino acid and nutrient transporters in response to IL-4R α signaling in AAM Φ (9). Transporter upregulation, in addition to the above-discussed effector functions, might allow macrophages to acquire essential nutrients needed for expansion. Thus, the anti-proliferative effect of AAM Φ on bystander cells *in vitro* might at least in part be a side effect of nutrient sequestration to which T cells are highly sensitive (175).

Critically, whether AAM Φ directly affect T-cell proliferation and polarization *in vivo* is not known. Although individual molecules expressed by AAM Φ (i.e. RELM α , YM1,

arginase) have been associated with immune suppression (58, 59, 87), it is not always clear whether it is expression by macrophages that imparts this effect. Indeed, our RNAseq data indicate a remarkable lack of chemokine receptor expression on nematode elicited macrophages (9), suggesting that these cells are prone to stay in the tissues and most likely do not migrate to the draining lymph nodes to direct immune response. Thus, the immune regulation observed *in vivo* (58, 59, 87) might rather be due to other cells, such as DCs (82), expressing AAM Φ -associated molecules or due to alternative activation of macrophages resident within the lymphoid structures (79). Undoubtedly AAM Φ do influence the local immune milieu in the tissues. Recruitment of eosinophils to the site of helminth infection is greatly enhanced by the presence of AAM Φ (176) and recruitment of monocytes and neutrophils is discouraged by IL-4 (9, 19). Thus, AAM Φ contribute to regulating cells in their immediate vicinity via various means and although it has yet to be documented, local T-cell proliferation may also be prevented. As described above in the context of complement, a major feature of AAM Φ may be to regulate the local immune environment and mediate key necessary effector functions, leaving more systemic immune regulation to other cells. In this context, the local expansion of the resident population and avoidance of pro-inflammatory

recruitment makes sense; shaping a 'safe' environment for the control of helminth infection or to mediate repair.

Plasticity

Classical macrophage activation, as found during bacterial infections, and IL-4R α -mediated alternative activation are two extremes of a wide spectrum of overlapping activation phenotypes (160). Although differentially activating stimuli will block or reduce opposing activation signals, macrophages are known to be highly plastic and rarely exhibit fixed phenotypes. Instead they form a continuum, adopting characteristics of activation as needed according to their environment (177). The lack of a fixed activation state is supported by data showing that once the initial stimulating agent is removed, macrophages quickly lose their activation phenotype (178, own unpublished data) and cells isolated from one activating environment subjected to an opposing stimulus will adopt characteristics of the newly activating environment (179). Indeed, macrophages isolated from *B. malayi* infected animals and subjected to stimulation with LPS/IFN γ readily express pro-inflammatory cytokines (e.g. IL-6, IL-10, TNF) and are able to control intracellular replication of *Leishmania mexicana* parasites (85). This flexibility in activation phenotype, demonstrated *in vitro*, is presumed to be important in the context of co-infections allowing macrophages to quickly adapt to new challenges.

Changes in macrophage activation phenotype *in vivo* have been demonstrated in the setting of wound healing, where initially pro-inflammatory macrophages eventually convert to a more anti-inflammatory phenotype (138). Conversely we have reported a change in cytokine expression by macrophages isolated from *B. malayi* infected mice early or late following parasite implantation, with an increase in pro-inflammatory cytokine production at later stages (180). However, *in vivo* macrophage activation, especially in the context of infection, is rarely homogenous, and only a subset of macrophages will show signs of activation (13, 32, 181). Thus, adaptation to environmental changes could at least in part be due to induction of differentially activated subsets of macrophages, rather than true plasticity. Moreover, waves of cellular recruitment/expansion of various macrophage populations might be favored over re-polarization of the existing already activated population. Thus, despite clear evidence of macrophage plasticity *in vitro*, macrophage plasticity during infection has to be re-evaluated. This is especially true in light of our new understanding of macrophage origins and proliferation (24) and the findings

by Gundra *et al.* (20). Macrophages derived from different cellular sources (i.e. blood monocytes versus tissue-resident macrophages) respond differently to the same stimulus (i.e. IL-4) with potentially different functional activation outcomes. Of note, functional differences between resident and monocyte-derived macrophages have been described previously. Uderhardt *et al.* (182) showed preferential uptake of apoptotic cells by resident peritoneal macrophages over inflammatory monocyte-derived cells. Disruption of this bias through genetic ablation of 12/15 lipoxygenase resulted in presentation and recognition of self-antigens leading to autoimmune reactions and lupus-like symptoms (182). Furthermore, although *B. malayi* derived AAM Φ readily upregulated iNOS and pro-inflammatory cytokine-production following stimulation with LPS/IFN γ , we were never able to detect release of IL-12 from these cells, normally a key feature of classical macrophage activation (85). Thus, it is likely that epigenetic or otherwise mediated determinants of macrophage phenotype are induced during development restricting the plasticity/variation in responses (178, 183).

Nonetheless, bone marrow-derived macrophages can adopt a resident-like phenotype and re-populate the tissues after lethal irradiation (24). Thus, these developmental restrictions may be imposed by the local environment rather than in the bone marrow or during embryogenesis. Local regulation of macrophage identity was highlighted in two recent studies, in which expression of the transcription factor GATA6 was shown to be essential for the resident peritoneal macrophage phenotype (184, 185). Furthermore, all-trans retinoic acid and omentum-derived soluble factors could drive expression of resident peritoneal macrophage signature genes in bone marrow derived macrophages *in vitro* (184). Thus, the environment of the peritoneal cavity seems necessary and sufficient to induce the resident phenotype. Although this does not occur in monocyte derived macrophages present in the peritoneal cavity under normal circumstances it highlights the context dependency and flexibility of macrophage phenotypes. In this context, the macrophages recruited to the site of *B. malayi* implant, as analyzed in our RNAseq analysis, are monocytic in origin (authors' unpublished data). Despite clear evidence of proliferation and adoption of a resident-like phenotype (i.e. F4/80^{high}, MHC-II^{low}) (13), data from protected irradiated chimeras suggest that an initial inflammatory influx, caused by the wounding of the surgical incision, leads to the accumulation of monocyte derived cells which are then converted to a resident like phenotype. Thus, in addition to

driving proliferative expansion of resident and monocyte-derived macrophages, IL-4 seems to enhance or enable the conversion of monocyte derived macrophages to a resident phenotype. Consistent with this hypothesis, macrophages from *B. malayi* implanted animals showed highly enhanced GATA6 expression as compared to thioglycollate elicited macrophages (9), and IL-4 induces GATA-6 expression on thioglycollate-elicited macrophages (20). The functional consequences of cellular source, proliferative expansion, and long-term residence of tissue macrophages on the resistance to helminth infection remain a wide open question.

The future

We are entering a new era in macrophage biology, particularly in light of our new understanding of cellular origins. New questions are emerging daily, and we have raised many

of them here in this review. For example, in the context of helminth infection, how do single molecules perform such diverse functions as parasite killing, immune suppression, and wound repair? More broadly, what is the extent of macrophage plasticity? Do recruited cells truly become fully functional resident cells? What are the epigenetic/transcriptional controls that regulate these functions both during development and locally in the tissues? New tools (186, 187) and collaborations with different disciplines from stem cell biology to systems biology will allow us to unravel some of the enormous complexity underlying macrophage function in health and disease. Studying these intricate dynamics in the context of infection with helminths, our ancient evolutionary partners, will further enhance our fundamental understanding of macrophage biology. The challenge will be to translate this new knowledge into the ability to fine tune responses to the benefit of infected or injured hosts.

References

- Hausdorf B. Early evolution of the bilateria. *Syst Biol* 2000;**49**:130–142.
- Allen JE, Wynn TA. Evolution of Th2 immunity: a rapid repair response to tissue destructive pathogens. *PLoS Pathog* 2011;**7**:e1002003.
- Hotez PJ, Brindley PJ, Bethony JM, King CH, Pearce EJ, Jacobson J. Helminth infections: the great neglected tropical diseases. *J Clin Invest* 2008;**118**:1311–1321.
- Lustigman S, et al. A research agenda for helminth diseases of humans: the problem of helminthiasis. *PLoS Negl Trop Dis* 2012;**6**:e1582.
- Crompton DWT, Nesheim MC. Nutritional impact of intestinal helminthiasis during the human life cycle. *Annu Rev Nutr* 2002;**22**:35–59.
- Allen JE, Maizels RM. Diversity and dialogue in immunity to helminths. *Nat Rev Immunol* 2011;**11**:375–388.
- Gause WC, Wynn TA, Allen JE. Type 2 immunity and wound healing: evolutionary refinement of adaptive immunity by helminths. *Nat Rev Immunol* 2013;**13**:607–614.
- Loke P, Nair MG, Parkinson J, Guiliano D, Blaxter M, Allen JE. IL-4 dependent alternatively-activated macrophages have a distinctive in vivo gene expression phenotype. *BMC Immunol* 2002;**3**:7.
- Thomas GD, Ruckerl D, Maskrey BH, Whitfield PD, Blaxter ML, Allen JE. The biology of nematode- and IL4R α -dependent murine macrophage polarization in vivo as defined by RNA-Seq and targeted lipidomics. *Blood* 2012;**120**:e93–e104.
- Gordon S, Martinez FO. Alternative activation of macrophages: mechanism and functions. *Immunity* 2010;**32**:593–604.
- Allen JE, et al. Of mice, cattle, and humans: the immunology and treatment of river blindness. *PLoS Negl Trop Dis* 2008;**2**:e217.
- Taylor MD, LeGoff L, Harris A, Malone E, Allen JE, Maizels RM. Removal of regulatory T cell activity reverses hyporesponsiveness and leads to filarial parasite clearance in vivo. *J Immunol* 2005;**174**:4924–4933.
- Jenkins SJ, et al. Local macrophage proliferation, rather than recruitment from the blood, is a signature of TH2 inflammation. *Science* 2011;**332**:1284–1288.
- Dai X-M, et al. Targeted disruption of the mouse colony-stimulating factor 1 receptor gene results in osteopetrosis, mononuclear phagocyte deficiency, increased primitive progenitor cell frequencies, and reproductive defects. *Blood* 2002;**99**:1111–1120.
- Hume DA, Pavli P, Donahue RE, Fidler IJ. The effect of human recombinant macrophage colony-stimulating factor (CSF-1) on the murine mononuclear phagocyte system in vivo. *J Immunol* 1988;**141**:3405–3409.
- MacDonald KPA, et al. An antibody against the colony-stimulating factor 1 receptor depletes the resident subset of monocytes and tissue- and tumor-associated macrophages but does not inhibit inflammation. *Blood* 2010;**116**:3955–3963.
- Tagliani E, Shi C, Nancy P, Tay C-S, Pamer EG, Erlebacher A. Coordinate regulation of tissue macrophage and dendritic cell population dynamics by CSF-1. *J Exp Med* 2011;**208**:1901–1916.
- Milner JD, et al. Sustained IL-4 exposure leads to a novel pathway for hemophagocytosis, inflammation, and tissue macrophage accumulation. *Blood* 2010;**116**:2476–2483.
- Jenkins SJ, et al. IL-4 directly signals tissue-resident macrophages to proliferate beyond homeostatic levels controlled by CSF-1. *J Exp Med* 2013;**210**:2477–2491.
- Gundra UM, et al. Alternatively activated macrophages derived from monocytes and tissue macrophages are phenotypically and functionally distinct. *Blood* 2014;**123**:e110–e122.
- Loke P, et al. Alternative activation is an innate response to injury that requires CD4⁺ T cells to be sustained during chronic infection. *J Immunol* 2007;**179**:3926–3936.
- Schulz C, et al. A lineage of myeloid cells independent of Myb and hematopoietic stem cells. *Science* 2012;**336**:86–90.
- Hashimoto D, et al. Tissue-resident macrophages self-maintain locally throughout adult life with minimal contribution from circulating monocytes. *Immunity* 2013;**38**:792–804.
- Sieweke MH, Allen JE. Beyond stem cells: self-renewal of differentiated macrophages. *Science* 2013;**342**:1242974.
- Ginhoux F, Jung S. Monocytes and macrophages: developmental pathways and tissue homeostasis. *Nat Rev Immunol* 2014;**14**:392–404.
- Jakubzick C, et al. Minimal differentiation of classical monocytes as they survey steady-state tissues and transport antigen to lymph nodes. *Immunity* 2013;**39**:599–610.
- Bain CC, Mowat AM. Macrophages in intestinal homeostasis and inflammation. *Immunol Rev* 2014;**260**:102–117.
- Anthony RM, et al. Memory T(H)2 cells induce alternatively activated macrophages to mediate protection against nematode parasites. *Nat Med* 2006;**12**:955–960.
- Davies LC, Rosas M, Smith PJ, Fraser DJ, Jones SA, Taylor PR. A quantifiable proliferative burst of tissue macrophages restores homeostatic macrophage populations after acute inflammation. *Eur J Immunol* 2011;**41**:2155–2164.

30. Davies LC, et al. Distinct bone marrow-derived and tissue-resident macrophage lineages proliferate at key stages during inflammation. *Nat Commun* 2013;**4**:1886.
31. Mohrs K, Harris DP, Lund FE, Mohrs M. Systemic dissemination and persistence of Th2 and type 2 cells in response to infection with a strictly enteric nematode parasite. *J Immunol* 2005;**175**:5306–5313.
32. Nascimento M, et al. Ly6Chi monocyte recruitment is responsible for Th2 associated host-protective macrophage accumulation in liver inflammation due to schistosomiasis. *PLoS Pathog* 2014;**10**:e1004282.
33. Girgis NM, Gundra UM, Ward LN, Cabrera M, Frevert U, Loke P. Ly6Chi^{high} monocytes become alternatively activated macrophages in schistosome granulomas with help from CD4⁺ cells. *PLoS Pathog* 2014;**10**:e1004080.
34. Finkelman FD, et al. Interleukin-4- and interleukin-13-mediated host protection against intestinal nematode parasites. *Immunol Rev* 2004;**201**:139–155.
35. Hewitson JP, et al. The secreted triose phosphate isomerase of *Brugia malayi* is required to sustain microfilaria production in vivo. *PLoS Pathog* 2014;**10**:e1003930.
36. van der Werf N, Redpath SA, Azuma M, Yagita H, Taylor MD. Th2 cell-intrinsic hypo-responsiveness determines susceptibility to helminth infection. *PLoS Pathog* 2013;**9**:e1003215.
37. Maizels RM, Yazdanbakhsh M. Immune regulation by helminth parasites: cellular and molecular mechanisms. *Nat Rev Immunol* 2003;**3**:733–744.
38. Graham AL, Taylor MD, Le Goff L, Lamb TJ, Magennis M, Allen JE. Quantitative appraisal of murine filariasis confirms host strain differences but reveals that BALB/c females are more susceptible than males to *Litomosoides sigmodontis*. *Microbes Infect* 2005;**7**:612–618.
39. Volkmann L, et al. Murine filariasis: interleukin 4 and interleukin 5 lead to containment of different worm developmental stages. *Med Microbiol Immunol* 2003;**192**:23–31.
40. Robinson M, Behnke JM, Williams DJ. Immunity to adult *Heligmosomoides polygyrus* (*Nematospiroides dubius*): survival or rejection of adult worms following transplantation to mice refractory to larval challenge. *J Helminthol* 1988;**62**:221–231.
41. Maizels RM, Balic A, Gomez-Escobar N, Nair M, Taylor MD, Allen JE. Helminth parasites—masters of regulation. *Immunol Rev* 2004;**201**:89–116.
42. Casaravilla C, et al. Unconventional maturation of dendritic cells induced by particles from the laminated layer of larval *Echinococcus granulosus*. *Infect Immun* 2014;**82**:3164–3176.
43. Babayan SA, Allen JE, Taylor DW. Future prospects and challenges of vaccines against filariasis. *Parasite Immunol* 2012;**34**:243–253.
44. Babayan SA, Attout T, Vuong PN, Le Goff L, Gantier J-C, Bain O. The subcutaneous movements of filarial infective larvae are impaired in vaccinated hosts in comparison to primary infected hosts. *Filaria J* 2005;**4**:3.
45. Urban JF, Katona IM, Finkelman FD. *Heligmosomoides polygyrus*: CD4⁺ but not CD8⁺ T cells regulate the IgE response and protective immunity in mice. *Exp Parasitol* 1991;**73**:500–511.
46. Esser-von Bieren J, et al. Antibodies trap tissue migrating helminth larvae and prevent tissue damage by driving IL-4R α -independent alternative differentiation of macrophages. *PLoS Pathog* 2013;**9**:e1003771.
47. Kreider T, Anthony RM, Urban JF, Gause WC. Alternatively activated macrophages in helminth infections. *Curr Opin Immunol* 2007;**19**:448–453.
48. Jenkins SJ, Allen JE. Similarity and diversity in macrophage activation by nematodes, trematodes, and cestodes. *J Biomed Biotechnol* 2010;**2010**:262609–262614.
49. Rajan TV, Ganley L, Paciorkowski N, Spencer L, Klei TR, Shultz LD. Brugian infections in the peritoneal cavities of laboratory mice: kinetics of infection and cellular responses. *Exp Parasitol* 2002;**100**:235–247.
50. Patel N, Kreider T, Urban JF, Gause WC. Characterisation of effector mechanisms at the host:parasite interface during the immune response to tissue-dwelling intestinal nematode parasites. *Int J Parasitol* 2009;**39**:13–21.
51. Filbey KJ, et al. Innate and adaptive type 2 immune cell responses in genetically controlled resistance to intestinal helminth infection. *Immunol Cell Biol* 2014;**92**:436–448.
52. Zhao A, et al. Th2 cytokine-induced alterations in intestinal smooth muscle function depend on alternatively activated macrophages. *Gastroenterology* 2008;**135**:217–225 e1.
53. Nakanishi H, Hori Y, Terashima K, Fujita K. Effect of macrophage blockade on the resistance to a primary *Brugia pahangi* infection of female BALB/c mice. *Trop Med Parasitol* 1989;**40**:75–76.
54. Rao UR, Vickery AC, Kwa BH, Nayar JK, Subrahmanyam D. Effect of carriage on the resistance of congenitally athymic nude and normal BALB/c mice to infective larvae of *Brugia malayi*. *Parasitol Res* 1992;**78**:235–240.
55. Bonne-Année S, et al. Extracellular traps are associated with human and mouse neutrophil and macrophage mediated killing of larval *Strongyloides stercoralis*. *Microbes Infect* 2014;**16**:502–511.
56. Bonne-Année S, et al. Human and mouse macrophages collaborate with neutrophils to kill larval *Strongyloides stercoralis*. *Infect Immun* 2013;**81**:3346–3355.
57. Hume DA. Applications of myeloid-specific promoters in transgenic mice support in vivo imaging and functional genomics but do not support the concept of distinct macrophage and dendritic cell lineages or roles in immunity. *J Leukoc Biol* 2011;**89**:525–538.
58. Pesce JT, et al. Arginase-1-expressing macrophages suppress Th2 cytokine-driven inflammation and fibrosis. *PLoS Pathog* 2009;**5**:e1000371.
59. Nair MG, et al. Alternatively activated macrophage-derived RELM- α is a negative regulator of type 2 inflammation in the lung. *J Exp Med* 2009;**206**:937–952.
60. Herbert DR, et al. Alternative macrophage activation is essential for survival during schistosomiasis and downmodulates T helper 1 responses and immunopathology. *Immunity* 2004;**20**:623–635.
61. Urban JF, Noben-Trauth N, Schopf L, Madden KB, Finkelman FD. Cutting edge: IL-4 receptor expression by non-bone marrow-derived cells is required to expel gastrointestinal nematode parasites. *J Immunol* 2001;**167**:6078–6081.
62. Michels CE, et al. Neither interleukin-4 receptor alpha expression on CD4⁺ T cells, or macrophages and neutrophils is required for protective immunity to *Trichinella spiralis*. *Immunology* 2009;**128**:e385–e394.
63. Grecis RK, Humphreys NE, Bancroft AJ. Immunity to gastrointestinal nematodes: mechanisms and myths. *Immunol Rev* 2014;**260**:183–205.
64. Taylor MD, et al. Early recruitment of natural CD4⁺Foxp3⁺ Treg cells by infective larvae determines the outcome of filarial infection. *Eur J Immunol* 2008;**39**:192–206.
65. Babayan SA, Read AF, Lawrence RA, Bain O, Allen JE. Filarial parasites develop faster and reproduce earlier in response to host immune effectors that determine filarial life expectancy. *PLoS Biol* 2010;**8**:e1000525.
66. Falcone FH, Loke P, Zang X, MacDonald AS, Maizels RM, Allen JE. A *Brugia malayi* homolog of macrophage migration inhibitory factor reveals an important link between macrophages and eosinophil recruitment during nematode infection. *J Immunol* 2001;**167**:5348–5354.
67. Ruckerl D, et al. Induction of IL-4R α -dependent microRNAs identifies PI3K/Akt signaling as essential for IL-4-driven murine macrophage proliferation in vivo. *Blood* 2012;**120**:2307–2316.
68. Sutherland TE, Maizels RM, Allen JE. Chitinases and chitinase-like proteins: potential therapeutic targets for the treatment of T-helper type 2 allergies. *Clin Exp Allergy* 2009;**39**:943–955.
69. Bussink AP, Speijer D, Aerts JMFG, Boot RG. Evolution of mammalian chitinase(-like) members of family 18 glycosyl hydrolases. *Genetics* 2007;**177**:959–970.
70. Zhu Z, et al. Acidic mammalian chitinase in asthmatic Th2 inflammation and IL-13 pathway activation. *Science* 2004;**304**:1678–1682.
71. Hartl D, et al. Acidic mammalian chitinase regulates epithelial cell apoptosis via a chitinolytic-independent mechanism. *J Immunol* 2009;**182**:5098–5106.
72. Seibold MA, et al. Chitotriosidase is the primary active chitinase in the human lung and is modulated by genotype and smoking habit. *J Allergy Clin Immunol* 2008;**122**:944–950 e3.
73. Nance JP, et al. Chitinase dependent control of protozoan cyst burden in the brain. *PLoS Pathog* 2012;**8**:e1002990.
74. Sutherland TE, et al. Analyzing airway inflammation with chemical biology: dissection of acidic mammalian chitinase function with a

- selective drug-like inhibitor. *Chem Biol* 2011;**18**:569–579.
75. Reese TA, et al. Chitin induces accumulation in tissue of innate immune cells associated with allergy. *Nature* 2007;**447**:92–96.
 76. Guo L, Johnson RS, Schuh JC. Biochemical characterization of endogenously formed eosinophilic crystals in the lungs of mice. *J Biol Chem* 2000;**275**:8032–8037.
 77. Marchesi F, et al. Eosinophilic crystals as a distinctive morphologic feature of a hyaline droplet nephropathy in a mouse model of acute myelogenous leukaemia. *J Vet Med A Physiol Pathol Clin Med* 2003;**50**:103–107.
 78. Harbord M, et al. Ym1 is a neutrophil granule protein that crystallizes in p47phox-deficient mice. *J Biol Chem* 2002;**277**:5468–5475.
 79. Nair MG, et al. Chitinase and Fizz family members are a generalized feature of nematode infection with selective upregulation of Ym1 and Fizz1 by antigen-presenting cells. *Infect Immun* 2005;**73**:385–394.
 80. Chang NC, et al. A macrophage protein, Ym1, transiently expressed during inflammation is a novel mammalian lectin. *J Biol Chem* 2001;**276**:17497–17506.
 81. Cai Y, Kumar RK, Zhou J, Foster PS, Webb DC. Ym1/2 promotes Th2 cytokine expression by inhibiting 12/15(S)-lipoxygenase: identification of a novel pathway for regulating allergic inflammation. *J Immunol* 2009;**182**:5393–5399.
 82. Cook PC, Jones LH, Jenkins SJ, Wynn TA, Allen JE, MacDonald AS. Alternatively activated dendritic cells regulate CD4+ T-cell polarization in vitro and in vivo. *Proc Natl Acad Sci USA* 2012;**109**:9977–9982.
 83. Arora M, et al. Simvastatin promotes Th2-type responses through the induction of the chitinase family member Ym1 in dendritic cells. *Proc Natl Acad Sci USA* 2006;**103**:7777–7782.
 84. Osborne LC, et al. Coinfection. Virus-helminth coinfection reveals a microbiota-independent mechanism of immunomodulation. *Science* 2014;**345**:578–582.
 85. Mylonas KJ, Nair MG, Prieto-Lafuente L, Paape D, Allen JE. Alternatively activated macrophages elicited by helminth infection can be reprogrammed to enable microbial killing. *J Immunol* 2009;**182**:3084–3094.
 86. El Kasmi KC, et al. Toll-like receptor-induced arginase 1 in macrophages thwarts effective immunity against intracellular pathogens. *Nat Immunol* 2008;**9**:1399–1406.
 87. Pesce JT, et al. Retnla (relmalpha/fizz1) suppresses helminth-induced Th2-type immunity. *PLoS Pathog* 2009;**5**:e1000393.
 88. Osborne LC, et al. Resistin-like molecule α promotes pathogenic Th17 cell responses and bacterial-induced intestinal inflammation. *J Immunol* 2013;**190**:2292–2300.
 89. Holcomb IN, et al. FIZZ1, a novel cysteine-rich secreted protein associated with pulmonary inflammation, defines a new gene family. *EMBO J* 2000;**19**:4046–4055.
 90. Munitz A, Seidu L, Cole ET, Ahrens R, Hogan SP, Rothenberg ME. Resistin-like molecule alpha decreases glucose tolerance during intestinal inflammation. *J Immunol* 2009;**182**:2357–2363.
 91. Patel SD, Rajala MW, Rossetti L, Scherer PE, Shapiro L. Disulfide-dependent multimeric assembly of resistin family hormones. *Science* 2004;**304**:1154–1158.
 92. Herbert D, et al. Intestinal epithelial cell secretion of RELM- β protects against gastrointestinal worm infection. *J Exp Med* 2009;**206**:2947–2957.
 93. Artis D, et al. RELM β /FIZZ2 is a goblet cell-specific immune-effector molecule in the gastrointestinal tract. *Proc Natl Acad Sci USA* 2004;**101**:13596–13600.
 94. Chen G, Chan AJ, Chung JI, Jang JC, Osborne LC, Nair MG. Polarizing the T helper 17 response in *Citrobacter rodentium* infection via expression of resistin-like molecule α . *Gut Microbes* 2014;**5**:24–25.
 95. Munitz A, Cole ET, Karo-Atar D, Finkelman FD, Rothenberg ME. Resistin-like molecule- α regulates IL-13-induced chemokine production but not allergen-induced airway responses. *Am J Respir Cell Mol Biol* 2012;**46**:703–713.
 96. Liu T, et al. FIZZ1 stimulation of myofibroblast differentiation. *Am J Pathol* 2004;**164**:1315–1326.
 97. Maizels RM, Balic A. Resistance to helminth infection: the case for interleukin-5-dependent mechanisms. *J Infect Dis* 2004;**190**:427–429.
 98. Steppan CM, et al. The hormone resistin links obesity to diabetes. *Nature* 2001;**409**:307–312.
 99. Lee M-R, et al. The adipokine Retnla modulates cholesterol homeostasis in hyperlipidemic mice. *Nat Commun* 2014;**5**:4410.
 100. Vats D, et al. Oxidative metabolism and PGC-1 β attenuate macrophage-mediated inflammation. *Cell Metab* 2006;**4**:13–24.
 101. Nguyen KD, et al. Alternatively activated macrophages produce catecholamines to sustain adaptive thermogenesis. *Nature* 2011;**480**:104–108.
 102. Yang Z, et al. Parasitic nematode-induced modulation of body weight and associated metabolic dysfunction in mouse models of obesity. *Infect Immun* 2013;**81**:1905–1914.
 103. Munder M, Eichmann K, Modolell M. Alternative metabolic states in murine macrophages reflected by the nitric oxide synthase/arginase balance: competitive regulation by CD4+ T cells correlates with Th1/Th2 phenotype. *J Immunol* 1998;**160**:5347–5354.
 104. Wijnands KAP, et al. Arginase-1 deficiency regulates arginine concentrations and NOS2-mediated NO production during endotoxemia. *PLoS ONE* 2014;**9**:e86135.
 105. Munder M, Choi B-S, Rogers M, Kropf P. L-arginine deprivation impairs *Leishmania* major-specific T-cell responses. *Eur J Immunol* 2009;**39**:2161–2172.
 106. Peranzoni E, et al. Role of arginine metabolism in immunity and immunopathology. *Immunobiology* 2007;**212**:795–812.
 107. Zhu X, et al. The central role of arginine catabolism in T-cell dysfunction and increased susceptibility to infection after physical injury. *Ann Surg* 2014;**259**:171–178.
 108. Obata-Ninomiya K, et al. The skin is an important bulwark of acquired immunity against intestinal helminths. *J Exp Med* 2013;**210**:2583–2595.
 109. Morris SM. Arginine metabolism: boundaries of our knowledge. *J Nutr* 2007;**137**:1602S–1609S.
 110. Taylor MJ, Cross HF, Mohammed AA, Trees AJ, Bianco AE. Susceptibility of *Brugia malayi* and *Onchocerca lienalis* microfilariae to nitric oxide and hydrogen peroxide in cell-free culture and from IFN gamma-activated macrophages. *Parasitology* 1996;**112**(Pt 3):315–322.
 111. Pfaff AW, Schulz-Key H, Soboslay PT, Geiger SM, Hoffmann WH. The role of nitric oxide in the innate resistance to microfilariae of *Litomosoides sigmodontis* in mice. *Parasite Immunol* 2000;**22**:397–405.
 112. Saeftel M, et al. Lack of interferon-gamma confers impaired neutrophil granulocyte function and imparts prolonged survival of adult filarial worms in murine filariasis. *Microbes Infect* 2001;**3**:203–213.
 113. Alonso-Trujillo J, Rivera-Montoya I, Rodríguez-Sosa M, Terrazas LI. Nitric oxide contributes to host resistance against experimental *Taenia crassiceps* cysticercosis. *Parasitol Res* 2007;**100**:1341–1350.
 114. Reynolds LA, Filbey KJ, Maizels RM. Immunity to the model intestinal helminth parasite *Heligmosomoides polygyrus*. *Semin Immunopathol* 2012;**34**:829–846.
 115. Langhans T. Ueber Riesenzellen mit wandständigen Kernen in Tuberkeln und die fibröse Form des Tuberkels. *Archiv für pathologische Anatomie und Physiologie und für Klinische Medizin* 1868;**42**:382–404.
 116. Anderson JM, Rodriguez A, Chang D. Foreign body reaction to biomaterials. *Semin Immunol* 2008;**20**:86–100.
 117. Moreno JL, Mikhailenko I, Tondravi MM, Keegan AD. IL-4 promotes the formation of multinucleated giant cells from macrophage precursors by a STAT6-dependent, homotypic mechanism: contribution of E-cadherin. *J Leukoc Biol* 2007;**82**:1542–1553.
 118. Helming L, Gordon S. The molecular basis of macrophage fusion. *Immunobiology* 2007;**212**:785–793.
 119. Kao WJ, McNally AK, Hiltner A, Anderson JM. Role for interleukin-4 in foreign-body giant cell formation on a poly(etherurethane urea) in vivo. *J Biomed Mater Res* 1995;**29**:1267–1275.
 120. Anthony RM, Rutitzky LI, Urban JF, Stadecker MJ, Gause WC. Protective immune mechanisms in helminth infection. *Nat Rev Immunol* 2007;**7**:975–987.
 121. Henson PM. The immunologic release of constituents from neutrophil leukocytes. I. The role of antibody and complement on nonphagocytosable surfaces or phagocytosable particles. *J Immunol* 1971;**107**:1535–1546.
 122. Zhao Q, Topham N, Anderson JM, Hiltner A, Lodoen G, Payet CR. Foreign-body giant cells and polyurethane biostability: in vivo correlation of cell adhesion and surface cracking. *J Biomed Mater Res* 1991;**25**:177–183.

123. Morgan BP, Gasque P. Extrahepatic complement biosynthesis: where, when and why? *Clin Exp Immunol* 1997;**107**:1–7.
124. Sahoo S, et al. Glyceraldehyde-3-phosphate dehydrogenase of the parasitic nematode *Haemonchus contortus* binds to complement C3 and inhibits its activity. *Parasite Immunol* 2013;**35**:457–467.
125. Giacomini PR, et al. The role of complement in innate, adaptive and eosinophil-dependent immunity to the nematode *Nippostrongylus brasiliensis*. *Mol Immunol* 2008;**45**:446–455.
126. Daffern PJ, Pfeifer PH, Ember JA, Hugli TE. C3a is a chemotaxin for human eosinophils but not for neutrophils. I. C3a stimulation of neutrophils is secondary to eosinophil activation. *J Exp Med* 1995;**181**:2119–2127.
127. Dai S, Rajaram MVS, Curry HM, Leander R, Schlesinger LS. Fine tuning inflammation at the front door: macrophage complement receptor 3—mediates phagocytosis and immune suppression for *Francisella tularensis*. *PLoS Pathog* 2013;**9**:e1003114.
128. Benoit ME, Clarke EV, Morgado P, Fraser DA, Tenner AJ. Complement protein C1q directs macrophage polarization and limits inflammasome activity during the uptake of apoptotic cells. *J Immunol* 2012;**188**:5682–5693.
129. Odegaard JI, et al. Macrophage-specific PPARgamma controls alternative activation and improves insulin resistance. *Nature* 2007;**447**:1116–1120.
130. Odegaard JI, Chawla A. Alternative macrophage activation and metabolism. *Annu Rev Pathol Mech Dis* 2011;**6**:275–297.
131. Hood MI, Skaar EP. Nutritional immunity: transition metals at the pathogen–host interface. *Nat Rev Microbiol* 2012;**10**:525–537.
132. Knubel CP, et al. Indoleamine 2,3-dioxygenase (IDO) is critical for host resistance against *Trypanosoma cruzi*. *FASEB J* 2010;**24**:2689–2701.
133. Sindrilaru A, Scharffetter-Kochanek K. Disclosure of the culprits: macrophages—versatile regulators of wound healing. *Adv Wound Care (New Rochelle)* 2013;**2**:357–368.
134. Lucas T, et al. Differential roles of macrophages in diverse phases of skin repair. *J Immunol* 2010;**184**:3964–3977.
135. Mantovani A, Biswas SK, Galdiero MR, Sica A, Locati M. Macrophage plasticity and polarization in tissue repair and remodelling. *J Pathol* 2013;**229**:176–185.
136. Maarsingh H, Zaagsma J, Meurs H. Arginase: a key enzyme in the pathophysiology of allergic asthma opening novel therapeutic perspectives. *Br J Pharmacol* 2009;**158**:652–664.
137. Eming SA, Krieg T, Davidson JM. Inflammation in wound repair: molecular and cellular mechanisms. *J Invest Dermatol* 2007;**127**:514–525.
138. Daley JM, Brancato SK, Thomay AA, Reichner JS, Albina JE. The phenotype of murine wound macrophages. *J Leukoc Biol* 2009;**87**:59–67.
139. Teng X, Li D, Champion HC, Johns RA. FIZZ1/REL α , a novel hypoxia-induced mitogenic factor in lung with vasoconstrictive and angiogenic properties. *Circ Res* 2003;**92**:1065–1067.
140. Li X, Yang Y, Fang J, Zhang H. FIZZ1 could enhance the angiogenic ability of rat aortic endothelial cells. *Int J Clin Exp Pathol* 2013;**6**:1847–1853.
141. Li D, Fernandez LG, Dodd-o J, Langer J, Wang D, Laubach VE. Upregulation of hypoxia-induced mitogenic factor in compensatory lung growth after pneumonectomy. *Am J Respir Cell Mol Biol* 2005;**32**:185–191.
142. Zhou Y, et al. Chitinase 3-like 1 suppresses injury and promotes fibroproliferative responses in mammalian lung fibrosis. *Sci Transl Med* 2014;**6**:240ra76.
143. Wynes MW, Riches DWH. Induction of macrophage insulin-like growth factor-I expression by the Th2 cytokines IL-4 and IL-13. *J Immunol* 2003;**171**:3550–3559.
144. Rom WN, Basset P, Fells GA, Nukiwa T, Trapnell BC, Cysal RG. Alveolar macrophages release an insulin-like growth factor I-type molecule. *J Clin Invest* 1988;**82**:1685–1693.
145. Toulon A, et al. A role for human skin-resident T cells in wound healing. *J Exp Med* 2009;**206**:743–750.
146. Chen F, et al. An essential role for TH2-type responses in limiting acute tissue damage during experimental helminth infection. *Nat Med* 2012;**18**:260–266.
147. Bayat A, McGrouther DA, Ferguson MWJ. Skin scarring. *BMJ* 2003;**326**:88–92.
148. Röcken M, Racke M, Shevach EM. IL-4-induced immune deviation as antigen-specific therapy for inflammatory autoimmune disease. *Immunol Today* 1996;**17**:225–231.
149. Goerdts S, Orfanos CE. Other functions, other genes: alternative activation of antigen-presenting cells. *Immunity* 1999;**10**:137–142.
150. Loke P, MacDonald AS, Allen JE. Antigen-presenting cells recruited by *Brugia malayi* induce Th2 differentiation of naïve CD4 (+) T cells. *Eur J Immunol* 2000;**30**:1127–1135.
151. Rodríguez-Sosa M, et al. Chronic helminth infection induces alternatively activated macrophages expressing high levels of CCR5 with low interleukin-12 production and Th2-biasing ability. *Infect Immun* 2002;**70**:3656–3664.
152. Fenton MJ, Buras JA, Donnelly RP. IL-4 reciprocally regulates IL-1 and IL-1 receptor antagonist expression in human monocytes. *J Immunol* 1992;**149**:1283–1288.
153. Mantovani A, Sica A, Sozzani S, Allavena P, Vecchi A, Locati M. The chemokine system in diverse forms of macrophage activation and polarization. *Trends Immunol* 2004;**25**:677–686.
154. Li MO, Wan YY, Sanjabi S, Robertson A-KL, Flavell RA. Transforming growth factor- β regulation of immune responses. *Annu Rev Immunol* 2006;**24**:99–146.
155. Gronert K, Maheshwari N, Khan N, Hassan IR, Dunn M, Laniado Schwartzman M. A role for the mouse 12/15-lipoxygenase pathway in promoting epithelial wound healing and host defense. *J Biol Chem* 2005;**280**:15267–15278.
156. Kühn H, O'Donnell VB. Inflammation and immune regulation by 12/15-lipoxygenases. *Prog Lipid Res* 2006;**45**:334–356.
157. Hung S-I, Chang AC, Kato I, Chang N-CA. Transient expression of Ym1, a heparin-binding lectin, during developmental hematopoiesis and inflammation. *J Leukoc Biol* 2002;**72**:72–82.
158. Haben I, et al. T-cell-derived but not B-cell-derived IL-10 suppresses antigen-specific T-cell responses in *Litomosoides sigmodontis*-infected mice. *Eur J Immunol* 2013;**43**:1799–1805.
159. Zigmund E, et al. Macrophage-restricted interleukin-10 receptor deficiency, but not IL-10 deficiency, causes severe spontaneous colitis. *Immunity* 2014;**40**:720–733.
160. Mosser DM, Edwards JP. Exploring the full spectrum of macrophage activation. *Nat Rev Immunol* 2008;**8**:958–969.
161. Shirey KA, et al. Control of RSV-induced lung injury by alternatively activated macrophages is IL-4R α -, TLR4-, and IFN- β -dependent. *Mucosal Immunol* 2010;**3**:291–300.
162. Cohen HB, Mosser DM. Extrinsic and intrinsic control of macrophage inflammatory responses. *J Leukoc Biol* 2013;**94**:913–919.
163. Williams M, Bruhns P, Saeys Y, Hammad H, Lambrecht BN. The function of Fc γ receptors in dendritic cells and macrophages. *Nat Rev Immunol* 2014;**14**:94–108.
164. MacDonald AS, Maizels RM, Lawrence RA, Dransfield I, Allen JE. Requirement for in vivo production of IL-4, but not IL-10, in the induction of proliferative suppression by filarial parasites. *J Immunol* 1998;**160**:4124–4132.
165. Taylor MD, Harris A, Nair MG, Maizels RM, Allen JE. F4/80+ alternatively activated macrophages control CD4+ T cell hyporesponsiveness at sites peripheral to filarial infection. *J Immunol* 2006;**176**:6918–6927.
166. Smith P, et al. *Schistosoma mansoni* worms induce energy of T cells via selective up-regulation of programmed death ligand 1 on macrophages. *J Immunol* 2004;**173**:1240–1248.
167. Terrazas LI, Montero D, Terrazas CA, Reyes JL, Rodríguez-Sosa M. Role of the programmed Death-1 pathway in the suppressive activity of alternatively activated macrophages in experimental cysticercosis. *Int J Parasitol* 2005;**35**:1349–1358.
168. Brys L, et al. Reactive oxygen species and 12/15-lipoxygenase contribute to the antiproliferative capacity of alternatively activated myeloid cells elicited during helminth infection. *J Immunol* 2005;**174**:6095–6104.
169. Michalek RD, Rathmell JC. The metabolic life and times of a T-cell. *Immunol Rev* 2010;**236**:190–202.
170. Vander Heiden MG, Cantley LC, Thompson CB. Understanding the Warburg effect: the metabolic requirements of cell proliferation. *Science* 2009;**324**:1029–1033.
171. van der Windt GJW, et al. Mitochondrial respiratory capacity is a critical regulator of CD8+ T cell memory development. *Immunity* 2012;**36**:68–78.

172. Pond CM, Mattacks CA. The source of fatty acids incorporated into proliferating lymphoid cells in immune-stimulated lymph nodes. *Br J Nutr* 2003;**89**:375–383.
173. Macqueen HA, Waights V, Pond CM. Vascularisation in adipose depots surrounding immune-stimulated lymph nodes. *J Anat* 1999;**194**(Pt 1):33–38.
174. Birsoy K, et al. Metabolic determinants of cancer cell sensitivity to glucose limitation and biguanides. *Nature* 2014;**508**:108–112.
175. Zheng Y, Delgoffe GM, Meyer CF, Chan W, Powell JD. Anergic T cells are metabolically anergic. *J Immunol* 2009;**183**:6095–6101.
176. Voehringer D, Van Rooijen N, Locksley RM. Eosinophils develop in distinct stages and are recruited to peripheral sites by alternatively activated macrophages. *J Leukoc Biol* 2007;**81**:1434–1444.
177. Díaz A, Allen JE. Mapping immune response profiles: the emerging scenario from helminth immunology. *Eur J Immunol* 2007;**37**:3319–3326.
178. Ishii M, et al. Epigenetic regulation of the alternatively activated macrophage phenotype. *Blood* 2009;**114**:3244–3254.
179. Stout RD, Jiang C, Matta B, Tietzel I, Watkins SK, Suttles J. Macrophages sequentially change their functional phenotype in response to changes in microenvironmental influences. *J Immunol* 2005;**175**:342–349.
180. Whyte CS, et al. Suppressor of cytokine signaling (SOCS)1 is a key determinant of differential macrophage activation and function. *J Leukoc Biol* 2011;**90**:845–854.
181. Dambuza I, et al. Efficacy of membrane TNF mediated host resistance is dependent on mycobacterial virulence. *Tuberculosis (Edinb)* 2008;**88**:221–234.
182. Uderhardt S, et al. 12/15-lipoxygenase orchestrates the clearance of apoptotic cells and maintains immunologic tolerance. *Immunity* 2012;**36**:834–846.
183. Satoh T, et al. The Jmjd3-Irf4 axis regulates M2 macrophage polarization and host responses against helminth infection. *Nat Immunol* 2010;**11**:936–944.
184. Okabe Y, Medzhitov R. Tissue-specific signals control reversible program of localization and functional polarization of macrophages. *Cell* 2014;**157**:832–844.
185. Rosas M, et al. The transcription factor Gata6 links tissue macrophage phenotype and proliferative renewal. *Science* 2014;**344**:645–648.
186. Chow A, Brown BD, Merad M. Studying the mononuclear phagocyte system in the molecular age. *Nat Rev Immunol* 2011;**11**:788–798.
187. Chattopadhyay PK, Gierahn TM, Roederer M, Love JC. Single-cell technologies for monitoring immune systems. *Nat Immunol* 2014;**15**:128–135.