



# THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### Mechanisms for CO<sub>2</sub> Leakage Prevention – A Global Dataset of Natural Analogues

**Citation for published version:**

McDermott, C 2013, 'Mechanisms for CO<sub>2</sub> Leakage Prevention – A Global Dataset of Natural Analogues' Energy Procedia, vol 40, pp. 320-328.

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Publisher final version (usually the publisher pdf)

**Published In:**

Energy Procedia

**Publisher Rights Statement:**

Open Access

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



European Geosciences Union General Assembly 2013, EGU

Division Energy, Resources & the Environment, ERE

## Mechanisms for CO<sub>2</sub> leakage prevention – a global dataset of natural analogues

Johannes M. Miocic <sup>a\*</sup>, Stuart M. V. Gilfillan <sup>a</sup>, Christopher McDermott <sup>a</sup>, R. Stuart Haszeldine <sup>a</sup>

<sup>a</sup>*School of Geosciences, University of Edinburgh, King's Buildings, West Mains Road, Edinburgh EH9 3JW, United Kingdom*

---

### Abstract

Natural CO<sub>2</sub> reservoirs have similar geological trapping mechanisms as required for CO<sub>2</sub> storage sites and often have held CO<sub>2</sub> for a geological period of time without any indication of leakage. Yet, migration of CO<sub>2</sub> from reservoirs to the surface is also common. 49 natural CO<sub>2</sub> reservoirs have been analysed to provide an overview of factors that are important for (1) retention of CO<sub>2</sub> in the subsurface and (2) leakage of CO<sub>2</sub> from the reservoir. Results indicate that overpressure of the overburden and the state of CO<sub>2</sub> in the reservoir influence the likelihood of migration and hence the performance of reservoirs.

© 2013 The Authors. Published by Elsevier Ltd.

Selection and peer-review under responsibility of the GFZ German Research Centre for Geosciences

*Keywords:* CCS; CO<sub>2</sub> Storage; Leakage; Natural Analogues; Overpressure

---

### 1. Introduction

Carbon Capture and Storage (CCS) is the only industrial scale technology currently planned to directly reduce CO<sub>2</sub> emissions from fossil fuelled power plants and large industrial point sources to the atmosphere [1]. CO<sub>2</sub> is captured at the source and transported to subsurface storage sites, such as depleted oil and gas fields or saline aquifers [2]. In order to have a reduction of emissions it is crucial that the amount of CO<sub>2</sub> leaking from storage sites to shallow aquifers or the surface remains very low (<0.1% per year, over 10.000 years) [3].

Therefore the long-term behaviour of CO<sub>2</sub> in the subsurface, including possible CO<sub>2</sub> migration pathways and CO<sub>2</sub>-brine-rock interactions, needs to be critically assessed for each storage site. This is ideally done with several methods: (1) geometric, structural and geochemical appraisal of the storage site; (2) planned monitoring strategies; (3) laboratory experiments on both reservoir and cap rocks; (4) geochemical and coupled modelling of CO<sub>2</sub> behaviour and (5) the study of natural CO<sub>2</sub> fields as analogues for storage sites [4]. Laboratory experiments help to understand how CO<sub>2</sub> may modify the reservoir and cap rocks during the first months to years after injection but fail to give insights into the long-term behaviour (100's to 1000's of years). Modelling approaches often use parameters obtained from laboratory experiments and/or simplify the complex subsurface. Additionally, the up-scaling of pore-scale processes to reservoir size needs significant computing power and new softwares. Natural CO<sub>2</sub> reservoirs have the advantage that CO<sub>2</sub> has

---

\* Corresponding author. Tel.: +44-131-650-5916; fax: +44-131-668-3184.  
E-mail address: [johannes.miocic@ed.ac.uk](mailto:johannes.miocic@ed.ac.uk).

interacted with the reservoir and cap rocks for a long period of time (up to 10s of millions of years). Leaking reservoirs offer the opportunity to study the mechanisms that lead to the migration of CO<sub>2</sub>. However, natural reservoirs are complex systems and detailed information on the subsurface is often rare. Here we present the results of a global study of natural CO<sub>2</sub> reservoirs with a focus on identifying possible leakage mechanisms.

## 2. Natural CO<sub>2</sub> reservoirs

Natural CO<sub>2</sub> reservoirs are widespread in sedimentary basins world-wide and have geological elements similar to hydrocarbon reservoirs. The CO<sub>2</sub> can originate from a number of sources such as mantle degassing, carbonate rock metamorphism or the degradation of organic matter [5]. The reservoirs are sometimes encountered during hydrocarbon exploration activities and regularly abandoned as they are not as profitable as hydrocarbon accumulations. However, because of the demand of CO<sub>2</sub> for enhanced oil recovery, several reservoirs are commercially exploited (e.g. on the Colorado Plateau, US).

Many natural CO<sub>2</sub> fields have been studied, often with focus on the origin of the CO<sub>2</sub> in order to avoid hydrocarbon exploration in areas where such reservoirs could occur [e.g. 6] and naturally do not focus on trapping mechanisms and possible leakage pathways. Other authors have focused on the impact the long-term residence of CO<sub>2</sub> has had on the mineralogy and geochemistry of reservoirs [e.g. 7, 8]. Detailed geological information including production data is only available for few reservoirs [e.g. 9]. Reviews and comparisons of natural CO<sub>2</sub> reservoirs exist on a regional scale, usually as analogue studies for carbon storage sites [10, 11]. Roberts [12] has completed a comparison of leaking and non-leaking reservoirs in Italy, but to date no global comparison of leaking and non-leaking reservoirs exists.

## 3. Creating a global dataset

We compiled a global dataset of 49 well described natural CO<sub>2</sub> reservoirs (Fig. 1; Appendix A). Well logs were analysed for additional information such as pressure gradients and temperature gradients. The quality of reports differs and only literature that had a certain scientific standard (peer-reviewed or well documented methodology and sources) was included. If in doubt about the reliability of a source, only reservoirs with several independent sources were incorporated.




Fig. 1. Locations and CO<sub>2</sub> content of natural CO<sub>2</sub> reservoirs included in this study.

The goal was to identify possible leakage mechanisms, consequently it was crucial to correctly differentiate between leaking and non-leaking reservoirs. The following criteria were used to identify leaking reservoirs:

- CO<sub>2</sub> showing at the surface spatially close to the reservoir. This includes CO<sub>2</sub> rich springs and diffusive degassing and indicates a present day leakage.

- Formation of carbonate rocks (travertines) at the surface spatially close to the reservoir. Even if there is no current precipitation, carbonates may indicate historical leakage.
- Gas chimneys identified on seismic data.
- Occurrence of CO<sub>2</sub> in an aquifer above the reservoir in lower concentration than in the actual reservoir.

While the first two points can be readily analysed by studying the relevant literature and maps, the latter two points can be harder identify. In addition, there is also a small chance that CO<sub>2</sub> is leaking from a reservoir into a shallower aquifer without being detected.

The dataset includes depth, temperature, pressure and CO<sub>2</sub> content for all reservoirs. For reservoirs for which in situ pressure and temperature data was not available missing pressure data has been calculated assuming a hydrostatic pressure gradient of 9.8 kPa/m. Missing temperature data was calculated using measured regional temperature gradients [13]. CO<sub>2</sub> state and density for each reservoir was calculated using the equation of state from Huang et al. [14]. Possible leakage pathways like faults were identified on structural cross-sections where available. Vertical pressure profiles for several analogues were created by using well- and mudlog data.

## 4. Results

### 4.1. Pressure controls on leakage

Natural CO<sub>2</sub> reservoirs follow in general “normal” depth-pressure trends, i.e. reservoir pressure is between hydrostatic and lithostatic pressure. Shallow reservoirs (<1200 m) are often underpressured with regards to hydrostatic pressure while deep reservoirs (>2000 m) tend to be overpressured. Leaking reservoirs are either shallow or near or above a fracture gradient for reservoir sandstone (Fig. 2).




Fig. 2. Depth-pressure plot of natural CO<sub>2</sub> reservoirs. Leaking reservoirs are either shallow or near/above a fracture gradient for reservoir rocks. Note that only reservoirs with in-situ pressure data are plotted. Main gas other than CO<sub>2</sub> for most reservoirs is CH<sub>4</sub>.

Vertical pressure profiles through several natural CO<sub>2</sub> reservoirs have been combined into a more general vertical profile (Fig. 3). Reservoirs that are underpressured with regards to the overburden are less likely to leak than reservoirs that are overpressured with regards to the overburden.




Fig. 3. Correlation between leakage from natural CO<sub>2</sub> reservoirs and vertical pressure profiles through the reservoirs and overburden (after [12]). Blue line represents a non-leaking natural analogue for which the pressure in the reservoir is lower than the pressure in the overburden. For leaking reservoirs the pressure in the overburden is lower than the pressure in the reservoir (red line). Thus leakage is less likely if the overburden is overpressured with regard to the reservoir.

#### 4.2. CO<sub>2</sub> state

The CO<sub>2</sub> state for all reservoirs has been calculated based on pressure and temperature data (fig. 4). Results show that reservoirs with gaseous CO<sub>2</sub> have a high likelihood of leakage (6 out of 9, 66%). Reservoirs with supercritical CO<sub>2</sub> have only a moderate to low chance of leaking (4 out of 40, 10%).




Fig. 4. CO<sub>2</sub> state diagram showing the temperatures and pressures in the natural CO<sub>2</sub> reservoirs. Blue lines (solid and dotted) indicate phase boundaries. Dashed lines are lines of equal density.

## 5. Discussion

### 5.1. Pressure controls on leakage

Vertical pressure profiles through the overburden and the reservoir are available for several fields. Comparison of those profiles from leaking and non-leaking reservoirs showed that reservoirs which are underpressured in regards to the overburden are less likely to leak CO<sub>2</sub> than reservoirs which are overpressured in regards to the overburden.

Overpressured reservoirs near or above fracture gradients have an increased probability of leakage as fractures in the reservoir rocks can propagate into the overburden and create fluid pathways through the sealing rock [15]. Additionally, elevated pore pressures due to the presence of natural gases such as CO<sub>2</sub> can lead to the reactivation of existing faults [16]. The depth dependency of leakage reflects the change of CO<sub>2</sub> state with rising pressures and temperatures (see section 4.2).

A positive pressure gradient from the reservoir to the overburden hampers the flow of fluids into the overburden and thus vertical leakage of CO<sub>2</sub>. This is illustrated by the fact that leakage seems less likely if the overburden is overpressured with regards to the reservoir. Other authors have suggested that this could be used to artificially overpressure the overburden to prevent CO<sub>2</sub> leakage [17].

### 5.2. CO<sub>2</sub> state controls on leakage

Gaseous CO<sub>2</sub> reservoirs are in shallow depths (generally <1000 m, depending on the local temperature and pressure gradients) while dense state CO<sub>2</sub> occurs in reservoirs with a depth greater than 1000 m. The fact that reservoirs with gaseous CO<sub>2</sub> are more likely to leak than reservoirs with dense state CO<sub>2</sub> can be partially attributed to the higher buoyancy of gaseous CO<sub>2</sub> compared to the buoyancy of supercritical CO<sub>2</sub> and thus higher stress on the overburden. Furthermore, recent laboratory experiments indicate that the flow of dense CO<sub>2</sub> through fractures in mudrock seals is impeded compared to the flow of gaseous CO<sub>2</sub> [18].

## 6. Conclusions

The analysis of a global dataset of 49 natural CO<sub>2</sub> reservoirs of which 10 are known to leak has helped to identify mechanisms that promote leakage of CO<sub>2</sub> from reservoirs: (1) shallow depth, (2) CO<sub>2</sub> in gas phase and (3) hydrostatic overburden pressure. Based on the results of this study sites for engineered containment of CO<sub>2</sub> are best where CO<sub>2</sub> is in the dense state, overburden is overpressured and the reservoir pressure is less than 50% of lithostatic pressure.

## Acknowledgements

This research has partly been funded by SCCS (Scottish Carbon Capture and Storage) and the EU FP7 research programme PANACEA.

## References

- [1] Scott V, Gilfillan S, Markusson N, Chalmers H, Haszeldine RS. Last chance for carbon capture and storage. *Nature Clim Change* 2013;3:105-11.
- [2] Metz B, Davidson O, de Coninck HC, Loose M, Meyer LA (Eds.). *IPCC Special report on Carbon Dioxide Capture and Storage*. Cambridge University Press, New York, USA, Cambridge, UK; 2005, pp. 431.
- [3] Song J, Zhang D. Comprehensive Review of Caprock-Sealing Mechanisms for Geologic Carbon Sequestration. *Env Sci & Tech* 2012;47:9-22.
- [4] Gaus I. Role and impact of CO<sub>2</sub>-rock interactions during CO<sub>2</sub> storage in sedimentary rocks. *Int J Greenhouse Gas Control* 2010;4:73-89.
- [5] Wycherley H, Fleet A, Shaw H. Some observations on the origins of large volumes of carbon dioxide accumulations in sedimentary basins. *Mar Pet Geol* 1999;16:489-94.
- [6] Zhang T, Zhang M, Bai B, Wang X, Li L. Origin and accumulation of carbon dioxide in the Huanghua depression, Bohai Bay Basin, China. *AAPG Bull* 2008;92:341-58.

- [7] Wilkinson M, Haszeldine RS, Fallick AE, Odling N, Stoker SJ, Gatliff RW. CO<sub>2</sub>-Mineral Reaction in a Natural Analogue for CO<sub>2</sub> Storage—Implications for Modeling. *J Sed Res* 2009;79:486-94.
- [8] Moore J, Adams M, Allis R, Lutz S, Rauzi S. Mineralogical and geochemical consequences of the long-term presence of CO<sub>2</sub> in natural reservoirs: An example from the Springerville-St. Johns Field, Arizona, and New Mexico, USA. *Chem Geol* 2005;217:365-85.
- [9] Becker TP, Lynds R. A geologic deconstruction of one of the world's largest natural accumulations of CO<sub>2</sub>, Moxa arch, southwest Wyoming. *AAPG Bull* 2012;96:1643-64.
- [10] Stevens SH, Pearce JM, Rigg AAJ. *Natural Analogs for Geologic Storage of CO<sub>2</sub>: An Integrated Global Research Program*. in: First National Conference on Carbon Sequestration U.S. Department of Energy, National Energy Technology Laboratory, Washington D.C.; 2001, pp. 12.
- [11] Pearce J, Czernichowski-Lauriol I, Lombardi S, Brune S, Nador A, Baker J, Pauwels H, Hatziyannis G, Beaubien S, Faber E. A review of natural CO<sub>2</sub> accumulations in Europe as analogues for geological sequestration. *Spec Pub Geol Soc London* 2004;233:29-41.
- [12] Roberts JJ. *Natural CO<sub>2</sub> fluids in Italy: Implications for the leakage of geologically stored CO<sub>2</sub>*. PhD, University of Edinburgh. 2013. pp. 229.
- [13] Yuan Y-S, Ma Y-S, Hu S-B, Guo T-L, Fu X-Y. Present day geothermal characteristics in South China. *Chin J Geophy* 2006;49:1005-14.
- [14] Huang F-H, Li M-H, Lee LL, Starling KE. An accurate equation of state for Carbon Dioxide. *J Chem Eng Japan* 1985;18:490-6.
- [15] Hawkes CD, Bachu S, McLellan PJ. Geomechanical factors affecting geological storage of CO<sub>2</sub> in depleted oil and gas reservoirs. *J Can Petrol Tech* 2005;44:52-61.
- [16] Wiprut D, Zoback MD. Fault reactivation and fluid flow along a previously dormant normal fault in the northern North Sea. *Geology* 2000;28:595-8.
- [17] Réveillère A, Rohmer J. Managing the risk of CO<sub>2</sub> leakage from deep saline aquifer reservoirs through the creation of a hydraulic barrier. *Energy Procedia* 2011;4:3187-94.
- [18] Edlmann K, Haszeldine RS, McDermott CI. Experimental investigation into the sealing capability of naturally fractured shale caprocks to supercritical carbon dioxide flow. *Env Earth Sci* 2013.
- [19] Stuart CA, Kozik HG. Geopressuring Mechanism of Smackover Gas Reservoirs, Jackson Dome Area, Mississippi. *J Petrol Tech* 1977;29:579-85.
- [20] Gilfillan SMV, Wilkinson M, Haszeldine RS, Shipton ZK, Nelson ST, Poreda RJ. He and Ne as tracers of natural CO<sub>2</sub> migration up a fault from a deep reservoir. *Int J Greenhouse Gas Control* 2011;5:1507-16.
- [21] Rauzi S. *Carbon Dioxide in the St. Johns-Springerville Area, Apache County, Arizona*. Arizona Geol Surv Open-File Report 99-2; 1999.
- [22] Kaszuba JP, Navarre-Sitchler A, Thyne G, Chopping C, Meuzelaar T. Supercritical carbon dioxide and sulfur in the Madison Limestone: A natural analog in southwest Wyoming for geologic carbon-sulfur co-sequestration. *Earth Planet Sci Lett* 2011;309:131-40.
- [23] Allis RG, Chidsey T, Gwynn W, Morgan C, White SP, Adams M, Moore J. *Natural CO<sub>2</sub> reservoirs on the Colorado Plateau and southern Rocky Mountains: Candidates for CO<sub>2</sub> sequestration*. AAPG meeting, Denver, Colorado; 2001.
- [24] Holloway S, Pearce JM, Hards VL, Ohsumi T, Gale J. Natural emissions of CO<sub>2</sub> from the geosphere and their bearing on the geological storage of carbon dioxide. *Energy* 2007;32:1194-201.
- [25] Lynch RD, McBride EJ, Perkins TK, Wiley ME. Dynamic Kill of an Uncontrolled CO<sub>2</sub> Well. *J Petrol Tech* 1985;37:1267-75.
- [26] Renfro JJ. Sheep Mountain CO<sub>2</sub> Production Facilities - A conceptual design. *J Petrol Tech* 1979;31:1462-8.
- [27] Allis R, Bergfeld D, Moore J, McClure K, Morgan C, Chidsey T, Heath J, McPherson B. Implications of results from CO<sub>2</sub> flux surveys over known CO<sub>2</sub> systems for long-term monitoring. in: *Conf Proc 4th Ann Conf CCS*; 2005.
- [28] Morgan CD, Chidsey T, Gordon Creek, Farnham Dome, and Woodside fields, Carbon and Emery Counties, Utah, in: Chidsey T (Ed.) *Geology of east-central Utah*, Utah Geol Assoc Pub, 1991, pp. 301-9.
- [29] Chidsey T, Chamberlain L, Gordon Creek, in: Hill BG, Bereskin SR (Eds.) *Oil and gas fields of Utah*, Utah Geol Assoc 1996.
- [30] Gerling CR, McElmo Dome Leadville carbon dioxide field, Colorado, in: Fasset JE (Ed.) *Oil and Gas Fields of the Four Corners area*, Four Corners Geol Soc, 1983, pp. 735-9.
- [31] Pearce JM, Holloway S, Wacker H, Nelis MK, Rochelle C, Bateman K. Natural occurrences as analogues for the geological disposal of carbon dioxide. *Energy Conv Manag* 1996;37:1123-8.
- [32] Broadhead RF. Carbon dioxide in Union and Harding Counties, in: *New Mexico Geol Soc Guidebook, 38th Field Conference, Northeastern New Mexico*, 1987, pp. 339-49.
- [33] Dubacq B, Bickle MJ, Wigley M, Kampman N, Ballentine CJ, Sherwood Lollar B. Noble gas and carbon isotopic evidence for CO<sub>2</sub>-driven silicate dissolution in a recent natural CO<sub>2</sub> field. *Earth Planet Sci Lett* 2012;341-344:10-9.
- [34] Ballentine CJ, Schoell M, Coleman D, Cain BA. 300-Myr-old magmatic CO<sub>2</sub> in natural gas reservoirs of the west Texas Permian basin. *Nature* 2001;409:327-31.
- [35] Gilfillan SMV, Lollar BS, Holland G, Blagburn D, Stevens S, Schoell M, Cassidy M, Ding Z, Zhou Z, Lacrampe-Couloume G, Ballentine CJ. Solubility trapping in formation water as dominant CO<sub>2</sub> sink in natural gas fields. *Nature* 2009;458:614-8.
- [36] Blann JR, Laville GM. Gas Lifting a Major Oil Field in Argentina with High CO<sub>2</sub> content associated gas. *SPE Prod Facil* 1997;12:41-5
- [37] Gachuz H, Berumen S, Alcazar LO, Ruodriguez JA. Quebrache, a Natural CO<sub>2</sub> Reservoir: A new source for EOR projects in Mexico. *Soc Petrol Engi*; 2007.

- [38] Gachuz H, Sanchez-Bujanos JL, Castro-Herrera I, Rodriguez-Pimentel JA. Quebrache Field: Evaluation to Date of the Natural CO<sub>2</sub> reservoir. in: SPE EUROPEC/EAGE Ann Conf, Soc Petrol Engi, Vienna, Austria; 2011.
- [39] Gaus I, Le Guern C, Pearce J, Pauwels H, Shepherd T, Hatziyannis G, Metaxas A. Comparison of long-term geochemical interactions at two natural CO<sub>2</sub>-analogues. Montmiral (Southeast basin, France) and Messokampos (Florina basin, Greece) case studies. in: Rubin ES, Keith DW, Gilboay CF (Eds.) *Conf Proc 7th Int Conf Greenhouse Gas Control Techn*, IEA Greenhouse Gas Programm, Cheltenham, UK; 2004, pp. 561-9.
- [40] Underhill JR, Lykakis N, Shafique S. Turning exploration risk into a carbon storage opportunity in the UK Southern North Sea. *Petrol Geosci* 2009;15:291-304.
- [41] Yielding G, Lykakis N, Underhill JR. The role of stratigraphic juxtaposition for seal integrity in proven CO<sub>2</sub> fault-bound traps of the Southern North Sea. *Petrol Geosci* 2011;17:193-203.
- [42] Schütze C, Sauer U, Beyer K, Lamert H, Bräuer K, Strauch G, Flechsig C, Kämpf H, Dietrich P. Natural analogues: a potential approach for developing reliable monitoring methods to understand subsurface CO<sub>2</sub> migration processes. *Envi Earth Sci* 2012;67:411-23.
- [43] Annunziatellis A, Beaubien SE, Bigi S, Ciotoli G, Coltella M, Lombardi S. Gas migration along fault systems and through the vadose zone in the Latera caldera (central Italy): Implications for CO<sub>2</sub> geological storage. *Int J Greenhouse Gas Control* 2008;2:353-72.
- [44] Fischer M, Botz R, Schmidt M, Rockenbauch K, Garbe-Schönberg D, Glodny J, Gerling P, Littke R. Origins of CO<sub>2</sub> in permian carbonate reservoir rocks (Zechstein, Ca<sub>2</sub>) of the NW-German Basin (Lower Saxony). *Chem Geol* 2006;227:184-213.
- [45] Doleschall S, Szittar A, Udvardi G. Review of the 30 years' experience of the CO<sub>2</sub> imported Oil Recovery Projects in Hungary. International Meeting on Petroleum Engineering, Soc Petrol Engi, Beijing; 1992.
- [46] Dai J, Song Y, Dai C. *Conditions governing the formation of Abiogenic Gas and Gas Pools in Eastern China*. Beijing: Science Press; 2000.
- [47] Anping H, Jinxing D, Chun Y, Qinghua Z, Yunyan N. Geochemical characteristics and distribution of CO<sub>2</sub> gas fields in Bohai Bay Basin. *Pet Explor Dev* 2009;36:181-9.
- [48] Dai J, Yang S, Chen H, Shen X. Geochemistry and occurrence of inorganic gas accumulations in Chinese sedimentary basins. *Org Geochem* 2005;36:1664-88.
- [49] Gong Y, Wang L, Liu S, Guo L, Cai J. Distribution Characteristics of the Geotemperature field in the Jiyan Depression, Shandong Province, North China *Chin J Geophy* 2003;46:993-42.
- [50] Li M, Wang T, Liu J, Lu H, Wu W, Gao L. Occurrence and origin of carbon dioxide in the Fushan Depression, Beibuwan Basin, South China Sea. *Mar Pet Geol* 2008;25:500-13.
- [51] Guang Y, Zhanyin Z, Mingli S. Formation of carbon dioxide and hydrocarbon gas reservoirs in the Changling fault depression, Songliao Basin. *Pet Expl Dev* 2011;38:52-8.
- [52] Huang B, Xiao X, Li X. Geochemistry and origins of natural gases in the Yinggehai and Qiongdongnan basins, offshore South China Sea. *Org Geochem* 2003;34:1009-25.
- [53] Bell Jr. RJ, Davis JM. Lost Circulation Challenges: Drilling Thick Carbonate Gas Reservoir, Natuna D-Alpha Block. in: SPE/IADC Drilling Conference, Soc Petrol Engin, New Orleans, Louisiana; 1987, pp. 947-54.
- [54] Watson MN, Zwingmann N, Lemon NM. The Ladbroke Grove–Katnook carbon dioxide natural laboratory: A recent CO<sub>2</sub> accumulation in a lithic sandstone reservoir. *Energy* 2004;29:1457-66.
- [55] Parker KA. The exploration and appraisal history of the Katnook and Ladbroke Grove gas fields, onshore Otway Basin, South Australia *APPEA* 1992;32:67-85.
- [56] Hortle A, deWijkerslooth C, Tenthorey E, Strand J, Giger S. Understanding the tuna field: An integrated approach to fault seal properties, natural CO<sub>2</sub> content & hydrodynamic analysis. *Energy Procedia* 2011;4:4732-8.
- [57] Hulston JR, Hilton DR, Kaplan IR. Helium and carbon isotope systematics of natural gases from Taranaki Basin, New Zealand. *Appl Geochem* 2001;16:419-36.
- [58] Webster M, O'Connor S, Pindar B, Swarbrick R. Overpressures in the Taranaki Basin: Distribution, causes, and implications for exploration. *AAPG Bull* 2011;95:339-70.
- [59] Lyon GL, Giggenbach WF, Sano Y. Variations in the chemical and isotopic composition of Taranaki gases and their possible causes. in: Proc NZ Petrol Conf 1996, pp. 171-8.

## Appendix A. List of fields and literature used in this study

| No. | Name of field  | Leakage | Source                 |
|-----|----------------|---------|------------------------|
| 1   | Jackson Dome   | No      | [10], [19]             |
| 2   | St. Johns Dome | Yes     | [8], [20], [21]        |
| 3   | Moxa Arch      | No      | [9], [22], [23]        |
| 4   | Sheep Mountain | No      | [23], [24], [25], [26] |



---

|    |                         |     |                        |
|----|-------------------------|-----|------------------------|
| 5  | Farnham Dome            | Yes | [23], [27], [28]       |
| 6  | Gordon Creek            | No  | [23], [29]             |
| 7  | McElmo Dome             | No  | [23], [30]             |
| 8  | Bravo Dome              | No  | [23], [31], [32], [33] |
| 9  | JM- Brown Bassett Field | No  | [34], [35]             |
| 10 | El Trapial Field        | No  | [36]                   |
| 11 | Quebrache Field         | No  | [37], [38]             |
| 12 | Montmiral               | Yes | [11], [39]             |
| 13 | Messokampos             | Yes | [39]                   |
| 14 | Fizzy Field             | No  | [7], [41], [42]        |
| 15 | Vorderrhön              | Yes | [42]                   |
| 16 | Cheb Basin              | Yes | [43]                   |
| 17 | Latera Caldera          | Yes | [44]                   |
| 18 | Benevento Field         | No  | [12]                   |
| 19 | Monte Taburno Reservoir | Yes | [12]                   |
| 20 | Muscillo Reservoir      | No  | [12]                   |
| 21 | Acerno Reservoir        | No  | [12]                   |
| 22 | Pieve Santo Stefano     | Yes | [12]                   |
| 23 | Frigento Field          | Yes | [12]                   |
| 24 | Wiehengebirgsvorland    | No  | [45]                   |
| 25 | Budafa Field            | No  | [46],                  |
| 26 | Mihalyi-Repcelak        | No  | [42]                   |
| 27 | Zaizhuangzi Field       | No  | [6], [47]              |
| 28 | You' aicun Field        | No  | [47], [48]             |
| 29 | Dazhongwang WG1         | No  | [6], [47], [48]        |
| 30 | Gaoqing Field           | NA  | [47], [49], [50]       |
| 31 | Ping Fang Wang Field    | No  | [47], [48], [50]       |
| 32 | Yang 25 Field           | No  | [47], [48], [50]       |
| 33 | Balipo Field            | No  | [47], [48]             |
| 34 | Pingnan                 | No  | [47], [48], [50]       |
| 35 | Hua 17 Field            | No  | [47], [48], [50]       |
| 36 | Huangquiao Field        | No  | [49]                   |
| 37 | Huanchang 3-4 Field     | No  | [51]                   |
| 38 | Wanjinta Field          | No  | [47], [49]             |
| 39 | Qian'an                 | No  | [47]                   |
| 40 | Nong'ancun Field        | No  | [47], [49]             |
| 41 | Changling Field         | No  | [52]                   |
| 42 | DF1-1 Field             | No  | [53]                   |
| 43 | LD28-1 Field            | No  | [53]                   |
| 44 | LD15-1 Field            | No  | [53]                   |
| 45 | Natuna D-Alpha Block    | No  | [54]                   |
| 46 | Ladbroke Grove Field    | No  | [55], [56]             |
| 47 | Tuna Field              | No  | [57]                   |
| 48 | Kapuni Field            | No  | [58], [59]             |

---

|    |                   |     |            |
|----|-------------------|-----|------------|
| 49 | New Plymouth Area | Yes | [58], [60] |
|----|-------------------|-----|------------|

---