

Blocking the k -holes of point sets in the plane

Javier Cano * Alfredo García † Ferran Hurtado ‡
Toshinori Sakai § Javier Tejel † Jorge Urrutia *

Abstract

Let P be a set of n points in the plane in general position. A subset H of P consisting of k elements that are the vertices of a convex polygon is called a k -hole of P , if there is no element of P in the interior of its convex hull. A set B of points in the plane blocks the k -holes of P if any k -hole of P contains at least one element of B in the interior of its convex hull. In this paper we establish upper and lower bounds on the sizes of k -hole blocking sets, with emphasis in the case $k = 5$.

1 Introduction

Let P be a set of n points in the plane in general position, i.e., such no three of them are collinear. All point sets considered in this paper are assumed to be in general position, and therefore this assumption is mentioned only occasionally hereafter. The *convex hull* of P , denoted as $CH(P)$, is the smallest convex set containing all of the elements of P . A set of points is in *convex position*, if its elements are the vertices of a convex polygon. A

¹Instituto de Matemáticas, UNAM, Mexico. j_cano@uxmcc2.iimas.unam.mx, urrutia@matem.unam.mx. Partially supported by projects MTM2006-03909 (Spain), and SEP-CONACYT of Mexico, Proyecto 80268.

²Departamento de Métodos Estadísticos, IUMA, Universidad de Zaragoza, Spain. olaverri@unizar.es, jtejel@unizar.es. Partially supported by projects E58-DGA, MICINN MTM2009-07242, MINECO MTM2012-30951, and ESF EUROCORES programme EuroGIGA, CRP ComPoSe: MICINN Project EUI-EURC-2011-4306.

³Departament de Matemàtica Aplicada II, UPC, Spain. Ferran.Hurtado@upc.edu. Partially supported by projects MICINN MTM2009-07242, MINECO MTM2012-30951, Gen. Cat. DGR2009SGR1040, and ESF EUROCORES programme EuroGIGA, CRP ComPoSe: MICINN Project EUI-EURC-2011-4306.

⁴Research Institute of Educational Development, Tokai University, Japan. sakai@tokai-u.co.jp.

16 subset H of P with k elements is called a k -hole of P if it is in convex
17 position, and no element of P belongs to the interior of $CH(H)$.

18 Counting and finding k -holes of point sets has been a very active area of
19 research since Erdős and Szekeres [9, 10] asked about the existence of k -holes
20 in planar point sets. Harborth proved that any point set with at least ten
21 points always contains at least one 5-hole [13]. Horton [14] proved that for
22 $k \geq 7$ there are point sets containing no k -holes. Recently Nicolás [17] and
23 independently Gerken [12] proved that any point set with sufficiently many
24 points contains at least one 6-hole.

25 Let $f_k(n)$ be the minimum number of k -holes that every point set has.
26 Katchalski and Meir [16] proved that $\binom{n}{2} \leq f_3(n) \leq kn^2$ for some $k < 200$;
27 see also Purdy [20]. Their lower bounds were improved by Dehnhardt [6] to
28 $n^2 - 5n + 10 \leq f_3(n)$, who also proved that $\binom{n-3}{2} + 6 \leq f_4(n)$. Point sets
29 with few k -holes for $3 \leq k \leq 6$ were obtained by Bárány and Valtr [4].

30 Chromatic variants of the Erdős-Szekeres problem were introduced by
31 Devillers, Hurtado, Károly, and Seara [7]. They proved among other results
32 that any bichromatic point set contains at least $\frac{n}{4} - 2$ compatible monochro-
33 matic empty triangles (i.e., having pairwise disjoint interiors). Aichholzer *et*
34 *al.* [1] proved that any bichromatic point set always contains $\Omega(n^{5/4})$ empty
35 monochromatic triangles; this bound was improved by Pach and Tóth [18]
36 to $\Omega(n^{4/3})$. For a thorough survey on this topic, the reader is referred to
37 B. Vogthenhuber's doctoral's thesis [2], where new variations on these and
38 other problems (e.g. dropping the convexity condition on holes) are studied.

39 In this paper we consider the problem of, given a point set P , finding a
40 second set of points, as small as possible, that *pierce*, *stab*, or *block* all the
41 holes of a certain size in P . More precisely: A point $q \notin P$ *blocks* a hole
42 H of P if it belongs to the interior of $CH(H)$. A set of points B such that
43 $B \cap P = \emptyset$ is called a *k-hole blocking set* of P , for short a *k-blocking set* of
44 P , if for any k -hole H of P , there at least one element of B in the interior
45 of $CH(H)$. In the rest of this paper, P will always be a point set in general
46 position with n elements, $n \geq 3$.

47 Given a point set P , let c_P be the number of elements of P on the bound-
48 ary of $CH(P)$. The problem of finding 3-blocking sets has been studied for
49 some time now. It is known that any point set P always has a 3-blocking
50 set with exactly $2n - c_P - 2$ elements, and since any triangulation of P
51 contains exactly $2n - c_P - 2$ elements, this bound is tight; see Katchalski
52 and Meir [16], and Czyzowicz, Kranakis and Urrutia [5].

53 Sakai and Urrutia proved in [21] that there are point sets for which
54 $2n - o(n)$ points are necessary to block all their 4-holes; as $2n - c_P - 2$
55 points are always sufficient to block all the 3-holes of any point set, and

56 thus its 4-holes, this bound is essentially tight. In fact, we believe that in
57 general, the number of points needed to block the 4-holes of any point set
58 P is essentially the same as the number of points needed to block the 3-
59 holes of P (i.e., that the asymptotically dominating terms are the same). In
60 Section 2, we prove that this is the case for point sets in convex position:
61 We prove that to block the 4-holes of any set of n points in convex position,
62 we need at least $n - O(\sqrt{n})$ points, while it is known that $n - 2$ points are
63 sufficient and necessary to block the 3-holes.

64 Remarkably, blocking the k -holes of a point set changes substantially for
65 $k \geq 5$, a problem that, to the best of our knowledge, had not been considered
66 before. In Section 3, the core of this paper, we show that there are point
67 sets, both in general and in convex position, for which the number of points
68 needed to block their 5-holes is as low as a fifth of the number of triangles
69 in a triangulation of the respective point set. We also prove the somehow
70 surprising fact that the number of points needed to block the 5-holes of
71 a point set depends on the geometry of the specific point set, unlike the
72 case of blocking its triangles which only depends on the number of points
73 in the convex hull: We show point sets of the same cardinality, with the
74 same number of points on their convex hulls, for which their 5-blocking sets
75 with minimum cardinality have different sizes. What is more, we show that
76 even for point sets in convex position the size of the 5-blocking sets may be
77 different and depends on the specific geometry.

78 Finally, in Section 4, we give results on blocking the k -holes of point sets
79 in convex position, for general values of k , and we conclude in Section 5 with
80 some observations and open problems.

81 As a final remark in this introduction, it is worth mentioning that the
82 case $k = 2$, *i.e.*, blocking the visibility between pairs of points, has also
83 received attention recently, see [19] and the references therein.

84 **2 Blocking the 4-holes of convex point sets**

85 It is well known that $n - 2$ points are sufficient and necessary to block the
86 3-holes of any set of n points in convex position [16, 5]. In this first section
87 we show that for 4-holes the same amount is essentially needed, in the sense
88 that $n - o(n)$ blocking points are always necessary. More precisely, our main
89 goal in this section is to prove the next result¹:

¹Another proof of this result has independently been found recently by P. Valtr, inspired by discussions during a meeting in Spain in May 2011 (personal communication).

90 **Theorem 2.1.** *Let P any set of n points in convex position. Then, any*
 91 *4-blocking set for P has at least $n - O(\sqrt{n})$ elements.*

92 To prove this, we use a result on the chromatic number of a certain geo-
 93 metric type Kneser graph. Araujo, Dumitrescu, Hurtado, Noy, and Urrutia
 94 [3] introduced the following graph: Let P be a set of n points in convex
 95 position. The *convex segment disjointness graph* of P , denoted by D_n , is
 96 the graph whose vertex set is the set of all line segments with endpoints
 97 in P , two of which are adjacent if they are disjoint. Clearly D_n does not
 98 depend on the choice of P .




Figure 1: Graph D'_7 .

99 Let $\chi(D_n)$ denote the chromatic number of D_n . A lower bound on this
 100 value was obtained by Fabila-Monroy and Wood in [11], while an upper
 101 bound was obtained by Dujmović and Wood in [8]. Both bounds combine
 102 into the following theorem:

Theorem 2.2 ([11, 8]).

$$n - \sqrt{2n + \frac{1}{4}} + \frac{1}{2} \leq \chi(D_n) < n - \sqrt{\frac{1}{2}n} - \frac{1}{2}(\log n) + 4.$$

103 Let D'_n be the graph obtained from D_n by removing the vertices of D_n
 104 corresponding to the edges of the convex hull of P , see Figure 1. Then D'_n
 105 has $\binom{n}{2} - n$ vertices. It is easy to see from the proof of Theorem 2.2 in [11],
 106 that the chromatic number of D'_n satisfies:

$$\chi(D'_n) \geq n - \sqrt{4n + \frac{1}{4}} + \frac{1}{2}.$$

107 We now use this bound to obtain a lower bound on the number of points
 108 blocking all the 4-holes of P that have two edges on the boundary of the
 109 convex hull of P . We call *2-quadrilateral of P* any convex quadrilateral
 110 having two sides that are non-consecutive edges of the convex hull of P (see
 111 Figure 2)




Figure 2: Two intersecting 2-quadrilaterals of P .

112 Let e_i be an edge in the convex hull of P , and m_i be its mid-point. Let
 113 P' the set of all mid-points of the edges of the convex hull of P . Let e_i
 114 and e_j be two *non-consecutive* edges of the convex hull of P . We denote
 115 by $Q(i, j)$ the 2-quadrilateral of P induced by e_i and e_j . It is obvious
 116 that $Q(i, j) \cap Q(r, s) \neq \emptyset$ if and only if the line segments $m_i m_j$ and $m_r m_s$
 117 intersect. Clearly, two 2-quadrilaterals of P can be simultaneously blocked
 118 by a point if and only if their interiors intersect.

119 Let $G'(P)$ be the graph whose vertex set is the set of the 2-quadrilaterals
 120 of P , two of which are adjacent if their interiors do not intersect. Observe
 121 that D'_n and $G'(P)$ are isomorphic graphs: if the elements of P are the
 122 points p_1, \dots, p_n , labelled as they appear clockwise ordered on the convex
 123 hull of P , diagonal $p_i p_j$ (with $j \neq i + 1$) corresponds to the 2-quadrilateral
 124 $Q(i, j)$ defined by the edges $e_i = p_i p_{i+1}$ and $e_j = p_j p_{j+1}$.

125 Suppose that we can block all the 4-holes of P using a set of points
 126 $S = \{q_1, \dots, q_t\}$ with less than $t < \chi(D'_n) = \chi(G'(P))$ points. For each

127 2-quadrilateral C of P , pick a point $q_r \in S$ that blocks C , and assign color r
128 to C . This induces a valid coloring of D'_n , and hence $t \geq n - \sqrt{4n + \frac{1}{4}} + \frac{1}{2}$.
129 Theorem 2.1 follows.

130 3 Blocking 5-holes

131 Given a set of n points P in general position, let us recall that we denote by
132 c_P the number of elements of P that are vertices of $CH(P)$. In this section
133 we study the problem of blocking the 5-holes of point sets in the plane. As
134 announced in the introduction, 5-holes behave, both for convex and general
135 position, quite differently than 4-holes and 3-holes do.

136 3.1 Point sets in convex position

137 3.1.1 Piercing the 5-holes

138 The main objective of this section is to prove the following result, which
139 requires several intermediate lemmas:

140 **Theorem 3.1.** $\frac{n}{2} - 2$ points are always necessary and sometimes sufficient
141 to block the 5-holes of a point set with n elements in convex position and
142 $n = 4k$.

143 We start by proving a more general result:

144 **Lemma 3.2.** Let P a set of n points in convex position. Then any 5-blocking
145 set for P has at least $2\lceil \frac{n}{4} \rceil - 3$ elements.

146 *Proof.* Let B be a 5-blocking set of P with r elements and \mathcal{M} a crossing-free
147 geometric matching of maximum cardinality of the elements of B ; that is, a
148 set of disjoint pairs of elements of B such that the line segments $\{\ell_1, \dots, \ell_{\lfloor \frac{r}{2} \rfloor}\}$
149 joining them do not intersect. Note that if r is odd, we are left with an
150 isolated element of B . One at a time, extend $\ell_1, \dots, \ell_{\lfloor \frac{r}{2} \rfloor}$ until they hit a
151 line segment in M or a previously extended segment. Observe that some
152 ℓ_i 's might be extended to semi-lines or lines. When r is odd, start with a
153 tiny line segment containing the unmatched element of B and extend it as
154 before; see Figure 3.

155 This process yields a decomposition of the plane into exactly $\lceil \frac{r}{2} \rceil + 1$
156 convex regions. If one of these regions contains five or more points, it would
157 contain a 5-hole of P not blocked by B . Thus each of these regions contains
158 at most 4 elements of P , and therefore $|B| = r \geq 2\lceil \frac{n}{4} \rceil - 3$. \square




Figure 3: Illustration of Theorem 3.2.

159 For $n = 4k$, we can improve slightly on the previous bound:

160 **Lemma 3.3.** *Let P a set of n points in convex position with $n = 4k$. Then*
 161 *any 5-blocking set for P has at least $\frac{n}{2} - 2$ elements.*

162 *Proof.* Suppose that we have a 5-blocking set B for P with $\frac{n}{2} - 3$ points and
 163 $n = 4k$. Obtain a decomposition of the plane as in the proof of Lemma 3.2
 164 by an almost perfect geometric matching of the elements of B . Clearly each
 165 cell of such decomposition contains exactly 4 elements of P . Since $|B|$ is
 166 odd, there is one element b of B unmatched and then, there is an edge ℓ of
 167 the decomposition that only contains b , rotate ℓ around b until it hits one
 168 element of P , now there are 5 points in one of the cells incident to ℓ that
 169 contains 5 elements of P in its closure, and clearly those 5 points define a
 170 5-hole that does not contains b in its interior, so we need at least one more
 171 point to block all the 5-holes of P . We conclude that any 5-blocking set of
 172 P contains at least $\frac{n}{2} - 2$ points. \square

173 A point set P is called *almost convex* if any triangle whose vertices are
 174 in P contains at most one element of P in its interior. Almost convex sets
 175 were introduced by Károlyi, Pach and Tóth in [15]. They constructed a
 176 family \mathcal{X}_j of almost convex point sets as follows.

177 Let Z_1 be the end-points of a horizontal line segment ℓ_1 of length two,
 178 and define $\mathcal{X}_1 = \mathcal{R}_1$. Let \mathcal{R}_2 be the set of endpoints of two vertical line
 179 segments ℓ_2 and ℓ_3 of length one whose mid-points are very close to the
 180 endpoints of ℓ_1 , and let $\mathcal{X}_2 = \mathcal{R}_1 \cup \mathcal{R}_2$. See Figure 4(a).




Figure 4: In (a) we show point set \mathcal{X}_2 , in (b) point set \mathcal{X}_3 .

181 Assume that we have already defined $\mathcal{R}_1, \dots, \mathcal{R}_j, \mathcal{X}_1, \dots, \mathcal{X}_j, j \geq 2$, such
 182 that they satisfy the following conditions:

- 183 (1) $\mathcal{X}_j := \mathcal{R}_1 \cup \dots \cup \mathcal{R}_j$ is in general position,
 184 (2) the vertices of $CH(\mathcal{X}_j)$ are the elements of \mathcal{R}_j , and
 185 (3) any triangle determined by three elements of \mathcal{R}_j contains precisely one
 186 point of \mathcal{X}_{j-1} in its interior.

187 Clearly \mathcal{X}_1 and \mathcal{X}_2 satisfy the preceding conditions. Observe that condi-
 188 tion (3), implies that \mathcal{X}_{j-1} is a 3-blocking set of $\mathcal{R}_j, j \geq 2$.

189 The set \mathcal{X}_{j+1} is constructed as follows. Let z_1, \dots, z_r denote the vertices
 190 of $CH(\mathcal{X}_j)$ in clockwise order around $CH(\mathcal{X}_j)$. For every $1 \leq i \leq r$, let ℓ_i
 191 denote the line through z_i orthogonal to the bisector of the angle of $CH(\mathcal{X}_j)$
 192 at z_i . Let z'_i and z''_i be the two points in ℓ_i at infinitesimal distance ε from
 193 z_i . Now move simultaneously z'_i and z''_i away from $CH(\mathcal{X}_j)$ in the direction
 194 orthogonal to ℓ_i by another infinitesimal distance δ , with $\varepsilon \gg \delta$, and denote
 195 the resulting points u'_i and u''_i , respectively.

196 It is proved in [15] that ε and δ can be chosen small enough such that
 197 $\mathcal{R}_{j+1} = \{u'_i, u''_i | i = 1, \dots, r\}$ and $\mathcal{X}_{j+1} := \mathcal{R}_1 \cup \dots \cup \mathcal{R}_{j+1}$ satisfy conditions
 198 1, 2, 3 above. See Figure 4(b).

199 With the preceding construction we are ready to prove:

200 **Lemma 3.4.** *There is a set P of n points in convex position with $n = 2^m$
 201 that has a 5-blocking set consisting of $\frac{n}{2} - 2$ elements.*

202 *Proof.* Let $P = \mathcal{R}_m$ and $B = \mathcal{X}_{m-2}$. Then $|P| = n$ and $|B| = \frac{n}{2} - 2$. We will
 203 show that B is a 5-hole blocking set for P . Suppose that B is not a 5-hole
 204 blocking set of P , then there is a 5-hole H of P such that no point of B lies

205 in the interior of the convex hull of H . Take a triangulation of H ; it will
 206 have three triangles of P . By construction, each of them contains exactly
 207 one element of \mathcal{X}_{m-1} , since $B = \mathcal{X}_{m-1} \setminus \mathcal{R}_{m-1}$. Then these three points
 208 have to be elements of \mathcal{R}_{m-1} and they form a triangle contained in H . By
 209 construction, such a triangle contains precisely one element q of \mathcal{X}_{m-2} . Thus
 210 q blocks H , which is a contradiction. Our result follows. \square

211 The proof of Theorem 3.1 follows now immediately from Lemmas 3.3
 212 and 3.4. \square

213

214 Theorem 3.1 is frankly surprising to us. We believed that a similar result
 215 to that obtained for blocking the 4-holes of point sets in convex position
 216 would also hold for 5-blocking sets, i.e., we thought that a 5-blocking set of
 217 any point set P in convex position would always have $n - o(n)$ elements.
 218 We have seen that that is not always the case yet, we still believe that for
 219 some point sets in convex position that may be the right answer. We pose
 220 explicitly a related open problem:

221 **Problem 3.5.** *Is it true that if P is the set of vertices of a regular polygon*
 222 *with n vertices, then any 5-blocking set of P has at least $n - o(n)$ elements?*

223 3.1.2 Blocking 5-holes of regular polygons

224 While a solution of Problem 3.5 remains elusive to us, we give in this section
 225 a proof for a special case, because the technique is used in Section 3.1.3, and
 226 we also hope that it may inspire a general solution.

227 Let $\mathcal{Q}_n = \{p_0, \dots, p_{n-1}\}$ be the vertices of a regular polygon \mathcal{R}_n with
 228 n vertices, given as they appear on the boundary in clockwise order. The
 229 arithmetic of their indices is done modulo n . A subset of \mathcal{Q}_n is called a
 230 *lateral k -hole* if its elements are k consecutive elements of \mathcal{Q}_n . To be more
 231 precise, we use the notation $S_{i,k} = \{p_i, \dots, p_{i+k-1}\}$ for the i -th lateral k -hole
 232 of \mathcal{Q}_n , with $0 \leq i \leq n-1$ and $3 \leq k \leq n$. The convex hull of $S_{i,k}$ is a convex
 233 k -gon, which we denote $R_{i,k}$. Abusing slightly the notation, we also say that
 234 $R_{i,k}$ is a *lateral k -hole* of \mathcal{R}_n .

235 **Lemma 3.6.** *Any 5 blocking set of \mathcal{Q}_{19} has at least eight elements.*

236 *Proof.* First, recall that according to Lemmas 3.2 and 3.3, to block the 5-
 237 holes of any convex polygon with 5, 8, 13, 16, 17, and 19 vertices, we need
 238 at least 1, 2, 5, 6, 7, and 7 points, respectively.




Figure 5: A regular 19-gon.

239 We prove now our claim by contradiction: Suppose that there is a 5-
 240 blocking set B of \mathcal{Q}_{19} consisting of seven points. Observe first that if we
 241 remove a lateral 4-hole $R_{i,4}$ from \mathcal{R}_{19} , we obtain a convex 17-gon, namely
 242 $R_{i+3,17}$. As mentioned in the preceding paragraph, to block the 5 holes
 243 of $R_{i+3,17}$ we need at least seven points. It follows that all the elements
 244 of B lie in the interior of $R_{i+3,17}$ and therefore, that no lateral 4-hole $R_{i,4}$
 245 contains any element of B . Let W_4 the union of these regions, i.e., $W_4 =$
 246 $\bigcup_{i=0,\dots,n-1} R_{i,4}$, a polygonal annulus that contains no point from B .

247 Let $R_{i,5}$ be a lateral 5-hole of \mathcal{R}_{19} , and $\widehat{R}_{i,5}$ the subset of $R_{i,5}$ obtained
 248 by removing from $R_{i,5}$ all the points that belong to some lateral 4-hole of
 249 \mathcal{R}_{19} : Equivalently, $\widehat{R}_{i,5} = R_{i,5} \setminus W_4$ (see Figure 5, upper part). Since the
 250 elements of B block all the 5-holes of \mathcal{Q}_{19} , every lateral 5-hole $R_{i,5}$ of \mathcal{R}_{19}
 251 contains at least one element of B , which must belong to $\widehat{R}_{i,5}$.

252 Observe that the polygonal region that complements $R_{i,5}$ in \mathcal{R}_{19} is pre-
 253 cisely $R_{i+4,16}$. As we know that we need at least six points to block the
 254 5 holes of the vertices of any convex polygon with 16 vertices, each lateral
 255 5-hole of \mathcal{R}_{19} must contain exactly one blocking point.

256 In a similar way, if we remove a lateral 8-hole $R_{i,8}$ from \mathcal{R}_{19} , we are
 257 left with a convex polygon $R_{i+7,13}$ with 13 vertices, and thus at least five
 258 elements of B belong to the interior of $R_{i+7,13}$. It follows that each lateral
 259 8-hole of Q contains exactly two elements of B .

260 Observe that for each lateral 8-hole $R_{i,8}$ of \mathcal{R}_{19} , there are exactly two
 261 lateral 5-holes of \mathcal{R}_{19} , namely $R_{i,5}$ and $R_{i+3,5}$, such that their corresponding
 262 regions $\widehat{R}_{i,5}$ and $\widehat{R}_{i+3,5}$ are disjoint and contained in $R_{i,8}$. Let $H_{i,8} = R_{i,8} \setminus$
 263 $(\widehat{R}_{i,5} \cup \widehat{R}_{i+3,5})$. The preceding discussion implies that the two blocking points
 264 of B in $R_{i,8}$ must be one in $\widehat{R}_{i,5}$ and the other one in $\widehat{R}_{i+3,5}$, and that $H_{i,8}$
 265 is empty of points from B .

266 Let R_B be the region obtained by removing from \mathcal{R}_{19} all the empty
 267 regions $H_{i,8}$ defined the lateral 8-holes $R_{i,8}$ of \mathcal{R}_{19} , with $0 \leq i \leq n-1$.
 268 All the points of B must lie in R_B . It is easy to see that R_B consists of a
 269 19-regular polygon $C_{19} = \bigcap_{i=0, \dots, 18} R_{i,13}$, with the same center than \mathcal{R}_{19} ,
 270 and 19 hexagons, which we call A_i , for $0 \leq i \leq n-1$, where we denote by
 271 A_i the hexagon that is closer to p_i . To be precise, $A_i = R_{i-3,5} \cap R_{i-1,5} \cap$
 272 $R_{i,12} \cap R_{i+1,17} \cap R_{i+2,17} \cap R_{i+7,12}$. The twenty connected components of R_B
 273 are shaded in yellow in Figure 5.

274 No point in the central 19-gon C_{19} can block any lateral 5-hole. In
 275 addition, putting a blocking point in one of the hexagonal regions A_i , we
 276 only block 3 lateral 5-holes, $R_{i-3,5}$, $R_{i-2,5}$ and $R_{i-1,5}$.

277 Therefore, to block the 19 lateral 5-holes of \mathcal{R}_{19} , we need to put the
 278 seven blocking points from B in the hexagonal regions. As every lateral
 279 5-hole contains three of these hexagons, one of the lateral 5-holes of \mathcal{R}_{19} will
 280 contain two blocking points, contradicting the fact that each lateral 5-hole
 281 of \mathcal{R}_{19} contains exactly one point in B . \square

282 3.1.3 Geometry matters

283 Lemmas 3.4 and 3.6 indicate that the geometry and distribution of the
 284 points has to be considered when finding 5-blocking sets for point sets, even
 285 in convex position. In this section we go deeper in that direction, and show
 286 two set of 11 points in convex position, for which their smallest 5-blocking
 287 point sets have different cardinalities.

288 Our first point set is \mathcal{Q}_{11} , the set of vertices of a regular polygon \mathcal{R}_{11} with
 289 eleven vertices. With an approach along the lines of the proof of Lemma 3.6
 290 it is easy to see that the 5-holes of \mathcal{Q}_{11} can be blocked with exactly three
 291 points, see Figure 6.

292 Our second point set, $\mathcal{S}_{11} = \{p_0, \dots, p_{10}\}$ is shown in Figure 7. First
 293 note that the four blue dots shown in Figure 7, block all the 5-holes of \mathcal{S}_{11} .
 294 We now prove that the 5-holes of \mathcal{S}_{11} cannot be blocked with three points.
 295 Let \mathcal{P}_{11} be the convex polygon with vertex set \mathcal{S}_{11} .

296 For any $0 \leq i \leq 10$ let T_i be the triangle bounded by the segments
 297 $p_i - p_{i+1}$, $p_i - p_{i+4}$, and $p_{i-3} - p_{i+1}$, addition taken *mod* 11. Observe that any




Figure 6: A regular 11-gon.

298 point of the plane can block at most four lateral 5-holes of \mathcal{S}_{11} , and that if it
 299 does, it must belong to some T_i , in which case it blocks the laterals 5-holes
 300 of \mathcal{S}_{11} with vertex sets $\{p_{i-3}, \dots, p_{i+1}\}$, $\{p_{i-2}, \dots, p_{i+2}\}$, $\{p_{i-1}, \dots, p_{i+3}\}$,
 301 and $\{p_i, \dots, p_{i+4}\}$. Suppose now that the 5-holes of \mathcal{S}_{11} can be blocked with
 302 a set of three points $\{x, y, z\}$. In particular $\{x, y, z\}$ also block the eleven
 303 lateral 5-holes of \mathcal{S}_{11} , and thus at least two points among x, y , and z cover
 304 four lateral 5-holes of \mathcal{S}_{11} , and the other point three or four. From this we
 305 can infer that two points among x, y , and z , say x and y , must belong to
 306 two triangles T_i and T_j such that $j = i + 4$ for some $0 \leq i \leq 10$, addition
 307 taken *mod* 11.

308 Since blocking the 5-holes of nine points in convex position requires at
 309 least three blocking points, all the lateral 4-holes of \mathcal{P}_S must be empty. Since
 310 T_1, T_2, T_4, T_7, T_9 and T_{10} are contained in lateral 4-holes of \mathcal{P}_S , they cannot
 311 contain any of the points x, y , or z . Then x and y are in T_0, T_3, T_5, T_6 , or
 312 T_8 .

313 But x and y must belong to some T_i and T_{i+4} , which is not possible:
 314 Therefore, to block the 5-holes of \mathcal{S}_{11} we need at least four points, as claimed.

315 Thus, we have proved:

316 **Theorem 3.7.** *There are two different sets of eleven points in convex posi-*
 317 *tion such that their smallest 5-blocking sets have different cardinalities.*




Figure 7: A set of 11 points in convex position that requires 4 points to block its 5-holes.

318 **3.2 Point sets in general position**

319 **3.2.1 Geometry matters**

320 As mentioned in the introduction, the number of points needed to block the
 321 set of triangles of a point set P , is exactly $2n - c_P - 2$, where $n = |P|$ and
 322 c_P is the number of elements from P that are vertices of $CH(P)$. A similar
 323 formula does not exist for blocking the 5-holes of a point set: We are next
 324 constructing point sets of the same cardinality, and having the same number
 325 of elements on their convex hulls, for which the number of points required
 326 to block their 5-holes are different.

327 In other words, we are giving here a result for points in general position,
 328 similar to Theorem 3.7, proving that the specific geometry and distribution
 329 of the points can change the size of the minimal 5-blocking sets.

330 We show first that there exist families of point sets with $4m$ elements,
 331 with $2m$ of them on the convex hull, such that all of their 5-holes can be
 332 blocked with $m - 2$ points.

333 **Lemma 3.8.** *For any m there is a point set P_{4m} in general position with*
 334 *$|P_{4m}| = n = 4m$ points and $c_P = 2m$, such that $m - 2$ points are sufficient*
 335 *and necessary to block all the 5-holes of P_{4m} .*




Figure 8: A point set in general position in which $\frac{n}{4} - 2$ points are sufficient and necessary to block all of its convex 5-holes. The image on the right is a close up look at each fat point of the regular m -gon at the left.

336 *Proof.* Let $\mathcal{R}_m = \{q_1, \dots, q_m\}$ be a regular m -gon. From the results in [5,
 337 16], we can choose $m - 2$ points $B = \{b_1, \dots, b_{m-2}\}$ such that any triangle
 338 with vertices in \mathcal{R}_m contains exactly an element of B in its interior. It is
 339 not hard to see that given such B , we can move the vertices of \mathcal{R}_m around
 340 some sufficiently small $\varepsilon > 0$, such that any triangle in the perturbed set
 341 contains exactly one element of B .

342 We construct a set P_{4m} with $4m$ points as follows. We substitute each
 343 vertex q_i of \mathcal{R}_m , $i = 1, 2, \dots, m$, by a set of 4 points $S_i = \{p_i^1, p_i^2, p_i^3, p_i^4\}$,
 344 each of them at distance no more than ε from q_i , and consider the set
 345 $P_{4m} = S_1 \cup \dots \cup S_m$. The replacement is as follows: Consider the bisector
 346 b_i of the internal angle of \mathcal{R}_m at q_i . Let ℓ_i be a line orthogonal to b_i that
 347 intersects the edges of \mathcal{R}_m , incident to q_i , infinitesimally enough to q_i . Let
 348 p_i^1 and p_i^4 be the points of intersection of ℓ_i with the circumcircle C of \mathcal{R}_m .
 349 Let p_i^2 and p_i^3 be two points equidistant to q_i , below ℓ_i , one on each of the
 350 edges of \mathcal{R}_m incident to q_i , and such that the angles $\angle p_i^1 p_i^2 p_i^3$ and $\angle p_i^4 p_i^3 p_i^2$
 351 are close to π , see Figure 8. With this replacement, the convex hull of P_{4m}
 352 has $2m$ vertices.

353 Observe that one can choose p_i^1 and p_i^4 such that one of the open half-
 354 planes bounded by the line passing through p_i^1 and p_i^3 (resp. p_i^4 and p_i^2)
 355 contains p_i^4 , (resp. p_i^1), and no other point of P_{4m} . See Figure 8.

356 Observe next, that no 5-hole can use more than two elements of S_i . It

357 follows now that any 5-hole has vertices in at least three different sets S_i ,
 358 S_j , S_k .

359 Moreover, since the elements of S_i are at distance no more than ε from
 360 q_i , any triangle containing a point in any three sets S_i , S_j , and S_k contains
 361 a point of B in its interior. Therefore the elements of B block all of the
 362 5-holes of P_{4m} .

363 Observe now that any 5-blocking set for P_{4m} can not have fewer points
 364 than $m - 2$. First, suppose that B' is a 5-blocking set for P_{4m} with at most
 365 $m - 3$ elements, then at least one triangle with vertices in \mathcal{R}_m that is not
 366 blocked (since the number of triangles in any triangulation of \mathcal{R}_m is $m - 2$).
 367 Assume that the vertices of one such triangle are q_i, q_j, q_k . Then, by taking
 368 two elements in S_i and S_j and one in S_k , we obtain a 5-hole of P_{4m} that is
 369 not blocked by any element of B' . Thus, P_{4m} requires $m - 2$ points in order
 370 to block all of its 5-holes. \square

371 We construct now point sets P'_{4m} with $4m$ elements, $2m$ on its convex
 372 hull, such that to block all of its 5 holes we need more than $2m$ points,
 373 roughly twice as many as for P_{4m} .

374 **Lemma 3.9.** *For every positive integer m divisible by 15 there is a point*
 375 *set P'_{4m} in general position with $|P'_{4m}| = n = 4m$ elements and $c_P = 2m$,*
 376 *such that more than $2m$ are points necessary to block all the 5-holes of P'_{4m} .*

377 *Proof.* Let P'_{4m} be a set with $4m = 30k$ points, with $15k$ on its convex
 378 hull forming the set of vertices of a regular $15k$ -gon. We consider on the
 379 boundary of $CH(P'_{4m})$ alternated subsets consisting of 10 and 5 vertices,
 380 yielding therefore k subsets of each class. For each of the subsets of 5
 381 vertices, we form a *block* connecting with a chord the first and last element
 382 and adding 15 points to the interior of the region, in such a way that the
 383 region can be decomposed into 11 convex 5-gons (the pattern corresponds to
 384 the classical plane drawing of the dodecahedron graph). See figure 9, where
 385 each block is labelled “a”.

386 The part of the convex hull of P'_{4m} that is not in the blocks is an empty
 387 convex polygon H with $12k$ vertices: $10k$ come from the subsets not used
 388 for the blocks and $2k$ come from gathering the first and last points of all the
 389 blocks.

390 By Lemma 3.3, H requires at least $12k/2 - 2$ points to block all of its
 391 5-holes, and for the pentagonized blocks we need at least $11k$ points. Thus,
 392 any 5-blocking set for P'_{4m} contains at least $(6k - 2) + 11k = 17k - 2$ points,
 393 which is larger than $2m = 15k$. \square




Figure 9:

394 Thus, combining Lemmas 3.8 and 3.9 we have proved:

395 **Theorem 3.10.** *There are two different sets of $n = 4m$ points in non-convex*
 396 *position, such that the number of vertices in the convex hull of each set $2m$,*
 397 *and such that their smallest 5-blocking sets have different cardinalities.*

398 3.2.2 Piercing the 5-holes of general point sets

399 We conclude this section with a general a lower bound on the number of
 400 points needed to block the 5-holes of any point set. We prove:

401 **Theorem 3.11.** *Let P be any set of n points in general position. Then any*
 402 *5-blocking set of P has at least $2\lceil \frac{n}{9} \rceil - 3$ points.*

403 *Proof.* Harborth [13] proved that any set of ten points in general position
 404 in the plane always contains a 5-hole. Let B be a 5-blocking set of P .
 405 Take a geometric planar matching of the elements of B , and decompose the
 406 plane into convex regions by extending the segments in the matching as in
 407 Lemma 3.2. Then any convex region in our decomposition cannot contain
 408 more than nine points, otherwise there would be a 5-hole of P not blocked
 409 by any element of B . It now follows, as in the proof of Lemma 3.2, that
 410 $B \geq 2\lceil \frac{n}{9} \rceil - 3$. \square

411 In view of the preceding results we conjecture:

412 **Conjecture 3.12.** *The number of points needed to block all the 5-holes of*
 413 *any point set with n elements is greater than or equal to $\frac{n}{4} \pm c$, where c is a*
 414 *constant.*

415 4 Blocking k -holes for points in convex position

416 In this last section before the concluding remarks, we consider the problem
 417 of blocking k -holes for larger values of k . As mentioned in the introduction,
 418 Horton [14] proved that for $k \geq 7$, there exist point sets that don't have
 419 any k -hole. Thus the question of finding *the minimum* number of blocking
 420 points is properly interesting only for some specific families of point sets
 421 always having k -holes; here we focus on point sets in convex position.

422 Let P be a set of n points in convex position. Using a similar argument
 423 as in the proof of Lemma 3.2, it can be verified that any k -blocking set for
 424 P has at least $2 \left\lceil \frac{n}{k-1} \right\rceil - 3$ elements. This bound is essentially tight for odd
 425 values of k , as we show next.

426 We construct a point set P as follows: Let \mathcal{R}_m , C , B , and ϵ as in the
 427 proof of Lemma 3.8, i.e., \mathcal{R}_m is a regular m -gon, C its circumcircle, B
 428 a set of $m - 2$ points blocking all the triangles of \mathcal{R}_m , and ϵ is the radius of
 429 infinitesimal disks centered at the vertices of \mathcal{R}_m in such a way that if these
 430 vertices are perturbed each to any position inside their associated disks, the
 431 set B keeps blocking all the triangles the perturbed vertices determine. Let
 432 $k = 2s + 1$. Replace each vertex p_i of \mathcal{R}_m with a set $S_i = \{p_1^i, \dots, p_s^i\}$ of s
 433 points on C within the circle of radius ϵ centered at p_i , see Figure 10. Let
 434 $P = S_1 \cup \dots \cup S_m$. Then P has $n = m \times s$ elements,




Figure 10: The general construction when $k = 11$.

435 Then any k -hole with vertices in P has vertices in at least three of the
 436 sets S_i , and thus the set B blocks all of the k -holes of P . But B has $m - 2$
 437 elements, and $2 \left\lceil \frac{n}{k-1} \right\rceil - 2 = 2 \left\lceil \frac{ms}{2s} \right\rceil - 2 = m - 2$.

438 Therefore, we have proved:

439 **Theorem 4.1.** $2 \left\lceil \frac{n}{k-1} \right\rceil - 3$ points are always necessary, and $2 \left\lceil \frac{n}{k-1} \right\rceil - 2$
 440 are sometimes sufficient to block the k holes of a point set with n elements
 441 in convex position.

442 Next we show that when k is even we can give a better lower bound.

443 **Proposition 4.2.** Let $P = \{p_1, \dots, p_n\}$ be a set of $n = mh$ points in convex
 444 position, with $h \geq 2$. Then, $\frac{n}{h} - O(\sqrt{(n)})$ points are necessary to block all
 445 the $2(h + 1)$ -holes of P .

446 *Proof.* Let us denote by P' the set of points $p_{i,h}$, for $i = 1, 2, \dots, m$. Then,
 447 the number of points in P' is $n/h = m$. Figure 11 shows the case with $h = 2$
 448 (6-holes), in which P' is the set of points with even indices.




Figure 11: Illustration for Proposition 4.2.

449 Take two arbitrary edges $p_{i,h} - p_{(i+1) \cdot h}$ and $p_{j,h} - p_{(j+1) \cdot h}$, with $i + 1 < j$,
 450 and let $H_{i,j}$ be the $2(h + 1)$ -hole of P determined by the set of points $\{p_{i,h},$
 451 $p_{i \cdot h+1}, \dots, p_{(i+1) \cdot h}, p_{j,h}, p_{j \cdot h+1}, \dots, p_{(j+1) \cdot h}\}$. Consider now the 4-hole $H'_{i,j}$
 452 of P' determined by $p_{i,h}, p_{(i+1) \cdot h}, p_{j,h}$ and $p_{(j+1) \cdot h}$. Observe that two edges

453 of this 4-hole are on the convex hull of P' and that the other two edges are
 454 diagonals (see Figure 11).

455 Therefore, we can define a bijection between the set Q' of 4-holes in
 456 P' defined by pairs of edges $p_{i \cdot h} p_{(i+1) \cdot h}$ and $p_{j \cdot h} p_{(j+1) \cdot h}$, and the set Q of
 457 $2(h+1)$ -holes $H_{i,j}$ defined above.

458 Now, take a set B of points blocking the $2h$ -holes of Q . Suppose that one
 459 of the blocking points x is inside the polygon with vertices $p_{i \cdot h}, p_{i \cdot h+1}, \dots, p_{(i+1) \cdot h}$
 460 (a triangle in the case $h = 2$). Let R be the set of $2(h+1)$ -holes of Q blocked
 461 only by x . Note that this point can only block the $2(h+1)$ -holes of Q formed
 462 using edge $p_{i \cdot h} p_{(i+1) \cdot h}$.

463 Then, we can remove x and we can add a point y very close to the
 464 midpoint of the edge $p_{i \cdot h} p_{(i+1) \cdot h}$, inside the convex hull of P' , such that y
 465 blocks at least the $2(h+1)$ -holes in R (see Figure 11).

466 Then we can assume that, for any set B blocking the $2(h+1)$ -holes of
 467 Q , all the blocking points are inside the convex hull of P' . In this case, note
 468 that, if a point z blocks a $2(h+1)$ -hole of Q , then its corresponding 4-hole
 469 in Q' is also blocked by z and vice versa.

470 Since there is a bijection between Q and Q' and since we need $\frac{n}{h} -$
 471 $O(\sqrt{n})$ points to block all the 4-holes in Q' (as proved in Section 2), it
 472 is impossible that the size of a $2(h+1)$ -blocking set for Q is smaller than
 473 $\frac{n}{h} - O(\sqrt{n})$, for otherwise we could block the 4-holes of Q' with less than
 474 $\frac{n}{h} - O(\sqrt{n})$ points. \square

475 5 Final remarks

476 Closing the gaps between the lower and upper bounds for this family of
 477 problems is obviously a main open problem for future research. Yet to be
 478 more specific, we would like to end this paper emphasizing the interest of
 479 bringing more light into two specific bounds.

480 As repeatedly mentioned in this paper, it is known that any point set S
 481 that blocks the set of triangles of any n -point set P in convex position, has
 482 at least $n - 2$ points; moreover, if $|S| = n - 2$, which is achievable, then any
 483 triangle with vertices in P has exactly one element of S in its interior. This
 484 gives a trivial upper bound on the number of elements sufficient to block
 485 the k -holes of P : Simply remove $k - 3$ elements from S . However, we do
 486 not know a better upper bound than that! In fact, we conclude with an
 487 apparently simpler question:

488 **Question 5.1.** *Is it true that to block all the k holes of the set of vertices*
 489 *of a regular n -gon, we need $n - c(k)$ points?*

490 We believe that the answer to the preceding question should be positive,
491 but a proof is still elusive to us.

492 References

- 493 [1] O. Aichholzer, R. Fabila-Monroy, D. Flores-Peñaloza, T. Hackl, C. Hue-
494 mer, and J. Urrutia. Empty Monochromatic Triangles, *Computational*
495 *Geometry, Theory and Applications*, **42** 9, (2009), 934–938.
- 496 [2] B. Vogtenhuber, Combinatorial aspects of colored point sets in the plane.
497 Doctoral Thesis at Graz University of Technology, Nov. 2011.
- 498 [3] G. Araujo, A. Dumitrescu, F. Hurtado, M. Noy, and J. Urrutia, On the
499 chromatic number of some geometric type Kneser graphs. *Computational*
500 *Geometry Theory and Applications*, 32(1):59–69, 2005.
- 501 [4] I. Bárány, and P. Valtr. Planar point sets with a small number of empty
502 convex polygons, *Studia Scientiarum Mathematicarum Hungarica*, bf 41,
503 2, (2004), 243-266.
- 504 [5] J. Czyzowicz, E. Kranakis, and J. Urrutia, Guarding the convex subsets
505 of a point set, in *12th Canadian Conference on Computational Geometry*,
506 Fredericton, New Brunswick, Canada, 2000, 47–50.
- 507 [6] K. Dehnhardt. Leere konvexe Vielecke in ebenen Punktmengen, *Dissert-*
508 *ation, TU Braunschweig*, (1987).
- 509 [7] O. Devillers, F. Hurtado, G. Károlyi, and C. Seara. Chromatic variants
510 of the Erdős-Szekeres Theorem, it *Computational Geometry Theory and*
511 *Applications*, **26**, 3, (2003) 193–208.
- 512 [8] V. Dujmović and D. Wood. Thickness and Antithickness, *Manuscript in*
513 *preparation*, (2011).
- 514 [9] P. Erdős. Some more problems on elementary geometry, *Austral. Math.*
515 *Soc. Gaz.* , **5** (1978), 52–54.
- 516 [10] P. Erdős, and G. Szekeres. A combinatorial problem in geometry, *Com-*
517 *positio Math.*, (1935), **2**, 463–470.
- 518 [11] R. Fabila-Monroy, and D.R. Wood, The chromatic number of the con-
519 vex segment disjointness graph. In *Proceedings of XIV Spanish Meeting*
520 *on Computational Geometry*, 2011.

- 521 [12] T. Gerken. Empty convex hexagons in planar point sets, *Discrete &*
522 *Computational Geometry*, **39**, 1-3, (2008), 239–272.
- 523 [13] H. Harborth, Konvexe, Fünfecke in ebenen Punktmengen. *Elemente*
524 *Math.*, **33** (1978), 116–118.
- 525 [14] J. D. Horton. *Sets with no empty convex 7-gons*. *Canad. Math. Bull.*,
526 **26**, 482 – 484 (1983).
- 527 [15] G. Károlyi, J. Pach, and G. Tóth, A modular version of the Erdős-
528 Szekeres theorem, *Studia Scientiarum Mathematicarum Hungarica* **51**
529 (2001), 245–260.
- 530 [16] M. Katchalski, and A. Meir, On empty triangles determined by points
531 in the plane, *Acta Math. Hungar* **51** (1988), 323–328.
- 532 [17] C.M. Nicolás. The empty hexagon theorem, *Discrete & Computational*
533 *Geometry*, **38** (2007), 389–397.
- 534 [18] J. Pach and G. Tóth. Monochromatic empty triangles in two-colored
535 point sets, *Geometry, Games, Graphs and Education: the Joe Malkevitch*
536 *Festschrift*, S. Garfunkel and R. Nath, eds, COMAP, Bedford, MA.,
537 (2008), 195–198.
- 538 [19] A. Pór, and D. R. Wood, On visibility and blockers, *Journal on Com-*
539 *putational Geometry* **1(1)** (2010), 29–40.
- 540 [20] G.B. Purdy. The minimum number of empty triangles, *AMS Abstracts*,
541 **3**, (1982), 318.
- 542 [21] T. Sakai, and J. Urrutia, Covering the convex quadrilaterals of point
543 sets, *Graphs and Combinatorics* **38** (2007), 343–358.