

Inorganics **2014**, *2*, 424-432; doi:10.3390/inorganics2030424

OPEN ACCESS

inorganics

ISSN 2304-6740

www.mdpi.com/journal/inorganics

Article

Gold Thione Complexes

Francesco Caddeo ^{1,2}, Vanesa Fernández-Moreira ¹, Massimiliano Arca ², Antonio Laguna ¹, Vito Lippolis ^{2,*} and M. Concepción Gimeno ^{1,*}

¹ Departamento de Química Inorgánica, Instituto de Síntesis Química y Catálisis Homogénea (ISQCH), CSIC-Universidad de Zaragoza, Zaragoza 50009, Spain;

E-Mails: suamaestailcaddeo@hotmail.it (F.C.); vanesa@unizar.es (V.F.-M.); alaguna@unizar.es (A.L.)

² Dipartimento di Scienze Chimiche e Geologiche, Università degli Studi di Cagliari, S.S. 554 Bivio per Sestu, Monserrato (CA) 09042, Italy; E-Mail: marca@unica.it

* Authors to whom correspondence should be addressed; E-Mails: lippolis@unica.it (V.L.); gimeno@unizar.es (M.C.G.); Tel.: +39-706-754-467 (V.L.); +34-976-762-291 (M.C.G.).

Received: 2 June 2014; in revised form: 16 July 2014 / Accepted: 18 July 2014 /

Published: 4 August 2014

Abstract: The reaction of the ligand Et₄todit (4,5,6,7-Tetrathiocino-[1,2-*b*:3,4-*b'*]-diimidazolyl-1,3,8,10-tetraethyl-2,9-dithione) with gold complexes leads to the dinuclear gold(I) complexes [$\{\text{Au}(\text{C}_6\text{F}_5)\}_2(\text{Et}_4\text{todit})$] and $[\text{Au}(\text{Et}_4\text{todit})_2(\text{OTf})_2]$, which do not contain any gold-gold interactions, or to the gold(III) derivative [$\{\text{Au}(\text{C}_6\text{F}_5)_3\}_2(\text{Et}_4\text{todit})$]. The crystal structures have been established by X-ray diffraction studies and show that the gold centers coordinate to the sulfur atoms of the imidazoline-2-thione groups.

Keywords: gold(I); gold(III); imidazole derivatives

1. Introduction

The coordination chemistry of organosulfur compounds featuring two or more 3-alkylimidazole-2-thione groups has not been studied in depth in spite of their potential structural coordination possibilities. The most intensely studied ligands are the derivatives that contain two 3-methylimidazoline-2-thione groups bounded by a $-(\text{CH}_2)_n-$ chain [$n = 1$ (Mbit), 2 (Ebit), 4 (Bbit)] and some complexes with nickel(II), cobalt(II) [1], rhodium(III) and iridium(III) [2], antimony(III),

bismuth(III) [3], tin(IV) [4], lead(II) [5], silver(I) [6,7] and group 11 elements [7] have been described in the literature with these types of ligands.


Tetrathiocino-diimidazolyl species also contains two C=S groups that can behave as donors towards Lewis acids [8–10]. In particular, the ligand Et₄todit (4,5,6,7-tetrathiocino-[1,2-*b*:3,4-*b'*]-diimidazolyl-1,3,8,10-tetraethyl-2,9-dithione) (Scheme 1) features very distant positions of the thiocarbonyl groups that make this molecule capable to act as bidentate bridging ligand. Some polymeric complexes with stoichiometry [M(Et₄todit)X₂] (M = Zn, Cd, Hg; X = Cl, Br) [11] or [Cu(Et₄todit)I] [12] have been reported. This notwithstanding, no one gold complex with this type of ligand has been obtained to date.

In this paper we report the synthesis and structural characterization of the dinuclear gold(I) complexes [{Au(C₆F₅)₂(Et₄todit)] and [Au(Et₄todit)]₂(OTf)₂ and of the gold(III) derivative [{Au(C₆F₅)₃]₂(Et₄todit)].

2. Results and Discussion


The Et₄todit ligand was prepared from the corresponding imidazoline-2-thione-4,5-dione by reaction with the Lawesson's reagent, as it was described previously [9,13,14]. The HOMO-1 and HOMO calculated at DFT level for Et₄todit are built up of the in-phase and out-of-phase combinations of the non-bonding lone pairs localized on the sulfur atoms, perpendicular to the imidazole plane, respectively, with remarkably negative NBO charges [9]. The terminal thiocarbonyl groups are potential donors not only towards molecular acids, [8–10] but also towards transition metal ions [11,12]. We have, therefore, studied the coordination properties of this ligand towards gold(I) and gold(III) derivatives. Scheme 1 represents the synthesized complexes.

Scheme 1. (i) 2[Au(C₆F₅)(tht)], (ii) [Au(tht)₂OTf], (iii) 2[Au(C₆F₅)₃(tht)].



The reaction of Et₄todit with [Au(C₆F₅)(tht)] (tht = tetrahydrothiophene, molar ratio = 1:2) leads to the dinuclear derivative [$\{\text{Au}(\text{C}_6\text{F}_5)\}_2(\text{Et}_4\text{todit})$] (**1**) (Scheme 1). It is a white solid that has been characterized by ¹H-NMR and ¹⁹F-NMR spectroscopy and mass spectrometry. Single crystals suitable for X-ray diffraction analysis were obtained by slow diffusion of hexane in a solution of **1** in 1,2-dichloroethane, and the molecular structure is depicted in Figure 1. The two five-membered rings of the bis(thiocarbonyl) donors are twisted about the C–C bond by 71.19°, similarly to what found in the crystal structure of Et₄todit (69.90°) and its derivatives, such as Et₄todit·2Br₂, where the four independent molecules of the asymmetric units show torsion values ranging between 67.5° and 79.7° [9,14]. The two gold centers present a distorted linear geometry and are bonded to the C=S groups of the thioimidazol rings and to the pentafluorophenyl groups. The S(1)–Au(1)–C(1) angle (174.8(1)°) and the Au(1)–C(1) (2.022(4) Å) and Au(1)–S(1) (2.333(1) Å) bond distances found in the solid structure are in good agreement with those reported in the literature [4,5]. No Au(I)–Au(I) interactions have been observed in the crystal structure. The complex is not luminescent in solid state or in solution.

Figure 1. Molecular structure of the compound **1**, with the atom labeling scheme (hydrogen atoms omitted for clarity).



A selection of bond lengths and angles for complex **1** are summarized in Table 1.


Table 1. Selected bond lengths (Å) and angles (°) for **1**.

Distances (Å)		Angles (°)	
Au(1)–S(1)	2.333(1)	S(1)–Au(1)–C(1)	174.8(1)
Au(2)–S(6)	2.309(1)	Au(1)–S(1)–C(7)	97.8(1)
Au(2)–C(21)	2.028(4)	Au(2)–S(6)–C(15)	113.7(1)
		Au(1)–C(1)–C(2)	121.3(3)
		Au(1)–C(1)–C(6)	124.1(3)

The ligand Et₄todit also reacts with [Au(tht)₂]OTf (OTf = triflate, CF₃SO₃) to give the dinuclear [Au(Et₄todit)]₂(OTf)₂ (**2**) (Scheme 1) as a yellow solid. It is soluble in dichloromethane and acetone

and insoluble in diethyl ether and hexane. It has been characterized by $^1\text{H-NMR}$ spectroscopy and mass spectrometry. Single crystals suitable for X-ray diffraction analysis were obtained by slow diffusion of hexane in a solution of **2** in 1,2-dichloroethane, and the molecular structure is depicted in Figure 2. The two five-membered rings of the bis(thiocarbonyl) donors are twisted about the C–C bond by 72.65° . The two gold centers present a distorted linear geometry and are bonded to two C=S groups of the thioimidazole of different Et_4todit ligands. The S(1)–Au(1)–S(2) angle ($174.8(1)^\circ$) and the Au(1)–S(1) ($2.289(1) \text{ \AA}$) and Au(1)–S(2) ($2.294(1) \text{ \AA}$) bond distances found in the solid structure are in good agreement with those reported in the literature [4,5]. No Au(I)–Au(I) intra- or inter-molecular interactions have been observed in the crystal structure. The complex is not luminescent in solid state or in solution.

Figure 2. Molecular structure of the compound **2**, with the atom labeling scheme (hydrogen atoms omitted for clarity).



A selection of bond lengths and angles for complex **2** are summarized in Table 2.


Table 2. Selected bond lengths (\AA) and angles ($^\circ$) for **2**.

Distances (\AA)		Angles ($^\circ$)	
Au(1)–S(1)	2.289(1)	S(1)–Au(1)–S(2)	174.8(1)
S(2)–C(10)	1.729(3)	Au(1)–S(1)–C(1)	111.5(1)
S(2)–Au(1)	2.294(1)		
S(1)–C(1)	1.709(2)		

With the gold(III) complex $[\text{Au}(\text{C}_6\text{F}_5)_3(\text{tht})]$ (molar ratio = 1:2) the dinuclear derivative $[\{\text{Au}(\text{C}_6\text{F}_5)_3\}_2(\text{Et}_4\text{todit})]$ (**3**) was obtained (Scheme 1). The yellow solid is soluble in dichloromethane, acetone or diethyl ether, but insoluble in hexane. Complex **3** has been characterized by $^1\text{H-NMR}$ and $^{19}\text{F-NMR}$ spectroscopy and mass spectrometry. Single crystals were obtained by slow diffusion of hexane in a solution of complex **3** in 1,2-dichloroethane. Crystal structure has been elucidated and the proposed stoichiometry was confirmed, although (crystal system, triclinic; space group, P-1; cell: a, 12.6131; b, 13.2567; c, 18.3299 \AA ; α , 78.727; β , 74.371; γ , 82.645 $^\circ$; V, 2885.41; Z, 2) the quality of the data was not good. Thus, no comparison of bond distances and angles with other

complexes can be made, although a molecular diagram is shown in Figure 3. The two gold centers present a distorted square-planar geometry and each one is bonded to the C=S groups of the thioimidazol rings and to three pentafluorophenyl groups.

Figure 3. Molecular structure of the compound **3** (hydrogen atoms omitted for clarity).



3. Experimental Section

3.1. Instrumentation

Mass spectra were recorded on a BRUKER (Bremen, Germany) ESQUIRE 3000 PLUS, with the electrospray (ESI) technique and on a BRUKER (Bremen, Germany) MICROFLEX (MALDI-TOF), with a Dithranol or a T-2-(3-(4-*t*-butyl-phenyl)-2-methyl-2-propenylidene)malononitrile matrix. Room-temperature NMR spectra were recorded with a Bruker (Karlsruhe,, Germany) ARX 400 spectrometer (^1H , 400 MHz and ^{19}F , 376.5 MHz). The chemical shifts are reported in ppm relative to the residual solvent peak [^1H (CD_3) $_2\text{CO}$: 2.05] and CFCl_3 , respectively.

3.2. Starting Materials

Et_4tudit [13,14], $[\text{Au}(\text{C}_6\text{F}_5)(\text{tht})]$ [15], $[\text{Au}(\text{tht})_2]\text{OTf}$ [16,17] and $[\text{Au}(\text{C}_6\text{F}_5)_3(\text{tht})]$ [15], were prepared according to literature procedures. Other starting materials and solvents were purchased from commercial suppliers and used as received unless otherwise stated.

3.3. General Procedure for the Synthesis of the Complexes 1–3

Synthesis of $[\{\text{Au}(\text{C}_6\text{F}_5)\}_2(\text{Et}_4\text{tudit})]$ (**1**). A solution of $[\text{Au}(\text{C}_6\text{F}_5)(\text{tht})]$ (83.7 mg, 0.18 mmol) in dichloromethane (5 mL) was added to a solution of Et_4tudit (40.4 mg, 0.09 mmol) in dichloromethane (5 mL). The mixture was stirred for two hours at room temperature and the solution was filtered through Celite. The complex **1** was obtained as white crystals by slow diffusion of hexane in the reaction mixture (92.8 mg, yield, 86.7%). ^1H -NMR (CD_3COCD_3 , 400 MHz): δ 1.33 (6H, t, CH_3 , $J(\text{HH})$ 7.1 Hz),

1.59 (6H, t, CH₃, $J(\text{HH})$ 7.1 Hz), 4.21 (2H, m, CH₂), 4.82 (4H, dt, CH₂, $J(\text{HH})$ 14.0 and 7.1 Hz), 4.98 (2H, m, CH₂). ¹⁹F-NMR (CD₃COCD₃, 377 MHz): δ -165.6 (4F, m, *m*-F), -163.3 (2F, t, *p*-F, $J(\text{FF})$ 19.8 Hz), -117.8 (4F, m, *o*-F). MS (MALDI⁺): m/z 1163.3 [(M)⁺ 0.8%, *calculated*: 1163.9], 1101.4 [(M - 2S + 2H)⁺ 4.0%], 933.3 [(M - 2S - C₆F₅)⁺ 36.8%], 569.2 [(M - 2S - C₁₂F₁₀Au)⁺ 100%]. Analytical data: C₂₆H₂₀Au₂F₁₀N₄S₆ (1163.92) requires C, 26.81; H, 1.73; N, 4.81; S, 16.52; found C, 26.72; H, 1.71; N, 4.66; S, 16.22.

Synthesis of [Au(Et₄todit)]₂(OTf)₂ (**2**). [Ag(tht)(OTf)] (51.2 mg, 0.15 mmol) was added to a solution of [AuCl(tht)] (47.6 mg, 0.15 mmol) in dichloromethane (25 mL). The mixture was stirred for three hours at room temperature and the AgCl precipitated was filtered off through Celite. To the filtered mixture a solution of Et₄todit (58.4 mg, 0.13 mmol) in dichloromethane (10 mL) was added dropwise and stirred for one hour at room temperature. The slightly yellow solution was concentrated under reduced pressure and solid was precipitated with hexane. The product was collected by filtration and recrystallized from dichloromethane/hexane (78.7 mg, yield: 75%). ¹H-NMR (400 MHz, CD₃COCD₃) δ 4.56 (4H, m, CH₂), 3.91 (4H, m, CH₂), 1.45 (6H, t, CH₃, $J(\text{HH})$ 7.1 Hz), 1.19 (6H, t, CH₃, $J(\text{HH})$ 7.1 Hz). MS (MALDI⁺): m/z 1414.9 [(M - CF₃SO₃)⁺ 4%] *calculated*: 1414.89, 1266 [(M - 2CF₃SO₃)⁺ 10%], *calculated*: 1265.94. Analytical data: C₃₀H₄₀Au₂F₆N₈O₆S₁₄ (1563.84) requires C, 23.02; H, 2.58; N, 7.16; S, 28.67; found C, 23.14; H, 2.44; N, 7.01; S, 28.17.

Synthesis of [{Au(C₆F₅)₃]₂(Et₄todit)] (**3**). [Au(C₆F₅)₃(tht)] (95.4 mg, 0.12 mmol) was added to a solution of Et₄todit (26.5 mg, 0.06 mmol) in dichloromethane (8 mL) and the mixture was stirred for two hours at room temperature. The slightly yellow solution was concentrated under reduced pressure and solid was precipitated with hexane. The yellow product was collected by filtration and recrystallized from dichloromethane/hexane (51.0 mg, yield: 46%). ¹H-NMR (400 MHz, CD₃COCD₃) δ 4.90 (2H, m, CH₂), 4.59 (4H, m, CH₂), 4.26 (2H, m, CH₂), 1.44 (12H, m, CH₃). ¹⁹F-NMR (377 MHz, CD₃COCD₃) δ -122.42 (4F, m, *o*-F), -122.57 (4F, m, *o*-F), -124.17 (4F, m, *o*-F), -159.42 (4F, t, $J(\text{FF})$ 19.7 Hz), -160.53 (2F, t, $J(\text{FF})$ 19.5 Hz), -163.04 (4F, m, *m*-F), -164.67 (8F, m, *m*-F). MS (MALDI⁺): m/z 1263.3 [(M - 3C₆F₅ - 2S)⁺ 10.1%] *calculated*: 1267.0, 1101.2 [(M - 4C₆F₅ - 2S)⁺ 8.7%], 933.3 [(M - 2S - 5C₆F₅)⁺ 11.1%], 736.2 [(M - 3S - 6C₆F₅ + 2H)⁺ 26.3%]. Analytical data: C₅₀H₂₀Au₂F₃₀N₄S₆ (1831.89) requires C, 32.76; H, 1.10; N, 3.06; S, 10.50; found C, 32.34; H, 1.44; N, 3.01; S, 10.17.

3.4. Crystallography

Crystals were mounted in inert oil on glass fibers and transferred to the cold gas stream of Xcalibur (Agilent Technologies, Waldbronn, Germany) Oxford Diffraction (**3**) diffractometer equipped with a low-temperature attachment. Data were collected using monochromated Mo K α radiation ($\lambda = 0.71073$ Å). Scan type ω . Absorption correction based on multiple scans were applied using spherical harmonics implemented in SCALE3 ABSPACK [18] scaling algorithm. The structures were solved by direct methods and refined on F^2 using the program SHELXL-97 [19], All non-hydrogen atoms were refined anisotropically. Refinements were carried out by full-matrix least-squares on F^2 for all data. Further details of the data collection and refinement are given in Table 3. CCDC-1006288 (**1**) and 1006289 (**2**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre [20].

Table 3. Details of Data Collection and Structure Refinement for Complexes **1** and **2**.

Compound	1	2·2CH₂Cl₂
Chemical Formula	C ₂₆ H ₂₀ Au ₂ F ₁₀ N ₄ S ₁₂	C ₃₂ H ₄₄ Au ₂ Cl ₄ F ₆ N ₈ O ₆ S ₁₄
Appearance	Colorless plate	Colorless needle
Crystal size/mm	0.42 × 0.23 × 0.06	0.20 × 0.08 × 0.06
Crystal system	Triclinic	Triclinic
Space group	P-1	P-1
<i>a</i> /Å	9.97740(10)	9.7616(2)
<i>b</i> /Å	12.1758(2)	12.9096(3)
<i>c</i> /Å	14.0916(2)	13.0537(3)
α /°	73.7730(10)	63.202(2)
β /°	82.3940(10)	77.979(2)
γ /°	88.3680(10)	76.346(2)
<i>U</i> /Å ³	1629.16(4)	1416.97(5)
<i>Z</i>	2	1
<i>D_c</i> /g cm ⁻³	2.374	2.034
<i>M</i>	1164.75	1735.33
<i>F</i> (000)	1096	844
<i>T</i> /°C	-173	-173
2 θ _{max} /°	51	51
μ (Mo- <i>K</i> α)/mm ⁻¹	9.465	5.943
Transmission	0.6005, 0.1092	0.7169, 0.3828
No. of reflections measured	29873	27048
No. of unique reflections	6032	5254
<i>R</i> _{int}	0.038	0.018
<i>R</i> ^a (<i>F</i> > 4 σ (<i>F</i>))	0.024	0.045
<i>wR</i> ² (<i>F</i> ² , all refl.)	0.066	0.175
No. of reflections used	6032	5254
No. of parameters	437	329
<i>S</i>	1.059	1.041
Max. $\Delta\rho$ /eÅ ⁻³	1.77	0.90

4. Conclusions

The reaction of the ligand Et₄todit with some gold(I) or gold(III) complexes leads to the dinuclear [$\{\text{Au}(\text{C}_6\text{F}_5)\}_2(\text{Et}_4\text{todit})$] (**1**), $[\text{Au}(\text{Et}_4\text{todit})_2(\text{OTf})_2]$ (**2**) or [$\{\text{Au}(\text{C}_6\text{F}_5)_3\}_2(\text{Et}_4\text{todit})$] (**3**). They represent some of the few examples of thione gold derivatives, and because the stability of the complexes the bond Au-S within these complexes may be strong. They do not contain intra- or inter-molecular gold-gold interactions, probably because steric effects of the ligand, and they are not luminescent in solid state or in solution.

Acknowledgments

The authors gratefully acknowledge the Ministry of Economy and Competitiveness (CTQ2013-48635-C2-1-P) and Aragon Government-European Social Funds (E77) for financial

support. Massimiliano Arca, Vito Lippolis and Francesco Caddeo also thank University of Cagliari for financial support.

Author Contributions

The synthesis and spectroscopic characterization of the new complexes were performed by Francesco Caddeo and Vanesa Fernández-Moreira. The synthesis of the ligand was carried out by Vito Lippolis and Massimiliano Arca. The X-ray structures were carried out by M. Concepción Gimeno. The expertise in gold chemistry and methodology was provided by M. Concepción Gimeno and Antonio Laguna. Data analysis and preparation of the manuscript were made by all the authors.

Conflicts of Interest

The authors declare no conflict of interest.

References

1. Jia, W.G.; Huang, Y.B.; Lin, Y.J.; Wang, G.L.; Jin, G.-X. Nickel complexes and cobalt coordination polymers with organochalcogen (S, Se) ligands bearing an *N*-methylimidazole moiety: Syntheses, structures, and properties. *Eur. J. Inorg. Chem.* **2008**, 4063–4073.
2. Jia, W.G.; Huang, Y.B.; Lin, Y.J.; Jin, G.-X. Syntheses and structures of half-sandwich iridium(III) and rhodium(III) complexes with organochalcogen (S, Se) ligands bearing *N*-methylimidazole and their use as catalysts for norbornene polymerization. *Dalton Trans.* **2008**, 5612–5620.
3. Williams, D.J.; vanDerveer, D.; Jones, R.L.; Menaldino, D.S. Main group metal halide complexes with sterically hindered thioureas XI. Complexes of antimony(III) and bismuth(III) chlorides with a new bidentate thiourea 1,1'-methylenebis(3-methyl-2H-imidazole-2-thione). *Inorg. Chim. Acta* **1989**, *165*, 173–178.
4. Bigoli, F.; Deplano, P.; Devillanova, F.A.; Lippolis, V.; Mercuri, M.L.; Pellinghelli, M.A.; Trogu, E.F. Synthesis, X-ray and spectroscopic characterization of $[\text{SnI}_2(\text{mbit})_2](\text{I}_3)_2 \cdot 2/3\text{I}_2$ obtained through the one-step reaction of $\text{mbit} \cdot 2\text{I}_2$ with tin metal powder ($\text{mbit} = 1,1'$ -bis(3-methyl-4-imidazoline-2-thione)methane). *Inorg. Chim. Acta* **1998**, *267*, 115–121.
5. Williams, D.J.; Shilatifard, A.; VanDerveer, D.; Lipscomb, L.A.; Jones, R.L. Main group metal halide complexes with sterically hindered thioureas XIII. Crystallographic study of a unique cross-linked polymeric dichlorolead(II) complex with 1,1'-methylenebis(3-methyl-2(3*H*)-imidazolethione). *Inorg. Chim. Acta* **1992**, *202*, 53–57.
6. Silva, R.M.; Smith, M.D.; Gardinier, J.R. Anion- and solvent-directed assembly in silver bis(thioimidazolyl)methane chemistry and the silver–sulfur interaction. *Inorg. Chem.* **2006**, *45*, 2132–2142.
7. Aroz, M.T.; Gimeno, M.C.; Kulcsar, M.; Laguna, A.; Lippolis, V. Group 11 complexes with imidazoline-2-thione or selone derivatives. *Eur. J. Inorg. Chem.* **2011**, 2884–2894.

8. Aragoni, M.C.; Arca, M.; Devillanova, F.A.; Isaia, F.; Lippolis, V.; Mancini, A.; Pala, L.; Slawin, A.M.Z.; Woollins, J.D. First example of an infinite polybromide 2D-network. *Chem. Commun.* **2003**, 2226–2227.
9. Mancini, A.; Aragoni, M.C.; Bricklebank, N.; Castellano, C.; Demartin, F.; Isaia, F.; Lippolis, V.; Pintus, A.; Arca, M. Formation of T-shaped *versus* charge-transfer molecular adducts in the reactions between bis(thiocarbonyl) donors and Br₂ and I₂. *Chem. Asian J.* **2013**, *8*, 639–647.
10. Mancini, A.; Aragoni, M.C.; Bingham, L.; Castellano, C.; Coles, S.L.; Demartin, F.; Hursthouse, M.B.; Isaia, F.; Lippolis, V.; Maninchedda, G.; *et al.* Reactivity of fluoro-substituted bis(thiocarbonyl) donors with diiodine: And XRD, FT-Raman, and DFT investigation. *Chem. Asian J.* **2013**, *8*, 2071–3078.
11. Bigoli, F.; Pellinghelli, M.A.; Deplano, P.; Trogu, E.F. Complexes of 4,5,6,7-tetrathiocino[1,2-*b*:3,4-*b'*]diimidazolyl-1,3,8,10-tetraethyl-2,9-dithione (Et₄todit) with group IIb metal halides. Crystal and molecular structure of (Cd(II)Et₄toditCl₂)_n. *Inorg. Chim. Acta* **1990**, *170*, 245–249.
12. Bigoli, F.; Pellinghelli, M.A.; Deplano, P.; Trogu, E.F. Preparation and characterization of polymeric compounds of copper(I) halogenides with Et₄todit = 4,5,6,7-tetrathiocino[1,2-*b*:3,4-*b'*]diimidazolyl-1,3,8,10-tetraethyl-2,9-dithione. Crystal and molecular structures of [Cu(I)(Et₄todit)I]_n and [Cu(I)(Et₄todit)I]_n·*n*/2Me₂CO. *Inorg. Chim. Acta* **1991**, *182*, 33–39.
13. Bigoli, F.; Pellinghelli, A.; Atzei, D.; Deplano, P.; Trogu, E.F. Synthesis of some 4,5,6,7-tetrathiocino[1,2-*b*:3,4-*b'*]diimidazolyl-1,3,8,10-tetrasubstituted-2,9-dithiones and crystal structure of the tetraethyl derivative. *Phosphorus Sulfur* **1988**, *37*, 189–194.
14. Aragoni, M.C.; Arca, M.; Demartin, F.; Devillanova, F.A.; Garau, A.; Isaia, F.; Lelj, F.; Lippolis, V.; Verani, G. New [M(R,R'timdt)₂] Metal-dithiolenes and related compounds (M = Ni, Pd, Pt; R,R'timdt = monoanion of disubstituted imidazolidine-2,4,5-trithiones): An Experimental and theoretical investigation. *J. Am. Chem. Soc.* **1991**, *121*, 7098–7107.
15. Usón, R.; Laguna, A. Polyaryl Derivatives of gold(I), silver(I) and gold(III). In *Organometallic Syntheses*; King, R.B., Eisch, J.J., Eds.; Elsevier: Amsterdam, Holland, 1986; Volume 3, pp. 322–342.
16. Usón, R.; Laguna, A.; Navarro, A.; Parish, R.V.; Moore, L.S. Synthesis and reactivity of perchlorate bis(tetrahydrothiophen)gold(I). ¹⁹⁷Au Mössbauer spectra of three-coordinate gold(I) complexes. *Inorg. Chim. Acta* **1986**, *112*, 295–208.
17. Usón, R.; Laguna, A.; Laguna, M.; Jiménez, J.; Gómez, M.P.; Sainz, A.; Jones, P.G. Gold complexes with heterocyclic thiones as ligands. X-ray structure determination of [Au(C₅H₅NS)₂]ClO₄. *J. Chem. Soc. Dalton Trans.* **1990**, 3457–3463.
18. *CrysAlisPro*, Version 1.171.35.11; Multi-scans absorption correction with SCALE3 ABSPACK scaling algorithm; Agilent Technologies: Waldbronn, Germany, 2011.
19. Sheldrick, G.M. *SHELXL-97, Program for Crystal Structure Refinement*; University of Göttingen: Göttingen, Germany, 1997.
20. The Cambridge Crystallographic Data Centre. Available online: http://www.ccdc.cam.ac.uk/data_request/cif (accessed on 2 June 2014).