



Proton pump inhibitor treatment and lower gastrointestinal bleeding: Balancing risks and benefits

Alberto Lué, Angel Lanas

Alberto Lué, Angel Lanas, Digestive Diseases Service, University Clinic Hospital Lozano Blesa, Avenida San Juan Bosco, 50009 Zaragoza, Spain

Alberto Lué, Angel Lanas, IIS Aragon, Avenida San Juan Bosco, 50009 Zaragoza, Spain

Angel Lanas, University of Zaragoza, Calle de Pedro Cerbuna, 50009 Zaragoza, Spain

Angel Lanas, CIBERehd, Av. Monforte de Lemos, 28029 Madrid, Spain

Author contributions: Lué A and Lanas A contributed to this paper.

Conflict-of-interest statement: Lué A and Lanas A declare no conflict of interest related to this publication.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Correspondence to: Angel Lanas, MD, DSc, Professor, Chairman, Digestive Diseases Service, University Clinic Hospital Lozano Blesa, Avenida San Juan Bosco, 50009 Zaragoza, Spain. alanas@unizar.es
 Telephone: +34-97-6768886
 Fax: +34-97-6768846

Received: August 27, 2016
 Peer-review started: August 31, 2016
 First decision: September 20, 2016
 Revised: September 28, 2016
 Accepted: November 12, 2016
 Article in press: November 13, 2016
 Published online: December 28, 2016

Abstract

Proton pump inhibitors (PPIs) represent a milestone in the treatment of acid-related diseases, and are the mainstay in preventing upper gastrointestinal bleeding in high-risk patients treated with nonsteroidal anti-inflammatory drugs (NSAIDs) or low-dose aspirin. However, this beneficial effect does not extend to the lower gastrointestinal tract. PPIs do not prevent NSAID or aspirin-associated lower gastrointestinal bleeding (LGB). PPIs may increase both small bowel injury related to NSAIDs and low-dose aspirin treatment and the risk of LGB. Recent studies suggested that altering intestinal microbiota by PPIs may be involved in the pathogenesis of NSAID-enteropathy. An increase in LGB hospitalization rates may occur more frequently in older patients with more comorbidities and are associated with high hospital resource utilization, longer hospitalization, and increased mortality. Preventive strategies for NSAID and aspirin-associated gastrointestinal bleeding should be directed toward preventing both upper and lower gastrointestinal damage. Future research should be directed toward identifying patients at low-risk for gastrointestinal events associated with the use of NSAIDs or aspirin to avoid inappropriate PPI prescribing. Alternatively, the efficacy of new pharmacologic strategies should be evaluated in high-risk groups, with the aim of reducing the risk of both upper and lower gastrointestinal bleeding in these patients.

Key words: Proton pump inhibitor; Small bowel; Small bowel; Lower gastrointestinal bleeding; Nonsteroidal anti-inflammatory drugs

© **The Author(s) 2016.** Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: Proton pump inhibitors (PPIs) reduce the risk of upper, but not lower gastrointestinal bleeding (LGB) in patients receiving nonsteroidal anti-inflammatory drugs (NSAIDs) or low-dose aspirin. PPIs could

exacerbate small bowel damage related to NSAIDs and low-dose aspirin, which contributes to an increased risk of LGB possibly related to pathological modifications of small bowel microbiota. LGB is a life-threatening condition, especially in older patients with comorbidities treated with NSAIDs, aspirin, or anticoagulants. No accepted treatments exist for decreasing the risk of LGB in these patients. Future research is needed on reducing inappropriate PPI use and evaluating possible pharmacologic interventions to decrease the risk of gastrointestinal bleeding.

Lué A, Lanás A. Protons pump inhibitor treatment and lower gastrointestinal bleeding: Balancing risks and benefits. *World J Gastroenterol* 2016; 22(48): 10477-10481 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v22/i48/10477.htm> DOI: <http://dx.doi.org/10.3748/wjg.v22.i48.10477>

INTRODUCTION

Proton pump inhibitors (PPIs) are one of the most prescribed drugs worldwide. PPIs are used to treat acid-related disorders like gastro-esophageal reflux disease, peptic ulcer, peptic ulcer bleeding, and *Helicobacter pylori* infection when combined with antibiotics. PPIs should also be considered in any patient with risk factors for gastrointestinal bleeding who receives treatment with nonsteroidal anti-inflammatory drugs (NSAIDs) or prophylaxis for cardiovascular events with aspirin or other antiplatelet agents^[1].

Treatment with NSAIDs, low-dose aspirin, other antiplatelet drugs, and anticoagulation are associated with an increased risk of both upper and lower gastrointestinal bleeding (LGB). Concomitant treatment with these compounds and PPIs has been associated with a decreased risk of upper gastrointestinal bleeding (UGB) but, obviously, not with an increased risk of LGB. The beneficial effect of PPIs is not expected to occur beyond the duodenum because NSAID-gastropathy, but not NSAID-enteropathy, is a pH-dependent phenomenon and gastroduodenal mucosal protection by PPIs is due mainly to their antisecretory effects^[2].

PROTON PUMP INHIBITOR TREATMENT AND RISK OF LOWER GASTROINTESTINAL BLEEDING

In a recent case-control study that included 1008 patients hospitalized for gastrointestinal bleeding, treatment with NSAIDs, low-dose aspirin, other antiplatelet agents, and anticoagulants increased the risk of both UGB and LGB. In this work, concomitant use of PPI was associated with a reduced risk of UGB but not LGB^[3]. In fact, there was a significant increase in the risk of LGB with PPI use, which has also been detected by other authors^[3-8]. Due to the intrinsic nature

of observational studies, the finding was explained as probably due to a "confounding by indication" bias. PPI use could be a marker of a group of patients that have an increased risk of LGB because of their clinical characteristics rather than the PPI use itself^[3]. In a previous study, we observed that among patients on dual antiplatelet treatment who received concomitant PPI treatment, gastrointestinal bleeding events were more frequent in the lower gastrointestinal tract (74% vs 26%)^[9]. There are several potential reasons explaining this change in the pattern of gastrointestinal events, which include (1) a very high protective effect of PPIs on the gastroduodenal mucosa associated with a profound decrease in the rate of UGB and a relative (but not absolute) increase in the proportion of LGB in low-dose aspirin users; (2) an absence of the mucosal protective effect of PPIs beyond the duodenum; (3) a direct harmful effect of low-dose aspirin on the small bowel; (4) promotion of bleeding of pre-existing lesions by these compounds; and (5) exacerbation of NSAID- and aspirin-enteropathy by PPI treatment^[10-14].

Lately, a hypothesis proposed that PPIs exacerbate NSAID and low-dose aspirin-associated small bowel injury by inducing changes in the intestinal microbiota^[15-17]. A multicenter, cross-sectional study including data collected from endoscopic capsule explorations from 205 patients treated with low-dose aspirin for 3 mo showed that 57.6% of patients had at least one mucosal lesion. In the multivariate analysis, concomitant PPI use (OR = 2.04; 95%CI: 1.05-3.97) and use of enteric-coated aspirin (OR = 4.05; 95%CI: 1.49-11.0) were independent risk factors for the presence of mucosal injury^[8]. Similar results were observed in patients receiving NSAIDs. In a recent randomized clinical trial, 57 healthy volunteers were allocated to receive celecoxib plus placebo or rabeprazole for 2 wk. The patients were evaluated by capsule endoscopy at the start and end of treatment. In the PPI group, the authors observed an increased rate of small bowel injury (44.7% vs 16.7%, $P = 0.04$)^[6].

Some authors suggest that acid gastric suppression induced by PPI treatment is related to changes in small bowel microbiota and explain, at least in part, NSAID-induced enteropathy. It is well known that in addition to bile acids^[18] and inhibition of cyclooxygenase activity^[19], bacterial mucosal translocation and the consequent activation of the innate inflammatory cascade are important factors in the pathogenesis of NSAID-enteropathy^[20,21]. In an animal model, Wallace *et al*^[17] demonstrated that concomitant treatment with PPIs and NSAIDs resulted in a higher rate of small bowel mucosal ulceration and bleeding. The authors observed that PPI treatment was related to a modification in the small bowel microbiota, consisting of a decrease in jejunal Actinobacteria and Bifidobacteria spp. In this study, the restoration of physiologic microbiota with probiotic treatment prevented intestinal injury^[17].

Similar findings have been observed in human subjects. A small randomized clinical trial, including 25

patients treated with low-dose enteric-coated aspirin and omeprazole randomized to either placebo or probiotic treatment with *Lactobacillus casei* for 3 mo, observed that patients in the probiotic group had a significant decrease in the number of mucosal breaks when evaluated by capsule endoscopy ($P = 0.039$)^[15].

All these data suggest that PPI treatment not only does not protect the lower gastrointestinal tract from NSAID- and aspirin-related injury, but that they may increase the intestinal mucosal damage and the risk of LGB. We observed that the LGB hospitalization rate has increased significantly in the past years^[9,12,13]. These LGB events occur more frequently in older patients, who usually have a higher number of comorbidities, and are associated with a longer hospital stay and higher mortality rates^[12]. These events are associated with greater use of hospital resources and also complicate the management of patients at discharge^[12]. A considerable percentage of patients did not resume prophylactic treatment with low-dose aspirin or anticoagulants after an acute episode of LGB, increasing the risk of serious cardiovascular events^[22]. A recent retrospective analysis including 295 patients with a previous diagnosis of LGB and low-dose aspirin observed that discontinuing low-dose aspirin is related to a decreased risk in LGB recurrence (6.9% vs 18.9%; $P = 0.007$), but increased the rates of mortality (26.7% vs 8.2%, $P = 0.001$) and cardiovascular events (36.5% vs 22.8%, $P = 0.017$)^[23]. This study is a clear example of how difficult it is to find the right balance between risks and benefits in these patients. On one side, there is a need to reduce or prevent cardiovascular events and on the other side there is an increased risk of LGB.

BALANCING RISK AND BENEFITS OF PROTON PUMP INHIBITOR TREATMENT

Current clinical practice faces several dilemmas in patients who need NSAIDs or aspirin, since damage to the upper gastrointestinal tract must be prevented without injuring the lower gastrointestinal tract. This clinical dilemma is especially important in patients who need, for example, preventive treatments against cardiovascular disease and who have had a previous LGB episode. Alternative therapeutic approaches with non-aspirin antiplatelet agents poses a similar risk of LGB as aspirin and many not be a valid option^[3]. The same is true for patients who need NSAIDs, although COX-2 selective NSAIDs have been shown to be less damaging to the lower gastrointestinal tract than traditional NSAIDs^[2,11,24]. The addition of a PPI to celecoxib increases intestinal mucosal damage^[6,10,17]. Therefore, future efforts must concentrate on finding strategies that help clinicians in the decision making process. The first relevant issue is to improve overall PPI prescribing and avoid use of PPIs in patients at low risk of gastrointestinal complications who take NSAIDs, aspirin, other antiplatelet agents, or anticoagulants.

PPIs are prescribed too often in routine clinical practice, where at least 50% of prescriptions are for non-approved indications^[25]. Therefore, the first step should be to promote the proper use of these drugs. Identifying patients at low risk of gastrointestinal events who may not benefit from treatment with PPIs is essential. Tools that could help physicians in the decision making process are now available^[26]. On the other hand, we have to reconsider whether current guidelines are valid or respond to problems in daily clinical practice. For example, the PPI treatment indication in patients receiving NSAIDs is based on data from clinical trials of chronic NSAID users. However, observational studies suggest that the risk of gastrointestinal bleeding may be higher when the drug use is recent, and most patients take these drugs for short periods of time^[11]. We also need more evidence to evaluate the risks of upper and lower gastrointestinal bleeding with new antiplatelet drugs, new oral direct anticoagulants, and the multiple combinations available in order to better assess the risks and benefits.

On the other hand, in patients at high risk for LGB, preventive strategies are needed. This group includes older patients, with a great number of comorbidities who need more complex management^[9,12,23]. As mentioned above, in these patients LGB represents a serious and life threatening complication. Currently, there is no clear effective pharmacological treatment to prevent LGB in NSAID, aspirin, or anticoagulant users. Some studies have evaluated the effect of the prostaglandin analogue misoprostol on small bowel injury induced by NSAIDs^[7,27]. Other mucosal protectants such as rebamipide^[28-31], irsogladine^[32,33], or geranylgeranylacetone^[34,35] have been tested and are only available in the Asia-Pacific region. Their potential beneficial effects in these patients still need to be evaluated in new studies.

In the same way, some studies have evaluated the efficacy of probiotics in preventing lower gastrointestinal tract injury among patients treated with NSAIDs or low-dose aspirin^[15,16]. The probiotic administered in these two trials was *Lactobacillus casei* in patient treated with low-dose aspirin, while patient treated with NSAIDs received the probiotic mixture VSL#3[®], that includes *Streptococcus thermophilus*, *Bifidobacterium breve*, *Bifidobacterium longum*, *Bifidobacterium infantis*, *Lactobacillus paracasei*, *Lactobacillus delbrueckii subsp. Bulgaricus*, *Lactobacillus acidophilus* and *Lactobacillus plantarum*^[15,16]. The results are promising and suggest that pharmacological prophylaxis could be provided by this strategy. However, more studies with a larger number of patients are also needed. At present, there is not sufficient evidence to recommend a specific probiotic strain to prevent lower gastrointestinal injury. However some bacteria have demonstrated antiinflammatory activity and a protective effect on intestinal mucosa. For example: (1) *Lactobacillus GG* protect mucosal cells from apoptosis and increase mucosal integrity; (2) *Lactobacillus plantarum*

promotes mucine from epithelial cells; (3) *Lactobacillus reuterii* and *Lactobacillus casei* have demonstrated to suppress tumor necrosis factor production *in vivo*; and (4) *Streptococcus thermophilus* and *Lactobacillus acidophilus* could prevent bacterial translocation^[36].

CONCLUSION

In conclusion, PPIs represent a milestone in the treatment of acid-related disease. Moreover, PPIs significantly reduce the risk of UGB in patients treated with NSAIDs, low-dose aspirin, other antiplatelet agents, or anticoagulants. This beneficial effect is not observed in the lower gastrointestinal tract. Furthermore, recent studies suggest that PPIs may in fact increase small bowel injury and contribute to the observed increase of LGB in the past years; however, results are controversial. Clinicians very often face clinical dilemmas and need to balance the risks and benefits of treatment. The use of aspirin in preventing cardiovascular disease and the risk of gastrointestinal bleeding is one of these situations. The dilemma is even more difficult to resolve when one of the drugs (PPI) can have a dual and opposite effect (beneficial to the upper GI tract but damaging to the lower GI tract). Clinicians have to evaluate carefully the risk of upper gastrointestinal bleeding in patients that have a previous episode of LGB in order to discontinue inappropriate administration of PPI. Future research should be focused on identifying patients at low risk for upper gastrointestinal bleeding to avoid inappropriate PPI treatment and to evaluate the efficacy of new pharmacologic strategies in high-risk groups. The aim of this strategy is to reduce the risk of LGB in these patients while at the same time providing the maximal beneficial effect to the upper gastrointestinal tract.

REFERENCES

- 1 **Bhatt DL**, Scheiman J, Abraham NS, Antman EM, Chan FK, Furberg CD, Johnson DA, Mahaffey KW, Quigley EM. ACCF/ACG/AHA 2008 expert consensus document on reducing the gastrointestinal risks of antiplatelet therapy and NSAID use: a report of the American College of Cardiology Foundation Task Force on Clinical Expert Consensus Documents. *Circulation* 2008; **118**: 1894-1909 [PMID: 18836135 DOI: 10.1161/circulationaha.108.191087]
- 2 **Scarpignato C**, Lanas A, Blandizzi C, Lems WF, Hermann M, Hunt RH. Safe prescribing of non-steroidal anti-inflammatory drugs in patients with osteoarthritis--an expert consensus addressing benefits as well as gastrointestinal and cardiovascular risks. *BMC Med* 2015; **13**: 55 [PMID: 25857826 DOI: 10.1186/s12916-015-0285-8]
- 3 **Lanas Á**, Carrera-Lasfuentes P, Arguedas Y, García S, Bujanda L, Calvet X, Ponce J, Perez-Aisa Á, Castro M, Muñoz M, Sostres C, García-Rodríguez LA. Risk of upper and lower gastrointestinal bleeding in patients taking nonsteroidal anti-inflammatory drugs, antiplatelet agents, or anticoagulants. *Clin Gastroenterol Hepatol* 2015; **13**: 906-912.e2 [PMID: 25460554 DOI: 10.1016/j.cgh.2014.11.007]
- 4 **Lanas A**, Sostres C. PPI therapy. The small bowel, low-dose aspirin and PPIs--should we be concerned? *Nat Rev Gastroenterol Hepatol* 2014; **11**: 458-460 [PMID: 25001974 DOI: 10.1038/nrgastro.2014.120]

- 5 **Nagata N**, Niikura R, Aoki T, Sakurai T, Moriyasu S, Shimbo T, Sekine K, Okubo H, Watanabe K, Yokoi C, Yanase M, Akiyama J, Uemura N. Effect of proton-pump inhibitors on the risk of lower gastrointestinal bleeding associated with NSAIDs, aspirin, clopidogrel, and warfarin. *J Gastroenterol* 2015; **50**: 1079-1086 [PMID: 25700638 DOI: 10.1007/s00535-015-1055-2]
- 6 **Washio E**, Esaki M, Machata Y, Miyazaki M, Kobayashi H, Ishikawa H, Kitazono T, Matsumoto T. Proton Pump Inhibitors Increase Incidence of Nonsteroidal Anti-Inflammatory Drug-Induced Small Bowel Injury: A Randomized, Placebo-Controlled Trial. *Clin Gastroenterol Hepatol* 2016; **14**: 809-815.e1 [PMID: 26538205 DOI: 10.1016/j.cgh.2015.10.022]
- 7 **Watanabe T**, Sugimori S, Kameda N, Machida H, Okazaki H, Tanigawa T, Watanabe K, Tominaga K, Fujiwara Y, Oshitani N, Higuchi K, Arakawa T. Small bowel injury by low-dose enteric-coated aspirin and treatment with misoprostol: a pilot study. *Clin Gastroenterol Hepatol* 2008; **6**: 1279-1282 [PMID: 18995219 DOI: 10.1016/j.cgh.2008.06.021]
- 8 **Endo H**, Sakai E, Taniguchi L, Kessoku T, Komiya Y, Ezuka A, Kawamura H, Taguri M, Higurashi T, Ohkubo H, Yamada E, Takahashi H, Inamori M, Maeda S, Sakaguchi T, Hata Y, Nagase H, Nakajima A. Risk factors for small-bowel mucosal breaks in chronic low-dose aspirin users: data from a prospective multicenter capsule endoscopy registry. *Gastrointest Endosc* 2014; **80**: 826-834 [PMID: 24830581 DOI: 10.1016/j.gie.2014.03.024]
- 9 **Casado Arroyo R**, Polo-Tomas M, Roncalés MP, Scheiman J, Lanas A. Lower GI bleeding is more common than upper among patients on dual antiplatelet therapy: long-term follow-up of a cohort of patients commonly using PPI co-therapy. *Heart* 2012; **98**: 718-723 [PMID: 22523056 DOI: 10.1136/heartjnl-2012-301632]
- 10 **Goldstein JL**, Eisen GM, Lewis B, Gralnek IM, Zlotnick S, Fort JG. Video capsule endoscopy to prospectively assess small bowel injury with celecoxib, naproxen plus omeprazole, and placebo. *Clin Gastroenterol Hepatol* 2005; **3**: 133-141 [PMID: 15704047]
- 11 **Lanas A**, García-Rodríguez LA, Arroyo MT, Gomollón F, Feu F, González-Pérez A, Zapata E, Bástida G, Rodrigo L, Santolaria S, Güell M, de Argila CM, Quintero E, Borda F, Piqué JM. Risk of upper gastrointestinal ulcer bleeding associated with selective cyclo-oxygenase-2 inhibitors, traditional non-aspirin non-steroidal anti-inflammatory drugs, aspirin and combinations. *Gut* 2006; **55**: 1731-1738 [PMID: 16687434 DOI: 10.1136/gut.2005.080754]
- 12 **Lanas A**, García-Rodríguez LA, Polo-Tomás M, Ponce M, Alonso-Abreu I, Perez-Aisa MA, Perez-Gisbert J, Bujanda L, Castro M, Muñoz M, Rodrigo L, Calvet X, Del-Pino D, Garcia S. Time trends and impact of upper and lower gastrointestinal bleeding and perforation in clinical practice. *Am J Gastroenterol* 2009; **104**: 1633-1641 [PMID: 19574968 DOI: 10.1038/ajg.2009.164]
- 13 **Lanas A**, García-Rodríguez LA, Polo-Tomás M, Ponce M, Quintero E, Perez-Aisa MA, Gisbert JP, Bujanda L, Castro M, Muñoz M, Del-Pino MD, Garcia S, Calvet X. The changing face of hospitalisation due to gastrointestinal bleeding and perforation. *Aliment Pharmacol Ther* 2011; **33**: 585-591 [PMID: 21205256 DOI: 10.1111/j.1365-2036.2010.04563.x]
- 14 **Smecoul E**, Pinto Sanchez MI, Suarez A, Argonz JE, Sugai E, Vazquez H, Litwin N, Piazuelo E, Meddings JB, Bai JC, Lanas A. Low-dose aspirin affects the small bowel mucosa: results of a pilot study with a multidimensional assessment. *Clin Gastroenterol Hepatol* 2009; **7**: 524-529 [PMID: 19249402 DOI: 10.1016/j.cgh.2008.12.019]
- 15 **Endo H**, Higurashi T, Hosono K, Sakai E, Sekino Y, Iida H, Sakamoto Y, Koide T, Takahashi H, Yoneda M, Tokoro C, Inamori M, Abe Y, Nakajima A. Efficacy of *Lactobacillus casei* treatment on small bowel injury in chronic low-dose aspirin users: a pilot randomized controlled study. *J Gastroenterol* 2011; **46**: 894-905 [PMID: 21556830 DOI: 10.1007/s00535-011-0410-1]
- 16 **Montalto M**, Gallo A, Curigliano V, D'Onofrio F, Santoro L, Covino M, Dalvai S, Gasbarrini A, Gasbarrini G. Clinical trial: the effects of a probiotic mixture on non-steroidal anti-inflammatory drug enteropathy - a randomized, double-blind, cross-over, placebo-

- controlled study. *Aliment Pharmacol Ther* 2010; **32**: 209-214 [PMID: 20384610 DOI: 10.1111/j.1365-2036.2010.04324.x]
- 17 **Wallace JL**, Syer S, Denou E, de Palma G, Vong L, McKnight W, Jury J, Bolla M, Bercik P, Collins SM, Verdu E, Ongini E. Proton pump inhibitors exacerbate NSAID-induced small intestinal injury by inducing dysbiosis. *Gastroenterology* 2011; **141**: 1314-1322 [PMID: 21745447 DOI: 10.1053/j.gastro.2011.06.075]
 - 18 **Petruzzelli M**, Vacca M, Moschetta A, Cinzia Sasso R, Palasciano G, van Erpecum KJ, Portincasa P. Intestinal mucosal damage caused by non-steroidal anti-inflammatory drugs: role of bile salts. *Clin Biochem* 2007; **40**: 503-510 [PMID: 17321514 DOI: 10.1016/j.clinbiochem.2007.01.015]
 - 19 **Takeuchi K**, Tanaka A, Kato S, Amagase K, Satoh H. Roles of COX inhibition in pathogenesis of NSAID-induced small intestinal damage. *Clin Chim Acta* 2010; **411**: 459-466 [PMID: 20074562 DOI: 10.1016/j.cca.2009.12.026]
 - 20 **Blackler RW**, Gemici B, Manko A, Wallace JL. NSAID-gastroenteropathy: new aspects of pathogenesis and prevention. *Curr Opin Pharmacol* 2014; **19**: 11-16 [PMID: 24929967 DOI: 10.1016/j.coph.2014.05.008]
 - 21 **Lanas A**, Scarpignato C. Microbial flora in NSAID-induced intestinal damage: a role for antibiotics? *Digestion* 2006; **73** Suppl 1: 136-150 [PMID: 16498262 DOI: 10.1159/000089789]
 - 22 **Sostres C**, Lanas A. Epidemiology of Low Dose Aspirin Damage in the Lower Gastrointestinal Tract. *Curr Pharm Des* 2015; **21**: 5094-5100 [PMID: 26369682]
 - 23 **Chan FK**, Leung Ki EL, Wong GL, Ching JY, Tse YK, Au KW, Wu JC, Ng SC. Risks of Bleeding Recurrence and Cardiovascular Events With Continued Aspirin Use After Lower Gastrointestinal Hemorrhage. *Gastroenterology* 2016; **151**: 271-277 [PMID: 27130815 DOI: 10.1053/j.gastro.2016.04.013]
 - 24 **Chan FK**, Lanas A, Scheiman J, Berger MF, Nguyen H, Goldstein JL. Celecoxib versus omeprazole and diclofenac in patients with osteoarthritis and rheumatoid arthritis (CONDOR): a randomised trial. *Lancet* 2010; **376**: 173-179 [PMID: 20638563 DOI: 10.1016/s0140-6736(10)60673-3]
 - 25 **Lanas A**. We Are Using Too Many PPIs, and We Need to Stop: A European Perspective. *Am J Gastroenterol* 2016; **111**: 1085-1086 [PMID: 27166129 DOI: 10.1038/ajg.2016.166]
 - 26 **Lanas A**, Polo-Tomás M, Casado-Arroyo R. The aspirin cardiovascular/gastrointestinal risk calculator--a tool to aid clinicians in practice. *Aliment Pharmacol Ther* 2013; **37**: 738-748 [PMID: 23413984 DOI: 10.1111/apt.12240]
 - 27 **Kojima Y**, Takeuchi T, Ota K, Harada S, Edogawa S, Narabayashi K, Nouda S, Okada T, Kakimoto K, Kuramoto T, Inoue T, Higuchi K. Effect of long-term proton pump inhibitor therapy and healing effect of irsogladine on nonsteroidal anti-inflammatory drug-induced small-intestinal lesions in healthy volunteers. *J Clin Biochem Nutr* 2015; **57**: 60-65 [PMID: 26236102 DOI: 10.3164/jcfn.15-32]
 - 28 **Niwa Y**, Nakamura M, Ohmiya N, Maeda O, Ando T, Itoh A, Hirooka Y, Goto H. Efficacy of rebamipide for diclofenac-induced small-intestinal mucosal injuries in healthy subjects: a prospective, randomized, double-blinded, placebo-controlled, cross-over study. *J Gastroenterol* 2008; **43**: 270-276 [PMID: 18458842 DOI: 10.1007/s00535-007-2155-4]
 - 29 **Kurata S**, Nakashima T, Osaki T, Uematsu N, Shibamori M, Sakurai K, Kamiya S. Rebamipide protects small intestinal mucosal injuries caused by indomethacin by modulating intestinal microbiota and the gene expression in intestinal mucosa in a rat model. *J Clin Biochem Nutr* 2015; **56**: 20-27 [PMID: 25834302 DOI: 10.3164/jcfn.14-67]
 - 30 **Kurokawa S**, Katsuki S, Fujita T, Saitoh Y, Ohta H, Nishikawa K, Sato Y, Sato Y, Ohira K, Yamada M, Kato M. A randomized, double-blinded, placebo-controlled, multicenter trial, healing effect of rebamipide in patients with low-dose aspirin and/or non-steroidal anti-inflammatory drug induced small bowel injury. *J Gastroenterol* 2014; **49**: 239-244 [PMID: 23595613 DOI: 10.1007/s00535-013-0805-2]
 - 31 **Watanabe T**, Takeuchi T, Handa O, Sakata Y, Tanigawa T, Shiba M, Naito Y, Higuchi K, Fujimoto K, Yoshikawa T, Arakawa T. A multicenter, randomized, double-blind, placebo-controlled trial of high-dose rebamipide treatment for low-dose aspirin-induced moderate-to-severe small intestinal damage. *PLoS One* 2015; **10**: e0122330 [PMID: 25874951 DOI: 10.1371/journal.pone.0122330]
 - 32 **Isomura Y**, Yamaji Y, Yamada A, Watanabe Y, Suzuki H, Kobayashi Y, Yoshida S, Watabe H, Hirata Y, Yoshida H, Koike K. Irsogladine improves small-intestinal injuries in regular users of nonsteroidal anti-inflammatory drugs. *Gastrointest Endosc* 2014; **80**: 118-125 [PMID: 24518124 DOI: 10.1016/j.gie.2013.12.030]
 - 33 **Kuramoto T**, Umegaki E, Nouda S, Narabayashi K, Kojima Y, Yoda Y, Ishida K, Kawakami K, Abe Y, Takeuchi T, Inoue T, Murano M, Tokioka S, Higuchi K. Preventive effect of irsogladine or omeprazole on non-steroidal anti-inflammatory drug-induced esophagitis, peptic ulcers, and small intestinal lesions in humans, a prospective randomized controlled study. *BMC Gastroenterol* 2013; **13**: 85 [PMID: 23672202 DOI: 10.1186/1471-230x-13-85]
 - 34 **Umegaki E**, Kuramoto T, Kojima Y, Nouda S, Ishida K, Takeuchi T, Inoue T, Tokioka S, Higuchi K. Geranylgeranylacetone, a gastro-mucoprotective drug, protects against NSAID-induced esophageal, gastroduodenal and small intestinal mucosal injury in healthy subjects: A prospective randomized study involving a comparison with famotidine. *Intern Med* 2014; **53**: 283-290 [PMID: 24531083]
 - 35 **Xiong L**, Huang X, Li L, Yang X, Liang L, Zhan Z, Ye Y, Chen M. Geranylgeranylacetone protects against small-intestinal injuries induced by diclofenac in patients with rheumatic diseases: a prospective randomized study. *Dig Liver Dis* 2015; **47**: 280-284 [PMID: 25660821 DOI: 10.1016/j.dld.2015.01.005]
 - 36 **Ganesh BP**, Versalovic J. Luminal Conversion and Immunoregulation by Probiotics. *Front Pharmacol* 2015; **6**: 269 [PMID: 26617521 DOI: 10.3389/fphar.2015.00269]

P- Reviewer: Hreinsson JP, Lakatos PL, Rabago L, Souza JLS, Tan HJ
S- Editor: Gong ZM **L- Editor:** A **E- Editor:** Liu WX





Published by **Baishideng Publishing Group Inc**

8226 Regency Drive, Pleasanton, CA 94588, USA

Telephone: +1-925-223-8242

Fax: +1-925-223-8243

E-mail: bpgoffice@wjgnet.com

Help Desk: <http://www.wjgnet.com/esps/helpdesk.aspx>

<http://www.wjgnet.com>



ISSN 1007-9327

