

ARRANGEMENTS OF HYPERSURFACES AND BESTVINA-BRADY GROUPS

ENRIQUE ARTAL BARTOLO, JOSÉ IGNACIO COGOLLUDO-AGUSTÍN, AND DANIEL MATEI

ABSTRACT. We show that quasi-projective Bestvina-Brady groups are fundamental groups of complements to hyperplane arrangements. Furthermore we relate other normal subgroups of right-angled Artin groups to complements to arrangements of hypersurfaces. We thus obtain examples of hypersurface complements whose fundamental groups satisfy various finiteness properties.

INTRODUCTION

An important open question in the topology of complex algebraic varieties is J.-P. Serre's problem to characterize fundamental groups of smooth algebraic varieties (cf. [31]): Which finitely presented groups are quasi-projective, i.e. isomorphic to the fundamental group of a smooth connected quasi-projective variety (the complement of a normal crossing divisor in a smooth, connected, complex projective variety)? One could also consider an analogous question for quasi-Kähler groups, that is, for fundamental groups of hypersurface complements in a compact Kähler manifold. On the other hand, one could explore particular versions of Serre's question by restricting either the ambient projective variety or the divisor, or both. For example: Which groups can be realized as fundamental groups of hypersurface (in particular hyperplane) arrangement complements in a projective space? or Which finitely presented groups are projective, i.e. isomorphic to fundamental groups of smooth connected projective varieties?

In this paper we are interested in quasi-projective groups from the point of view of their finiteness properties. It is well known that such groups are finitely presented, but much less is known about their higher dimensional finiteness properties. In [34], C.T.C. Wall introduced general finiteness properties of groups and CW-complexes. A group G is said to be of type F_n if it has an Eilenberg-MacLane complex $K(G, 1)$ with finite n -skeleton. Clearly G is finitely generated if and only if it is F_1 and finitely presented if and only if it is F_2 . An interesting example of a finitely generated group which is not finitely presented was given by Stallings

2010 *Mathematics Subject Classification.* 14F35, 20F65, 14F45, 20F36, 55R55, 14N20, 14J70.

Key words and phrases. Bestvina-Brady quasi-projective groups, Artin kernels, hyperplane and toric arrangements, quasifibrations.

Partially supported by MTM2010-21740-C02-02. The third author is also partially supported by grant CNCSIS PNII-IDEI 1188/2008 and FMI 53/10 (Gobierno de Aragón).

in [33]. While Wall's property F_n has a substantial geometric meaning, it is not easily detectable by homological algebraic methods.

In [5], Bieri introduced homological counterparts of Wall's finiteness conditions. A group G is said to be of type FP_n if the $\mathbb{Z}G$ -module \mathbb{Z} admits a projective resolution which is finitely generated in dimensions $\leq n$. The condition FP_n is clearly weaker than F_n . Note also that G is of type FP_1 if and only if it is finitely generated, and that G is of type FP_2 if it is finitely presented. But, as shown by Bestvina and Brady [3] FP_2 does not imply finite presentation. Nevertheless, if G is F_2 then G is FP_n if and only if it is F_n .

The first example of a group which is finitely presented but not of type FP_3 was given by Stallings in [33]. Afterwards Bieri [4, 5] generalized Stallings' example to an infinite family. In a nutshell, let $\mathbb{F}_2 \times \cdots \times \mathbb{F}_2$ be the direct product of $r + 1$ free groups, each of rank 2. Then the kernel N of the map taking each generator to $1 \in \mathbb{Z}$ is F_r but not F_{r+1} . Stallings considered the cases $r \leq 2$. In [3], Bestvina and Brady generalized the Bieri-Stallings family by presenting a systematic way of constructing groups N of type F_r , but not of type F_{r+1} . This construction was further extended in [22, 6], and revisited in [29]. The input is a finite graph Γ , together with a character χ of its associated right-angled Artin group A_Γ . The generalized Bestvina-Brady group N_Γ^χ , or Artin kernel, is then defined as the kernel of the homomorphism $\chi : A_\Gamma \rightarrow \mathbb{Z}$, see [29]. The Bestvina-Brady groups N of [3] are recovered as $N_\Gamma := N_\Gamma^\chi$ for the diagonal character $\chi : A_\Gamma \rightarrow \mathbb{Z}$ sending each generator to 1.

The class of Bestvina-Brady groups N_Γ which are quasi-projective was determined by Dimca, Papadima and Suciu in [12] as those corresponding to either trees or special multipartite complete graphs Γ , i.e. $N_\Gamma = \mathbb{F}_{n_0} \times \cdots \times \mathbb{F}_{n_r}$ with either $r \geq 2$ or $n_i = 1$ for some i . In [12], it was proved in particular that the Stallings-Bieri groups are quasi-projective. It was noticed by Matei and Suciu in 2004, and pointed out in [27], that the Stallings group may be realized as the fundamental group of the complement of a complex line arrangement in \mathbb{P}^2 , first considered by Arvola in unpublished work from 1992, see [30]. In [20] and [1] we reported on our generalization of this observation. More precisely, we exhibited in [20], for each quasi-projective Bestvina-Brady group N_Γ , a complex line arrangement in \mathbb{P}^2 having N_Γ as the fundamental group of the complement, and in [1] higher dimensional hyperplane arrangements realizing just the Stallings-Bieri groups. In this work we will present the geometric construction that was announced in [20, 1] (see Theorem 3.2 for more details).

Theorem 1. *Let Γ be a graph on v vertices, and suppose N_Γ is a quasi-projective Bestvina-Brady group. Then N_Γ is the fundamental group of the complement of a hyperplane arrangement \mathcal{A}_Γ in \mathbb{P}^r , for some $r < v$ depending only on Γ .*

Theorem 1 answers negatively [2, Question 2.10] in Bestvina's problem list. Indeed, the fundamental group $G = N_\Gamma$ of the complement of \mathcal{A}_Γ cannot have a finite $K(G, 1)$, as long as $H_{r+1}(N_\Gamma)$ has infinite rank. Moreover, as we shall see later in Corollary 3.3, \mathcal{A}_Γ may be given by

real equations, and by taking a generic plane section of \mathbb{P}^r one can realize N_Γ as the fundamental group of the complement to a real line arrangement in \mathbb{P}^2 . For another approach to this result we refer to the work of Cohen, Falk and Randell [8].

Our geometric realization of the quasi-projective Bestvina-Brady group N_Γ as the fundamental group of an arrangement complement M , gives more than just the fundamental group of $K = K(N_\Gamma, 1)$. In fact M has the homotopy type of the r -skeleton of K .

Theorem 2. *Let \mathcal{A}_Γ be a hyperplane arrangement in \mathbb{P}^r with complement $M = M_\Gamma$ such that $N_\Gamma = \pi_1(M)$ is a quasi-projective Bestvina-Brady group. Then M has the homotopy type of the r -skeleton of $K(N_\Gamma, 1)$. Suppose N_Γ is not the direct product of free groups. Then we have that*

- (1) *the homology groups $H_i(M)$ and $H_i(N_\Gamma)$ are isomorphic for $i \leq r$.*
- (2) *the homotopy groups $\pi_i(M)$ vanish in the range $1 < i < r$.*
- (3) *the homotopy group $\pi_r(M)$ does not vanish, in fact its group of coinvariants $\pi_r^*(M)$ under the π_1 -action is isomorphic to $H_{r+1}(N_\Gamma)$.*

One could compare this with the results in [13] where examples are constructed of smooth projective manifolds M of dimension $r \geq 2$ with $\pi_1(M)$ having the finiteness properties of the Bestvina-Brady groups, and $\pi_r(M)$ the next non-vanishing higher homotopy group.

The results in Theorems 1 and 2 extend to quasi-projective Artin kernels beyond the Bestvina-Brady groups. We will show that an Artin kernel N_Γ^χ is quasi-projective if Γ is a multipartite complete graph with N_Γ quasi-projective, and the character χ of the Artin group $A_\Gamma = \mathbb{F}_{n_0} \times \cdots \times \mathbb{F}_{n_r}$ sends each free factor \mathbb{F}_{n_i} to a non-zero integer. More precisely, N_Γ^χ will be realized as the fundamental group of an arrangement of hypersurfaces in an r -dimensional weighted projective space whose weights are given by the images under χ of the generators of A_Γ . In practice though we will work not in the weighted projective space, but in its associated torus. Thus all Artin kernel N_Γ^χ realized by our construction turn out to be fundamental groups of complements to arrangements of hypertori in an ambient algebraic torus. The basic properties of such toric arrangements are similar to those of hyperplane arrangements, see monographs [10] and [25].

The description of the quasi-projective Bestvina-Brady groups provided by Theorems 1 and 2 yields a large class of hyperplane arrangements that exhibit new topological features. More precisely, the affine arrangements in \mathbb{C}^{r+1} induced by the projective arrangements \mathcal{A}_Γ , cannot be hypersolvable since their groups are not of type F. Thus the techniques described in [26] can no longer be used to determine the homotopy and homology groups of these arrangement complements. Compare also with the earlier results of [30], obtained using the Lefschetz hyperplane theorem. This suggests the following open problem.

Problem. Characterize the quasiprojective Artin kernels.

In the following we will explain the motivation and the basic steps for the geometric construction that will produce quasi-projective Artin kernels. Suppose X is a smooth (quasi)-projective

variety. Then any finite covering of X is again a (quasi)-projective manifold (see for instance Griffiths and Harris [15, p. 192] for the projective case and Namba [24, Lemma 2.1.2] for the general case using the Grauert-Remmert extension theorem of unramified coverings of complex spaces).

In general, the (quasi)-projectivity property is lost when passing to infinite coverings. Suppose $\pi_1(X)$ admits an epimorphism $\nu : \pi_1(X) \rightarrow \mathbb{Z}$. Let Y be the regular connected infinite cyclic covering of X associated with ν . We are interested in deciding whether or not Y is a (quasi)-projective manifold. The first obstruction would be the finite presentability of $\pi_1(Y)$. We may wonder what other properties of X or $\pi_1(X)$ survive after infinite cyclic coverings. For example the $K(\pi, 1)$ property for X , or other finiteness properties of $\pi_1(X)$ such as F_n , F , or FP_n .

In this context, the general problem that motivates our construction is the following:

Problem. Identify (quasi)-projective manifolds X admitting infinite cyclic coverings which have the homotopy of a (quasi)-projective manifold. Alternatively, identify quasi-projective groups G having normal subgroups N , such that $G/N \cong \mathbb{Z}$, which are again quasi-projective.

A very natural idea is to start with a polynomial mapping $f : \mathbb{C}^n \rightarrow \mathbb{C}$, and analyze the restriction $f : X \rightarrow \mathbb{C}^*$, where $X = \mathbb{C}^n \setminus V(f)$. The bifurcation set $\Lambda = \Lambda_f$ of f is the minimal set to be removed so that $f : X \setminus f^{-1}(\Lambda) \rightarrow \mathbb{C}^* \setminus \Lambda$ is a locally trivial fibration. Then Λ contains $C = C_f$ the set of critical values of f (in general only as a proper subset).

Now suppose that f has finitely many singular points, and it behaves like a proper map, in particular the critical set is the bifurcation set of f . If $n \geq 3$, then one can obtain, by standard arguments, an exact sequence of groups

$$1 \rightarrow \pi_1(F_a) \rightarrow \pi_1(X) \rightarrow \mathbb{Z} \rightarrow 1,$$

for any fiber $F_a = f^{-1}(a)$, $a \in \mathbb{C}^*$ of f , whether smooth or singular, see [12, 32].

In the present paper a different approach will be presented. Suppose f and g are two polynomial mappings $\mathbb{C}^n \rightarrow \mathbb{C}$, and consider the restrictions $f|_T : T \rightarrow \mathbb{C}^*$, where $T = \mathbb{C}^n \setminus V(f)$, and $f|_X : X \rightarrow \mathbb{C}^*$, where $X = \mathbb{C}^n \setminus V(g)$.

If $f|_T : T \rightarrow \mathbb{C}^*$ is a trivial fiber bundle, and $V(g)$ is transverse (in the stratified sense) to the fibers $f|_T^{-1}(a)$ with $a \in \mathbb{C}^* \setminus \Lambda$, then $f|_{X_\Lambda} : X_\Lambda \rightarrow \mathbb{C}^* \setminus \Lambda$, $X_\Lambda := X \setminus f^{-1}(\Lambda)$, is again a fiber bundle (Λ is the bifurcation set of $f : X \rightarrow \mathbb{C}^*$).

Let Σ be the union of the strata in $V(g)$ which are not transverse to $f^{-1}(\Lambda)$, and denote its dimension by $s := \dim \Sigma$. Then, using Dold-Thom fibration theory one can see that the map $f : X \rightarrow \mathbb{C}^*$ behaves like a fibration up to dimension $n - s - 1$ (details will be given in Section 5).

Now suppose X is aspherical and $s = 0$. It follows that all the fibers $F_a = f^{-1}(a)$ share the same homotopy type up to dimension $(n - 2)$, in particular, if $n \geq 4$, the same fundamental group. Furthermore, if the monodromy of $f : X \rightarrow \mathbb{C}^*$ about the points in Λ is trivial, we then have an exact sequence $1 \rightarrow \pi_1(F) \rightarrow \pi_1(X) \rightarrow \mathbb{Z} \rightarrow 1$. We will show that indeed that is the

case for the constructions of interest to us, by analyzing the homotopy short exact sequence $1 \rightarrow \pi_1(F) \rightarrow \pi_1(X_\Lambda) \rightarrow \pi_1(\mathbb{C}^* \setminus \Lambda) \rightarrow 1$.

The paper is organized as follows: in the first two sections the main objects are introduced. More specifically the group theoretical objects, such as Artin groups, Artin kernels, and their presentations, are described in Section 1. The geometrical objects such as hyperplane and toric arrangements are described in Section 2 together with their homological properties. The main results extending Theorems 1 and 2 are stated in Section 3 and their proofs will be shown in the following two sections. Section 4 describes the arrangements realizing the quasi-projective groups of Section 1. Finally, the homotopy and homology properties of these groups are discussed in Section 5 in terms of the arrangements using quasifibration theory.

1. ARTIN KERNELS

Let Γ be a finite simplicial graph, i.e., $\Gamma := (V_\Gamma, E_\Gamma)$, where V_Γ is a finite set, say $\{0, 1, \dots, s\}$, and $E_\Gamma \subset \{A \subset V_\Gamma \mid \#A = 2\}$; let L_Γ be the *flag simplicial complex* generated by Γ , where $J := \{j_1, \dots, j_k\}$ is a simplex of L_Γ if and only if the complete graph on the vertices J is a subgraph of Γ . The *right-angled Artin group* A_Γ associated with Γ is the group with generators $\sigma_1, \dots, \sigma_s$ and relations $\sigma_i \sigma_j = \sigma_j \sigma_i$, one for each edge $\{i, j\} \in E_\Gamma$ of Γ .

The Eilenberg-MacLane space $K(A_\Gamma, 1)$ has two remarkable incarnations as toric spaces. First, consider the real $(s+1)$ -torus $T = (\mathbb{S}^1)^{s+1}$ with its standard CW-complex structure, where k -cells are in bijection with k -subsets of $\{0, 1, \dots, s\}$. From a simplicial complex $L = L_\Gamma$ one constructs the subcomplex T_L of T , by retaining only the cells corresponding to the simplices of L . It is well known that T_L is a $K(A_\Gamma, 1)$ -space [7]. In the language of [11], T_L is the *moment-angle complex* $Z_L(\mathbb{S}^1)$, obtained by gluing subtori of T according to the simplicial structure of L . A similar gluing process, but using the complex $(s+1)$ -torus $\mathbb{T}^{s+1} = (\mathbb{C}^*)^{s+1}$ and complex subtori accordingly, produces $Z_L(\mathbb{C}^*)$, which we denote by \mathbb{T}_L .

The properties of the construction Z_L , see [11], ensure that T_L and \mathbb{T}_L have the same homotopy type. It also follows that $T_{L*M} = T_L \times T_M$, and respectively $\mathbb{T}_{L*M} = \mathbb{T}_L \times \mathbb{T}_M$, where $L * M$ is the join of L and M .

Example 1.1. Let us consider two standard examples of graphs on $s+1$ vertices, namely, K_{s+1} the complete graph, and \bar{K}_{s+1} the graph with no edges. The flag complexes are respectively an s -dimensional simplex and \bar{K}_{s+1} itself. As for the associated right-angled Artin groups, $A_{K_{s+1}} \cong \mathbb{Z}^{s+1}$ and $A_{\bar{K}_{s+1}} \cong \mathbb{F}_{s+1}$, the free group of rank $s+1$. The toric complexes are $T_{K_{s+1}} = T \simeq \mathbb{T} = \mathbb{T}_{K_{s+1}}$ and $T_{\bar{K}_{s+1}} = \bigvee^{s+1} \mathbb{S}^1 \simeq \bigvee^{s+1} \mathbb{C}^* = \mathbb{T}_{\bar{K}_{s+1}}$ respectively. Note that $\bigvee^{s+1} \mathbb{C}^*$ has the homotopy type of \mathbb{C} punctured $s+1$ times. We denote by \mathbb{C}_{s+1}^* an $(s+1)$ -punctured \mathbb{C} . Thus we have $\mathbb{T}_{\bar{K}_{s+1}} \simeq \mathbb{C}_{s+1}^*$.

Example 1.2. Let $\bar{\mathbf{n}} := (n_0, \dots, n_r)$ be an $(r+1)$ -tuple of positive integers, $n_i \geq 1$. Denote by $\bar{K}_{\bar{\mathbf{n}}}$ the multipartite graph defined by the join $\bar{K}_{n_0} * \dots * \bar{K}_{n_r}$. The associated flag complex

$L = L_{\bar{K}_{\mathbf{n}}}$ has the homotopy type of a wedge of $m := \prod_{i=0}^r (n_i - 1)$ spheres $\bigvee^m \mathbb{S}^r$. The Artin group $A_{\bar{K}_{\mathbf{n}}}$ is the product $\mathbb{F}_{n_0} \times \cdots \times \mathbb{F}_{n_r}$ of free groups, and $\mathbb{T}_L \simeq \mathbb{C}_{n_0}^* \times \cdots \times \mathbb{C}_{n_r}^*$.

We will consider here a class of subgroups of the Artin group A_Γ , generalizing the Bestvina-Brady groups of [3]. An *Artin kernel* is nothing but the kernel of an epimorphism $\chi : A_\Gamma \rightarrow \mathbb{Z}$, see [29]. More precisely, given an epimorphism $\chi : A_\Gamma \rightarrow \mathbb{Z}$ such that $\chi(\sigma_i) \in \mathbb{Z}$, we denote its kernel by N_Γ^χ .

Theorem 1.3 ([3, 22, 6, 29]). *The Artin kernel N_Γ^χ is finitely presented if and only if L_Γ^χ is simply-connected. Moreover, N_Γ^χ is of type F_r (respectively FP_r) if and only if the flag complex L_Γ^χ is $(r-1)$ -connected (respectively $(r-1)$ -acyclic). In addition, N_Γ^χ is of type FP_r if and only if $\dim_{\mathbb{K}} H_{\leq r}(N_\Gamma^\chi, \mathbb{K}) < \infty$, for any field \mathbb{K} .*

In this paper we will always assume that all $\chi(\sigma_i)$ are non-zero. This implies that L_Γ^χ in Theorem 1.3 is the usual flag complex L_Γ on Γ . The particular case where χ is the epimorphism $\chi : A_\Gamma \rightarrow \mathbb{Z}$ defined by $\chi(\sigma_i) := 1$, the kernel $N_\Gamma := \ker \chi$ is known as the *Bestvina-Brady group* associated with Γ . As noted, it is finitely presented if and only if L is simply-connected. In that case, N_Γ admits a commutator-relators presentation, see [28]. As a consequence, one immediately obtains that $b_1(N_\Gamma) = |V_\Gamma| - 1$ and $b_2(N_\Gamma) = |E_\Gamma| - |V_\Gamma| + 1$.

The Artin kernels that interest us most here are those associated with the multipartite graph $\bar{K}_{\mathbf{n}}$. For this special case, we introduce now some notation. After a suitable labeling, we denote the set of vertices of $\bar{K}_{\mathbf{n}}$ by $\bigcup_{k=0}^r \{(k, i) \mid 1 \leq i \leq n_k\}$ where (k, i) is a vertex of \bar{K}_{n_k} . The corresponding generators of $A_{\bar{K}_{\mathbf{n}}}$ are denoted by $\sigma_{k,i}$.

Let $\mathbf{d} := (d_0, \dots, d_r) \in \mathbb{Z}^r$ and consider the character $\chi_{\mathbf{d}} : A_{\bar{K}_{\mathbf{n}}} \rightarrow \mathbb{Z}$ defined by $\chi_{\mathbf{d}}(\sigma_{k,i}) := d_k$.

Corollary 1.4. *The Artin kernel $N_{\mathbf{n}}^{\mathbf{d}} := N_{\bar{K}_{\mathbf{n}}}^{\chi_{\mathbf{d}}}$ is of type F_r , but not FP_{r+1} , if all $n_i > 1$.*

Proof. Since $L_{\bar{K}_{\mathbf{n}}} \simeq \bigvee^m \mathbb{S}^r$ is $(r-1)$ -connected, but $H_r(L_{\bar{K}_{\mathbf{n}}}) \neq 0$, it follows from Theorem 1.3 that $N_{\mathbf{n}}^{\mathbf{d}}$ is of type F_r , but not FP_{r+1} , if all $n_i > 1$. \square

Note that, as soon as $r \geq 2$, the Artin kernel $N_{\mathbf{n}}^{\mathbf{d}}$ is finitely presented and thus the F_m and FP_m properties are equivalent. For $N = N_{\bar{K}_{\mathbf{n}}}$ one obtains:

$$b_1(N) = \sum_{i=0}^r n_i - 1, \quad b_2(N) = \sum_{0 \leq i < j \leq r} n_i n_j - \sum_{i=0}^r n_i + 1.$$

Recall the definition of the Poincaré series as the generating function of the Betti numbers, that is, $P_G(t) := \sum_{k=0}^{\infty} b_k(G)t^k$. From the results of [29] it follows that the truncated Poincaré polynomials of A_Γ and N_Γ are related as

$$P_{A_\Gamma}(t) \equiv (1+t)P_{N_\Gamma}(t) \pmod{t^{r+1}}.$$

It is easily seen that the r -th truncation of the Poincaré polynomial of $A = A_{\bar{K}_{\bar{n}}}$ is given by

$$P_A(t) \equiv \prod_{i=0}^r (1 + n_i t) \pmod{t^{r+1}}.$$

This gives the Betti numbers b_k of $N = N_{\bar{K}_{\bar{n}}}$, in the range $0 \leq k < r$:

$$b_k(N) = \sum_{p=0}^k (-1)^{k-p} \sum_{0 \leq i_1 < \dots < i_p \leq r} n_{i_1} \dots n_{i_p}.$$

1.5. Finite presentations of N_Γ . Assume Γ is a finite simplicial graph with $s + 1$ vertices. As a general fact, an infinite presentation for N_Γ can be obtained by the Reidemeister-Schreier algorithm. A set of generators can be taken to be

$$\alpha_{i;m} := \sigma_0^m \sigma_i \sigma_0^{-m-1}, \quad 0 < i \leq s, m \in \mathbb{Z}.$$

In order to simplify the notations, we will also consider $\alpha_{0,m} := 1$. Then a complete set of relators is given by:

$$\alpha_{i;m} \alpha_{j;m+1} = \alpha_{j;m} \alpha_{i;m+1}, \quad \{i, j\} \in E_\Gamma, \quad m \in \mathbb{Z}.$$

Suppose now that $\Gamma = \bar{K}_{\bar{n}}$ where $2 \leq n_0 \leq \dots \leq n_r$. If $r \geq 2$ then the flag complex L is simply-connected and Bestvina-Brady work ensures that the countable presentation above can be reduced to a finite one; for $r = 1$ the group will be finitely generated. In the following we approach this question in a more direct way.

The generators of A_Γ are denoted by $\sigma_{k,i}$, for $1 \leq i \leq n_k$ and $0 \leq k \leq r$. An infinite set of generators of N_Γ is given by

$$\alpha_{k,i;m} := \sigma_{0,1}^m \sigma_{k,i} \sigma_{0,1}^{-m-1}, \quad 0 \leq k \leq r, \quad m \in \mathbb{Z},$$

with the trivial relations $\alpha_{0,1;m} = 1, \forall m \in \mathbb{Z}$. An additional set of complete relators is given by:

$$(1.1) \quad \alpha_{k,i;m} \alpha_{\ell,j;m+1} = \alpha_{\ell,j;m} \alpha_{k,i;m+1}, \quad 0 \leq k < \ell \leq r, \quad 1 \leq i \leq n_k, \quad 1 \leq j \leq n_\ell, \quad m \in \mathbb{Z}.$$

We consider three different cases separately. If $r = 0$ then no relation exists and N_Γ is a free group in $\alpha_{0,i;m}$, where $2 \leq i \leq n_0$ and $m \in \mathbb{Z}$. As expected, this group is not finitely generated.

For $r = 1$, the relations (1.1) for $(k, i) = (0, 1)$ allow to conclude that $\alpha_{1,j;m}$ does not depend on m . The relations (1.1) for $\ell = 1$ imply that

$$(1.2) \quad \alpha_{0,i;m} = \alpha_{0,i;0}^m, \quad 2 \leq i \leq n_0, \quad m \in \mathbb{Z}, \quad 1 \leq j \leq n_1.$$

Using $j = 1$ in (1.2) allows one to show that the group is generated by $\alpha_{0,i;0}, \alpha_{1,j;0}$ for $2 \leq i \leq n_0$ and $1 \leq j \leq n_1$; for simplicity, we drop the index 0. The remaining relations in (1.2) are rewritten as:

$$(1.3) \quad [\alpha_{0,i}, \alpha_{1,j}^m \alpha_{1,1}^{-m}] = 1, \quad 2 \leq i \leq n_0, \quad 2 \leq j \leq n_1, \quad m \in \mathbb{Z}.$$

This infinite set of relators cannot be reduced.

For $r \geq 2$ we proceed as above; the set

$$(1.4) \quad \{\alpha_{0,i;0}\}_{i=2}^{n_0} \cup \bigcup_{k=1}^r \{\alpha_{k,j;0}\}_{j=1}^{n_k}$$

is a generator system for N_Γ ; as before, we will drop the subindex 0.

To the relations (1.3) we add the relations (1.1) for $1 \leq k \leq \ell$:

$$(1.5) \quad [\alpha_{k,i}, \alpha_{\ell,j}] = 1, \quad 1 \leq k < \ell \leq r, \quad 1 \leq i \leq n_k, \quad 1 \leq j \leq n_\ell,$$

We add now the relations (1.1) for $k = 0$ and $\ell \geq 2$ and we obtain:

$$(1.6) \quad [\alpha_{0,i}, \alpha_{\ell,j}^m \alpha_{1,1}^{-m}] = 1, \quad 2 \leq \ell \leq r, \quad 2 \leq i \leq n_0, \quad 1 \leq j \leq n_\ell, \quad m \in \mathbb{Z}.$$

For $\ell \geq 2$, one has $\alpha_{\ell,j}^m \alpha_{1,1}^{-m} = (\alpha_{\ell,j} \alpha_{1,1}^{-1})^m$ and hence (1.3) and (1.6) are consequences of

$$(1.7) \quad [\alpha_{0,i}, \alpha_{k,j} \alpha_{1,1}^{-1}] = 1, \quad 1 < i \leq n_0, \quad (k = 1, \quad 2 < j \leq n_1) \text{ and } (2 \leq k \leq r, \quad 1 \leq j \leq n_k).$$

A finite presentation of N_Γ is given by

$$\left\langle (1.4) \middle| (1.5), (1.7) \right\rangle$$

1.6. Finite presentations of N^χ . We now consider presentations for $N^\chi = N_{\bar{\mathbf{n}}}^{\mathbf{d}}$, where $r \geq 2$ and $n_0, \dots, n_r \geq 2$. Fix $a_k \in \mathbb{Z}$ so that $\sum_{k=0}^r a_k d_k = 1$ and define $\tau := \prod_{k=0}^r \sigma_{k,1}^{a_k}$; note that $\chi(\tau) = 1$. Since $\sigma_{k,1}, \sigma_{\ell,1}$ commute whenever $k \neq \ell$, note that $\tau^m := \prod_{k=0}^r \sigma_{k,1}^{m a_k}$, $m \in \mathbb{Z}$. We can add τ to the generators of $A_{\bar{K}_{\bar{\mathbf{n}}}}$, adding the following relations (some of them are redundant but useful):

$$(1.8) \quad \tau = \prod_{k=0}^r \sigma_{k,1}^{a_k}, \quad [\tau, \sigma_{k,1}] = 1, \quad k = 0, \dots, r.$$

Applying the Reidemeister-Schreier method, a set of generators of N^χ is given by

$$\alpha_{k,i;m} := \tau^m \sigma_{k,i} \tau^{-m-d_k}, \quad 0 \leq k \leq r, \quad 1 \leq i \leq n_k, \quad m \in \mathbb{Z}.$$

The commutativity relations in (1.8) imply that $\alpha_{k,1;m}$ does not depend on m and hence we denote them by β_k . The remaining relation in (1.8) gives the following one for N^χ :

$$(1.9) \quad \prod_{k=0}^r \beta_k^{a_k} = 1.$$

The relations $[\sigma_{k,i}, \sigma_{\ell,j}] = 1$, $k \neq \ell$ give the following ones in the kernel N^χ

$$(1.10) \quad \alpha_{k,i;m} \alpha_{\ell,j;m+d_k} = \alpha_{\ell,j;m} \alpha_{k,i;m+d_\ell}, \quad \text{if } k \neq \ell.$$

If $i = j = 1$, (1.10) simplifies as:

$$(1.11) \quad [\beta_k, \beta_\ell] = 1, \quad k \neq \ell.$$

If $i = 1 < j$, (1.10) simplifies as:

$$(1.12) \quad \alpha_{\ell,j;m+d_k} = \beta_k^{-1} \alpha_{\ell,j;m} \beta_k, \quad \forall m \in \mathbb{Z}, \quad k \in \{0, \dots, r\}, \quad \ell \neq k, \quad j \in \{2, \dots, n_\ell\}.$$

Let us denote $e_\ell := \gcd\{d_k \mid k \neq \ell\}$ and choose integers $a_{k,\ell}$, $a_{k,k} := 0$, such that

$$e_\ell = \sum_{k=0}^r a_{k,\ell} d_k.$$

Clearly, using relations (1.12), we can reduce the infinite set of generators $\{\alpha_{k,i,m} \mid m \in \mathbb{Z}, k \in \{0, \dots, r\}, 1 \leq i \leq n_k\}$ to the finite subset

$$(1.13) \quad \{\beta_k \mid 1 \leq k \leq r\} \cup \{\alpha_{k,j;p} \mid 0 \leq p < e_k, 0 \leq k \leq r, 1 < j \leq n_k\}.$$

Let us denote $\gamma_\ell := \prod_{k=0}^r \beta_k^{a_{k,\ell}}$. Then, we obtain:

$$(1.14) \quad \alpha_{\ell,j;m+e_\ell} = \gamma_\ell^{-1} \alpha_{\ell,j;m} \gamma_\ell, \quad \forall m \in \mathbb{Z}, \ell \in \{1, \dots, r\}, j \in \{2, \dots, n_\ell\}.$$

Denote $f_{k,\ell} := \frac{d_k}{e_\ell}$, $k \neq \ell$. Combining (1.12)-(1.14), we obtain:

$$(1.15) \quad [\alpha_{\ell,j;m}, \gamma_\ell^{f_{k,\ell}} \beta_k^{-1}] = 1 \quad k \in \{0, \dots, r\}, \ell \neq k, j \in \{2, \dots, n_\ell\}, m \in \{0, \dots, e_k - 1\}.$$

Consider now the relations (1.10) for $i, j > 1$:

$$(1.16) \quad \alpha_{k,i;m} \beta_k^{-1} \alpha_{\ell,j;m} \beta_k = \alpha_{\ell,j;m} \beta_\ell^{-1} \alpha_{k,i;m} \beta_\ell, \text{ if } k \neq \ell.$$

Given $k \neq \ell$, fix $m \neq k, \ell$ such that d_m is minimal; this is possible since $r \geq 2$. Then it is enough to consider relations (1.16) for $m \in \{0, \dots, d_m - 1\}$.

The presentation of N^χ admits (1.13) as a system of generators and (1.9), (1.11), (1.15), and (1.16) as a system of relators.

Example 1.7. If $d_0 = 1$, we can choose $\tau = \sigma_{0,1}$, i.e. $a_0 = 1$ and $a_k = 0$ if $k > 0$. We get $\beta_0 = 1$, $e_k = 1$ if $k > 0$, and hence $\gamma_k = 1$ if $k > 0$. Then N^χ is generated by $\alpha_{k,i;m} = \sigma_{0,1}^m \sigma_{k,i} \sigma_{0,1}^{-m-d_k}$, for

$$\{(0, i; m) \mid 1 < i \leq n_0, 0 \leq m < e_0\} \cup \{(k, i; 0) \mid 0 < k \leq r, 1 \leq i \leq n_k\}.$$

Let us denote $\mu_{i,m} := \alpha_{0,i;m}$ and $\nu_{k,i} := \alpha_{k,i;0}$ ($k > 0$). Note also that if $k \neq \ell$ and both are different from 0 then, we can choose $m = 1$. The relations are:

$$\begin{aligned} [\nu_{k,1}, \nu_{\ell,1}] &= 1, & 0 < k < \ell, \\ [\mu_{i,m}, \gamma_0^{\frac{d_k}{e_0}} \nu_{k,1}^{-1}] &= 1, & 1 < i \leq n_0, 0 \leq m < e_0, 0 < k \leq r, \\ [\nu_{k,i}, \nu_{\ell,j}] &= 1, & 0 < k, \ell \leq r, k \neq \ell, 1 \leq i \leq n_k, 1 \leq j \leq n_\ell, \\ [\mu_{j,m}, \nu_{k,i} \nu_{k,1}^{-1}] &= 1 & 0 < k \leq r, 1 \leq i \leq n_k, 1 < j \leq n_0, 0 \leq m < d_0, \end{aligned}$$

where $\mu_{j,m+e_1} := \gamma_0^{-1} \mu_{j,m} \gamma_0$.

2. HYPERSURFACE ARRANGEMENTS AND THEIR COMPLEMENTS

In this section we introduce the arrangements of hypersurfaces that will be needed in the sequel. We will consider two classes: hyperplane arrangements and toric arrangements. In each case, we will explain how the homology of complements to such arrangements may be computed in terms of their combinatorics. We will end with computations of the Poincaré polynomial for the arrangements that we will later relate with Artin kernels.

2.1. Hyperplane arrangements.

An arrangement of hyperplanes is a collection \mathcal{A} of hyperplanes in a projective, or affine space. An arrangement \mathcal{A} in \mathbb{P}^r may be viewed as a central arrangement in \mathbb{C}^{r+1} . If M is the complement of \mathcal{A} in \mathbb{P}^r , and M' the complement in \mathbb{C}^{r+1} , we have that M' is homeomorphic to $M \times \mathbb{C}^*$. It follows that $P(M', t) = (1+t)P(M, t)$, and $\pi_1(M') = \pi_1(M) \times \mathbb{Z}$.

The Poincaré polynomial of a complement to a hyperplane arrangement \mathcal{A} is completely determined by its intersection poset $L(\mathcal{A})$, which consists of all non-empty intersections among hyperplanes in \mathcal{A} ordered by reversed inclusion and ranked by codimension.

More precisely, if \mathcal{A} is an affine hyperplane arrangement, the characteristic polynomial of the intersection poset $L(\mathcal{A})$ is defined by

$$\chi(\mathcal{A}, q) := \sum_{S \in L(\mathcal{A})} \mu(S) q^{\dim S},$$

where $\mu : L(\mathcal{A}) \rightarrow \mathbb{Z}$ is the Möbius function of the ranked poset $L(\mathcal{A})$, see [25].

Proposition 2.1. *Let \mathcal{A} be a hyperplane arrangement in \mathbb{C}^{r+1} with complement M . We then have*

$$P(M, t) = t^{r+1} \chi(\mathcal{A}, -t^{-1}).$$

The best understood arrangements are those of supersolvable type, see [25]. The complement M of such an arrangement sits at the top of a tower of linear fiber bundles with fiber a punctured complex plane. In particular M is a $K(G, 1)$ space for $G := \pi_1(M)$. A more general class, comprising both supersolvable and generic arrangements, is that of the hypersolvable arrangements, see [19]. The group G of a hypersolvable arrangement \mathcal{A} is still of type F, as it is also a group of supersolvable arrangement $\hat{\mathcal{A}}$ to which \mathcal{A} deforms.

A class of arrangements that contains both hypersolvable and non-hypersolvable examples is that of graphic arrangements. We refer to [26, 27] for the details of the discussion. One associates to a graph $\mathcal{G} = (\mathcal{V}, \mathcal{E})$ on a set of vertices $\mathcal{V} := \{0, \dots, r\}$, the arrangement $\mathcal{A}_{\mathcal{G}}$ in \mathbb{C}^{r+1} of all hyperplanes $H_{i,j} = \{w_i = w_j\}$ for $\{i, j\} \in \mathcal{E}$ an edge of \mathcal{G} . Any $\mathcal{A}_{\mathcal{G}}$ is a sub-arrangement of the braid arrangement, which corresponds to the complete graph K_{r+1} . Denote by c_p the number of complete subgraphs K_{p+1} of \mathcal{G} . Thus c_1 is the number of edges and c_2 is the number of triangles of \mathcal{G} .

An arrangement $\mathcal{A}_{\mathcal{G}}$ is supersolvable if and only if \mathcal{G} is a chordal graph (i.e. every circuit of length ≥ 4 in \mathcal{G} has a chord). For example, an arrangement $\mathcal{A}_{\mathcal{G}}$ with $c_2 = 0$ is hypersolvable. Suppose \mathcal{G} contains no complete graph K_4 as a subgraph, that is $c_3 = 0$. A sufficient condition for such an arrangement $\mathcal{A}_{\mathcal{G}}$ to be non-hypersolvable is that $0 < c_1 \leq 2c_2$.

Example 2.2. Let $\mathcal{G} = W_r$ be the (wheel) graph on $r+1$ vertices obtained by coning an r -cycle, hence \mathcal{A}_{W_r} is defined, after the linear change of coordinates $z_0 = w_0, z_i = w_i - w_0, 1 \leq i \leq r$ by

$$z_1 \dots z_r (z_1 - z_2) \dots (z_{r-1} - z_r) (z_r - z_1).$$

Then $c_1 = 2r$, $c_2 = r$, and $c_3 = 0$, if $r > 3$. Thus $c_1 = 2c_2$, and so \mathcal{A}_{W_r} is decomposable and non-hypersolvable. We shall see that the group of the arrangement \mathcal{A}_{W_r} is not of type F_r .

2.2. Toric arrangements.

Let ν be a character of the torus $\mathbb{T}^{r+1} = (\mathbb{C}^*)^{r+1}$ and $a \in \mathbb{C}^*$. The pair (ν, a) defines a translated 1-codimensional subtorus $H_{\nu, a} = \{\mathbf{x} \in \mathbb{T}^{r+1} \mid \nu(\mathbf{x}) = a\}$. Note that if $\nu(\mathbf{x}) = x_0^{d_0} \cdot \dots \cdot x_r^{d_r}$, then $H_{\nu, a}$ has $d := \gcd(d_0, \dots, d_r)$ connected components.

A finite set \mathcal{A} of such hypertori in \mathbb{T}^{r+1} is called a toric arrangement. Much of the combinatorial and topological theory of such arrangements parallels that of hyperplane arrangements, see [10]. The de Rham cohomology ring of the complement was determined in [9]. Moreover, under certain conditions the de Rham algebra is formal and generated by logarithmic 1-forms.

Furthermore, the Poincaré polynomial of the complement $M = \mathbb{T}^{r+1} \setminus V(\mathcal{A})$ essentially coincides with the characteristic polynomial of the intersection poset $L(\mathcal{A})$ of \mathcal{A} , as shown in [23].

Proposition 2.3. *Let \mathcal{A} be a toric arrangement in a $(r+1)$ -torus with complement M . We then have*

$$P(M, t) = (-t)^{r+1} \chi(\mathcal{A}, -t^{-1} - 1).$$

Here $L(\mathcal{A})$ consists of all connected components of all non-empty intersections among hypertori in \mathcal{A} ordered by reversed inclusion and ranked by codimension. As before, $\chi(\mathcal{A}, q) := \sum_{S \in L(\mathcal{A})} \mu(S) q^{\dim S}$.

Remark 2.4. Let \mathcal{T} be a toric arrangement in \mathbb{T}^{r+1} which is obtained by the restriction a hyperplane arrangement \mathcal{H} in \mathbb{C}^{r+1} defined by equations of the form $z_i - az_j = 0$, with $a \in \mathbb{C}^*$.

Then the complement of \mathcal{T} in \mathbb{T}^{r+1} may be seen as the complement in \mathbb{C}^{r+1} to the hyperplane arrangement \mathcal{A} obtained by adding the coordinate hyperplanes to \mathcal{H} . The equality of Poincaré polynomials gives the following relation between characteristic polynomials

$$\chi(\mathcal{A}, q) = (-1)^{r+1} \chi(\mathcal{T}, q - 1).$$

Remark 2.5. Note that the formula of Proposition 2.3 can be as well applied to an arrangement of subtori in an r -torus $\mathbb{T}_{\nu, a} \subset \mathbb{T}^{r+1}$ defined as $\mathbb{T}_{\nu, a} = \{\nu(x) = a\}$ for some $a \in \mathbb{C}^*$ and some character ν of \mathbb{T}^{r+1} with $\gcd\{d_0, \dots, d_r\} = 1$.

2.3. General position toric arrangements.

We focus now on a particular class of toric arrangements, which consists of general position toric arrangements and some mild deformations of them. Our goal is to compute the Poincaré polynomials of such arrangements.

We start by analyzing an example illustrating all the features of the general case.

Example 2.6. Let $\mathcal{A}_a = \{H_{0,a}, H_1, \dots, H_r\}$ be the toric arrangement in \mathbb{T}^r defined by the hypertori $H_0 = H_{0,a} = \{x_1 \cdots x_r = a\}$ and $H_i = \{x_i = 1\}$, $1 \leq i \leq r$, for some $a \in \mathbb{C}^*$.

Note that the hypertori in \mathcal{A}_a intersect in general position, except maybe if we consider the intersection of all $r+1$ of them. Indeed, if H_I denotes the intersection $\bigcap_{i \in I} H_i$, and I is a proper subset of $\{0, \dots, r\}$, then H_I is a connected torus of codimension $|I|$. Now $\bigcap_{i=0}^r H_i$ is either empty if $a \neq 1$, or the point $(1, \dots, 1) \in \mathbb{T}$. It follows that the intersection poset $L(\mathcal{A}_a)$ is either the truncation of the lattice of subsets of $\{0, \dots, r+1\}$ (maximal element removed), if $a \neq 1$, or the obvious collapsing of this truncation for $a = 1$. More precisely, the poset $L(\mathcal{A}_{a \neq 1})$ differs from $L(\mathcal{A}_1)$ only in the maximal elements: there are $r+1$ of them, if $a \neq 1$, each with Möbius function $\mu = (-1)^r$, respectively only one, if $a = 1$, with $\mu = (-1)^{r+1}$.

The calculation of the characteristic polynomial of \mathcal{A}_a is then immediate:

$$\chi(\mathcal{A}_a, q) = \begin{cases} \sum_{k=0}^r (-1)^k \binom{r+1}{k} q^{r-k} & \text{if } a \neq 1, \\ \sum_{k=0}^r (-1)^k \binom{r+1}{k} q^{r-k} - (-1)^{r+1} & \text{if } a = 1. \end{cases}$$

After evaluating at $q = -\frac{1+t}{t}$, we obtain the Poincaré polynomials

$$P(M(\mathcal{A}_a), t) = \begin{cases} \frac{(2t+1)^{r+1} - t^{r+1}}{t+1} & \text{if } a \neq 1, \\ (2t+1) \frac{(2t+1)^r - t^r}{t+1} & \text{if } a = 1. \end{cases}$$

Note that the complements $M(\mathcal{A}_{a \neq 1})$ differ from $M(\mathcal{A}_1)$ only in their top Betti number b_r , as $P(M(\mathcal{A}_a), t) - P(M(\mathcal{A}_1), t) = t^r$. In fact the complements $M(\mathcal{A}_a)$ are all homeomorphic if $a \neq 1$, and, as we shall see later, they share their $(r-1)$ -skeleton, up to homotopy, with $M(\mathcal{A}_1)$.

We now treat the general case. We work in \mathbb{T}^{r+1} and fix a character ν such that $\nu(\mathbf{x}) = x_0^{d_0} \cdots x_r^{d_r}$ with $\gcd(d_0, \dots, d_r) = 1$. Consider the connected r -torus $\mathbb{T}_{\nu,a} = \{\nu(x) = a\}$ in \mathbb{T}^{r+1} for $a \in \mathbb{C}^*$. Let $\mathbf{n} := (n_0, n_1, \dots, n_r)$ and $m_i := n_i - 1$ as before. We want to compute the Poincaré polynomial of the complement in $\mathbb{T}_{\nu,a}$ of the arrangement $\mathcal{A}_{\nu,a}^{\mathbf{n}}(\boldsymbol{\alpha})$ consisting of the hypertori $H_i^j = \{x_i = \alpha_{i,j}\}$, $0 \leq i \leq r$, $1 \leq j \leq m_i$. As we shall see, $\mathcal{A}_{\nu,a}^{\mathbf{n}}(\boldsymbol{\alpha})$ depends on the data $\boldsymbol{\alpha} := (\alpha_{i,j})$. Denote by $\Lambda := \{\nu(\alpha_{0,j_0}, \dots, \alpha_{r,j_r}) \mid 1 \leq j_i \leq m_i, 0 \leq i \leq r\}$.

Proposition 2.7. *Let $M_a := M(\mathcal{A}_{\nu,a}^{\mathbf{n}}(\boldsymbol{\alpha}))$ be the complement of $\mathcal{A}_{\nu,a}^{\mathbf{n}}(\boldsymbol{\alpha})$ in $\mathbb{T}_{\nu,a}$. Suppose $a \notin \Lambda$. Then the Poincaré polynomial of the complement is equal to*

$$P(M_a, t) = \sum_{k=0}^r t^k (1+t)^{r-k} \sum_{I \subseteq [r], |I|=k} d_{\bar{I}} m_I.$$

In particular, the i -th Betti number of M_a is

$$b_i(M_a) = \sum_{l=r-i}^r \binom{l}{r-i} \sum_{|I|=r-l} d_{\bar{I}} m_I.$$

Furthermore, if $\lambda \in \Lambda$ then $P(M_\lambda, t) \neq P(M_a, t)$. More precisely, we have

$$b_i(M_a) = b_i(M_\lambda) \quad \forall i < r, \quad \text{and} \quad b_r(M_a) > b_r(M_\lambda).$$

Proof. We compute the characteristic polynomial $\chi(\mathcal{A}, q)$ of the toric arrangement $\mathcal{A} = \mathcal{A}_{\nu,a}^{\mathbf{n}}(\boldsymbol{\alpha})$. We claim that the hypertori H_i^j intersect in general position, except maybe if we consider intersections of $r+1$ of them.

First note that $H_i^{j'}$ and $H_i^{j''}$ never intersect if $j' \neq j''$. Then denote by $H_I^{\mathbf{j}}$ the intersection

$$H_I^{\mathbf{j}} = \mathbb{T}_{\nu,a} \cap \bigcap_{i \in I} H_i^{j_i}, \quad \text{for } I \subseteq [r], \quad \mathbf{j} \in \prod_{i \in I} [m_i],$$

where $[r] := \{0, \dots, r\}$. If $|I| \leq r$, then it can be easily seen that $H_I^{\mathbf{j}}$ is non-empty and of codimension $r - |I|$, and it consists of $d_{\bar{I}}$ connected components. Indeed, if $I = \{i_1, \dots, i_k\}$, $\mathbf{j} = (j_1, \dots, j_k)$ then $H_I^{\mathbf{j}}$ has $d_{\bar{I}}$ connected components

$$H_I^{\mathbf{j}} = \left\{ \prod_{i \notin I} x_i^{d_i} = a \prod_{p=1}^k \alpha_{i_p, j_p}^{-d_{i_p}}, \quad x_{i_p} = \alpha_{i_p, j_p}, \quad 1 \leq p \leq k \right\}.$$

If $a \notin \Lambda$, then $H_{[r]}^{\mathbf{j}}$ is empty for all \mathbf{j} , and so it does not appear in $L(\mathcal{A})$. Moreover, the Möbius function is readily computed as $\mu(H_I^{\mathbf{j}}) = (-1)^{|I|}$.

Now if we fix $k = |I|$ then clearly there are $\sum_{I, |I|=k} m_I$ intersections $H_I^{\mathbf{j}}$, and all these are mutually disjoint. Summing up the Möbius function contributions for all the connected components of $H_I^{\mathbf{j}}$ gives the coefficient of q^{r-k} in the characteristic polynomial $\chi(\mathcal{A}, q)$, and we obtain:

$$\chi(\mathcal{A}, q) = \sum_{k=0}^r (-1)^k q^{r-k} \sum_{I \subseteq [r], |I|=k} d_{\bar{I}} m_I.$$

Which, after the substitution $q = -t^{-1} - 1$, gives the Poincaré polynomial of the M_a .

The formula for the Betti numbers follows after a routine rearrangement of the sum. More precisely, if

$$\chi(q) = \sum_{k=0}^r (-1)^k c_{r-k} q^{r-k}, \quad P(t) = \sum_{i=0}^r b_i t^i, \quad \text{and} \quad P(t) = (-t)^r \chi(-t^{-1} - 1)$$

then $b_i = \sum_{l=r-i}^r \binom{l}{r-i} c_l$. Note that $b_r = \sum_{l=0}^r c_l$ and that b_i does not depend on c_0 if $i < r$.

Consider now the case $\lambda \in \Lambda$. It is readily seen that positive dimensional intersections $H_I^{\mathbf{j}}$ have the same description as in the generic case. Thus the coefficient of q^{r-k} in $\chi(\mathcal{A}, q)$ will stay the same if $k < r$. The only difference may appear in the free term. To that coefficient only $H_I^{\mathbf{j}}$ with $|I|$ equals r or $r+1$ may contribute.

Let us start with the case $|I| = r$; let $p \in [r] \setminus I$. We distinguish two types for the intersections $H_I^{\mathbf{j}}$. In the first type there exists $j_p \in \{1, \dots, m_p\}$ such that $\lambda = \nu(\alpha_{0,j_0}, \dots, \alpha_{r,j_r})$. In the second type, $\forall j_p \in \{1, \dots, m_p\}$ we have $\lambda \neq \nu(\alpha_{0,j_0}, \dots, \alpha_{r,j_r})$. Then, we have

$$H_I^{\mathbf{j}} = \begin{cases} \left\{ \left\{ x_p^{d_p} = \alpha_{p,j_p}^{d_p}, x_s = \alpha_{s,j_s}, s \neq p \right\} \right. & \text{in the first type,} \\ \left. \left\{ x_p^{d_p} = a \prod_{s \neq p} \alpha_{s,j_s}^{-d_s}, x_s = \alpha_{s,j_s}, s \neq p \right\} \right. & \text{in the second one.} \end{cases}$$

Both types consist of d_p distinct points. The conditions on the value of ν immediately imply that intersections of the first and second type are disjoint. Moreover two different intersections of the second type are disjoint. Finally if in the second type we choose \mathbf{j}' such that $j_i = j'_i$ if and only if $i \neq p$, then $H_{[r]}^{\mathbf{j}'}$ is empty. It remains to consider what happens with the intersection of two pairs of the first type.

We fix notations. Let $H_I^{\mathbf{j}}, H_{I'}^{\mathbf{j}'}$ be two intersections of the first type ($|I| = |I'| = r$, $p \in [r] \setminus I$, $p' \in [r] \setminus I'$). Let $j_p, j_{p'}$ be such that $\nu(\alpha_{0,j_0}, \dots, \alpha_{r,j_r}) = \lambda = \nu(\alpha_{0,j'_0}, \dots, \alpha_{r,j'_r})$. Then $H_I^{\mathbf{j}}$ and $H_{I'}^{\mathbf{j}'}$ will intersect if and only if either

$$p = p', j_s = j'_s \text{ for } s \neq p, \text{ and } \alpha_{p,j_p}^{d_p} = \alpha_{p',j'_p}^{d_p},$$

or

$$p \neq p', j_s = j'_s \text{ for } s \neq p, p', \text{ and } \alpha_{p,j_p}^{d_p} = \alpha_{p',j'_p}^{d_p}, \alpha_{p',j'_p}^{d_{p'}} = \alpha_{p',j'_p}^{d_{p'}}.$$

We add an additional hypothesis: for any $p \in [r]$ and $1 \leq j_p < j'_p$ then

$$(2.1) \quad \alpha_{p,j_p}^{d_p} \neq \alpha_{p,j'_p}^{d_p}.$$

Under this hypothesis $H_I^{\mathbf{j}}$ and $H_{I'}^{\mathbf{j}'}$ are disjoint.

Now we consider the case $|I| = r+1$, i.e. $I = [r]$. Let \mathbf{j} such that $H_{[r]}^{\mathbf{j}} \neq \emptyset$, i.e. it consists of a single point $\{x_s = \alpha_{s,j_s}, 0 \leq s \leq r\}$. For any $p \in [r]$, this point lies also in $H_p^{\mathbf{j}^p}$ (where \mathbf{j}^p is the r -tuple obtained by forgetting j_p). Note also that the other $(d_p - 1)$ points in $H_p^{\mathbf{j}^p}$ only belong to this set. Hence, for the point in $H_{[r]}^{\mathbf{j}} \neq \emptyset$ it follows that $\mu = (-1)^r$ while for the other points, we have $\mu = r(-1)^r$.

Now if we sum up the contributions of all connected components of zero-dimensional intersections, grouping them into the two types, we obtain a free term contribution with sign $(-1)^r$ and whose absolute value is at most $\sum_{p \in [0,r]} d_p m_{\bar{p}}$, the free term in the generic case. Indeed,

the free term writes as $(-1)^r$ times

$$\sum_{J \text{ type 1}} \left(\sum_{p=0}^r (d_p - 1) + r \right) + \sum_{K \text{ type 2}} \left(\sum_{p=0}^r d_p \right) = \sum_{p=0}^r d_p m_{\bar{p}} - \#\{J \text{ type 1}\}.$$

Finally, in case condition (2.1) is not satisfied, then the free term clearly drops even more in absolute value. \square

Let us fix $\mathbf{n} := (n_0, n_1, \dots, n_r)$. We denote $m_i := n_i - 1$. For $0 \leq i \leq r$, fix arbitrary sets of m_i non-zero complex numbers $\{\alpha_{i,j}\}_{j=1}^{m_i}$. Let us denote $H_i^j = \{x_i = \alpha_{i,j}\}$, if $i > 0$, and $H_{\nu,a}^j = \{\nu(x) = a\alpha_{0,j}^{-1}\}$, for some $a \in \mathbb{C}^*$. Fix $a \in \mathbb{C}^*$ and denote $H_0^j := H_{\nu,a}^j$. We will consider the following toric arrangement $\mathcal{A}_{\nu,a}^{\mathbf{n}} := \bigcup_{k=0}^r \{H_k^j\}_{j=1}^{m_k}$.

3. MAIN THEOREMS

In this section we collect the main results of the paper. The proofs will be given in the last two sections. The quasi-projective Bestvina-Brady groups were determined in [12].

Theorem 3.1 ([12]). *Let Γ be a finite graph. Suppose the Bestvina-Brady group N_{Γ} associated to Γ is quasi-projective. Then Γ is either a tree, or a complete multipartite graph K_{n_0, \dots, n_r} , with either some $n_i = 1$ or all $n_i \geq 2$ and $r \geq 2$.*

We shall make this statement more precise, by showing that all quasi-projective groups N_{Γ} are in fact fundamental groups of hyperplane arrangement complements in \mathbb{P}^r .

Theorem 3.2. *Any quasi-projective Bestvina-Brady group N_{Γ} is a hyperplane arrangement group for an arrangement in \mathbb{P}^r .*

In the first case, N_{Γ} is simply a free group of rank $v - 1$ (v is the number of vertices of the tree), which is known to be the fundamental group of the complement in \mathbb{P}^1 of v points.

The second case will interest us most. Let $\Gamma = K_{\mathbf{n}}$, with $\mathbf{n} := (n_0, \dots, n_r)$. The first part is again easy. Assume $n_i = 1$ for $0 \leq i \leq s$ and $n_i > 1$ for $s < i \leq r$. Then it is readily seen that $N_{K_{\mathbf{n}}} \cong \mathbb{Z}^s \times \mathbb{F}_{n_{s+1}} \times \dots \times \mathbb{F}_{n_r}$. This group is clearly the fundamental group of the complement of a hyperplane arrangement in $\mathbb{C}^r = \mathbb{P}^r \setminus \{z_0 = 0\}$ with equation

$$z_i = \alpha_i, \quad 1 \leq i \leq s, \quad z_i = \alpha_{i,j}, \quad s+1 \leq i \leq r, \quad 1 \leq j \leq n_i,$$

for generic choices of $\alpha_i, \alpha_{i,j}$. Taking a generic plane section it can be seen as the fundamental group of the complement of a line arrangement: Consider $r - s$ distinct directions in \mathbb{C}^2 and take n_i lines parallel to the $s + i^{\text{th}}$ direction, and s other lines in general position.

So only the case $N_{K_{\mathbf{n}}}$, with all $n_i \geq 2$ and $r \geq 2$, is left. We will show in the next section that $N_{K_{\mathbf{n}}}$ is the fundamental group of the complement in \mathbb{P}^r to the hyperplane arrangement $\mathcal{A}_{\nu,a}^{\mathbf{n}}$

defined by the polynomial

$$z_0 \cdots z_r \cdot \prod_{j=1}^{n_0-1} (az_0 - \alpha_{0,j}z_1) \cdot \left(\prod_{i=1}^{r-1} \prod_{j=1}^{n_i-1} (z_i - \alpha_{i,j}z_{i+1}) \right) \cdot \prod_{j=1}^{n_r-1} (z_r - \alpha_{r,j}z_0),$$

A generic plane section provides a line arrangement with a precise combinatorics.

Corollary 3.3. *The group $N_{K_{\mathbf{n}}}$ can be realized as $\pi_1(\mathbb{P}^2 \setminus \mathcal{A}_{\mathbf{n}})$, where $\mathcal{A}_{\mathbf{n}}$ is a line arrangement in \mathbb{P}^2 consisting of $(n_0 + 1) \cdots (n_r + 1)$ lines, formed by r lines in general position cutting out a polygon with $r + 1$ sides which have at vertex i another $n_i - 1$ lines that intersect transversally among themselves.*

If $\Gamma = K_{n_0, \dots, n_r}$ then the right-angled Artin group A_{Γ} is the product of free groups $\mathbb{F}_{n_0} \times \cdots \times \mathbb{F}_{n_r}$. Using the same arguments, one can extend this result to other subgroups of A_{Γ} , namely to the generalized Bestvina-Brady group $N_{\mathbf{n}}^{\mathbf{d}}$, where $\mathbf{n} = (n_0, \dots, n_r)$, and $\mathbf{d} = (d_0, \dots, d_r)$ with $\gcd(d_i) = 1$.

Theorem 3.4. *Any generalized Bestvina-Brady group $N_{\mathbf{n}}^{\mathbf{d}}$ is the fundamental group of the complement to an algebraic hypersurface in the weighted projective space $\mathbb{P}^r(\mathbf{d})$.*

In fact, it is more convenient to view this complement space as obtained from an algebraic r -torus by removing $\sum_{i=0}^r (n_i - 1)$ hypertori. More precisely, we will consider the toric arrangement $\mathcal{A}_{\mathbf{n}, \mathbf{a}}^{\mathbf{d}}(\alpha)$ from Section 2. In certain cases, as we shall see later, it is still possible to realize this toric arrangement complement as a hypersurface complement in \mathbb{P}^r , but in general this hypersurface is not the union of hyperplanes. Now recall that a $K(\pi, 1)$ space for $N_{\mathbf{n}}^{\mathbf{d}}$ may be chosen to be the infinite cyclic cover $\mathbb{T}_{K_{\mathbf{n}}}^{\chi^{\mathbf{d}}}$ of the product $\mathbb{T}_{K_{\mathbf{n}}}$ of tori. Then, by putting together the geometric realizability of $N_{\mathbf{n}}^{\mathbf{d}}$ from Section 4 and its higher dimensional consequences from Section 5, we obtain the following.

Theorem 3.5. *Let $M = M(\mathcal{A}_{\mathbf{n}, \mathbf{a}}^{\mathbf{d}}(\alpha))$ be the toric arrangement complement, where $r \geq 2$, $n_i > 1$, and $\mathbf{a} \notin \Lambda$. Then we have the following*

- (1) *the r -skeleton of $\mathbb{T}_{K_{\mathbf{n}}}^{\chi^{\mathbf{d}}}$ has the homotopy type of M .*
- (2) *the Artin kernel $N_{\mathbf{n}}^{\mathbf{d}}$ is isomorphic to $\pi_1(M)$.*
- (3) *the homotopy groups $\pi_i(M)$ vanish in the range $1 < i < r$.*
- (4) *the homology groups $H_i(M)$ and $H_i(N_{\mathbf{n}}^{\mathbf{d}})$ are isomorphic for $i \leq r$.*

4. A GEOMETRIC CONSTRUCTION

Suppose $\mathbb{T}^{r+1} \setminus D$ is a hypersurface complement in a complex torus of dimension $r + 1$. If $f: \mathbb{T}^{r+1} \setminus D \rightarrow \mathbb{C}^*$ is a polynomial function then let $\Lambda_f \subset \mathbb{C}^*$ be the bifurcation set of f , that is the smallest set Λ such that f is a locally trivial fibration over the complement of Λ .

In this section we will determine the bifurcation set Λ_f for certain monomial maps defined on $\mathbb{T}^{r+1} \setminus D$ for which D is a union of hypertori of \mathbb{T}^{r+1} , that is a toric arrangement complement.

More precisely, given $(r+1)$ -tuples $\mathbf{n} := (n_i)_{i=0}^r$ and $\mathbf{d} := (d_i)_{i=0}^r$ of positive integers, with $\gcd \mathbf{d} = 1$, let $p := p^{\mathbf{d}}$ denote the multiplication map:

$$p : \mathbb{T}^{r+1} \rightarrow \mathbb{C}^*, \quad p(x_0, \dots, x_r) := x_0^{d_0} \cdot \dots \cdot x_r^{d_r}.$$

Now, let $f := f_{\mathbf{n}}^{\mathbf{d}} : X \rightarrow \mathbb{C}^*$ denote the restriction of p to the complement $X := \mathbb{T}^{r+1} \setminus D$ of the hypersurface $D := \bigcup_{(i,j) \in B} D_{i,j}$ in the torus \mathbb{T}^{r+1} , where $D_{i,j} := \{x_i = \alpha_{i,j}\}$, $\alpha_{i,j} \in \mathbb{C}^*$ and $B := \{(i,j) \mid 1 \leq j < n_i, 0 \leq i \leq r\}$. Note that X is homeomorphic to $\mathbb{C}_{n_0}^* \times \dots \times \mathbb{C}_{n_r}^*$, the toric complex $\mathbb{T}_{K_{\mathbf{n}}}$ associated to the complete multipartite graph $K_{\mathbf{n}}$; the analytic structure of X depend on the sets of $n_i - 1$ points removed from each \mathbb{C}^* factor. In fact, $f = f_{\mathbf{n}}^{\mathbf{d}}$ also depends on them, but they are omitted for notational simplicity.

Lemma 4.1. *The map $f = f_{\mathbf{n}}^{\mathbf{d}} : X \rightarrow \mathbb{C}^*$ induces the group homomorphism $\chi := \chi_{\mathbf{n}}^{\mathbf{d}} : A_{\Gamma} \rightarrow \mathbb{Z}$.*

Let $a \in \mathbb{C}^*$. Then the fiber $p^{-1}(a) = \{x_0^{d_0} \cdot \dots \cdot x_r^{d_r} = a\}$ is a connected hypertorus ambient isomorphic to the hypertorus $\{y_0 = a\}$, via a monomial automorphism $x_j \mapsto \prod_{i=0}^r y_i^{a_{ij}}$, determined by a unimodular integer matrix $A = (a_{ij})_{0 \leq i, j \leq r}$. Clearly the map p is a trivial fiber bundle with fiber a connected torus. Let $C := \{\mathbf{j} = (j_0, \dots, j_r) \mid 1 \leq j_i < n_i, 0 \leq i \leq r\}$.

Proposition 4.2. *The bifurcation set Λ_f consists of the values $\{a_{\mathbf{j}} := \alpha_{0,j_0} \cdot \dots \cdot \alpha_{r,j_r}\}_{\mathbf{j} \in C}$. For generic points $\{\alpha_{i,j}\}$ the map f has exactly $m := (n_0 - 1) \cdot \dots \cdot (n_r - 1)$ distinct special fibers $f^{-1}(a_{\mathbf{j}})$.*

Proof. The hypersurface D determines a stratification of the ambient torus \mathbb{T}^{r+1} as follows. For each $I \subset B$ consider the intersection $D_I = \bigcap_{(i,j) \in I} D_{i,j}$; the strata are defined as differences of these closed sets. Then the top stratum is X itself and the positive codimension strata are the intersections D_I away from lower dimensional $D_{I'}$.

Now $f : X \rightarrow \mathbb{C}^*$ is a restriction of p which is a trivial fiber bundle. In order to apply the Thom Isotopy Lemma, it is enough to compactify X as a subspace of (normalization of)

$$Z := \{([x_0 : \dots : x_r : y], t) \in \mathbb{P}^{r+1} \times \mathbb{C}^* \mid x_0^{d_0} \cdot \dots \cdot x_r^{d_r} = ty^{d_{r+1}}\}, \quad d_{r+1} = \sum d_i,$$

and extend p to a proper map of $Z \rightarrow \mathbb{C}^*$. Note that p defines a trivial fibration at the strata at infinity.

In order to make f into a fiber bundle we only have to remove those values $a \in \mathbb{C}^*$ (and their preimages from X) such that $p^{-1}(a)$ does not intersect transversally at least one stratum of the above stratification of \mathbb{T}^{r+1} . It is immediate that the fibers $p^{-1}(a)$ intersect D_I transversally away from lower dimensional $D_{I'}$, unless D_I are already zero-dimensional. In that case, the stratum is simply a point which may lie or not on $p^{-1}(a)$. More precisely, D_I is a point if and only if $I = I_{(j_0, \dots, j_r)} = \{(0, j_0), \dots, (r, j_r)\}$ for $(j_0, \dots, j_r) \in C$. Note that $I_{(j_0, \dots, j_r)} \subset f^{-1}(\alpha_{0,j_0} \cdot \dots \cdot \alpha_{r,j_r})$ and hence, Λ_f is determined.

Finally, if $\{\alpha_{i,j}\}_{(i,j) \in B}$ are generic, then the values $a = \alpha_{0,j_0} \cdot \dots \cdot \alpha_{r,j_r}$ are pairwise distinct. \square

For the sake of simplicity, we will first treat the situation where $d_i = 1$ for all i . In that case we just write $f = f_{\mathbf{n}}$ and $p : \mathbb{T}^{r+1} \rightarrow \mathbb{C}^*$, $p(\mathbf{x}) = x_0 \cdot \dots \cdot x_r$. The fibers $p^{-1}(a) = \{\mathbf{x} \in \mathbb{T}^{r+1} \mid x_0 \cdot \dots \cdot x_r = a\}$ of the product map are isomorphic to the r -torus \mathbb{T}^r under the identifications

$$(4.1) \quad (x_0, \dots, x_r) \mapsto (x_1, \dots, x_r), \quad (x_1, \dots, x_r) \mapsto \left(\frac{a}{x_1 \cdot \dots \cdot x_r}, x_1, \dots, x_r \right).$$

Moreover one can identify \mathbb{T}^r with $\mathbb{P}^r \setminus \{z_0 \cdot \dots \cdot z_r = 0\}$, via

$$(4.2) \quad (x_1, \dots, x_r) \mapsto \left[1 : \prod_{j=1}^r x_j : \prod_{j=2}^r x_j : \dots : x_{r-1} x_r : x_r \right], \quad [z_0 : \dots : z_r] \mapsto \left(\frac{z_1}{z_2}, \dots, \frac{z_r}{z_0} \right).$$

Thus, composing the inverse of the identifications (4.1)-(4.2), we obtain a diffeomorphism from $\mathbb{P}^r \setminus \{z_0 \cdot \dots \cdot z_r = 0\}$ to $p^{-1}(a)$ provided by

$$(4.3) \quad [z_0 : \dots : z_r] \mapsto \left(a \frac{z_0}{z_1}, \frac{z_1}{z_2}, \dots, \frac{z_r}{z_0} \right).$$

Lemma 4.3. *Consider the map $f = f_{\mathbf{n}} : X \rightarrow \mathbb{C}^*$ as defined above. The fiber $F = f^{-1}(a)$ over $a \in \mathbb{C}^*$ is homeomorphic to the complement in \mathbb{P}^r of the hyperplane arrangement \mathcal{A}_a defined by the polynomial*

$$z_0 \cdot \dots \cdot z_r \cdot \prod_{j=1}^{n_0-1} (az_0 - \alpha_{0,j} z_1) \cdot \left(\prod_{i=1}^{r-1} \prod_{j=1}^{n_i-1} (z_i - \alpha_{i,j} z_{i+1}) \right) \cdot \prod_{j=1}^{n_r-1} (z_r - \alpha_{r,j} z_0),$$

Proof. Under the identification $\{\mathbf{x} \in \mathbb{T}^{r+1} \mid x_0 \cdot \dots \cdot x_r = a\} \rightarrow \mathbb{P}^r \setminus \{x_0 \cdot \dots \cdot x_r = 0\}$ described in (4.3) the fiber $F = f^{-1}(a) = \{\mathbf{x} \in \mathbb{T}^{r+1} \mid x_0 \cdot \dots \cdot x_r = a, x_i \neq \alpha_{i,j}\}$ is sent to

$$\{[z_0 : z_1 : \dots : z_r] \in \mathbb{P}^r \mid z_i \neq 0, az_0 \neq \alpha_{0,j} z_1, z_i \neq \alpha_{i,j} z_{i+1}, z_r \neq \alpha_{r,j} z_0\}. \quad \square$$

Remark 4.4. Because of Proposition 4.2, we know $m = \prod_{i=0}^r (n_i - 1) \geq m' := \#\Lambda_f$. Let us denote

$$\mathbb{C}_{m'+1}^* = \mathbb{C}^* \setminus \Lambda_f^*, \quad E = \prod_{i=0}^r \mathbb{C}_{n_i}^* \setminus f^{-1}(\Lambda_f^*).$$

The restriction $f = f_{\mathbf{n}} : E \rightarrow \mathbb{C}_{m'+1}^*$ is a locally trivial fibration. For generic values, $m = m'$ as stated in Proposition 4.2. For non-generic values of the $\alpha_{i,j}$'s the bifurcation set could very well have smaller cardinality than m .

For example the map $f = f_{3,\dots,3} : (\mathbb{C}^* \setminus \{\pm 1\})^{r+1} \rightarrow \mathbb{C}^*$ will have $\Lambda_f = \{\pm 1\}$, thus $m' = 2 < 2^{r+1} = m$. More generally, one may take the set of $\{\alpha_{i,j}\}_{1 \leq j < n_i}$ to consist of the subgroup μ_{n_i-1} of roots of unity of order $n_i - 1$ in \mathbb{C}^* .

We assume from now that $m = m'$. The exact sequence in homotopy of the fibration $f = f_{\mathbf{n}} : E \rightarrow \mathbb{C}_{m+1}^*$ gives a short exact sequence of groups

$$(4.4) \quad 1 \rightarrow \pi_1(F) \rightarrow \pi_1(E) \rightarrow \pi_1(\mathbb{C}_{m+1}^*) \rightarrow 1.$$

Identify $\pi_1(\mathbb{C}_{m+1}^*)$ with the free group of rank $m + 1$ written as $\mathbb{Z} * \mathbb{F}_m$, where \mathbb{Z} is generated by the class t of a meridian around the origin, and \mathbb{F}_m is the free group on a_j (coming from meridians around the points in Λ_f), $\mathbf{j} \in C$.

Note that $\pi_1(E)$ surjects onto the Artin group $A = \prod_{i=0}^r \mathbb{F}_{n_i}$, and the epimorphism is induced by the inclusion $E \hookrightarrow \prod_{i=0}^r \mathbb{C}_{n_i-1}^*$. The inclusion $\mathbb{C}^* \setminus \Lambda_f \hookrightarrow \mathbb{C}^*$ induces the obvious epimorphism $\mathbb{Z} * \mathbb{F}_m \rightarrow \mathbb{Z}$. Comparing the short exact sequence of the fiber bundle with the one defining Bestvina-Brady group, we obtain a commutative diagram whose vertical arrows are epimorphisms. In fact we have the following.

Lemma 4.5. *The arrangement group $\pi_1(F)$ is isomorphic to the Bestvina-Brady group $N = N_{\mathbf{n}}$.*

Remark 4.6. The particular case $f = f_{2,\dots,2} : (\mathbb{C}^* \setminus \{1\})^{r+1} \rightarrow \mathbb{C}^*$ leads to interesting hyperplane arrangements. The fiber $F = f^{-1}(a)$ over $a \neq 1$ is homeomorphic to the complement of the hyperplane arrangement \mathcal{A}_a in \mathbb{P}^r defined by $z_0 z_1 \dots z_r (a z_0 - z_1)(z_1 - z_2) \dots (z_{r-1} - z_r)(z_r - z_0)$. The bifurcation set Λ_f is just $\lambda = 1$. If $r > 2$ then it is readily seen that the fundamental group of the complement to \mathcal{A}_a does not change even when $a = 1$. Lemma 4.5 will imply that $N_{2,\dots,2}$ is the group of the graphic arrangement $z_0 z_1 \dots z_r (z_0 - z_1)(z_1 - z_2) \dots (z_{r-1} - z_r)(z_r - z_0)$, $r > 2$.

Proof of Lemma 4.5. We have the following commutative diagram, where the rows are exact and the vertical arrows are surjective:

$$\begin{array}{ccccccc} 0 & \rightarrow & \pi_1(F) & \rightarrow & \pi_1(E) & \rightarrow & \pi_1(\mathbb{C}_{m+1}^*) \cong \mathbb{Z} * \mathbb{F}_m \rightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \rightarrow & N & \rightarrow & A & \rightarrow & \pi_1(\mathbb{C}^*) \cong \mathbb{Z} \rightarrow 0 \end{array}$$

Let us fix $\mathbf{j} := (j_0, \dots, j_r) \in C$. Let γ_j be a lift of a_j to $\pi_1(E)$ that becomes trivial in the Artin group A . Let G be the subgroup of $\pi_1(E)$ normally generated by γ_j , $\mathbf{j} \in C$. If we identify $\pi_1(F)$ as a subgroup of $\pi_1(E)$, then $K := \ker(\pi_1(F) \rightarrow N) = G \cap \pi_1(F)$. It can be easily checked that K is normally generated by $[g, \gamma_j]$, $g \in \pi_1(F)$, $\mathbf{j} \in C$. The goal is to prove that K is trivial and this fact holds if $g^{\gamma_j} = g$ for all g in $\pi_1(F)$ and $\mathbf{j} \in C$.

The crucial observation is that the conjugation by γ_j is obtained by means of the monodromy action of a_j on $\pi_1(F)$; this is due to the choice of the lift γ_j to be trivial in A . In order to understand the action, we need to understand the behavior of f near the bifurcation value α_j . The only stratum not transversal to $f^{-1}(\alpha_j)$ is the stratum I_j which consists of a point $p_j := (\alpha_{0,j_0}, \dots, \alpha_{r,j_r})$. Let \mathbb{D}_j be a small closed disk around α_j and let us denote $\mathbb{D}_j^* := \mathbb{D}_j \setminus \{\alpha_j\}$. The disk is chosen in order to ensure the following facts:

- There is a *small* Milnor ball $\mathbb{B}_j \subset p^{-1}(\mathbb{D}_j)$ for p around p_j .
- $X_j := \overline{\phi^{-1}(\mathbb{D}_j)} \setminus \overline{\mathbb{B}_j}$ fibers trivially over \mathbb{D}_j .

We have realized a decomposition $f^{-1}(\mathbb{D}_j^*) = X_j^* \cup \check{\mathbb{B}}_j$ for $X_j^* := X_j \cap E$ and $\check{\mathbb{B}}_j := \mathbb{B}_j \cap E$. The intersections $Y_j := X_j \cap \check{\mathbb{B}}_j$ $Y_j^* := X_j^* \cap \check{\mathbb{B}}_j$ are arcwise connected.

Since the statement deals with a conjugation equality, we may choose as F any non-special fiber. Let us fix $\tilde{\alpha}_j \in \partial\mathbb{D}_j$ and fix $F := f^{-1}(\tilde{\alpha}_j)$; for later use, let us denote $F_j := f^{-1}(\alpha_j)$, which is a special fiber. Let us denote $M := F \cap \mathbb{B}_j$, $M_j := F \cap \check{\mathbb{B}}_j$, $\check{F} := F \cap X_j$, $\check{F}_j := F_j \cap X_j$, $F^\partial = M \cap \check{F}$ and $F_j^\partial = M_j \cap \check{F}_j$.

By the above construction:

$$X_j \cong \check{F} \times \mathbb{D}_j \cong \check{F}_j \times \mathbb{D}_j, \quad X_j^* \cong \check{F} \times \mathbb{D}_j^*$$

and these homeomorphisms restrict to:

$$Y_j \cong F^\partial \times \mathbb{D}_j^* \cong F_j^\partial \times \mathbb{D}_j, \quad Y_j^* \cong F^\partial \times \mathbb{D}_j^*;$$

moreover the pairs (\check{F}, F^∂) and $(\check{F}_j, F_j^\partial)$ are isotopic in X_j . We can choose a representative of γ_j in Y_j^* ; the above homeomorphism imply that the action of γ_j by conjugation on $\pi_1(\check{F})$ is trivial. Hence, applying Seifert-van Kampen theorem, it is enough to check that the action of γ_j by conjugation on $\pi_1(M)$ is also trivial.

This is done as follows. Using the Taylor expansion of p around p_j we can choose analytic coordinates t for \mathbb{D}_j (centered at α_j) and (y_0, \dots, y_r) for \mathbb{B}_j (centered at p_j) such that $p(y_0, \dots, y_r) = y_0 + \dots + y_r$.

Then, M has the homotopy type of the complement of the arrangement of equation $z_1 \dots z_r \cdot (z_1 + \dots + z_r - b) = 0$ in \mathbb{C}^r , for some $b \in \mathbb{C}^*$. This group is generated by the meridians of the hyperplanes and it is abelian (since $r \geq 2$). Let μ be the meridian of one of the hyperplanes; the element μ^{γ_j} is constructed geometrically using the monodromy around a_j and it is again a meridian of the same hyperplane. Since the group is abelian we obtain that $\mu^{\gamma_j} = \mu$; since the group $\pi_1(M)$ is generated by these meridians we obtain that γ_j acts trivially on $\pi_1(M)$ as required. \square

Proof of Theorem 3.2. It is an immediate consequence of Lemmas 4.3 and 4.5. \square

Proof of Corollary 3.3. In order to obtain the arrangement \mathcal{A}_n in \mathbb{P}^2 we only need to take a generic 2-dimensional slice of the hyperplane arrangement \mathcal{A}_a in \mathbb{P}^r . \square

We now look at the general case of $f = f_{\mathbf{n}}^{\mathbf{d}}$, where $d_i \geq 1$ are integers whose greatest common divisor is 1. The fibers of the product map $p = p^{\mathbf{d}} : \mathbb{T}^{r+1} \rightarrow \mathbb{C}^*$ are still tori isomorphic to \mathbb{T}^r , but the identification map does not have an explicit form in terms of the d_i 's. In fact, the torus \mathbb{T}^r naturally lives as $\mathbb{P}^r(\mathbf{d}) \setminus \{z_0 \dots z_r = 0\}$ inside the weighted projective space $\mathbb{P}^r(\mathbf{d})$. Therefore the fibers of f can be identified with complements of hypersurfaces in $\mathbb{P}^r(\mathbf{d})$.

Remark 4.7. In case one of the d_i 's is equal to 1, say $d_0 = 1$, we can still realize the fibers of f as complements of hypersurfaces in \mathbb{P}^r . Indeed, the fiber $p^{-1}(a) = \{x_0 \cdot x_1^{d_1} \dots x_r^{d_r} = a\} \subset \mathbb{T}^{r+1}$ is homeomorphic to $\mathbb{P}^r \setminus \{z_0 \dots z_r = 0\}$ via the mapping: $x_i = z_i z_{i+1}^{-1}$ for $1 \leq i \leq r$, and $x_0 = a z_0^{d_r} z_1^{-d_1} z_2^{d_1 - d_2} \dots z_r^{d_{r-1} - d_r}$.

Example 4.8. Consider the map $f = f_{2,\dots,2}^{\mathbf{d}} : (\mathbb{C}^* \setminus \{1\})^{r+1} \rightarrow \mathbb{C}^*$ and assume $d_0 = 1$. The bifurcation set is $\Lambda_f = \{1\}$. The fiber $F = f^{-1}(a)$ is homeomorphic to the complement of the arrangement of hypersurfaces \mathcal{A}_a in \mathbb{P}^r defined by the polynomial

$$z_0 \cdots z_r \cdot \prod_{i=1}^r (z_i - z_{i+1}) \cdot \prod_{j=1}^{n_0-1} (az_0^{d_r} - z_1^{d_1} z_2^{d_2-d_1} \cdots z_r^{d_r-d_{r-1}}).$$

Proof of Theorem 3.4. The proof follows the same ideas as the proof of Theorem 3.2. We have seen above that the generic fibers of f are in a natural way hypersurfaces of in some weighted projective space. The special fibers are obtained as in Proposition 4.2; they correspond to some 0-dimensional non-zero strata. If we take out the special fibers we obtained a short exact sequence like (4.4). The local behavior around the *bad* 0-dimensional strata are like in the proof of Lemma 4.5 and the result follows. \square

5. FIBRATIONS AND HOMOTOPY GROUPS

We will show here that the map $f : X = \mathbb{T}^{r+1} \setminus D \rightarrow \mathbb{C}^*$ considered in the previous section is in fact *like a fibration* in codimension 1. Also, in order to investigate the homology of the Artin kernels N_{Γ}^{χ} , we construct geometric approximations of their $K(\pi, 1)$ complexes.

Recall that $f : f^{-1}(B) = X_{\Lambda} \rightarrow \mathbb{C}^* \setminus \Lambda = B$ is a fiber bundle. The bifurcation set Λ consists of the $a \in \mathbb{C}^*$ for which the fiber $F_a = f^{-1}(a)$ is not transverse to all the strata of the divisor D . Also recall from the proof of Proposition 4.2 that the union Σ of those non-transversal strata is a finite set of points.

5.1. Quasifibrations.

In order to describe the objects and concepts mentioned above we need some definitions. In what follows we will consider $f : X \rightarrow C$ surjective maps where C is path-connected.

Definition 5.1 ([14]). A map $f : X \rightarrow C$ is a *quasifibration* (resp. up to dimension n) if it induces a weak homotopy equivalence (resp. n -equivalence) $f : (X, F_c) \rightarrow (C, c)$ for all $c \in C$, $F_c = f^{-1}(c)$, that is, if for any $x \in F_c$ the morphism

$$f_* : \pi_i(X, F_c, x) \rightarrow \pi_i(C, c)$$

is an isomorphism for all $i \geq 0$ (resp. for all $i < n$ and a surjection for $i = n$). We call a quasifibration up to dimension n an *n -quasifibration* for short.

A characterization of n -quasifibrations is the following (cf. [21, 16]): A map $f : X \rightarrow C$ is an n -quasifibration if and only if the inclusion $F_c \rightarrow H_c$ of any fiber $F_c = f^{-1}(c)$ into the homotopy theoretic fiber H_c is a (weak) homotopy n -equivalence.

As a consequence of this, all the fibers of an n -quasifibration have the same weak homotopy type up to dimension n .

5.2. Monomial quasifibrations.

We will show that the monomial maps $f : X = \mathbb{T}^{r+1} \setminus D \rightarrow \mathbb{C}^*$ considered in the previous section have a quasifibration structure. For the rest of the section we assume $r \geq 3$ unless otherwise stated.

Proposition 5.2. *The map $f : X = \mathbb{T}^{r+1} \setminus D \rightarrow \mathbb{C}^*$ defined by $f(\mathbf{x}) = x_0^{d_0} \cdots x_r^{d_r}$ is an r -quasifibration.*

Proof. The proof will use the tools for quasifibrations devised in the proof of the Dold-Thom theorem as outlined in Hatcher [17, Chapter 4, Section 4.K]. The restriction $f : X_\Lambda \rightarrow \mathbb{C}^* \setminus \Lambda = B$ is a fiber bundle, and thus a fibration and a quasifibration. Choose neighborhoods B_j of the points q_j in the finite set Λ so that $B \cup \bigcup_j B_j$ is a cover of \mathbb{C}^* . The result will follow from the gluing theorem in [17, Lemma 4K.3] (see also [16, Theorem 2.3]) as soon as we prove that the restrictions $f : f^{-1}(B_j) \rightarrow B_j$ are r -quasifibrations.

Recall that to each $q_j \in \Lambda$ it corresponds a non-transversality point p_j in the intersection of D with the fiber $f^{-1}(\lambda)$ of $f : \mathbb{T}^{r+1} \rightarrow \mathbb{C}^*$. Using [17, Corollary 4K.2], it is enough to show the following local statement: For any $q_j \in \Lambda$, there exist neighborhoods U_j of p_j in \mathbb{T}^{r+1} and V_j of q_j in \mathbb{C}^* such that $f : X_j = U_j \cap X \rightarrow V_j$ is an r -quasifibration.

As in Lemma 4.5, in local coordinates centered at p_j , respectively q_j , we have to consider the map $f : \mathbb{T}^{r+1} \rightarrow \mathbb{C}$ given by $f(z) = d_0 z_0 + \cdots + d_r z_r$. The fibers of this map $F_c = f^{-1}(c)$ are complements to hyperplane arrangements:

$$F_c = \mathbb{C}^r \setminus \{z_1 \dots z_r (d_1 z_1 + \cdots + d_r z_r - c) = 0\}.$$

The arrangements involved are in general position as in the work of Hattori [18]. The results there give that F_c has the following homotopy type

$$F_c \simeq \begin{cases} (T^r)^{(r)} & \text{if } c \neq 0, \\ (T^{r-1})^{(r-1)} & \text{if } c = 0, \end{cases}$$

where T^k is the real k -dimensional torus, and $(T^k)^{(l)}$ is its l -skeleton. That yields $\pi_1(F_c) = \mathbb{Z}^r$ and

$$\pi_i(F_c) = 0 \text{ if } \begin{cases} 1 < i < r & \text{and } c \neq 0, \\ 1 < i < r - 1 & \text{and } c = 0. \end{cases}$$

The long exact sequence of the pair (\mathbb{T}^{r+1}, F_c) provides the vanishing and surjection needed to obtain that the map $f : (\mathbb{T}^{r+1}, F_c) \rightarrow (\mathbb{C}, c)$ is an r -equivalence. Hence $f : \mathbb{T}^{r+1} \rightarrow \mathbb{C}$ is an r -quasifibration, and the local statement is proved. \square

Corollary 5.3. *Let F be any fiber of the map $f : X = \mathbb{T}^{r+1} \setminus D \rightarrow \mathbb{C}^*$. Then $\pi_i(F) = 0$ for $1 < i < r - 1$.*

Proof. The result follows from the long exact sequence of f which is an r -quasifibration with aspherical base and total space. \square

Proposition 5.4. *Let F be the general fiber of the map $f : X = \mathbb{T}^{r+1} \setminus D \rightarrow \mathbb{C}^*$, $r \geq 2$. Then $\pi_{r-1}(F) = 0$.*

Proof. Note that $f : E \rightarrow B$ is a fiber bundle with fiber F and aspherical base $B = \mathbb{C}^* \setminus \Lambda$. Thus we are done if we show that $\pi_{r-1}(E) = 0$.

Suppose the map $f : X = \mathbb{T}^{r+1} \setminus D \rightarrow \mathbb{C}^*$ defined by $f(\mathbf{x}) = x_0^{d_0} \cdots x_r^{d_r}$ has m special fibers $\Lambda = \{a_1, \dots, a_m\}$ (see Proposition 4.2), thus $E = X \setminus \bigcup_{j=1}^m \{x_0^{d_0} \cdots x_r^{d_r} = a_j\}$.

Consider now the divisor $\bar{D} = D \cup \bigcup_{j=1}^m \{x_{r+1} - aa_j^{-1} = 0\}$ in \mathbb{T}^{r+2} , for some $a \in \mathbb{C}^*$. It is readily seen that E can be identified with the special fiber $\bar{F}_a = \bar{f}^{-1}(a)$ of the map $\bar{f} : \bar{X} = \mathbb{T}^{r+2} \setminus \bar{D} \rightarrow \mathbb{C}^*$ given by $\bar{f}(\mathbf{x}, x_{r+1}) = x_0^{d_0} \cdots x_r^{d_r} x_{r+1}$. But $\pi_{r-1}(\bar{F}_a) = 0$ from Corollary 5.3. \square

Let F be the general fiber of the map $f : X = \mathbb{T}^{r+1} \setminus D \rightarrow \mathbb{C}^*$ defined by $f(\mathbf{x}) = x_0^{d_0} \cdots x_r^{d_r}$, where $D = \bigcup_{i=0}^r \bigcup_{j=1}^{n_i-1} \{x_i = \alpha_{i,j}\}$. Then X is clearly homotopy equivalent to $T_\Gamma \subset T^{r+1}$ the toric complex associated with the complete graph $\Gamma = K_{\mathbf{n}}$. Let T_Γ^χ be the infinite cyclic cover of T_Γ associated with the epimorphism $\chi : A_\Gamma \rightarrow \mathbb{Z}$ defined by $\chi(\sigma_{i,j}) = d_i$. The multiplication map $f : T_\Gamma \rightarrow \mathbb{S}^1$, now seen in the real context, is a fibration with fiber T_Γ^χ . Recall that T_Γ^χ is an Eilenberg-MacLane space for the Artin kernel $N_\Gamma^\chi = \ker \chi$.

Corollary 5.5. *For $r \geq 2$, the general fiber F is a smooth quasi-projective variety of dimension r that has the homotopy type of an r -dimensional CW-complex. The r -skeleton of the infinite cyclic cover T_Γ^χ has the homotopy type of F . In particular*

$$H_i(N_\Gamma^\chi) \cong \begin{cases} H_i(F) & \text{for } i \leq r, \\ \pi_r^*(F), & \text{for } i = r + 1, \end{cases}$$

where π_i^* are the coinvariants of π_i under the π_1 -action.

Proof. We know that $\pi_1(F) = N_\Gamma^\chi$. From Corollary 5.3 and Proposition 5.4 we have that $\pi_i(F) = 0$ for $1 < i < r$. That ensures $(T_\Gamma^\chi)^{(r)} = F$. \square

We obtain that the first non-vanishing homotopy group of the general fiber $\pi_r(F)$ is of infinite rank, as $\pi_r^*(F)$ is so.

All the special fibers F_s are smooth r -dimensional quasi-projective varieties sharing the homotopy type of an r -dimensional CW-complex. If $r \geq 3$, the $(r-1)$ -skeletons of F_s and of the general fiber F have the same homotopy type.

Corollary 5.6. *The special fiber F_s is a smooth r -dimensional quasi-projective variety. If $r = 2$, the special fiber F_s is aspherical and $\pi_1(F_s)$ is not isomorphic to N_Γ^χ . If $r \geq 3$, we have that*

$$\pi_i(F_s) \begin{cases} = N_\Gamma^\chi, & \text{if } i = 1, \\ = 0, & \text{if } 1 < i < r - 1, \\ \neq 0 & \text{if } i = r - 1, \end{cases} \quad H_i(N_\Gamma^\chi) \cong H_i(F_s) \text{ for } i < r$$

and an exact sequence

$$(5.1) \quad H_r(F_s) \rightarrow H_r(N_\Gamma^\chi) \rightarrow \pi_{r-1}^*(F_s) \rightarrow 0.$$

In other words, if $r \geq 3$, then $\pi_{r-1}^*(F_s)$ is a finitely generated abelian group of rank $b_r(F_s) - b_r(F)$.

Proof. We know that $\pi_1(F_s) = N_\Gamma^\chi$, if $r \geq 3$. From Corollary 5.3 we have that $\pi_i(F_s) = 0$ for $1 < i < r$. That ensures $F_s^{(r-1)} = F^{(r-1)} = (T_\Gamma^\chi)^{(r-1)}$. The exact sequence (5.1) comes from the long exact sequence in homology obtained from the classifying map $F_s \rightarrow T_\Gamma^\chi = K(N_\Gamma^\chi, 1)$. \square

REFERENCES

- [1] E. Artal, J.I. Cogolludo-Agustín, and D. Matei, *Quasi-projectivity, Artin-Tits groups, and pencil maps*, Topology of algebraic varieties and singularities, Contemp. Math., vol. 538, Amer. Math. Soc., Providence, RI, 2011, pp. 113–136.
- [2] M. Bestvina, *Questions in geometric group theory*, available at <http://www.math.utah.edu/~bestvina/eprints/questions-updated.pdf>, manuscript, 2000, (updated 2004).
- [3] M. Bestvina and N. Brady, *Morse theory and finiteness properties of groups*, Invent. Math. **129** (1997), no. 3, 445–470.
- [4] R. Bieri, *Normal subgroups in duality groups and in groups of cohomological dimension 2*, J. Pure Appl. Algebra **7** (1976), no. 1, 35–51.
- [5] ———, *Homological dimension of discrete groups*, second ed., Queen Mary College Mathematical Notes, Queen Mary College Department of Pure Mathematics, London, 1981.
- [6] K.-U. Bux and C. González, *The Bestvina-Brady construction revisited: geometric computation of Σ -invariants for right-angled Artin groups*, J. London Math. Soc. (2) **60** (1999), no. 3, 793–801.
- [7] R. Charney and M.W. Davis, *Finite $K(\pi, 1)$ s for Artin groups*, Prospects in topology (Princeton, NJ, 1994), Ann. of Math. Stud., vol. 138, Princeton Univ. Press, Princeton, NJ, 1995, pp. 110–124.
- [8] D.C. Cohen, M. Falk, and R. Randell, *Discriminantal bundles, arrangement groups, and subdirect products of free groups*, Preprint available at [arXiv:1008.0417](https://arxiv.org/abs/1008.0417) [math.AG].
- [9] C. De Concini and C. Procesi, *On the geometry of toric arrangements*, Transform. Groups **10** (2005), no. 3-4, 387–422.
- [10] ———, *Topics in hyperplane arrangements, polytopes and box-splines*, Universitext, Springer, New York, 2011.
- [11] G. Denham and A.I. Suciú, *Moment-angle complexes, monomial ideals and Massey products*, Pure Appl. Math. Q. **3** (2007), no. 1, Special Issue: In honor of Robert D. MacPherson. Part 3, 25–60.
- [12] A. Dimca, Ş. Papadima, and A.I. Suciú, *Quasi-Kähler Bestvina-Brady groups*, J. Algebraic Geom. **17** (2008), no. 1, 185–197.
- [13] ———, *Non-finiteness properties of fundamental groups of smooth projective varieties*, J. Reine Angew. Math. **629** (2009), 89–105.

- [14] A. Dold and R. Thom, *Quasifaserungen und unendliche symmetrische Produkte*, Ann. of Math. (2) **67** (1958), 239–281.
- [15] P. Griffiths and J. Harris, *Principles of algebraic geometry*, Wiley-Interscience [John Wiley & Sons], New York, 1978, Pure and Applied Mathematics.
- [16] M.A. Guest, M. Kwieciński, and B.-W. Ong, *Pseudo vector bundles and quasifibrations*, Hokkaido Math. J. **29** (2000), no. 1, 159–170.
- [17] A. Hatcher, *Algebraic topology*, Cambridge University Press, Cambridge, 2002.
- [18] A. Hattori, *Topology of C^n minus a finite number of affine hyperplanes in general position*, J. Fac. Sci. Univ. Tokyo Sect. IA Math. **22** (1975), no. 2, 205–219.
- [19] M. Jambu and Ş Papadima, *Deformations of hypersolvable arrangements*, Topology Appl. **118** (2002), no. 1-2, 103–111, Arrangements in Boston: a Conference on Hyperplane Arrangements (1999).
- [20] D. Matei, *Arrangements of hypersurfaces and Bestvina-Brady groups*, Oberwolfach Rep. **4** (2007), no. 3, 2347–2348.
- [21] J. P. May, *Weak equivalences and quasifibrations*, Groups of self-equivalences and related topics (Montreal, PQ, 1988), Lecture Notes in Math., vol. 1425, Springer, Berlin, 1990, pp. 91–101.
- [22] J. Meier, H. Meinert, and L. VanWyk, *Higher generation subgroup sets and the Σ -invariants of graph groups*, Comment. Math. Helv. **73** (1998), no. 1, 22–44.
- [23] L. Moci, *A Tutte polynomial for toric arrangements*, Trans. Amer. Math. Soc. **364** (2012), no. 2, 1067–1088.
- [24] M. Namba, *Branched coverings and algebraic functions*, Pitman Research Notes in Mathematics Series, vol. 161, Longman Scientific & Technical, Harlow, 1987.
- [25] P. Orlik and H. Terao, *Arrangements of hyperplanes*, Grundlehren der Mathematischen Wissenschaften, vol. 300, Springer-Verlag, Berlin, 1992.
- [26] Ş. Papadima and A.I. Suciuc, *Higher homotopy groups of complements of complex hyperplane arrangements*, Adv. Math. **165** (2002), no. 1, 71–100.
- [27] ———, *When does the associated graded Lie algebra of an arrangement group decompose?*, Comment. Math. Helv. **81** (2006), no. 4, 859–875.
- [28] ———, *Algebraic invariants for Bestvina-Brady groups*, J. Lond. Math. Soc. (2) **76** (2007), no. 2, 273–292.
- [29] ———, *Toric complexes and Artin kernels*, Adv. Math. **220** (2009), no. 2, 441–477.
- [30] R. Randell, *Homotopy and group cohomology of arrangements*, Topology Appl. **78** (1997), no. 3, 201–213.
- [31] J.-P. Serre, *Sur la topologie des variétés algébriques en caractéristique p* , Symposium internacional de topología algebraica International symposium on algebraic topology, Universidad Nacional Autónoma de México and UNESCO, Mexico City, 1958, pp. 24–53.
- [32] I. Shimada, *Fundamental groups of algebraic fiber spaces*, Comment. Math. Helv. **78** (2003), no. 2, 335–362.
- [33] J. Stallings, *A finitely presented group whose 3-dimensional integral homology is not finitely generated*, Amer. J. Math. **85** (1963), 541–543.
- [34] C.T.C. Wall, *Finiteness conditions for CW-complexes*, Ann. of Math. (2) **81** (1965), 56–69.

DEPARTAMENTO DE MATEMÁTICAS, IUMA, UNIVERSIDAD DE ZARAGOZA, C/ PEDRO CERBUNA 12, E-50009 ZARAGOZA SPAIN

E-mail address: artal@unizar.es, jicogo@unizar.es

INSTITUTE OF MATHEMATICS OF THE ROMANIAN ACADEMY, P.O. BOX 1-764, RO-014700 BUCHAREST ROMANIA

E-mail address: Daniel.Matei@imar.ro

URL: <http://www.imar.ro/~dmatei>