

# Accepted Manuscript

Assessment of the variability of atmospheric pollution in National Parks of mainland Spain

M. Escudero, A. Lozano, J. Hierro, O. Tapia, J. del Valle, A. Alastuey, T. Moreno, J. Anzano, Xavier Querol



PII: S1352-2310(16)30176-5

DOI: [10.1016/j.atmosenv.2016.03.006](https://doi.org/10.1016/j.atmosenv.2016.03.006)

Reference: AEA 14488

To appear in: *Atmospheric Environment*

Received Date: 31 July 2015

Revised Date: 17 February 2016

Accepted Date: 4 March 2016

Please cite this article as: Escudero, M., Lozano, A., Hierro, J., Tapia, O., Valle, J.d., Alastuey, A., Moreno, T., Anzano, J., Querol, X., Assessment of the variability of atmospheric pollution in National Parks of mainland Spain, *Atmospheric Environment* (2016), doi: 10.1016/j.atmosenv.2016.03.006.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Author's data requested for the accepted paper AEA14488:**

Name: Xavier Querol

Institution: Institute of Environmental Assessment and Water Research (IDAEA-CSIC),  
Jordi Girona 18-26, E-08034 Barcelona, SPAIN

Email: xavier.querol@idaea.csic.es

ACCEPTED MANUSCRIPT

# Assessment of the variability of atmospheric pollution in National Parks of mainland Spain

M. Escudero<sup>a,\*</sup>, A. Lozano<sup>a</sup>, J. Hierro<sup>a</sup>, O. Tapia<sup>a</sup>, J. del Valle<sup>a</sup>, A. Alastuey<sup>b</sup>, T. Moreno<sup>b</sup>, J. Anzano<sup>c</sup>

<sup>a</sup>Centro Universitario de la Defensa (CUD) de Zaragoza, Academia General Militar, Ctra. de Huesca s/n, E-50090 Zaragoza, SPAIN

<sup>b</sup>Institute of Environmental Assessment and Water Research (IDAEA-CSIC), Jordi Girona 18-26, E-08034 Barcelona, SPAIN

<sup>c</sup>Laser and Environment Laboratory, Department of Analytical Chemistry, Faculty of Sciences, University of Zaragoza, Pedro Cerbuna 12, E-50009 Zaragoza, SPAIN

---

## Abstract

Air quality in nine National Parks in mainland Spain was assessed analysing  $SO_2$ ,  $NO_x$ ,  $O_3$ ,  $PM_{10}$  and  $PM_{2.5}$  data from background stations. As emissions in and around parks are limited, the levels of primary pollutants are low. Concentrations of secondary pollutants are high especially in summer due to photochemical production. The geographical variability of pollutants responds to regional emission patterns and the dominant circulation regimes in different regions resulting in west-east gradients for  $O_3$  and  $PM$ . Seasonal variability of pollutants was also interpreted in virtue of transport scenarios, changes in photochemical activity and emissions variability.  $NO_x$  and  $SO_2$  maximize in winter due to higher emissions while  $O_3$  and  $PM$  do it in summer due to photochemical production, lower precipitation and, in the case of  $PM$ , the occurrence of African dust outbreaks. The diurnal evolution was interpreted in virtue of variability in emissions and changes in the Planetary Boundary Layer height.

*Keywords:*  $O_3$ ,  $NO_x$ ,  $SO_2$ ,  $PM_{10}$ ,  $PM_{2.5}$ , National Parks, Spain.

---

## 1. INTRODUCTION

Atmospheric pollutants are gases and particles present in the atmosphere which exert negative effects on health (WHO, 2006, 2008), climate (IPCC, 2013), ecosystems and vegetation (UNECE, 1988; Irwin et al., 2002; Fenn et al., 2003; Paoletti, 2006) and materials (Kucera and Fitz, 1995; de Leeuw, 2000; Screpanti and Marco, 2009). Regarding to their epidemiological and environmental effects, those classically monitored in air quality networks are sulfur dioxide ( $SO_2$ ), nitrogen oxides ( $NO_x = NO + NO_2$ ), ozone ( $O_3$ ) and atmospheric particulate matter ( $PM$ ).

---

\*Corresponding author  
Email address: [mescu@unizar.es](mailto:mescu@unizar.es) (M. Escudero)

Natural sources of  $SO_2$  include volcanoes and terrestrial or oceanic biogenic sources that emit significant amounts of precursor species such as dimethyl sulfide ( $(CH_3)_2S$ ) which interact chemically to produce  $SO_2$  (Andreae and Crutzen, 1997) while anthropogenic sources are mainly related with the use of fossil fuels (coal, fuel oil or wood). In recent years in Europe and North America, there has been a clear reduction in the use of these fuels but they are still used in power stations, ships and aircraft engines and in the residential sector (Querol et al., 2014).  $NO_x$  are generally emitted in combustion processes in the form of  $NO$  that, after interaction with other species such as  $O_3$  and volatile organic compounds ( $VOC_s$ ), is transformed to  $NO_2$ . The anthropogenic sources of  $NO_x$  include traffic, biomass burning and industry. Lightning, forest fires and bacterial processes in soils that convert ammonia ( $NH_3$ ) into  $NO_x$  are the major natural sources. Ozone is a trace gas with secondary origin which, except for a small proportion coming from stratospheric intrusions, is formed in the troposphere by means of photochemical reactions involving  $NO_x$  and  $VOC_s$  (Atkinson, 2000). Finally, the atmospheric aerosols or  $PM$  are solid and liquid particles in the atmosphere with a wide variety of sizes, compositions and origins. Size is essential to evaluate  $PM$  toxicity because finer particles have a greater capacity to penetrate into the respiratory tract (Dockery et al., 1993). Following epidemiological criteria, two size fractions are generally monitored:  $PM_{10}$  and  $PM_{2.5}$  (particles with less than 10 and 2.5  $\mu m$  of diameter respectively). The heterogeneity of  $PM$  explains the large number of potential sources (EC, 2004). Oceans, soils, volcanoes and vegetation are major natural sources of primary  $PM$  and of gaseous precursors ( $SO_2$ ,  $NO_x$ ,  $COV_s$ ,  $NH_3$  among others). The main anthropogenic sources are traffic, industry, the residential sector and agricultural and livestock activities.

The National Parks (NP) are areas with significant natural and environmental features which are legally recognised by a nation in order to protect their resources and biodiversity with the objective of preserving them for future generations. The classification of a region as a NP represents the maximum figure of environmental protection granted in national legislation. In North America the importance of monitoring and controlling levels of atmospheric pollutants in National Parks has been largely recognised. Monitoring networks for different atmospheric parameters have been active for over 25 years (Percy and Karnosky, 2007) controlling gaseous pollutants such as  $SO_2$ ,  $NO_x$  and  $O_3$ , speciation of  $PM$  and visibility and deposition. NP also implements field campaigns with the objective of obtaining a deeper insight on certain pollutants such as  $O_3$  using passive samplers or mobile monitors and other photochemical oxidant pollutants. These data have been employed in a considerable number of research studies. The first efforts focused on the study of reactive gases ( $SO_2$  and  $NO_x$ ) and  $PM$  and their link with atmospheric visibility (Eatough et al., 1997; Pitchford et al., 2000; Malm et al., 2005; McMeeking et al., 2005; Engling et al., 2006). Excess nitrogen deposition in NP in the United States has been addressed by several authors (Burns, 2003; Fenn et al., 2003; Beem et al., 2010; Benedict et al., 2013; Ellis et al., 2013; Prenni et al., 2014) with emphasis on high elevated areas such as the

Rocky Mountains. The variety of ecological effects of  $O_3$  has motivated several studies (Temple, 1989; Ray, 2001; Burley and Ray, 2007; Preisler et al., 2010; Burley et al., 2014) highlighting the high levels of  $O_3$  existing in the parks located at the western regions of the US and the important impact of wildfires increasing  $O_3$  levels in NP (Preisler et al., 2010).

In Spain vigilance of air quality in the NP network has not been regular and has been mostly based on specific studies. Few studies have addressed changes in deposition in NP (Camarero and Aniz, 2010; García-Gómez et al., 2014; Barberán et al., 2014) revealing the importance of the monitoring of reactive nitrogen species because of the risk of exceedances of critical loads for different types of vegetation. Regarding pollutant gases, Sanz et al. (2007) conducted a study on  $O_3$  levels in different NP based on dosimetry campaigns, Villanueva et al. (2014) studied the levels of ozone and  $VOC_s$  in Cabañeros NP while Adame et al. (2014) performed a complete field campaign of atmospheric photo-oxidants in Doñana NP. Finally, other studies have evaluated levels of PM components in particular NP such as Doñana (de la Campa et al., 2009; González-Castanedo et al., 2015). Despite these efforts, it is still necessary to make an assessment of the levels of air quality in NP in Spain to provide consistent and comparable information which allows the detection of the potential problems and environmental hazards associated with air pollutants. The main objective of this study is to conduct such assessment seeking to quantify and interpret the differences in pollutant levels in NP with different geographical locations and topographical features and to determine seasonal behavioral patterns of atmospheric pollutants.

## 2. MATERIALS AND METHODS

### 2.1. Stations and data representativity

One of the limitations for the planning of this study was the scarcity of long term data on air quality parameters available inside the NP of Spain. The main reason is that most of NP do not have access to the national electrical grid due to preservation policies. Therefore, an initial selection of stations belonging to operational air quality networks was made searching for regional background or remote sites which could assure the adequate representativity for the parks. Requirements such as maximum possible proximity to the NP (always  $< 40$  km and in most cases less than 20 km) and similarity with the nearby park in aspects such as environmental conditions, altitude and topography (Table 1) were imposed.

Different authors have performed studies to evaluate the representativeness for average measurements of rural stations as the ones used in this work. Larssen et al. (1999) offered a definition of the area of representativeness of a monitoring station as the region in which the concentration does not differ from the concentration measured at the station by more than a specified amount. Based on this

definition, the authors estimate the radius of the area of representativeness for regional background stations ranges between 25 and 150 km while for remote stations it is 200-500 km. More recently, Henne et al. (2010) estimated the area of representativeness of rural stations for  $O_3$ ,  $NO_2$  and  $CO$  across Europe by evaluating the catchment areas for different advection periods. For the shortest advection period tested in that work (12 h), catchment areas with radii between 29 and 195 km were determined. Based on the relationship between annual concentrations of  $NO_2$  with land use characteristics, Janssen et al. (2012) also found large areas of representativeness (radius > 10 km) for rural stations in Belgium. Summarising, all these references demonstrate that the radius of the area of representativity of rural stations is larger than the distance from the NP and the sites selected for this study. Moreover, the analysis performed in this study deals with the average levels of pollutants and not with specific events of hourly or daily durations in which representativity is difficult to be assured. In consequence, it can be assumed that the data employed in this study is closely indicative of the air quality standards in the NP.

Based on the data from the selected stations, 9 of the 10 NP located in mainland Spain (Figure 1 and Table 1) have been characterised: Islas Atlánticas (IAT), Picos de Europa (PEU), Ordesa y Monte Perdido (ORD), Aigüestortes y lago de Sant Maurici (AIG), Sierra de Guadarrama (GUA), Monfragüe (MON), Cabañeros (CAB), Doñana (DON) and Sierra Nevada (SNE).

Station code	Name	Network	Lat. (°N)	Long. (°E)	Alt. (m.a.s.l.)	Nat. Park (dist.)	Data
ES0005R	Noia	EMEP	42.72	-8.92	685	IAT (~ 20 km)	$SO_2$ , $NO_x$ , $O_3$ , $PM_{10}$
ES1989A	Lario	Castilla y León	43.04	-5.09	1140	PEU1 (~ 15 km)	$SO_2$ , $NO_x$ , $O_3$
ES0008R	Niembro	EMEP	43.44	-4.85	115	PEU (~ 17 km)	$SO_2$ , $NO_x$ , $O_3$ , $PM_{10}$ , $PM_{2.5}$
ES1883A	Torrelisa	Aragón	42.46	0.18	1005	ORD (~ 10 km)	$SO_2$ , $NO_x$ , $O_3$ , $PM_{10}$
ES1982A	Montsec	Catalunya + IDAEA-CSIC	42.05	0.73	1570	AIG (~ 40 km)	$SO_2$ , $NO_x$ , $O_3$ , $PM_{10}$ , $PM_{2.5}$
ES1802A	El Atazar	Madrid	40.91	-3.47	940	GUA (~ 20 km)	$SO_2$ , $NO_x$ , $O_3$ , $PM_{10}$ , $PM_{2.5}$
ES0001R	SP de los Montes	EMEP	39.55	-4.35	917	CAB (~ 15 km)	$SO_2$ , $NO_x$ , $O_3$ , $PM_{10}$ , $PM_{2.5}$
ES1616A	Monfragüe	Extremadura	39.85	-5.94	376	MON (-)	$SO_2$ , $NO_x$ , $O_3$ , $PM_{10}$
ES0017R	Doñana	EMEP	37.05	-6.56	5	DON (~ 1 km)	$SO_2$ , $NO_x$ , $O_3$ , $PM_{10}$
ES0007R	Víznar	EMEP	37.24	-3.53	1230	SNE (~ 10 km)	$SO_2$ , $NO_x$ , $O_3$ , $PM_{10}$ , $PM_{2.5}$

Table 1: Monitoring stations and their related NP employed in this study.

The features of the 9 NP analysed in this study present strong differences. Five of them are high-mountain ecosystems from which 3 are Atlantic (PEU, ORD and AIG) and 2 are Mediterranean (GUA and SNE), 2 more are located in Mediterranean medium-mountain terrain (CAB and MON) while IAT and DON represent low altitude terrestrial-maritime and wetland ecosystems respectively. Although anthropogenic emissions from inside NP are practically non-existent, external sources located in the surroundings can be relevant in some cases. Important cities are present near IAT (Vigo), GUA (Madrid and Segovia), DON (Seville, Huelva, Jerez de la frontera and Cádiz) and SNE (Granada). Regarding traffic sources, major roads run in the vicinity of GUA, DON, SNE and PEU while important




Figure 1: Location of the NP and the monitoring stations used in this study.

industrial estates can be found in Huelva and Gibraltar which may influence Doñana NP air quality.

The parameters used in this study are regularly monitored in air quality stations ( $SO_2$ ,  $NO_x$ ,  $O_3$ ,  $PM_{10}$  and  $PM_{2.5}$ ). Data was collected making use of standard methods as reflected in the EU Directive 2008/50/EC:  $SO_2$  was measured by UV fluorescence,  $NO_x$  by chemiluminescence and  $O_3$  by UV photometry in the three cases by means of automatic monitors. Daily data of  $PM_{10}$  and  $PM_{2.5}$  were obtained by the standard gravimetric method in all cases except in El Atazar where automatic equipment based on  $\beta$  attenuation was used applying a correction factor of, obtained after intercomparison with the gravimetric method (1.0 for 2008-2010 and 0.73 for 2011-2012). Complete time series from 2008 to 2012 were always available except in Montsec where monitoring began in 2010. High data availability was obtained in all cases ( $> 90\%$  for automatic data and  $> 80\%$  for  $PM$ ). A particular consideration on the characteristics of  $NO_x$  and  $SO_2$  monitors must be mentioned here since trace level analyzers were available in all the stations with the exception of Lario, Torrelisa, El Atazar and Monfragüe. The detection limit of a regular monitor is around 0.5 ppb (equivalent to 1.31

$\mu\text{g}/\text{m}^3$  for  $\text{SO}_2$  and  $0.94 \mu\text{g}/\text{m}^3$  for  $\text{NO}_x$ ), approximately one order of magnitude higher than in trace  
115 level ones. When ambient concentrations are below detection limit (a situation that can often occur  
in background stations), the measurement is usually set at a fixed level generally corresponding to  
50% of the detection limit, generating an upward bias in stations equipped with regular monitors with  
respect to those with trace level devices.

## 2.2. Meteorological analyses

120 In order to characterise the origin of air masses, 48-hours back-trajectories were computed once a  
day for the period 2008-2012 for each NP with HYSPLIT4 (Draxler and Hess, 1998) model using GDAS  
0.5-deg meteorological data. Clustering analyses of the back-trajectories were also performed with  
HYSPLIT4 to quantify the occurrence of transport regimes over each NP. A similar clustered trajectory  
analysis was employed before for characterising atmospheric transport in three NP in western US in  
125 a study by Hafner et al. (2007). In this work it was demonstrated that the prediction of precipitation  
and fine particle concentrations in remote areas improved for short period trajectory clustering with  
respect to longer periods. This result suggests that the chosen 48-h clusters are adequate although the  
importance of long range transport can be slightly underestimated.

An analysis of planetary boundary layer (PBL) heights, every three hours, was also performed. It  
130 should be noticed that GFS/GDAS code suffered a major upgrade on July 27th 2010, which corresponds  
to the studied period. This upgrade affects the reported values of PBL heights, with a clear reduction  
in maximum PBL heights after the upgrade, in all the studied locations. This lack of homogeneity in  
the available GDAS PBL heights is the reason why ERA-Interim data with 0.75-deg resolution were  
used as a source of PBL heights instead of GDAS 0.5-deg data. Nevertheless, a comparison between  
135 PBL heights as obtained from ERA-Interim and GDAS during the period considered was checked  
in Figure 2 for four specific locations: Doñana (ES0017R) which is close to sea in the south of the  
Iberian Peninsula, Vízcar (ES0007R) which is a southern mean mountain location, Montsec (ES1982A)  
which is a northern mean mountain location and Monfragüe (ES1616A) which lies in a southwestern  
hilly area. A simple moving average of the PBL height at 15:00 UTC was obtained from a 29 days  
140 sample centered on the considered date; during the first and the last period of 29 days the sample was  
truncated so as to fit into the 2008-2012 period <sup>1</sup>.

An analysis of figure 2 reveals the reduction of PBL GDAS heights after the code upgrade. In  
locations where GDAS data were higher than ERA-Interim data, this reduction has improved the  
agreement between both data sources; however, in locations where GDAS data was either lower or

---

<sup>1</sup>It means, for instance, that the first value is the moving average of the 15 first days, whereas the last value in the  
moving average of the 15 last days.






Figure 2: Number of days since Jan 1st 2008 in the X axis versus moving average of PBL height at 15:00 UTC, Y axis, in four locations: a) DON, b) SNE, c) AIG and d) MON. The reported value of PBL height is a simple mean of the actual values during the 29 days centered around the considered date, at 15:00 UTC every day. ERA-Interim values versus GDAS ones.

145 similar to ERA-Interim data before the upgrade, this reduction has increased the discrepancies between both sources. The particular case of Víznaar would be worth of a more refined study.

### 3. RESULTS

#### 3.1. Average levels

Due to their distant location from major sources, the average concentrations of primary pollutants in the NP are low (Figure 3 and Table 2). This occurs for  $SO_2$  with mean levels in the range  
 150  $1\text{-}2 \mu\text{g}/\text{m}^3$ . However, differences are observed between the stations equipped with trace level analyzers ( $0.44\text{-}1.08 \mu\text{g}/\text{m}^3$ ) and the rest, equipped with regular monitors ( $1.55\text{-}2.28 \mu\text{g}/\text{m}^3$ ) reflecting the upward bias of regular monitors explained above.

		ES1989A-PEU	ES0008R-PEU	ES1802A-GUA	ES1616A-MON	ES0001R-CAB	ES1883A-ORD	ES0017R-DON	ES0005R-IAT	ES1982A-AIG	ES0007R-SNE
$SO_2$	Mean	1.72	1.08	2.28	1.55	0.44	1.60	0.75	0.76	0.96	0.74
	Std	2.05	1.61	1.51	1.49	0.57	1.62	1.13	1.42	1.16	0.98
	P90	3.80	2.19	4.00	3.56	0.82	4.00	1.47	1.78	2.20	1.63
$NO_x$	Mean	4.54	5.20	7.38	9.26	3.35	7.63	6.31	3.10	1.87	6.84
	Std	10.10	4.83	8.42	6.53	3.28	4.99	5.91	3.17	1.35	9.92
	P90	14.00	10.54	12.00	15.65	6.31	13.44	13.17	6.17	3.30	15.00
$O_3$	Mean	61.69	67.61	82.58	68.12	83.92	73.92	63.07	71.98	89.53	83.96
	Std	23.75	18.07	25.15	30.11	20.70	20.49	28.56	22.54	18.83	19.83
	P90	91.00	91.14	114.00	107.60	111.30	102.00	100.20	99.36	116.00	109.80
$PM_{10}$	Mean	—	13.35	19.32	—	—	13.35	—	—	—	—
	Std	—	8.15	13.10	—	—	8.15	—	—	—	—
	P90	—	22.21	34.00	—	—	22.21	—	—	—	—
$PM_{2.5}$	Mean	—	7.89	8.28	—	—	7.89	—	—	—	—
	Std	—	4.12	5.45	—	—	4.12	—	—	—	—
	P90	—	12.95	15.00	—	—	12.95	—	—	—	—

Table 2: Average, standard deviation and 90<sup>th</sup> percentile of the concentrations of  $SO_2$ ,  $NO_x$ ,  $O_3$ ,  $PM_{10}$  and  $PM_{2.5}$  for the period 2008-2012 in the 10 monitoring stations used in this study.

Average  $NO_x$  levels in the study period ranged between 1.9 and 9.3  $\mu g/m^3$  with the highest concentrations in MON, ORD and GUA although in these sites it may be reflected again the use of regular analyzers with higher detection limits. The proximity to traffic ways has a clear impact on  $NO_x$  concentrations as observed in DON, SNE or PEU (5.2-6.3  $\mu g/m^3$ ) compared with sites located further from important roads such as CAB, IAT or AIG (1.9-3.4  $\mu g/m^3$ ). Moreover, in the case of PEU the presence of power plants in the region may also increase  $NO_x$  levels (Querol et al., 2014).

The Directive EU/50/2008 establishes critical levels for protection of the vegetation for  $SO_2$  (20  $\mu g/m^3$  for both the of the annual and winter period averages) and  $NO_x$  (30  $\mu g/m^3$  for the annual mean) that were not surpassed during the study period in the analysed sites.

$PM_{10}$  and  $PM_{2.5}$  concentrations ranged in 7-19  $\mu g/m^3$  and 6-11  $\mu g/m^3$  respectively. The highest levels corresponded to locations eventually affected by anthropogenic sources such as densely populated areas as Madrid (GUA) and Granada (SNE), or important traffic ways like PEU and DON. The impact of African dust intrusion may also increase  $PM$  levels especially in southern areas as SNE. The remaining stations are less exposed to anthropogenic emissions and, consequently, present lower  $PM$  levels.

As it is the case of other rural areas across the Mediterranean basin (Sicard et al., 2013), mean  $O_3$  levels in the analysed stations reached considerable values in the study period (61.7-89.5  $\mu g/m^3$ ). The strong emissions of precursors occurring in the highly industrialised and populated Mediterranean flank help to increase  $O_3$  levels in this area (73.9-89.5  $\mu g/m^3$ ) in contrast with western Iberia (61.7-72.0  $\mu g/m^3$ ) although climatic effects also are important.

Apart from the variability associated with emission patterns, climatic conditions are essential to an understanding of the spatial changes in pollutants levels. The atmospheric circulation over each




Figure 3: Average levels of atmospheric pollutants in NP of mainland Spain for the period 2008-2012

NP has been studied considering five source areas: Atlantic Ocean (ATL), European Continent (EU), Mediterranean Sea (MED), Northern African Continent (NAF) and Iberian Peninsula (PIB). As shown in Figure 4, the transport of Atlantic air masses was the most common situation in all NP but the frequency was higher over west of the PIB (53-80% of the days) than in the remaining NP (30-51%).

180 ATL episodes imply the arrival of clean oceanic air masses sometimes with precipitation associated with frontal systems. This generate ventilation and, in occasions, atmospheric wash-out.

PIB events occur when lack of prevalent advective conditions were observed. Typically, anticyclonic conditions over the Iberian Peninsula give rise to this scenario and, in consequence, precipitation is scarce. During winter PIB episodes, near-ground inversion layers develop leading to the accumulation of




Figure 4: Proportion of air mass transport scenarios over NP in mainland Spain based on clustering 48-h back-trajectories at surface level in the period 2008-2012

185 pollutants in urban and industrial locations. Summer PIB episodes are generated by the displacement of the Azores high over the western Mediterranean basin and the development of a thermal low over the central plateau of the Iberian Peninsula. The convergence of winds from the coastal areas, through natural passes, towards the centre of the Peninsula (Millán et al., 1996) and the enhanced mesoscale circulation yield to recirculation and aging of air masses. As shown in Figure 4, PIB transport is  
 190 frequent over NP at eastern and central Iberia (33-48% of the days), and less frequent over northern and western parks (14-26% of the days).

EU transport is rare in parks located in the centre, east, southwest and southeast (1-5% of the days) and more frequent in those located near the European continent (7-13% of the days). MED episodes are less frequent in northern NP (1-9%) than in southern ones (11-19%) mainly because the same  
 195 synoptic situations inferring EU transport over the north often generate MED events at the southern part of the peninsula. Precipitation is common under MED transport due to the moist character of Mediterranean air masses.

The occurrence of NAF episodes increase in frequency from north to south and from west to east (3-5% on the south and the east, and 1-2% on the north). These proportions are low compared with  
 200 previous studies (Escudero et al., 2005; Pey et al., 2013) due to two main reasons. Firstly, in order

to avoid excessive computational effort in the clustering analyses, 48-h back-trajectories have been computed so long range transport such as NAF can be underestimated. Secondly, the origin of air masses has been elucidated here exclusively from back-trajectories clustering while in previous studies sources of information such as model outputs or satellite imagery were used for detecting dust.

205  $O_3$  and, in a lesser extent,  $PM$  show a west-east gradient closely related with the variability of the occurrence of ATL episodes and opposite to the variability of the occurrence of PIB events. The greater ventilation and precipitation provided by ATL transport in the western flank of the Iberian Peninsula help to keep  $PM$  and  $O_3$  lower than in oriental and central areas. In DON,  $NO$  emissions from the industrial estates located near this NP may have resulted in lowering  $O_3$  levels. Moreover, 210 the specific orography of the eastern side of the Iberian peninsula characterized by mountain ranges parallel to the coast interspersed by valleys, generates regional circulation and aging of air masses resulting in elevated  $O_3$  and  $PM$  levels (especially the secondary fraction) in background locations (Millán et al., 1996; Rodríguez et al., 2002). Finally, the higher frequency of NAF episodes in south and eastern NP contribute to increase  $PM$  levels in those parks (Escudero et al., 2005, 2007).

### 215 3.2. Seasonal variability

The seasonal variability of pollutant levels in NP (Figure 5) was associated with emission patterns and with changes in ambient conditions. The concentrations of certain pollutants, such as  $SO_2$ , exhibit very low variability between summer-spring and autumn-winter. Only in some NP (PEU and IAT) light winter increases were observed associated with greater emissions from the residential sector and 220 coal power plants.

$NO_x$  concentrations showed higher levels in autumn-winter (20-60 % higher than in spring-summer). Winter maxima reflect the increment of emissions with respect to summer months. In addition, the reduced photochemical activity during winter results in lower  $NO_2$  destruction by photolysis. Finally, lower  $O_3$  concentrations during autumn-winter also inhibits the elimination of  $NO$  by tritiation. Ex- 225 ceptions were found in ORD, DON and SNE where maxima in summer is probably associated with an regional increment in traffic during summer vacations.

$O_3$  levels present strong spring-summer increases associated with enhanced photochemical activity. Average concentrations in early spring (March), are comparable to summer levels in most NP. Western parks (IAT, PEU and DON) show maxima in April decreasing slightly during summer linked with an 230 earlier decline of solar radiation in that sector than in the remaining NP where maxima concentrations are observed during summer (CEAM, 2014).

The highest  $PM_{10}$  and  $PM_{2.5}$  levels in NP are generally recorded in spring-summer but the differences between summer and winter are greater in  $PM_{10}$ . Spring-summer concentrations increase in 20-80% for  $PM_{10}$  and 10-40% for  $PM_{2.5}$ , with the weakest increments being registered in the




Figure 5: Monthly mean concentrations of atmospheric pollutants in NP of mainland Spain for the period 2008-2012

235 northwestern locations (IAT and PEU). This is explained by the greater frequency of African dust intrusions over the Iberian Peninsula (Escudero et al., 2005, 2007), the lower precipitation reducing atmospheric washout and favouring *PM* resuspension, the enhanced photochemical production of secondary aerosols and the stronger vertical development of the mixing layer which eases the arrival towards NP of pollutants emitted in urban and industrial spots. Another maximum, secondary in the case of *PM*<sub>10</sub> but of the same importance or even greater than the spring-summer peak for *PM*<sub>2.5</sub>, is

240

observed in February-March. These late winter maxima are associated with two processes. Firstly, the effect of winter African dust intrusions over the Iberian Peninsula with a outstanding impact on  $PM$  levels because, contrarily to summer African dust events, transport occurs at surface level (Escudero et al., 2005). Moreover, the occurrence of anticyclonic situations in late winter with enough radiation may activate mountain breezes facilitating the arrival of pollutants to natural areas (Pey et al., 2010). These two processes combined with the milder summer conditions in northwestern NP (PEU and IAT) explain why average winter concentrations are higher than in summer in those sites.

### 3.3. Daily variability

One of the key parameters to understand the variability of the levels of pollutants during the day in remote areas is the evolution of the altitude and stability of the PBL. A strong vertical development of an unstable PBL aided by the enhancement of coastal and mountain breezes (Pey et al., 2010) facilitate a rapid transport of pollutants from urban and industrial areas towards rural sites. An unstable PBL may be decomposed into three sublayers: a surface layer, the lowest part of the PBL where averaged winds show a logarithmic profile versus elevation above the surface with strong turbulence production, a mixed layer, the core zone of the PBL where meteorological variables are well mixed and turbulence decays, and an entrainment or inversion layer, a transition zone between the PBL and the free troposphere where there is a mixture of air masses coming from both zones accompanied by strong variations of the meteorological variables.

Before realizing any analysis based on the height and stability of the PBL, it should be pointed out that there is not a unique definition of these properties. With regard to the estimation of the PBL height, see for instance (Seidel et al., 2012) and references therein, where ten different definitions were compared. Concerning the stability of the PBL, most studies are based on either the bulk Richardson number at the surface layer, or the Obukhov length (which is related to the height at which the TKE production of turbulence because of buoyancy equals its production because of shear flow), or the value of the surface heat flux (Zhang et al., 2014). In this work, PBL height values were taken from ERA-Interim data. According to the documentation of the IFS/ECWMF numerical code, the PBL height is defined as the level where the bulk Richardson number, based on the differences between quantities at that level and the lowest model level, reaches a critical value of 0.25. The definition is a slight modification of the one proposed by (Troen and Mahrt, 1986): virtual dry static energy is used instead of virtual potential temperature.

With regard to the stability criterion, IFS/ECWMF uses the sign of the Obukhov length,  $L_O < 0$  for unstable conditions,  $L_O > 0$  for stable conditions and  $L_O = 0$  for neutral conditions. In its turn, the sign of the Obukhov length is the opposite of the sign of the virtual temperature flux, see eqn.(1), in the surface layer, where positive flux is assumed to go upwards. The virtual temperature flux depends

on heat and humidity fluxes and it may go to zero only if there is not momentum flux. Heat and momentum fluxes are provided by ERA-Interim.

$$Q_{0v} = \frac{J_{sh}}{\rho C_p} + \varepsilon T_n \frac{J_{lh}}{\rho L_v} \quad (1)$$

where  $J_{sh}$  is the sensible heat flux,  $J_{lh}$  is the latent heat flux,  $\rho$  is moist air density,  $C_p$  is the heat capacity at constant pressure for dry air,  $\varepsilon \approx 0.6063$  is a constant used to compute virtual temperatures,  $T_n$  is an absolute reference temperature taken at a point close to the surface ( $T$  at 2m of elevation is provided by ERA-Interim) and  $L_v$  is the latent heat of vaporization of water. Since, the sign of  $Q_{0v}$  is the only relevant quantity, one may simply check the sign of  $\rho Q_{0v} = (J_{sh}/C_p) + \varepsilon T_n (J_{lh}/L_v)$  without the need to compute air density near surfaces.

Figure 6 shows the average diurnal cycle of the PBL height for each month in the studied NP for 2008-2012. Every diurnal cycle is divided into eight steps with a three hour separation between them. PBL altitude maximizes in all sites around 15 h UTC coinciding with the maximum surface warming. Three different groups of sites can be differentiated regarding the PBL daily cycles. In coastal locations (IAT, DON and PEU1), the influence of marine winds, intensified in the warm season, limits the vertical extent of the PBL. In continental locations in the North of the Iberian Peninsula (PEU, ORD and AIG), the PBL has a strong development in spring-summer reaching averages around 2000 metres. This effect, which is due to an intense surface heating in peninsular areas isolated from maritime influence, is even stronger in continental locations in the South and the Center of the Iberian Peninsula (MON, GUA, CAB and SNE) with averages above 2500 metres. During the cold season the vertical extent of the mixing layer is lower due to the reduced surface heating (maximum averages between 500 and 1000 metres) and the main difference between continental and coastal sites is the much lower amplitude of the diurnal cycle in coastal sites.

This average behaviour of the PBL can be affected sporadically during African dust outbreaks as explained by Pandolfi et al. (2014). According to these authors, the PBL height decreases during African dust episodes due to two major factors related with the presence of the dust layer. Firstly, the decay in incoming solar radiation by dispersion and backscattering caused by mineral aerosols and the generation of a low-level inversion layer at the altitude where the African dust layer ends. Nevertheless, this a phenomenon that takes place under the influence of African dust transport which, as detailed above, is not dominant among transport scenarios so the actual impact on the average behaviour of the PBL (which is analysed in this paper) is not expected to be decisive.

Figure 7 shows the diurnal cycle of the probability that the PBL be unstable for each month in three selected stations (DON, AIG and SNE) during the studied period with a three hour time step. No more locations were selected because inland sites showed an intermediate behaviour between that of SNE and that of AIG, whereas coastal locations behaved like DON. For a given location, month






Figure 6: Mean daily cycles of the PBL height for each month in NP of Spain during the period 2008-2012

and hour, the distribution of the stability is rather sharp: there is a clear dominance of one of the two conditions and transitions towards the opposite situation are fast. The most striking feature is the high probability of stable conditions during the central hours of the day in coastal locations. The reason being that the sea surface keeps much cooler than the surrounding air what enforces a net sensible heat flux directed downwards (towards the sea) which is associated with the formation of a stable PBL. In the morning and in the late afternoon, differences of temperature between the sea and the surrounding air layer are not so high and the PBL turns unstable. An analysis of momentum fluxes revealed that there were no exact neutral conditions in any studied point at any considered time.

Figure 8 shows the average daily cycles of the concentrations of pollutants separately for winter (October-March) and summer (April-September).  $SO_2$  levels increase slightly in the central hours of the day although levels are really close to detection limit so those increments are in the same order of measurements errors. These midday increases are also detected in  $NO_x$  concentrations and may be associated with the PBL height evolution presented above. The development of an unstable PBL with increasing altitude at the central hours facilitates the transport of pollutants from urban and industrial areas towards background regions which lie in the direction of the prevailing winds at the top of the PBL. In some parks (SNE, PEU and DON) two  $NO_x$  maxima are observed one in the morning and another in the evening. These are associated with traffic emissions in the rush hours indicating exposure to traffic sources due to the proximity to major roads or cities. Other stations




Figure 7: Mean daily cycles of the probability that the PBL be unstable for DON, AIG and SNE during the period 2008-2012.

320 (ORD, GUA or AIG) just present a single peak in the concentrations of  $NO_x$  during the afternoon. The direct impact of traffic sources on these sites is limited so it is the development of the PBL which progressively allows the arrival of well mixed air masses loaded with  $NO_x$  giving rise to a single concentration maximum.

The diurnal evolution of  $NO_x$  also present seasonal differences. As stated before, winter levels are 325 higher than summer concentrations due to three major factors associated with wintertime: Higher emissions, reduction in the photochemical activity which inhibits  $NO_2$  photolysis and the reduction of  $NO$  titration limited by low  $O_3$  concentrations. Another noticeable feature is that, in the sites where two daily maxima are detected, the morning peak occurs earlier during spring-summer than during autumn-winter. The stronger vertical extent of the PBL in summer along with the intensification of 330 mountain breezes cause by higher temperature and radiation, facilitates the rapid arrival of  $NO_x$  from emission sources to the natural areas while in winter a more limited and slower growth of the mixing layer, which may be absent under thermal inversion conditions, delays the arrival of  $NO_x$ .

The mean daily cycles of  $NO_x$  have also been studied disaggregating weekdays from weekends (Figure 9). This approach was used in Escudero et al., (2014) in order to discriminate the impact of 335 anthropogenic activity on the levels of  $NO_x$  given that emissions derived from human activities fall drastically during weekends. In some NP (PEU, GUA, DON and SNE) present  $NO_x$  increments in




Figure 8: Winter and summer mean daily cycles of the concentrations of gaseous pollutants in NP of Spain for the period 2008-2012

weekdays with respect to weekends. These are especially relevant in the hours when concentrations reach their maximum values in the central hours or during rush hours. This must be interpreted as the direct or semi-direct impact of anthropogenic sources. In the remaining sites, with a higher degree of isolation from anthropogenic sources, daily cycles present very limited variations both in magnitude and morphology.

The daily variability of  $O_3$  is conditioned by meteorological factors and precursor gases emissions




Figure 9: Weekdays and weekends mean daily cycles of  $NO_x$  concentrations in National parks of Spain for the period 2008-2012

patterns. The concentration peaks are observed in the afternoon following the maximum photochemical activity. The reduction in concentrations during the night is not particularly marked in some sites  
 345 responding to processes involving air masses, previously loaded with  $O_3$  produced locally or regionally during daytime, which suffer recirculation and generate notable ozone concentrations in remote areas at night (Escudero et al., 2014).

#### 4. CONCLUSIONS

This work presents a comprehensive assessment of the levels of atmospheric pollutants in 9 Spanish  
 350 NP located in the Iberian Peninsula. This has been done working with data ( $SO_2$ ,  $NO_x$ ,  $PM_{10}$ ,  $PM_{2.5}$  and  $O_3$ ) corresponding to the period 2008-2012 obtained from regional background stations belonging to operative monitoring networks. These stations were selected after imposing different criteria in order to guarantee the representativeness with respect to the NP.

The natural and remote characteristics of NP, generally far from major emission sources, and the  
 355 progressive decline in the use of  $S$ -rich fuels result in low  $SO_2$  concentrations in all NP.  $NO_x$  levels were low although, in some parks (DON, SNE or PEU), the influence of main traffic pathways and certain cities generate higher concentrations. The seasonal variability of  $NO_x$  is characterised by higher winter levels due to greater emissions and reduced photochemical activity which reduces  $NO_2$  photolysis and, as consequence of lower  $O_3$  levels, the destruction of  $NO$  by titration. The analysis  
 360 of daily variability demonstrate the direct influence of traffic and other anthropogenic sources on the 3 NP mentioned above where two concentration peaks associated with traffic emissions in rush hours were observed. At these sites, marked reductions were detected for weekends compared to weekdays

confirming the impact of anthropogenic sources.

$PM_{10}$  and  $PM_{2.5}$  levels were moderate but, as in the case of  $NO_x$ , the sites with a higher presence  
365 of anthropogenic sources around them (GUA, SNE, PEU and DON) presented higher concentrations. In SNE and DON the influence of African dust outbreaks can also be a factor for those higher levels.  $PM$  variability is marked with higher summer levels due to higher frequency of African events, lower precipitation and increased photochemical production of secondary particles. Moreover, the high vertical extent of the PBL in summer facilitates the arrival of pollutants from urban and industrial  
370 areas to NP. A secondary maximum of  $PM$  is detected in February-March by the effect of African dust outbreaks and the enhancement of breeze dynamics.

Ozone concentrations are significantly high in NP in Spain. A west-east gradient is associated with climatic differences as the western Iberian Peninsula is more exposed to the Atlantic ventilation than the east. Moreover, the specific orography of the eastern side of the Iberian Peninsula aids to the  
375 regional circulation and aging of air masses resulting in elevated  $O_3$ . The photochemical origin of  $O_3$  results in a very marked seasonal variability with much higher levels in the spring-summer period.

The conclusions drawn in this work should motivate new research in this field. It is essential to continue collecting data on atmospheric pollutants in NP especially in cases like  $O_3$  due to the elevated levels of this pollutant in background areas and the difficulties for the implementation and evaluation  
380 of mitigation plans (Tapia et al., 2016). For this purpose it is basic to use trace level monitors for  $SO_2$  and  $NO_x$  and incorporate online  $PM$  measurements for describing time evolution. In order to complete the present work focused on long term variability, efforts describing spatial variability or vertical profiles (for example in mountainous regions) of pollutants in each NP should be carried out using dosimetry which is a useful and economically affordable technique. Deposition should be  
385 monitored in order to obtain information on incoming fluxes of relevant species such as trace metals or nitrogen.  $PM$  speciation and source apportionment studies can offer excellent estimation of major atmospheric sources affecting NP. Finally, studies dedicated to evaluate the impact of particular sources (traffic, industrial, agriculture o livestock) in certain NP may be useful to provide a better overview of air quality in these relatively remote but ecologically vulnerable locations.

## 390 5. ACKNOWLEDGEMENTS

This work was founded by the Fundación Biodiversidad (project AQ-NAT), the CUD (project CUD 2013/18) and the Government of Department of Research, Innovation and University of the Aragón Regional Government and the European Social Found through projects E15 and E75. Air quality data provision is acknowledge to the following entities: MAGRAMA, AEMET, IDAEA-CSIC and the  
395 Governments of Aragón, Extremadura, Catalunya, Madrid and Castilla y León. The authors gratefully

acknowledge the NOAA-ARL for GDAS data provision and for the use of HYSPLIT4 model. Finally, we would like to thank the two blind reviewers for their useful comments.

## References

- Adame JA, Martínez M, Sorribas M, Hidalgo PJ, Harder H, Diesch JM, Drewnick F, Song W, Williams  
400 J, Sinha V, Hernández-Ceballos MA, Vilà-Guerau de Arellano J, Sander R, Hosaynali-Beygi Z,  
Fischer H, Lelieveld J, De la Morena B. Meteorology during the domino campaign and its connection  
with trace gases and aerosols. *Atmospheric Chemistry and Physics* 2014;14(5):2325–42. doi:10.5194/  
acp-14-2325-2014.
- Andreae MO, Crutzen PJ. Atmospheric aerosols: Biogeochemical sources and role in atmospheric  
405 chemistry. *Science* 1997;276(5315):1052–8.
- Atkinson R. Atmospheric chemistry of {VOCs} and {NOx}. *Atmospheric Environment*  
2000;34(1214):2063–101. doi:http://dx.doi.org/10.1016/S1352-2310(99)00460-4.
- Barberán A, Henley J, Fierer N, Casamayor EO. Structure, inter-annual recurrence, and global-scale  
connectivity of airborne microbial communities. *Science of The Total Environment* 2014;487:187  
410 –95. doi:http://dx.doi.org/10.1016/j.scitotenv.2014.04.030.
- Beem KB, Raja S, Schwandner FM, Taylor C, Lee T, Sullivan AP, Carrico CM, McMeeking GR, Day  
D, Levin E, Hand J, Kreidenweis SM, Schichtel B, Malm WC, Collett JL. Deposition of reactive  
nitrogen during the Rocky Mountain Airborne Nitrogen and Sulfur (RoMANS) study. *Environmental  
Pollution* 2010;158(3):862–72. doi:http://dx.doi.org/10.1016/j.envpol.2009.09.023.
- 415 Benedict KB, Chen X, Sullivan AP, Li Y, Day D, Prenni AJ, Levin EJT, Kreidenweis SM, Malm  
WC, Schichtel BA, Collett JL. Atmospheric concentrations and deposition of reactive nitrogen in  
Grand Teton National Park. *Journal of Geophysical Research: Atmospheres* 2013;118(20):11875–87.  
doi:10.1002/2013JD020394; 2013JD020394.
- Burley JD, Bytnerowicz A, Ray JD, Schilling S, Allen EB. Surface ozone in Joshua Tree National Park.  
420 *Atmospheric Environment* 2014;87(0):95–107. doi:http://dx.doi.org/10.1016/j.atmosenv.  
2013.12.043.
- Burley JD, Ray JD. Surface ozone in Yosemite National Park. *Atmospheric Environment*  
2007;41(28):6048–62. doi:http://dx.doi.org/10.1016/j.atmosenv.2007.03.021.
- Burns DA. Atmospheric nitrogen deposition in the Rocky Mountains of Colorado and southern  
425 Wyoming: a review and new analysis of past study results. *Atmospheric Environment* 2003;37(7):921  
–32. doi:http://dx.doi.org/10.1016/S1352-2310(02)00993-7.

- Camarero L, Aniz M. El sistema de seguimiento de las aguas en el nodo LTER-Aigestortes: tendencias e indicadores de los impactos de la deposición atmosférica de N y S. *Revista Ecosistemas* 2010;19(2).
- de la Campa AS, Pio C, de la Rosa J, Querol X, Alastuey A, González-Castanedo Y. Characterization and origin of {EC} and {OC} particulate matter near the Doñana national park (sw Spain). *Environmental Research* 2009;109(6):671–81. doi:<http://dx.doi.org/10.1016/j.envres.2009.05.002>.
- CEAM . Informe Final Memoria Técnica Proyecto CONOZE- CONTaminación por OZono en España. Technical Report; Fundación Centro de Estudios Ambientales del Mediterráneo; 2014.
- Dockery DW, Pope CA, Xu X, Spengler JD, Ware JH, Fay ME, Ferris BG, Speizer FE. An association between air pollution and mortality in six U.S. cities. *New England Journal of Medicine* 1993;329(24):1753–9. doi:[10.1056/NEJM199312093292401](https://doi.org/10.1056/NEJM199312093292401). arXiv:<http://dx.doi.org/10.1056/NEJM199312093292401>; PMID: 8179653.
- Draxler RR, Hess GD. An overview of the HYSPLIT4 modelling system for trajectories, dispersion, and deposition. *Australian meteorological magazine* 1998;47(4):295–308.
- Eatough DJ, Du A, Joseph JM, Caka FM, Sun B, Lewis L, Mangelson NF, Eatough M, Rees LB, Eatough NL, Farber RJ, Watson JG. Regional source profiles of sources of  $SO_x$  at the Grand Canyon during project MOHAVE. *Journal of the Air & Waste Management Association* 1997;47(2):101–18. doi:[10.1080/10473289.1997.10464421](https://doi.org/10.1080/10473289.1997.10464421).
- EC . Second position paper on particulate matter. Technical Report; CAFE Working Group on Particulate Matter, European Commission; 2004.
- Ellis RA, Jacob DJ, Sulprizio MP, Zhang L, Holmes CD, Schichtel BA, Blett T, Porter E, Pardo LH, Lynch JA. Present and future nitrogen deposition to national parks in the United States: critical load exceedances. *Atmospheric Chemistry and Physics* 2013;13(17):9083–95. doi:[10.5194/acp-13-9083-2013](https://doi.org/10.5194/acp-13-9083-2013).
- Engling G, Herckes P, Kreidenweis SM, Malm WC, Collett JL. Composition of the fine organic aerosol in Yosemite National Park during the 2002 Yosemite aerosol characterization study. *Atmospheric Environment* 2006;40(16):2959–72. doi:<http://dx.doi.org/10.1016/j.atmosenv.2005.12.041>.
- Escudero M, Castillo S, Querol X, Avila A, Alarcón M, Viana MM, Alastuey A, Cuevas E, Rodríguez S. Wet and dry african dust episodes over eastern Spain. *Journal of Geophysical Research: Atmospheres* 2005;110(D18). doi:[10.1029/2004JD004731](https://doi.org/10.1029/2004JD004731); d18S08.

- Escudero M, Lozano A, Hierro J, del Valle J, Mantilla E. Urban influence on increasing ozone concentrations in a characteristic Mediterranean agglomeration. *Atmospheric Environment* 2014;99(0):322–32. doi:<http://dx.doi.org/10.1016/j.atmosenv.2014.09.061>.
- Escudero M, Querol X, Ávila A, Cuevas E. Origin of the exceedances of the European daily {PM} limit value in regional background areas of Spain. *Atmospheric Environment* 2007;41(4):730–44. doi:<http://dx.doi.org/10.1016/j.atmosenv.2006.09.014>.  
460
- Fenn ME, Baron JS, Allen EB, Rueth HM, Nydick KR, Geiser L, Bowman WD, Sickman JO, Meixner T, Johnson DW, Neitlich P. Ecological effects of nitrogen deposition in the Western United States. *BioScience* 2003;53(4):404–20. doi:10.1641/0006-3568(2003)053[0404:EEONDI]2.0.CO;2.
- García-Gómez H, Garrido J, Vivanco M, Lassaletta L, Rábago I, Avila A, Tsyro S, Sánchez G, Ortiz AG, González-Fernández I, Alonso R. Nitrogen deposition in Spain: Modeled patterns and threatened habitats within the Natura 2000 network. *Science of The Total Environment* 2014;485486(0):450–60. doi:<http://dx.doi.org/10.1016/j.scitotenv.2014.03.112>.  
465
- González-Castanedo Y, Sánchez-Rodas D, de la Campa AS, Pandolfi M, Alastuey A, Cachorro V, Querol X, de la Rosa J. Arsenic species in atmospheric particulate matter as tracer of the air quality of doana natural park (sw spain). *Chemosphere* 2015;119:1296–303. doi:<http://dx.doi.org/10.1016/j.chemosphere.2014.09.093>.  
470
- Hafner W, Solorzano N, Jaffe D. Analysis of rainfall and fine aerosol data using clustered trajectory analysis for National Park sites in the Western {US}. *Atmospheric Environment* 2007;41(14):3071–81. URL: <http://www.sciencedirect.com/science/article/pii/S1352231006012003>. doi:<http://dx.doi.org/10.1016/j.atmosenv.2006.11.049>.  
475
- Henne S, Brunner D, Folini D, Solberg S, Klausen J, Buchmann B. Assessment of parameters describing representativeness of air quality in-situ measurement sites. *Atmospheric Chemistry and Physics* 2010;10(8):3561–81. doi:10.5194/acp-10-3561-2010.
- IPCC . Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University press, 2013.  
480
- Irwin JG, Campbell G, Vincent K. Trends in sulphate and nitrate wet deposition over the United Kingdom: 1986/1999. *Atmospheric Environment* 2002;36(17):2867–79. doi:[http://dx.doi.org/10.1016/S1352-2310\(02\)00096-1](http://dx.doi.org/10.1016/S1352-2310(02)00096-1).  
485



- Janssen S, Dumont G, Fierens F, Deutsch F, Maiheu B, Celis D, Trimpeneers E, Mensink C. Land use to characterize spatial representativeness of air quality monitoring stations and its relevance for model validation. *Atmospheric Environment* 2012;59:492 – 500. doi:<http://dx.doi.org/10.1016/j.atmosenv.2012.05.028>.
- 490 Kucera V, Fitz S. Direct and indirect air pollution effects on materials including cultural monuments. *Water, Air, and Soil Pollution* 1995;85(1):153–65. doi:10.1007/BF00483697.
- Larssen S, Sluyter R, Helmis C. Criteria for EUROAIRNET. The EEA Air Quality Monitoring and Information Network. Technical Report; EEA; 1999.
- de Leeuw FA. Trends in ground level ozone concentrations in the European Union. *Environmental Science & Policy* 2000;3(4):189 –99. doi:[http://dx.doi.org/10.1016/S1462-9011\(00\)00090-3](http://dx.doi.org/10.1016/S1462-9011(00)00090-3).
- 495 Malm WC, Day DE, Carrico C, Kreidenweis SM, Collett JL, McMeeking G, Lee T, Carrillo J, Schichtel B. Intercomparison and closure calculations using measurements of aerosol species and optical properties during the Yosemite aerosol characterization study. *Journal of Geophysical Research: Atmospheres* 2005;110(D14). doi:10.1029/2004JD005494; d14302.
- 500 McMeeking GR, Kreidenweis SM, Carrico CM, Collett JL, Day DE, Malm WC. Observations of smoke-influenced aerosol during the Yosemite aerosol characterization study: 2. Aerosol scattering and absorbing properties. *Journal of Geophysical Research: Atmospheres* 2005;110(D18). doi:10.1029/2004JD005624; d18209.
- Millán M, Salvador R, Mantilla E, nano BA. Meteorology and photochemical air pollution in Southern Europe: Experimental results from {EC} research projects. *Atmospheric Environment* 505 1996;30(12):1909 –24. doi:[http://dx.doi.org/10.1016/1352-2310\(95\)00220-0](http://dx.doi.org/10.1016/1352-2310(95)00220-0).
- Pandolfi M, Tobias A, Alastuey A, Sunyer J, Schwartz J, Lorente J, Pey J, Querol X. Effect of atmospheric mixing layer depth variations on urban air quality and daily mortality during Saharan dust outbreaks. *Science of The Total Environment* 2014;494495:283 –9. doi:<http://dx.doi.org/10.1016/j.scitotenv.2014.07.004>.
- 510 Paoletti E. Impact of ozone on Mediterranean forests: A review. *Environmental Pollution* 2006;144(2):463 –74. doi:<http://dx.doi.org/10.1016/j.envpol.2005.12.051>.
- Percy K, Karnosky D. Air quality in natural areas: Interface between the public, science and regulation. *Environmental Pollution* 2007;149(3):256 –67. doi:<http://dx.doi.org/10.1016/j.envpol.2007.05.023>; air Pollution and Vegetation Effects Research in National Parks and Natural Areas: Implications for Science, Policy and Management.

- Pey J, Pérez N, Querol X, Alastuey A, Cusack M, Reche C. Intense winter atmospheric pollution episodes affecting the western mediterranean. *Science of The Total Environment* 2010;408(8):1951–9. doi:<http://dx.doi.org/10.1016/j.scitotenv.2010.01.052>.
- 520 Pey J, Querol X, Alastuey A, Forastiere F, Stafoggia M. African dust outbreaks over the mediterranean basin during 2001-2011: Pm<sub>10</sub> concentrations, phenomenology and trends, and its relation with synoptic and mesoscale meteorology. *Atmospheric Chemistry and Physics* 2013;13(3):1395–410. URL: <http://www.atmos-chem-phys.net/13/1395/2013/>. doi:10.5194/acp-13-1395-2013.
- Pitchford M, Green M, Kuhns H, Farber RJ. Characterization of regional transport and disper-  
525 sion using project MOHAVE tracer data. *Journal of the Air & Waste Management Association* 2000;50(5):733–45. doi:10.1080/10473289.2000.10464119.
- Preisler HK, Zhong SS, Esperanza A, Brown TJ, Bytnerowicz A, Tarnay L. Estimating contribution of wildland fires to ambient ozone levels in National Parks in the Sierra Nevada, California. *Environmental Pollution* 2010;158(3):778–87. doi:<http://dx.doi.org/10.1016/j.envpol.2009.10.007>.
- 530 Prenni A, Levin E, Benedict K, Sullivan A, Schurman M, Gebhart K, Day D, Carrico C, Malm W, Schichtel B, Jr. JC, Kreidenweis S. Gas-phase reactive nitrogen near Grand Teton national park: Impacts of transport, anthropogenic emissions, and biomass burning. *Atmospheric Environment* 2014;89(0):749–56. doi:<http://dx.doi.org/10.1016/j.atmosenv.2014.03.017>.
- Querol X, Alastuey A, Pandolfi M, Reche C, Pérez N, Minguillón M, Moreno T, Viana M, Escudero  
535 M, Orío A, Pallarés M, Reina F. 20012012 trends on air quality in Spain. *Science of The Total Environment* 2014;490(0):957–69. doi:<http://dx.doi.org/10.1016/j.scitotenv.2014.05.074>.
- Ray JD. Spatial distribution of tropospheric ozone in national parks of California: Interpretation of passive-sampler data. *TheScientificWorldJOURNAL* 2001;1(0):483–97. doi:doi:10.1100/tsw.2001.83.
- 540 Rodríguez S, Querol X, Alastuey A, Mantilla E. Origin of high summer {PM<sub>10</sub>} and {TSP} concentrations at rural sites in eastern Spain. *Atmospheric Environment* 2002;36(19):3101–12. doi:[http://dx.doi.org/10.1016/S1352-2310\(02\)00256-X](http://dx.doi.org/10.1016/S1352-2310(02)00256-X).
- Sanz M, Sanz F, Calatayud V, na GSP. Ozone in Spain's national parks and protected forests. *TheScientificWorldJOURNAL* 2007;7(0):67–77. doi:doi:10.1100/tsw.2007.8.
- 545 Screpanti A, Marco AD. Corrosion on cultural heritage buildings in Italy: A role for ozone? *Environmental Pollution* 2009;157(5):1513–20. doi:<http://dx.doi.org/10.1016/j.envpol.2008.09.046>; special Issue Section: Ozone and Mediterranean Ecology: Plants, People, Problems.

- Seidel D, Zhang Y, Beljaars A, Golaz JC, Jacobson A, Medeiros B. Climatology of the planetary boundary layer over the continental united states and europe. *Journal of Geophysical Research: Atmospheres* 2012;117(D17). doi:10.1029/2012JD018143; d17106.
- Sicard P, De Marco A, Troussier F, Renou C, Vas N, Paoletti E. Decrease in surface ozone concentrations at Mediterranean remote sites and increase in the cities. *Atmospheric Environment* 2013;79:705–15.
- Tapia O, Escudero M, Lozano , Anzano J, Mantilla E. New classification scheme for ozone monitoring stations based on frequency distribution of hourly data. *Science of The Total Environment* 2016;544:1–9. doi:http://dx.doi.org/10.1016/j.scitotenv.2015.11.081.
- Temple PJ. Oxidant air pollution effects on plants of joshua tree national monument. *Environmental Pollution* 1989;57(1):35 – 47. doi:http://dx.doi.org/10.1016/0269-7491(89)90128-0.
- Troen I, Mahrt L. A simple model of the atmospheric boundary layer; sensitivity to surface evaporation. *Boundary Layer Meteorology* 1986;37:129–48.
- UNECE . Final draft report of ECE Critical Levels Workshop. Technical Report; United Nations Economic Commission for Europe; 1988.
- Villanueva F, Tapia A, Notario A, Albaladejo J, Martínez E. Ambient levels and temporal trends of  $voc_s$ , including carbonyl compounds, and ozone at Cabañeros national park border, Spain. *Atmospheric Environment* 2014;85(0):256 –65. doi:http://dx.doi.org/10.1016/j.atmosenv.2013.12.015.
- WHO . Air Quality Guidelines for particulate matter, ozone, nitrogen dioxide and sulfur dioxide: Global Update 2005. Copenhagen, Denmark: WHO Regional Office for Europe, 2006.
- WHO . Health risks of ozone from long-range transboundary air pollution. Copenhagen, Denmark: WHO Regional Office for Europe, 2008.
- Zhang Y, Gao Z, Li D, Li Y, Zhang N, Zhao X, Chen J. On the computation of planetary boundary-layer height using the bulk richardson number method. *Geoscientific Model Development* 2014;7:2599–611. doi:10.5194/gmd-7-2599-2014.

First comprehensive assessment of the levels of atmospheric pollutants in National Parks in Spain

Regional sources and the influence of transport scenarios affect pollutants levels in National Parks

Emissions patterns and changes in the PBL height explain seasonal variability of pollutants

Ozone exhibit high levels in National Parks increasing concentrations towards the east of Iberia

Analysis of daily variability reveal impact of anthropogenic sources in some parks