

ISSN 2089 - 2136

Tropical Medicine Journal

Volume 03, No. 1, 2013

- Risk Factor of HIV Infection Among Young Age in Voluntary Counseling Testing (VCT) Clinics of Yogyakarta
- Evaluation of the Performance of Malaria Microscopist in Primary Health Center and Cross Checker in Belu East Nusa Tenggara
- The Kinetics of White Blood Cells in Acute Dengue Infection
- The Effect of *Pandanus conoideus* Lamik Extract to the Serum Level of TNF- α , IL-10 and Parasitemia of *Plasmodium berghei* Infected in Mice
- Comparison of Immunochromatography Method and Immunocytochemistry Method in Rapid Detection of NS-1 Antigen in Dengue Infection
- Filariasis Bancrofti Epidemiology Post Mass Drug Administration in Waris District Keerom Regency Province of Papua
- The Relationship of Behavior and Environment to the Incidence of Malaria in the Work Area of Desao Public Health Center (PHC) of East Kupang Sub-District of Kupang District in 2013
- The Red Fruit (*Pandanus Conoideus* Lam) Ethanol Extract Decrease Tumor Necrosis Factor-Alpha (TNF-Alpha) Level and Intercellular Adhesion Molecule-1 (ICAM-1) Expression of *Plasmodium berghei* Infected Swiss Mice Malaria Model
- Training of Sputum Microscopy Improves the Smear Quality and Slide Positivity Rate for Pulmonary Tuberculosis Diagnosis
- Integrated and Comprehensive Action to Reduce and Control Dengue Hemorrhagic Fever: A Survey in Pekalongan City, Central Java

TMJ	Volume 03	Number 01	Page 1 - 93	ISSN 2089 - 2136
-----	--------------	--------------	----------------	---------------------

Center for Tropical Medicine, Faculty of Medicine, Universitas Gadjah Mada
in collaboration with Indonesian Society of Tropical Medicine and Infectious Disease (PETRI)

Editor-in-chief

Prof. dr. Supargiyono, DTM&H., SU., Ph.D, S.Park

Managing Editor

Dr. dr. Mahardika Agus Wijayanti, M.Kes

Associate Editors

Prof. Dr. Mustofa, M.Kes., Apt.

dr. Yodhi Mahendradhata, M.Sc, Ph.D

Dr. Dra. Erna Kristin, MSi, Apt.

dr. Rinis Andono Ahmad, MPH., Ph.D

dr. Doni Priambodo, SpPD-KPTI

Editorial Advisory Board

Dr. Tedjo Sasmono, Bsc

dr. Din Syafruddin, Ph.D

Prof. dr. Ni Made Mertaniasih, MS, Sp.MK

Prof. Dr. dr. Arie Mansyur, SpPD-KPTI

dr. Subagyo Loeheeri, SpPD

Dr. dr. Budiman Bela, Sp.MK (K)

All right reserved. No part of this publication may be reproduced, stored or transmitted in any form or by any mean, electronic or mechanical, without written permission from the publisher.

Address : Tropical Medicine Journal, PAU Building, Jalan Teknika Utara Berek, Yogyakarta

Universitas Gadjah Mada, Yogyakarta 55281, Phone : +62-274-588483, E-mail: tropmedjournal@gmail.com

TROPICAL MEDICINE JOURNAL

ISSN : 2089-2136

Center for Tropical Medicine, Faculty of Medicine, Universitas Gadjah Mada in collaboration with
Indonesian Society of Tropical Medicine and Infectious Disease (PETRI)

Volume 03, Number 01

CONTENTS

- 1-15 Risk Factor of HIV Infection Among Young Agein *Voluntary Counseling Testing* (VCT) Clinics of Yogyakarta
Ismael Saleh, Sumardi, Lutfan Lazuardi
- 16-28 Evaluation of the Performance of Malaria Microscopist in Primary Health Center and Cross Checker in Belu East Nusa Tenggara
Fridolina Mau, Supargiyono, Elsa Herdiana Murhandarwati
- 29-38 The Kinetics of White Blood Cells in Acute Dengue Infection
Mohd Nasrul Bin Mohd Ghazali, Umi Solekhah Intansari, Ida Safitri Laksanawati
- 39-47 The Effect of *Pandanus conoideus* Lamk Extract to the Serum Level of TNF- α , IL-10 and Parasitemia of *Plasmodium berghei* Infected in Mice
Zeth Robeth Felle, Mahardika Agus Wijayanti, Supargiyono.
- 48-56 Comparison of Immunochromatography Method and Immunocytochemistry Method in Rapid Detection of NS-1 Antigen in Dengue Infection
How Tien Jack, Sitti Rahmah Umniyati, Elsa Herdiana Murhandarwati
- 57-63 Filariasis Bancrofti Epidemiology Post Mass Drug Administration in Waris District Keerom Regency Province of Papua
Korinus Suweni, Soeyoko, Sri Sumarni
- 64-70 The Relationship of Behavior and Environment to the Incidence of Malaria in the Work Area of Oesao Public Health Center (PHC) of East Kupang Sub-District of Kupang District in 2013
Titik Yuliaty, Yayi S. Prabandari, Tri Baskoro T. Satoto
- 71-80 TThe Red Fruit (*Pandanus Conoideus* Lam) Ethanol Extract Decrease Tumor Necrosis Factor-Alpha (TNF-Alpha) Level and Intercellular Adhesion Molecule-1 (ICAM-1) Expression of *Plasmodium berghei* Infected Swiss Mice Malaria Model
Demianus Tafor, Mujur, Achmad Djunaidi, Widya Wasityastuti, Eti Nurwening Sholikhah
- 81-84 Training of Sputum Microscopy Improves the Smear Quality and Slide Positivity Rate for Pulmonary Tuberculosis Diagnosis
Dede Kurniawan, Ning Rintiswati. Dibyong Pramono
- 85-93 Integrated and Comprehensive Action to Reduce and Control Dengue Hemorrhagic Fever: A Survey in Pekalongan City, Central Java
Nur Siyam

Risk Factor of HIV Infection Among Young Agein Voluntary Counseling Testing (VCT) Clinics of Yogyakarta

Ismael Saleh^{1*}, Sumardi², Lutfan Lazuardi³

¹Postgraduate Programme, Majoring in Tropical Health, Faculty of Medicine, Universitas Gadjah Mada, Yogyakarta, Indonesia; ²Division of Internist Education Program, Faculty of Medicine, Universitas Gadjah Mada, Yogyakarta, Indonesia; ³Public Health Department, Faculty of Medicine, Universitas Gadjah Mada, Yogyakarta, Indonesia.

Corresponding author: ismael_irmawan@yahoo.com

ABSTRACT

Introduction: HIV/AIDS is a health problem in the world. The Highest HIV/ AIDS cases in Indonesia were reported in young age groups, including in Province of Yogyakarta (DIY). Knowing the determinant and risk factors of HIV is necessary to determine effective prevention programs, especially in young age which one of the vulnerable populations, either because of individual and family characteristics, media influence, peer pressure, substance use and risky behavior.

Objectives: To know the relationship between risk factors for HIV infection among young age in VCT clinic of Yogyakarta.

Methods: This study used a cross-sectional design and the population was young clients at VCT clinics of Yogyakarta. Samples were obtained from VCT clinics of Griya Lentera PKBI Yogyakarta and Edelweis Clinics of Dr Sardjito Hospital. Variable risk factors were obtained by questionnaire and interviews than HIV test results was obtained from VCT clinics.

Results: Of the 62 respondents, 74.2% were female, age average were 24,4 years old, 41.9% level education were junior high school, 82.3% had migration, 62.9% received less monitoring from families, 61.3% ever use the TV / HP for pornography, 56.5% getting high sexual pressure from peers, 62.9% had been smoking and alcohol consumption, 98.4% had ever sexual intercourse, 95.2% had sexual intercourse in the last 6 months, 93.5% risky sexual intercourse, 32.3% HIV positive and 55% through heterosexual.

Conclusion: HIV infection among young people in VCT Clinic of Yogyakarta, by bivariate analysis correlated with gender, residency status, education level, relationship with parents, TV / mobile and internet used to watch pornography. However, according to multivariate analysis, HIV infection correlated with education level at senior high school and college, residing with their parents/ family and in the last 6 months are still using alcohol and illicit drugs. It is recommended to enhance more comprehensive prevention for young people in accordance with the factors associated with HIV risk behaviour.

Keywords: HIV infection, young age, individual characteristics, family, peers, media usage, substance use, risky behavior.

INTISARI

Pendahuluan: HIV/AIDS merupakan masalah kesehatan dunia. Kasus HIV/ AIDS tertinggi di Indonesia dilaporkan pada kelompok umur muda, termasuk di Propinsi Daerah Istimewa Yogyakarta (DIY). Mengetahui faktor penentu dan faktor risiko pada pasien HIV. Pemahaman tentang faktor penentu dan faktor risiko dari HIV diperlukan untuk menentukan program pencegahan yang efektif, terutama pada kelompok usia muda sebagai satu dari kelompok yang rentan, baik karena karakteristik individual dan keluarga, pengaruh media, tekanan kelompok pergaulan, dan perilaku bermasalah.

Tujuan: Untuk mengetahui hubungan antara faktor risiko infeksi HIV di antara kelompok usia muda pada klinik VCT di Yogyakarta.

Metode: Penelitian ini menggunakan rancangan potong lintang dengan populasi berupa klien muda di klinik VCT dari Griya LenteraPKBI Yogyakarta dan klinik Edelweis dari Rumah Sakit Dr Sardjito. Faktor risiko diperoleh melalui kuesioner dan wawancara sedangkan hasil tes HIV diperoleh dari klinik VCT.

Hasil: Dari 62 responden, 74.2% wanita, rerata umur 24,4 tahun, 41.9% memiliki tingkat pendidikan Sekolah Menengah Pertama (SMP), 82.3% mengalami pindahan, 62.9% kurang terpantau oleh keluarganya, 61.3% pernah menggunakan TV/HP untuk pornografi, 56.5% mendapat tekanan seksual yang tinggi dari kelompok sebaya, 62.9% mengkonsumsi alkohol dan merokok, 98.4% pernah berhubungan seksual, 95.2% pernah berhubungan seksual dalam 6 bulan terakhir, 93.5% berhubungan seksual yang berisiko, 32.3% positif mengidap HIV dan 55% melalui heteroseksual.

Simpulan: Infeksi HIV di antara orang muda di Klinik VCT Yogyakarta, berdasarkan analisis bivariat, berkorelasi dengan jenis kelamin, status tempat tinggal, level pendidikan, hubungan dengan orang tua, pemakaian TV/telepon seluler dan internet untuk melihat pornografi. Namun, menurut analisis multivariat, infeksi HIV berkorelasi dengan level pendidikan SMA dan perguruan tinggi, tinggal dengan orang tua/keluarga meredakan dalam 6 bulan terakhir masih mengkonsumsi alkohol dan obat terlarang. Dianjurkan untuk meningkatkan pencegahan yang lebih komprehensif untuk orang muda mengikuti faktor-faktor yang berhubungan dengan perilaku yang berisiko terinfeksi HIV

Kata kunci: infeksi HIV, usia muda, karakteristik individual, keluarga, kelompok sebaya, pemakaian media, pemakaian narkoba, perilaku berisiko

INTRODUCTION

Human immunodeficiency virus (HIV) and acquired immunodeficiency syndrome (AIDS) is currently the largest health problem in the world. Based on data published by the Joint United Nations Programme on HIV / AIDS (UNAIDS) at the end of 2010, an estimated 34 million people were living with HIV worldwide. The number of people dying of AIDS-related causes fell to 1.8 million and there were 2.7 million new infection in 2010¹.

In Indonesia, since first reported in 1987 until June 2011, were reported 66.693 cases of HIV, 26.483 cases of AIDS and 5.056 people died among it². Most cases of AIDS occurred in the age group 20-29 years (46.4%), recent data showed highest rate in the same age group at 36.4%. In Province of Yogyakarta (DIY), HIV cases were infected 1.217 people until June 2011 and AIDS prevalence were reported 18,87 per 100.000 population (higher than the national rate 11,09 per 100,000)². Furthermore, the data of Voluntary Counseling Testing (VCT) from the health department officer of DIY in 2011 showed

that 9.7% of new incidence were among 15-24 years old.

AIDS, caused by viruses, but it also a behavior disease, most infected get it through sexual intercourse (70-80%) and drug abuse (10%)³. Among young people, HIV spreads almost exclusively in two ways: unsafe sex (between males and females, and between males) and sharing injecting drug equipment⁴.

It is probable that most of the individuals in their twenties who have AIDS, is contracted HIV during their teenage years⁵. Environmental factors such as social, cultural, economic and political forces such as poverty, migration, urbanization, war, civil disturbance caused the vulnerable of adolescents infected by HIV^{6,7} rather than information and communication technology that enables them to access the things that drives the desire for deviant behavior⁸. Yogyakarta has a large number of young people, high level of urbanization⁹, easy internet access and an increase in unemployment rate¹⁰ that might cause the young driven to the risky behavior.

Knowing the determinant and risk factors for HIV is necessary to determine effective prevention programs, especially in young people which one of the most vulnerable groups with regard to HIV/AIDS, either because of characteristics of individual and family, media influence, peer pressure, substance use and risky behavior.

The aim of this study is to know the relationship between risk factors for HIV infection among young age in VCT clinic in Yogyakarta.

MATERIALS AND METHODS

Participants and data collection procedures

This research is a cross-sectional survey, conducted from September to October 2012. Population are young VCT clients (18-29 years old) in Yogyakarta, while the sample subject is VCT clients in two locations: Griya Lentera PKBI Yogyakarta and Edelweis Clinics of Dr. Sardjito Hospital. Data collection from Griya Lentera PKBI assisted by counselors who had been trained in filling out the questionnaire, while at the Edelweis clinic of Dr. Sardjito hospital directly conducted by researchers when clients visited the clinic. All the participants were willing to follow the informed consent form.

Parameter Measurements

This study participants filled out questionnaires anonymously and guided its contents by the researcher or team. Variables were asked in the questionnaire include individual and family characteristics, peer pressure, media usage, substance usage, risky behaviors and HIV status. Individual characteristics consisted of age, gender, level of education, status of residence and migration status. Residency status consisted of the kind of house where they lived, categorized as a boarding house, rent, parents and family house. Migration status were asked the history of their movement from one place to another.

Family characteristics include level of relationships with parents and families monitoring to respondent behavior. Relationships with parents consisted of level relationships and a sense of excitement in their interactions with parents. Family monitoring consisted of the informing family about where and with whom they were when far from home.

Peer pressure consisted of the number of their peers who did and invite to risky behavior, both sexual and drug used. Influence of the media consisted of frequency of television or cell phones and internet used to watch movies/porn scenes. Substance usage consisted of the history and continuity of substance usage including cigarettes, alcohol and illicit drugs.

Risky behavior consisted of the frequency and type of risky behavior that vulnerable to HIV infection. The questions include risky sexual behavior, injecting drug use and other behaviors such as tattoos and blood donor recipient. Risky sexual behavior includes involvement in a sexual commercial (sex worker or customers), ever having anal sex (sex among males), ever diagnosed STDs (sexual transmitted diseases) both respondents and partner, do not always use condom when having sex, number of sexual partners and frequency of sexual intercourse in last 6 months¹¹⁻¹⁴.

Data Analysis

Data analysis was performed with 3 steps. The first step, we conducted descriptive analysis of each variable. The second step of analysis, we conducted bivariate analysis using appropriate correlation test to determine the relationship between each independent with dependent variable. In the third step of analysis, we conducted multivariate analysis using logistic regression to see the effect of independent variables on the dependent variable simultaneously.

RESULT AND DISCUSSION

Univariate analysis

Table 1 Individual and Family Characteristic among Young Client at VCT clinic of Yogyakarta in 2012

Variable	n (%)	Variable	n (%)
Age (mean)	24,4	Education	
Sex		Primary	13 (21)
Male	16 (25,8)	Junior High School	26 (41,9)
Female	46 (74,2)	Senior High School	16 (25,8)
Residency Status		Diploma	3 (4,8)
Board/ Dormitory	38 (61,3)	Graduate/ Post	4 (6,5)
Rent house	9 (14,5)	Relationship with Parent	
Family House	3 (4,8)	Less	16 (25,8)
Parent house	12 (19,4)	Good	46 (74,2)
Migration		Family Monitoring	
Ever	51 (82,3)	Less	39 (62,9)
Never	11 (17,7)	Good	23 (37,1)

Of the 62 respondent, average age was 24.4 years, 74.2% were female, most junior high school (41.9%), residing in board/ dormitory(61.3 %) and ever migration (82.3%). Of the 62 respondents, 46

(74.2%) of respondents have good relationship with her parents, but claimed to have received less monitoring from his family (62.9%).

Table 2. Peer Pressure, Usage of Media and Substance Use among Young Client at VCT clinic of Yogyakarta in 2012

Variable	n (%)	Variable	n (%)
Peer Pressure (in sexual intercourse)		Smoking Continued (in last 30 days)	
High	35 (56,5)	Has Used	31 (79,5)
Low	27 (43,5)	Sometime	5 (12,8)
Peer Pressure (in illicitdrugs used)		Hasn't use	3 (7,7)
High	8 (12,9)	Alcohol consumption	
Low	54 (87,1)	Ever	39 (62,9)
TV/ HP for Pornography		never	24 (37,1)
Often	5 (8,1)	Continuity of alcohol (in last 6 month)	
Rarely	33 (53,2)	Has Used	20 (51,3)
Never	24 (38,7)	Sometime	8 (20,5)
Internet for Pornography		Hasn't use	11 (28,2)
Often	2 (3,2)	Illicit drug used	
Rarely	16 (25,8)	Ever	9 (14,5)
Never	44 (71)	Never	53 (85,5)
Cigarette		Continuity of IDUs (in last 6 month)	
Ever	39 (62,9)	Has Used	4 (44,4)
Never	23 (37,1)	Hasn't use	5 (55,6)

About peer pressure, 56.5% respondents had high pressure related-sexual and 12.9% for drug use. Most respondents were rarely (53.2%) using the television/ cell phone media and only 29% had used internet to watch porn scenes. Of 62 respondent, 39 respondents (62.9%) had smoked and 79.5% of them had continued in the past 30 days. The majority (62.9%) of respondents had used alcohol and 71.8% of them still using it in last 6 months. Only 9 respondents (14.5%) ever used illicit drugs and 4

respondents (44.4%) were still using it in last 6 months.

Of the 62 respondents, 58 respondents having risk sexual intercourse (93.5%). Injecting drug use were only done by 4 respondents (6.5%), 1 of them do by syringe sharing. Other risk factors, only 12 respondents (19.4%) had having tattoos and 3 respondents (4.8%) had ever been blood donor recipients.

Table 3. Risk behavior and HIV status among Young Client at VCT clinic of Yogyakarta in 2012

Variable	n (%)	Variable	n (%)
Sexual Intercourse		HIV status	
Risky	58 (93,5)	Positive	20 (32,3)
Not risky	4 (6,5)	Negative	42 (67,7)
IDUs		Transmission type	
Used	4 (6,5)	Bisexual	3 (15)
Not used	58 (93,5)	Homosexual (MSM)	5 (25)
Other risk		Heterosexual	11 (55)
Tattoo	12 (19,4)	IDUs	1 (5)
Be blood recipient	3 (4,8)		

Based on results of HIV antibody test, 20 respondents (32.3%) were HIV-positive and 67.7% negative. Respondents were mostly positive (55%) infected through heterosexual and 25% homosexual (male).

About sexual behavior, 32.8% of respondents had first sexual intercourse at under 16 years, 86.9% had sexual intercourse with the opposite sex, 47.6% do not know the history of sexual transmitted

diseases (STD) infection of the sexual partner, 49.2% claimed never infected with STDs and 42.6% did not know the presence of other sexual partner of partner. Sexual intercourse in the last 6 months had been done by 59 (95.2%) respondents, 42 (71.2%) having more than two sexual partners, 76.3% had sexual intercourse more than 12 times and 55.9% sometimes using condom when having sex.

Table 4. Sexual behavior among Young Client at VCT clinic of Yogyakarta in 2012

Variable	n (%)	Variable	n (%)
First sexual intercourse		Sex. intercourse	
Before or 16 years	20 (32,8)	(in last 6 month)	
After 16 years	41 (67,2)	Has Had	59 (95,2)
Type of Sex. Intercourse		Hasn't Had	3 (4,8)
Homosex	5 (8,2)	Number of Sexual partner	
Bisex	3 (4,9)	1	14 (23,7)
Heterosex	53 (86,9)	2	3 (5,1)
STDs History of Partner		More than 2	42 (71,2)
Ever	7 (11,5)	Frequency of sexual intercourse	
Don't know	27 (44,3)	Less than 6	6 (10,2)
Never	27 (44,3)	6 – 12	8 (13,6)
STDs History of respondent		More than 12	45 (76,3)
Ever	12 (19,7)	Condom used	
Don't know	19 (31,1)	Always	18 (30,5)
Never	30 (49,2)	Sometime	33 (55,9)
Other sexual partner		Never	8 (13,6)
Has Had	17 (27,9)		
Don't know	25 (41,0)		
Hasn't had	19 (31,1)		

Bivariate analysis

Age in this study showed no significantly correlation ($p = 0.487$) with HIV status. This is due to the subjects in this study were VCT clients who are mostly at-risk groups that have risk behaviors did not differ at various age levels, as seen in the data of studies showing no difference by age of respondents risky behavior. Other study with different result show that older age had significantly association with

HIV infection^{11,15}. It can happen because older young agemore actively in sexually activity¹⁶.

Gender in this study showed significantly correlation ($p = 0.000$) with HIV infection. Young women tend to not infected by HIV. This happens because young men was more likely to have risky sex than women¹⁶.The this study similar with case of HIV/AIDS in Indonesia, which also showed the majority of AIDS patients were male (72.3%).

Table 5. Bivariat analysis of individual and family characteristic related with HIV infection among Young Client at VCT clinic of Yogyakarta in 2012

Variable	HIV +	HIV –	p	r
Age	-	-	0,487	0,090
Sex			0,000	0,460
Male	11 (68,8)	5 (31,2)		
Female	9 (19,6)	37 (80,4)		
Residence status			0,000	-0,496
Board/Dormitory&Rent house	9 (15,1)	38 (80,9)		
Family &Parent house	11 (73,3)	4 (26,7)		
Education			0,000	-0,470
Primary - Junior High School	6 (15,4)	33 (84,6)		
Senior High School – College	14 (60,9)	9 (39,1)		
Migration			0,311	-0,131
Ever	15 (29,4)	36 (70,6)		
Never	5 (45,5)	6 (54,5)		
Relationship with Parent			0,017	0,303
Less	9 (56,2)	7 (43,8)		
Good	11 (23,9)	35(76,1)		
Family Monitoring			0,174	0,173
Less	15 (38,5)	24 (61,5)		
Good	5 (21,7)	18 (78,3)		

Residency status of respondents showed significantly correlation ($p = 0.000$) with HIV infection. Respondent who reside in board or dormitory or rent house tend to not infected by HIV. Young age who living in board or dormitory or not living with parent or family, tend to have good relationship with their parents. Its way for them to get messages and advice to avoid risky behaviors. Different with other similar research that showed HIV infection among young age was related to not living with their parents¹³. In addition, survey result showed that parent communication and monitoring may play an important role in reaching youth early with preventon messages¹⁷. Related to these findings, the parents and families need to always maintain communication and supervision of young age, especially related attitudes and behaviors that may increase the risk of contracting HIV.

Education level of respondents showed significantly correlation ($p = 0.006$) with HIV infection. Primary – Junior high school leveleducation tend to not infected by HIV. Other similar research showed different result, as low education level was one of the predictors of HIV infection in women¹³ and young men with low levels of education-related HIV risk behaviors¹⁸. In this research, higher education (SMA-PT) had tend to get higher peer pressure, more frequently in media us to access pornography, had a history of smoking and alcohol use and riskier sexual behavior than lower education level. Its increased respondents to behave at risk of contracting HIV.

Migration status in this study showed no significant correlation ($p = 0.311$) with HIV infection. It happened because most of respondents (82.3%) have ever migration so that it was not comparable to other variable. Nevertheless, the data showed that

respondents who had migrated tend to have risky sexual behavior, although not associated with HIV infection. Other study had similar result showed that young people who leave school later migrated with his parents to Beijing, had high vulnerability to HIV infection¹⁹.

Data in this study indicate that there were at least 50% of the respondents infected by HIV outside of Yogyakarta. This suggests that Yogyakarta is not potentially location of HIV transmission, however, Yogyakarta were the favourite treatment location because only 7 respondents (35%) coming from Yogyakarta, the other come from outside Yogyakarta.

Relationships with parents in this study correlated with HIV infection ($p = 0.017$). Good

relationship with parent can be an important means to avoid young people by the contagion environment. Good communication with parents causes young would not tend to engage in sexual relationships²⁰. Therefore, parents need to keep good relationship with the child and to follow the developments in the dynamics of his life.

Family monitoring in this study showed no significant correlation ($p = 0.174$) with HIV infection. Although, this study found that lacking of monitoring family of young people tend to risky sexual behavior. Other study with similar result showed that youth with high risk behavior tend to be done by those who lack monitoring from caregivers / parents^{16,21}.

Table 6. Bivariate analysis of Peer pressure and media usage related with HIV infection among Young Client at VCT clinic of Yogyakarta in 2012

Variable	HIV +	HIV -	<i>p</i>	<i>r</i>
Peer Pressure (Sexual Intercourse)			0,697	0,049
High	12 (34,3)	23 (65,7)		
Low	8 (29,6)	19 (70,4)		
Peer Pressure (illicit drugs used)			0,418	0,146
High	4 (50)	4 (50)		
Low	16 (29,6)	38 (70,4)		
TV/ HP for Pornography			0,037	0,265
Often&Rarely	16 (42,1)	22 (57,9)		
Never	4 (16,7)	20 (83,3)		
Internet for Pornography			0,002	0,395
Often&Rarely	11 (61,1)	7 (38,9)		
Never	9 (20,5)	35 (79,5)		

Peer pressure (related-sexual intercourse) in this study showed no significant correlation ($p = 0.697$) with HIV status. It happened because peer pressure does not directly influence to HIV infection, but effect on risk behavior. The data of this study indicate higher peer pressure (sexual intercourse) in young

men associated ($p = 0.013$) with risky sexual behavior. It be done because influence of peer pressure are more likely at risk of contracting HIV, because they would take drugs and have multiple sex partners and unsafe sex²². A similar result study found that higher proportion of close friends (peers)

who started the sexual intercourse, will improve their chances to emulate, including the number of sexual partners²³.

Peer pressure (related-illicit drug use) showed no correlation ($p = 0.250$) with HIV status. It happened because peer pressure does not directly influence on HIV infection, but effect on risk behavior. Other similar result study showed that peers were strongly associated with substance used, but not directly related to the use of condoms in sexual intercourse²⁴. Peer pressure (related-illicit drugs) was associated with injecting drug use and other risky behaviors (Tattoos donor and recipient). Peers not only can establish the behavior of injecting drug use, but also be powerful predictor of use of non-sterile needles²⁵. This happens because peer group can influence and

reinforce norms and values †, shaping social and cultural identity and provide a model of behavior, but in the perspective of health, peer influence can be protective factors and risk factors²⁶.

Media use, both TV/HP ($p = 0.037$) and internet ($p = 0.002$) showed significant correlation with HIV status. Frequency HP and internet related with sexual sensation seeking⁸ which would increase desire have sexual intercourse. It would increased the risk to infected by HIV because young people who frequently access to internet and social networks were more widely used to seeking for sexual partners²⁷ and it has a significantly higher risk of STDs and HIV infection²⁸ because it can increase their chances to get the infected individuals.

Table 7. Bivariat test of substance used related with HIV infection among Young Client at VCT clinic of Yogyakarta in 2012

Variable	HIV +	HIV –	p	r
Cigarette			0,814	0,030
Ever	13 (33,3)	26 (66,7)		
Never	7 (30,4)	16 (69,6)		
Smoking Continued (last 30 days)			0,375	- 0,113
Has Continued	10 (27,8)	26 (72,2)		
Hasn't Continued	10 (38,5)	16 (61,5)		
Alcohol consumption			0,744	- 0,041
Ever	12 (30,8)	27 (69,2)		
never	8 (34,8)	15 (65,2)		
Continuity of alcohol (in last 6 month)			0,098	-0,210
Has Continued	6 (21,4)	22 (78,6)		
Hasn't Continued	14 (41,2)	20 (58,8)		
Illicit drug used			0,133	0,205
Ever	5 (55,6)	4 (44,4)		
Never	15 (28,3)	38 (71,7)		
Continuity of drug used (in last 6 month)			0,095	0,240
Has Continued	3 (75)	1 (25)		
Hasn't Continued	17 (29,3)	41 (70,7)		

Tobacco use (history, $p = 0.814$ and continuity, $p = 0.375$) in this study showed no correlation with HIV status. Smoking does not directly influence on HIV infection, but effect on risky behaviors. This study showed association between continuity smoked in the past 30 days with history of drug use. Other similar result study showed that smoking was not associated with HIV infection, but related to drug abuse²⁹. Related to this, the efforts to stop smoking and drug abuse needs to be done at young people.

Alcohol consumption (history, $p = 0.744$) and continuity, $p = 0.098$) in this study showed no correlation with HIV status. Alcohol consumption does not directly influence on HIV infection, but effect on risky behaviors. This study also showed that history of alcohol consumption ($p = 0.002$) in young

men associated with not using condom during sexual intercourse in last 6 months. It happened because alcohol can influence on the increase of sexual desire and it may contibuted to HIV risk behavior³⁰.

Illicit drugs used (history, $p = 0.133$ and continuity $p = 0.095$) in this study showed no correalation with HIV status. It happened because illicit drugs used does not directly influence on HIV infection, but on risky behaviors. These results contrast with the results of a national survey of risk behaviors in the United States found that young people who use hazardous materials on a whim or for years, they are more likely to engage in high-risk behaviors such as unprotected sex. Other study showed that one of HIV risk factor in young men, were the use of drugs¹⁸.

Table 8. Bivariat test of risk behavior related with HIV infection among Young Client at VCT clinic of Yogyakarta in 2012

Variable	HIV +	HIV -	<i>p</i>	R
Sexual Intercourse			1,000	0,041
Risky	19 (32,8)	39 (67,2)		
Not risky	1 (25)	3 (75)		
IDUs			1,000	-0,041
Used	1 (25)	3 (75)		
Not used	19 (32,8)	39 (67,2)		
Other risk			0,349	0,122
Present	6 (42,9)	8 (57,1)		
Not present	14 (15,5)	34 (70,8)		

Risky sexual intercourse ($p = 0.748$) in this study showed no correlation with HIV infection. Its caused weakness of this study that categorizes risk sex based on many criteria, so that almost all respondents (93.5%) categorized risk in sexual intercourse, while the main risk factors of transmission, the use of condoms is not a major question items, but included

in question of sexual intercourse in the last 6 months.

Injection drug user (IDU) in this study showed no correlation ($p = 0.748$) with HIV infection. Nevertheless, the one IDU users who share syringes in this study were infected by HIV. This can happen because the HIV virus can be transmitted through shared needles used to inject drugs³¹.

Table 9. Bivariat analysis of sexual behavior related with HIV infection among Young Client at VCT clinic of Yogyakarta in 2012

Variable	HIV +	HIV –	p	R
First sexual intercourse			0,746	- 0,041
Before or at 16 years	6 (30)	14 (70)		
After 16 years	14 (34,1)	27 (65,9)		
Type of Sex. Intercourse			0,000	0,556
Homosex – bisex	8 (100)	0 (0)		
Heterosex	12 (22,6)	41 (77,4)		
STDs History of Partner			0,674	0,077
Ever	3 (42,9)	4 (57,1)		
Don't know – never	17 (31,5)	37 (68,5)		
STDs History of resp.			0,001	0,445
Ever	9 (75)	3 (25)		
Don't know – Never	11 (22,4)	38 (77,6)		
Other sex. partner of part			0,040	0,325
Has Had	8 (50)	8 (50)		
Don't know	10 (38,5)	16 (61,5)		
Hasn't had	2 (10,5)	17 (89,5)		

Age of first sexual intercourse in this study showed no correlation ($p = 0.746$) with HIV infection. This is because the age of first sexual intercourse is not directly lead to HIV infection, but if the risky sexual behavior with partner who has been infected. Related to this, the data showed that respondents who have sex the first time before 16 years old, were engage torisky sexual intercourse. Other study showed the similar result that sexual intercourse in younger people carried at risk of contracting STDs, including HIV³². Therefore, for young people who have not started sexual activity, delayed sexual intercourse were an important strategy to be applied³². Delayed sexual intercourse among young people can be done by high of parental monitoring³², good communication between parent and child²⁰ and parents state the disagreement to sexual intercourse in young age³³.

Type of sexual intercourse in this study showed significantly correlation ($p = 0.000$) with HIV status with tend to same-sex sexual intercourse. It happened because sexual behavior of homosexual

groups (of men) may cause anal injury and it can more easilyinfected by HIV. National survey of young risk behaviors in the United States was also getting the same thing, that young people who have sexual relations among men, especially for certain racial groups have high number of new HIV infections¹⁷.

STDs history of sexual partner showed no correlation ($p = 0.674$) with HIV infection. This can occur due to the weakness of the study that justused questionnaire. Information bias may be occurred because ignorance by respondent about STDs of his sexual partnerthat sometimes do not show physical symptoms. These results was contrast to Dorjgochoo et al. (2009) which states that HIV infection at young age were more likely to respondents who had sexual partner with STDs infection history.

Respondents with history of STDs showed significantly correlation ($p = 0.001$) with HIV infection. Respondents who do not know and have never been infected STDs tend to not infected by HIV. It can be explained that occurred STD infection, indicatedthat sexual intercourse be done, were

unsafe or unprotected, this increases the risk contracting by HIV. These risks can occur due to unprotected sexual behavior and genital injuries due to STDs facilitate HIV infection¹³.

The presence of other sexual partner of respondent partner in this study showed significantly correlation ($p=0.040$) with HIV status. This finding is

similar with Dorjgochoo et al. (2009) which states that HIV infection in young women one of which relates to the suspicion that a sex partner has another partner. It can be explained that the presence of sexual partner other than the respondent partner increases the chances of contracting HIV and then can transmit to the respondent.

Table 10. Bivariate analysis of sexual behavior in last 6 month related with HIV infection among Young Client at VCT clinic of Yogyakarta in 2012

Variabel	HIV +	HIV -	p	r
Number of Sexual partner (in last 6month)			0,000	-0,669
2 or More	6 (13,3)	39 (86,7)		
1	12 (85,7)	2 (14,3)		
Sexual Frequency (in last 6month)	12 (22,6)	41 (77,4)	0,001	-0,508
6 - 12 and More 12	6 (100)	0 (0)		
Less than 6			0,360	0,119
Condom used (in last 6month)	14 (34,1)	27 (65,9)		
Sometime & Never	4 (22,2)	14 (12,5)		
Always				

Number of sexual partner showed significantly correlation ($p = 0.000$) with HIV infection. Cross tabulation showed that respondent who had 2 or more sexual partner tend to not infected by HIV. This may occur because sexual intercourse that be done were safe, protected or sexual partner not infected by HIV. This finding contrasts with results of study¹¹ which states that HIV infection at young age relates to the number of their sexual partners. Although other study showed that has a lot of number of sexual partners did not increase the risk of contracting HIV¹³, having multiple sexual partners or even having a new partner puts adolescents at risk for STI because of a higher chance of exposure to an infected individual, but also because prevention behavior (e.g., condom use) changes with new or different partners³².

Frequency of sexual intercourse in last 6 month showed significantly correlation ($p=0.001$) with HIV infection. Cross tabulation showed that fewer sexual intercourse tend to infected by HIV. It happened because the limitation of research that could not uncover the truth of respondent answered. Its caused some secret and privacy answered or its wrong answered because could not remembered. This result different with general theory that revealed HIV infected correlation with higher sexual intercourse³¹.

Condom used in this study showed no correlation ($p = 0.360$) with HIV status. It happened because measures of condom used only in sexual intercourse last 6 months and researchers can not confirm the honesty of respondents' answers related to the use of condoms as only using questionnaire.

Multivariate analysis

Logistic regression test showed that HIV infection among young age may be related with: residing, continuity of alcohol use, education level and continuity of drug use. It can be explained that young age with senior high school-college level education, typically getting trust from parents than lower education, so they would get less monitoring. Higher Education level at young age also generally have better skills in using media as they wish, but the frequency of use of media among young age associated with sexual sensations seeking⁸, which can increase sex desire and associated also with looking for sexual partners through these media²⁷. This increases their exposure to HIV risk behaviors. In addition, continuity of alcohol and drugs consumption will increase their risk for engaging in risky sexual behavior HIV^{18,21,34}.

Model of HIV infection among young people that was obtained from this study can not be used to good predict. Although, the probabilities amounted to 68.76%, but its AUC value were not good and getting bad in calibration test results.

The limitations of our cross-sectional study include the temporal ambiguity inherent in assessing causal or predictive inferences from these associations. We may have encountered social response bias in some subjects because highly sensitive sexual behavioral information relied on self-report. Although the questionnaire was administered with limited interviewed, its not sure to eliminate information bias. Other limitation were small sample size and subjects may not represent the general population because they may have come for HIV testing for a variety of reasons.

Based on these results, it is recommended to improve information dissemination on HIV, increasing instruction to the risk group especially homosex community, substance use (cigarettes, alcohol and illicit drugs) control of among young age

through facilitating and creation of conducive environment.

CONCLUSION

HIV infection among young people in VCT Clinic of Yogyakarta, by bivariate analysis correlated with gender, residency status, education level, relationship with parents, TV / mobile and internet used to watch pornography. However, according to multivariate analysis, residing with their parents/ family, higher education level and in last 6 months are still using alcohol and illicit drugs.

ACKNOWLEDGEMENT

In this occasion the author acknowledge to Mr. dr. Sumardi, Sp.PD (KP), dr. Lutfan Lazuardi, Ph.D, all of the lecturers and staff of tropical medicine of Gadjah Mada University for their direction and support, and all parties who already helped so this study able to run finely. The author also acknowledge to Novan (PKBI DIY), Zainal Abidin (Edelweis Sardjito), who already helped the author to get the responden and also my friends in Tropical Medicine of Gadjah Mada University of 2010 academic year for their support during a study.

REFERENCE

1. UNAIDS. UNAIDS World AIDS Day Report 2011. Geneva: Joint United Nations Programme on HIV/AIDS, 2011.
2. Ditjen P2PL. Statistik Kasus HIV/AIDS di Indonesia sampai Juni 2011. Jakarta: Kementerian Kesehatan RI, 2011.
3. Djoerban Z. Membidik AIDS Ikhtiar Memahami HIV dan ODHA. 1st ed. Yogyakarta: Galang Press Yogyakarta bekerjasama dengan Yayasan Memajukan Ilmu Penyakit Dalam, 1999.
4. UNAIDS. HIV / AIDS and young people Hope for tomorrow. Geneva: Joint United Nations Programme on HIV/AIDS, 2003.

5. Boyer C, Kegeles S. AIDS risk and prevention among adolescents. *Soc. Sci. Med*, 1991; 33(1):11-23.
6. Borek N, Allison S, Caceres CF. Involving Vulnerable Populations of Youth in HIV Prevention Clinical Research. *J. Acquir. Immune. Defic. Syndr*, 2010;54(Supplement 1): 43–9.
7. Aggleton P, Rivers K. Interventions for Adolescents. In: Gibney L, DiClemente RJ, Vermund SH, editors. *Preventing HIV in Developing Countries Biomedical and Behavioral Approaches*. New York, Boston, Dordrecht, London, Moscow: Kluwer Academic Publishers, 2002;231-55.
8. Whiteley L, Brown L, Swenson R, Romer D, DiClemente R, Salazar L, et al. African American adolescents and new media: associations with HIV/STI risk behavior and psychosocial variables. *Ethn. Dis*, 2011; 21(2): 216–22.
9. Widyatmoko DS. Proses urbanisasi perdesaan di Daerah Istimewa Yogyakarta dan urbanisasi di Indonesia yang melatarbelakanginya (abstrak) [dissertation]. Yogyakarta: Universitas Gadjah Mada, 2007.
10. Yulianingsih. Pengangguran di Yogyakarta Terus Meningkat [Internet]. *Republika*, Selasa, 23 November 2010 23:31 WIB, 2010 [cited 2012 Mar 15]; Available from: <http://www.republika.co.id/berita/breaking-news/nusantara/10/11/23/148299-pengangguran-di-yogyakarta-terus-meningkat>
11. Amornkul PN, Vandenhoudt H, Nasokho P, Odhiambo F. HIV Prevalence and Associated Risk Factors among Individuals Aged 13-34 Years in Rural Western Kenya. *PLoS one*, 2009;4(7):1-11.
12. Ennett ST, Federman EB, Bailey SL, Ringwalt CL, Hubbard ML. HIV-Risk Behaviors Associated With Homelessness Characteristics in Youth. *Journal of Adolescent Health*, 1999; 25(5):344-53.
13. Dorjgochoo T, Noel F, Deschamps MM, Theodore H, Dupont W, Wright PF, et al. Risk Factors for HIV Infection Among Haitian Adolescents and Young Adults Seeking Counseling and Testing in Port-au-Prince. *J. Acquir. Immune. Defic. Syndr*, 2009; 52(4):498-508.
14. Marshall BDL, Kerr T, Shoveller JA, Patterson TL, Buxton JA, Wood E. Homelessness and unstable housing associated with an increased risk of HIV and STI transmission among street-involved youth. *Health & Place*, 2009;15:783-90.
15. Sturdevant MS, Belzer M, Weissman G, Friedman LB, Sarr M, Muenz LR, et al. The Relationship of Unsafe Sexual Behavior and the Characteristics of Sexual Partners of HIV Infected and HIV Uninfected Adolescent Females. *Journal of Adolescent Health*, 2001;29(3s):64-71.
16. Puffer E, Meade C, Drabkin A, Broverman S, Ogwang-Odhiambo R, Sikkema K. Individual and family level psychosocial correlates of HIV risk behavior among youth in rural Kenya. *AIDS Behav*, 2011;15(6):1264-74.
17. CDC. HIV among Youth. USA: Departement of Health and Human Service, 2011.
18. Simbayi L, Kalichman S, Jooste S, Cherry C, Mfecane S, Cain D. Risk factors for HIV-AIDS among youth in Cape Town, South Africa. *AIDS Behav*, 2005;9(1):53-61.
19. Ma Y-H, Ding S-Q, Wang C. Vulnerability to HIV Infection and Related Health Risk Behaviors of the Out of School Adolescents Migrated with Their Parents to Beijing. *Biomedical and Environmental Sciences*, 2008;21:404-10.
20. Karofsky PS, Zeng L, Kosorok MR. Relationship Between Adolescent–Parental Communication and Initiation of First Intercourse by Adolescents. *Journal of Adolescent Health*, 2000;28:41-5.

21. Peltzer K, Pengpid S. Prevalence and Social Correlates of Sexual Intercourse among School-Going Adolescents in Thailand. *The Scientific World Journal*, 2011;11:1812-20.
22. Singh S, Saini S. HIV risk perception in relation to peer pressure and drug abuse behavior among adolescents. *Indian J. Sex Transm. Dis*, 2007;28(1):53-4.
23. Ali MM, Dwyer DS. Estimating peer effects in sexual behavior among adolescents. *Journal of Adolescence*, 2011;34(1):183-90.
24. Elkington KS, Bauermeister JA, Zimmerman MA. Do parents and peers matter? A prospective socio-ecological examination of substance use and sexual risk among African American youth. *Journal of Adolescence*, 2011;34(5):1035-47.
25. Bailey SL, Ouellet LJ, Mackesy-amiti ME, Golub ET, Hagan H, Hudson SM, et al. Perceived risk, peer influences, and injection partner type predict receptive syringe sharing among young adult injection drug users in five U.S. cities. *Drug and Alcohol Dependence*, 2007;91S:18-29.
26. Settertobulte W, Matos MG de. Peer. In: Currie C, Roberts C, Morgan A, Smith R, Settertobulte W, Samdal O, et al., editors. *Young people's health in context. Health Behaviour in School-aged Children (HBSC) study: international report from the 2001/2002 survey*. Copenhagen: WHO Regional Office for Europe, 2004. p. 34-52.
27. Rice E, Monro W, Barman-adhikari A, Young SD. Internet Use, Social Networking, and HIV/AIDS Risk for Homeless Adolescents. *Journal of Adolescent Health*, 2010;47(6):610-3.
28. Mcfarlane M, Bull SS, Rietmeijer CA. Young Adults on the Internet: Risk Behaviors for Sexually Transmitted Diseases and HIV. *Journal of Adolescent Health*, 2002;31(1):11-6.
29. Marshall MM, Kirk GD, Caporaso NE, McCormack MC, Merlo CA, Hague JC, et al. Tobacco use and nicotine dependence among HIV-infected and uninfected injection drug users. *Addictive Behaviors [Internet]*. 2011;36(1-2):61-7. Available from: <http://dx.doi.org/10.1016/j.addbeh.2010.08.022>
30. Drumright LN, Colfax GN. HIV risk and prevention for non-injection substance users. In: Mayer KH, Pizer HF, editors. *HIV Prevention a comprehensive approach*. London: Academic Press, 2009. p. 344-6.
31. Gunung IK, Sumantera IGM, Sawitri AAS, Wirawan DN, editors. *Buku Pegangan Konselor HIV/AIDS*. Denpasar: Yayasan Kerti Praja, Yayasan Burnet Indonesia dan Australia NGO Cooperation Program (ANCP-AusAID), 2003.
32. Ethier KA, Orr DP. Behavioral Interventions for Prevention and Control of STDs Among Adolescents. In: Aral SO, Lipshutz JA, Douglas JM, editors. *Behavioral Interventions for Prevention and control of Sexual Transmitted Diseases*. New York: Springer Science+Business Media, LLC, 2007. p. 277-309.
33. Ford CA, Pence BW, Miller WC, Resnick MD, Bearinger LH, Pettingell S, et al. Predicting Adolescents' Longitudinal Risk for Sexually Transmitted Infection: Results From the National Longitudinal Study of Adolescent Health. *Archives of Pediatrics and Adolescent Medicine*, 2005;159:657-64.
34. Yan AF, Chiu Y-wen, Stoesen CA, Wang MQ. STD-/HIV-Related Sexual Risk Behaviors and Substance Use among U.S. Rural Adolescents. *Journal of The National Medical Association*, 2007;99(12):1386-94.

Tropical Medicine Journal

PAU Building
Jl. Teknik Utara, Berek, Yogyakarta 55281
0274-588483, email: tropmedjournal@gmail.com
Published by Faculty of Medicine, Universitas Gadjah Mada

Instructions to the Authors

Tropical Medicine Journal is a journal devoted to the publication of original articles in all field of basic, tropical medicine and tropical medical biotechnology.

This journal is a journal in tropical medical sciences that used as the media for dissemination of original research, innovative, ideas and new hypotheses in biomedicine, both for medical development, education and application. It also welcomes perspectives articles, biomedical history abridged articles, and reviews.

Statements and opinions expressed in the articles herein are those of author(s) responsibility and not necessary those of the Editor(s), the Faculty of Medicine, or Universitas Gadjah Mada.

Tropical Medicine Journal is published in June and December by the Faculty of Medicine, Universitas Gadjah Mada.

Submission of papers

Articles should be submitted in both hard copy and soft copy forms or in electronic form through e-mails as attachment to: The Editor-in-Chief, Tropical Medicine Journal, Faculty of Medicine, Universitas Gadjah Mada, Sekip Utara, Yogyakarta 55281, Phone: 0274-588483, Fax: 0274-588483
E-mail: tropmedjournal@gmail.com

Basic requirements for articles submitted to Tropical Medicine Journal are: a) original work; b) have not been previously published and not under consideration for publication elsewhere and if accepted will not be published elsewhere; c) should have obtained approval from the Ethics Committee; d) must obtained signed informed

consent from subjects for articles involving human subjects.

Referee suggestions

Upon submission, the author should provide one cover letter. In the covering letter, authors should suggest names and addresses (including e-mail) of at least three experts in the field for evaluation of article. The choice of referees will however remain with the editorial board.

Language

Tropical Medicine Journal will publish the articles in English. Editors encourage authors to submit their articles in English. Even so, when a language barrier is encountered, editors allow authors to submit their article in Bahasa Indonesia and it will be translated in English by in-house translator.

Typescripts

Articles should be neatly typed in Times New Roman, 12 pt, double-spaced on A4 format with 3 cm on all margins. Receipt of papers will be acknowledged. Authors will be informed of the referee's comments.

Article types

Three types of articles may be submitted: a) Original research article (maximum: 25 pages, 35 references); b) Review article (maximum: 40 pages, 100 references); c) Case Report article (maximum: 10 pages, 20 references)

Proofs and Reprints

Proofs of manuscript will be sent to the author for approval prior to publication. Page proofs are considered to be the final version of the manuscript. With the exception of typographical or minor clerical errors, no changes will be made in the manuscript at the proof stage. Corrections should be returned to the Editor within one week. Authors of accepted article will receive 10 free off prints of their articles and can place order for additional off prints or hard copy of the journal after the acceptance of the articles.

Copyright

Submission of an article for publication implies to the transfer of the copyright from the author(s) to the publisher upon acceptance. Accepted articles become the permanent property of Tropical Medicine Journal and may not be reproduced by any means without the written consent of the Editor-in-Chief.

Manuscript preparation

The format of the typescript should be as follows:

- a. Title and authors:** The title should be a brief phrase describing the contents of the article. The title page should include the author's full names and affiliation that marked Arabic number. The name of the corresponding author should be indicated with postal adresse, phone, fax and e-mail information.
- b. Abstract:** The author should provide two abstract, in Indonesian and English language. All articles should be provided with an sbstract of between 200-300 words in one spacing. The abstract should be written in simple language with structured abstract style. Abstract should describe of the study using below headings: Introduction, Objectives, Methods, Results and Discussion, and Conclusion. Standard nomenclature should be used and abbreviations should be avoided.

- c. Keywords:** A maximum of 5 keywords must be given at the end of the abstract.
- d. Introduction:** The Introduction should provide the problem statement clearly, the relevant literature on the subject, and the proposed approach or solution.
- e. Materials and methods:** The materials and methods should be clear enough to allow experiments to be reproduced. Previously published research procedure should be cited, and important modifications of it should be mentioned briefly. If the conducted research involved the use of human subjects or animal laboratory, it should be stated that the clearance from the Research Ethics Committee was obtained. The Editor may request a copy of the clearance document or informed consent form for verification.
- f. Results and Discussion:** The Results should be presented with clarity and precision and explained without referring to the literature. The original and important findings should be stated. The Results should be illustrated with figures or tables where necessary but these should be kept to the minimum. The Discussion should interpret the findings in view of the results obtained against the background of existing knowledge. The Discussion should highlight what is new in the paper. Any assumption on which conclusions are made must be stated clearly
- g. Conclusions:** State the Conclusions in a few sentences at the end of the paper.
- h. Acknowledgments:** The Acknowledgments should be presented at the end of the text and before the references. Technical assistance, financial support and advice may be acknowledged.
- i. Tables:** The tables should be kept to a minimum and be designed to be as simple as possible. Each table should be numbered consecutively in Arabic numerals and supplied

with a heading and a legend. Tables should be self-explanatory without reference to the text.

j. Figure: The figures should be numbered consecutively with Arabic numerals. Graphics should be prepared using applications capable of generating high resolution GIF, TIFF, JPEG or Powerpoint before pasting in the Microsoft Word manuscript file. The figures should be constructed in such a manner that they can be understood without reading the text. Appropriate symbols should be used on graphs and explained in the legends. Graphs should not duplicate results presented in tables. Title and comments of the figures and photographs should be provided on separate page using MS Word.

k. References: References should be numbered consecutively in the order in which they are first mentioned in the text (Vancouver style). Identify references by Arabic number as superscript in order of appearance. A number must be used even if the author(s) is named in the text. The original number assigned to the reference is reused each time the reference is cited in the text, regardless of its previous position in the text. For example :

..... it has been reported¹

..... according to Sardjito²

..... Winstein & Swartz³ conducted

..... by Avon *et al.*⁴

Authors are responsible for the accuracy and the completeness of their references. References should be listed numerically (Vancouver style) at the end of the text and in the same order that they have been cited in the text. For citation references with six or less authors, all authors should be listed, when seven or more authors only first three authors should be listed followed by *et al.* Journal names are abbreviated according to Index Medicus and Index of Indonesia Learned Periodicals (PDIN 1974). References to journal articles, books, chapters in books, theses, etc. should be listed as given in Sample References.

Sample References

Scientific Journal

1. *Standard journal article*

You CH, Lee KY, Chey RY, Menguy R. Electro-gastro-graphic study of patients with unexplained nausea, bloating and vomiting. *Gastroenterology* 1980; 79(2):311-14.

Goate AM, Haynes AR, Owen MJ, Farral M, James LA, Lai LY, et al. Predisposing locus for Alzheimer's disease on chromosome 21. *Lancet* 1989;1:352-55.

2. *Organization as author*

The Royal Marsden Hospital Bone-marrow Transplantation. Team. Failure of syngeneic bone-marrow graft without preconditioning in post-hepatitis marrow aplasia. *Lancet* 1977;2:742-44.

3. *No author given*

Coffee drinking and cancer of the pancreas [editorial]. *BMJ* 1981;283-628.

4. *Article not in English*

Massone L, Borghi S, Pestarino A, Piccini R, Gambini C. Localisations palmaires purpuriques de la dermatite herpetiforme. *Ann Dermatol Venereol* 1987;114:1545-47.

5. *Volume with supplement*

Magni F, Rossoni G, Berti F, BN-52021 protects guinea-pig from heart anaphylaxis. *Pharmacol Res Commun* 1988;20 Suppl 5:75-78.

6. *Issue with supplement*

Gardos G, Cole JO, Haskell D, Marby D, Paine SS, Moore P. The natural history of tardive dyskinesia. *J Clin Psychopharmacol* 1988;8(4 Suppl):31S-37S.

7. *Volume with part*

Hanly C. Metaphysics and innateness: a psychoanalytic perspective. *Int J Psychoanal* 1988;69(Pt 3):389-99.

8. *Issue with part*

Edwards L, Meyskens F, Levine N. Effect of oral isotretinoin on dysplastic nevi. *J Am Acad Dermatol* 1989;20(2 Pt 1):257-60.

9. *Issue with no volume*
Baumeister AA. Origins and control of stereotyped movements. *Monogr Am Assoc Ment Defic* 1978; (3):353-84.
10. *No issue or volume*
Danoek K. Skiing in and through the history of medicine. *Nord Midicinhist Arsb* 1982;86-100.
11. *Pagination in roman numerals*
Ronne Y. Ansvarfall. Bloodtransfusion till fel patients. *Vard-facket* 1989;13:XXVI-XXVII.
12. *Type of article indicated as needed*
Spargo PM, Manners JM, DDAVP and open heart surgery [letter]. *Anaesthesia* 1989;44:363-64.
Fuhrman SA, Joiner KA. Binding of the third component of complement C3 by *Toxoplasma gondii* [abstract]. *Clin Res* 1987; 35:475A.
13. *Article containing retraction*
Shishido A. Retraction notice: Effect of platinum compounds on murine lymphocyte mitogenesis [Retraction of Alsabti EA, Ghalib ON, Salem MH. In: *Jpn J Med Sci Biol* 1979; 32:53-65). *Jpn J Med Sci Biol* 1980;33:235-37.
14. *Article retracted*
Alsabti EA, Ghalib ON, Salem Mh. Effect of platinum compounds on murine lymphocyte mitogenesis [Retracted by Shishido A. In: *Jpn J Med Sci Biol* 1980;33:235-7]. *Jpn J Med Sci Biol* 1979;32:53-65.
15. *Article containing comment*
Piccoli A, Bossatti A. Early steroid therapy in IgA neuropathy: still open question [comment]. *Nephron* 1989;51:289-91.
16. *Article in comment*
Kobayashi Y, Fujii K, Hiki Y, Tateno S, Kurokawa A, Kamiyama M. Steroid therapy in IgA nephropathy: a retrospective study in heavy proteinuric cases [see comments]. *Nephron* 1988;48:12-7. Comment in: *Nephron* 1989;51:289-91.
17. *Article with published erratum*
Schofield A. The CAGE questionnaire and psychological health [published erratum

appears in *Br J Addict* 1989;84:701]. *Br J Addict* 1988;83:761-64.

Books and Other Monographs

18. *Personal author(s)*
Colson JH, Armour WJ. Sports injuries and their treatment. 2nd rev. ed. London: S. Paul, 1986.
19. *Editor(s) as author*
Diener HC, Wilkinson M, editors. Drug-induced headache. New York: Springer-Verlag, 1988.
20. *Organization(s) as author*
Virginia Law Foundation. The medical and legal implications of AIDS. Charlottesville: The Foundation, 1987.
21. *Chapter in a book*
Winstein L, Swartz MN. Pathologic properties of invading microorganisms. In: Sodeman WA Jr, Sodeman WA, editors. *Pathologic Physiology, mechanisms of disease*. Philadelphia: Saunders, 1974:457-72.
22. *Conference proceedings*
Vivian VL, editor. Child abuse and neglect: a medical community response. Proceedings of the First AMA National Conference on Child Abuse and Neglect; 1984 Ma 30-31; Chicago. Chicago: American Medical Association, 1985.
23. *Conference paper*
Harley NH. Comparing radon daughter dosimetric and risk models. In: Gammage RB, Kaye SV, editors. *Indoor air and human health. Proceedings of the Seventh Life Sciences Symposium*; 1984 Oct 29-31; Knoxville (TN). Chelsea (MI):Lewis, 1985:69-78
24. *Scientific or technical report*
Akutsu T. Total heart replacement device. Bethesda (MD): National Institutes of Health. National Heart and Lung Institute; 1974 Apr. Report No.:NIH-NIHI-69-2185-4.
Disertasi Youssef NM. School adjustment of children with congenital heart disease [dissertation]. Pittsburg (PA): Univ. of Pittsburg, 1988.

25. *Dissertation*
 Kay JG. Intracellular cytokine trafficking and phagocytosis in macrophages [Dissertation]. St Lucia, Qld: University of Queensland; 2007.
26. *Patent*
 Harred JF, Knight AR, McIntyre JS, inventors. Dow Chemical Company, assignee. Epoxidation process. US patent 3,654,317, 1972 Apr 4.

Other Published Material

27. *Newspaper article*
 Resberger B, Specter B. CFCs may be destroyed by natural process. The Washington Post 1989 Aug 7;Sect. A:2(col. 5).
28. *Audiovisual material*
 AIDS epidemic: the physician's role [video-recording]. Cleveland (OH): Academy of Medicine of Cleveland, 1987.
29. *Computer program*
 Renal system [computer program]. MS-DOS version. Edwardsville (KS): Medi-Sim, 1988.
30. *Legal material*
 Toxic Substances Control Act: Hearing on S. 776 Before the Subcomm. on the Environment of the Senate Comm. on Commerce, 94th Cong., 1st Sess. 343(1975).
31. *Map*
 Scotland [topographic map]. Washington: National Geographic Society (US), 1981.
32. *Dictionary or Encyclopaedia*
 Ectasia. Dorland's illustrated medical dictionary. 27th ed. Philadelphia: Saunders, 1988: 527.
33. *Classic material*
 The Winter's Tale: act 5, scene I, lines 13-16. The complete works of William Shakespeare. London: Rex, 1973.
34. *In press*
 Lillywhite HB, Donald JA. Pulmonary blood flow regulation in an aquatic snake. Science. In press.

Electronic Material

35. *Journal article in the internet*
 Morse SS. Factors in the emergence of infectious diseases. Emerg Infect Dis [serial online] 1995 Jan-Mar [cited 1996 Jun 5];1(1):[24 screens]. Available from: URL: <http://www.cdc.gov/ncidod/EID/eid.htm>
36. *Monograph in electronic format*
 CDI, clinical dermatology illustrated [monograph on CD-ROM]. Reeves JRT, Maibach H. CMEA Multimedia Group, producers. 2nd ed. Version 2.0 San Diego: CMEA; 1995.
37. *Computer program*
 Hemodynamics III: the ups and downs of hemodynamics [computer program]. Version 2.2. Orlando (FL): Computerized Educational System; 1993.

We thank to the reviewers of this edition:

dr. Abu Tholib, M.Sc., Ph.D., Sp.MK

dr. Ahmad Hamim Sadewa, Ph.D

dr. Arta Firmawati, Ph.D

dr. Elizabeth Henny Heningtyas, M.Si, Ph.D

dr. Hanggoro Tri Rinonce, Ph.D

Prof. Dr. Mustofa, Apt., M.Kes

Dra. Ning Rintiswati, M.Kes

dr. Titik Nuryastuti, Ph.D, Sp.MK