

ISSN 2089 - 2136

Tropical Medicine Journal

Volume 03, No. 2, 2013

- The Effects of Curcumin Against Dengue-2 Virus Based on Immunocytochemistry Technique
- Risk Factors Analysis of Typhoid Fever Occurrence of Inpatient in Kebumen Public Hospital in 2013
- Knowledge, Attitude and Practice on Dengue Fever Transmission Among Urban and Periurban Residents of Dhaka City, Bangladesh
- Geographic Information System (GIS) for Dengue Research in Indonesia: A Review
- Risk Factors of Pneumonia Among Under Five Children in Purbalingga District, Central Java Province
- Factors Associated with Delayed Diagnosis Among Tuberculosis Patient in Kebumen District
- Effication Test of Srikaya Seeds Extract (*Annona squamosa* L.) to Kill *Aedes aegypti* Larvae in Laboratory
- Immune Response against Hepatitis B Virus after Vaccination among Low Birth Weight and Preterm Newborns: A Retrospective Cohort Study in Magelang District Central Java
- Tumor Necrosis Factor-Alpha (TNF-Alpha) and Intercellular Adhesion Molecule-1 (ICAM-1) Expression of *Plasmodium berghei* Infected Swiss Mice Treated with Red Fruit (*Pandanus Conoideus* Lam) Ethanol Extract
- Validity of p-LDH/HRP2-Based Rapid Diagnostic Test for the Diagnosis of Malaria on Pregnant Women in Maluku
- Comparing the Sensitivity and Specificity of Zinc Sulphate Flotation Method to Formal Ether Sedimentation Method in Identifying Intestinal Protozoa's Cysts
- The Effect of Anticoagulant in Blood Meal Source on the *Aedes aegypti* Reproductive Ability in Laboratory

TMJ

Volume
03Number
02Page
95 - 195ISSN
2089 - 2136

Center for Tropical Medicine, Faculty of Medicine, Universitas Gadjah Mada
in collaboration with Indonesian Society of Tropical Medicine and Infectious Disease (PETRI)

Editor-in-chief

Prof. dr. Supargiyono, DTM&H., SU., Ph.D, S.Park

Managing Editor

Dr. dr. Mahardika Agus Wijayanti, M.Kes

Associate Editors

Prof. Dr. Mustofa, M.Kes., Apt.
dr. Yodhi Mahendradhata, M.Sc, Ph.D
Dr. Dra. Erna Kristin, MSi, Apt.
dr. Riris Andono Ahmad, MPH., Ph.D
dr. Doni Priambodo, SpPD-KPTI

Editorial Advisory Board

Dr. Tedjo Sasmono, Bsc
dr. Din Syafruddin, Ph.D
Prof. dr. Ni Made Mertaniasih, MS, Sp.MK
Prof. Dr. dr. Arie Mansyur, SpPD-KPTI
dr. Subagyo Loehoeri, SpPD
Dr. dr. Budiman Bela, Sp.MK (K)

All right reserved. No part of this publication may be reproduced, stored or transmitted in any form or by any mean, electronic or mechanical, without written permission from the publisher.

Address : Tropical Medicine Journal, PAU Building, Jalan Teknik Utara Barek, Yogyakarta

Universitas Gadjah Mada, Yogyakarta 55281, Phone : +62-274-588483, E-mail: tropmedjournal@gmail.com

TROPICAL MEDICINE JOURNAL

ISSN : 2089-2136

Center for Tropical Medicine, Faculty of Medicine, Universitas Gadjah Mada in collaboration with
Indonesian Society of Tropical Medicine and Infectious Disease (PETRI)

Volume 03, Number 02

CONTENTS

- 95 - 102 The Effects of Curcumin Against Dengue-2 Virus Based on Immunocytochemistry Technique
Dewi Marbawati, Sitti Rahmah Umniyati
- 103- 109 Risk Factors Analysis of Typhoid Fever Occurrence of Inpatient in Kebumen Public Hospital in 2013
Rina Hidayani, Hari Kusnanto, Rizka Humardewayanti, Trisno Agung W
- 110 - 120 Knowledge, Attitude and Practice on Dengue Fever Transmission Among Urban and Periurban Residents of Dhaka City, Bangladesh
Muhammad Sohel Rana, Mohammad Syaket Ahmed Shakil
- 121 - 127 Geographic Information System (GIS) for Dengue Research in Indonesia: A Review
Adnanto Wiweko
- 128- 135 Risk Factors of Pneumonia Among Under Five Children in Purbalingga District, Central Java Province
Ni Kadek Nira, Dibylo Pramono, Roni Naning
- 136 - 141 Factors Associated with Delayed Diagnosis Among Tuberculosis Patient in Kebumen District
Edwin Sovvan Aritonang, Ning Rintiswati, Riris Andono Ahmad
- 142 - 148 Effication Test of Srikaya Seeds Extract (*Annona squamosa* L.) to Kill *Aedes aegypti* Larvae in Laboratory
Eny Sofiyatun, Joko Malis Sunarno
- 149 - 154 Immune Response against Hepatitis B Virus after Vaccination among Low Birth Weight and Preterm Newborns: A Retrospective Cohort Study in Magelang District Central Java
Muhardison, Hari Kusnanto, Nenny Sri Mulyani
- 155 - 165 Tumor Necrosis Factor-Alpha (TNF-Alpha) and Intercellular Adhesion Molecule-1 (ICAM-1) Expression of *Plasmodium berghei* Infected Swiss Mice Treated with Red Fruit (*Pandanus Conoideus* Lam) Ethanol Extract
Demianus Tafor, Achmad Djunaidi, Widya Wasityastuti, Eti Nurwening Sholikhah
- 166 - 175 Validity of p-LDH/HRP2-Based Rapid Diagnostic Test for the Diagnosis of Malaria on Pregnant Women in Maluku
Vebiyanti, E. Elsa Herdiana Murhandarwati, Bambang Udji Jokorianto
- 176 - 183 Comparing the Sensitivity and Specificity of Zinc Sulphate Flotation Method to Formol Ether Sedimentation Method in Identifying Intestinal Protozoa's Cysts
Dini Alyani, Elsa Herdiana Murhandarwati, Sri Sumarni, Ernaningsih
- 184 - 195 The Effect of Anticoagulant in Blood Meal Source on the *Aedes aegypti* Reproductive Ability in Laboratory
Novyan Lusiyana, Budi Mulyaningsih, Sitti Rahmah Umniyati

Factors Associated with Delayed Diagnosis Among Tuberculosis Patient in Kebumen District

Edwin Sovvan Aritonang^{1*}, Ning Rintiswati², Riris Andono Ahmad³

¹Sibolga City Health Department, North Sumatera, Indonesia; ²Department of Microbiology, Faculty of Medicine, Universitas Gadjah Mada, Yogyakarta, Indonesia; ³Department of Public Health, Faculty of Medicine, Universitas Gadjah Mada, Yogyakarta, Indonesia.

Corresponding author: edwinsbg@yahoo.co.id

ABSTRACT

Introduction: Tuberculosis is a major global health problem. It is estimated that almost 9 million new cases detected in 2011 and 1.4 million died because of TB. Early diagnosis and effective treatment are the key elements of the TB control program. Delayed diagnosis associated with a longer duration of infectiousness, increase the severity of the disease, more severe complication and even death.

Objectives: To determine the factors associated with delayed diagnosis among tuberculosis patients in Kebumen District.

Methods : A cross sectional study among newly diagnosed TB patients in April to June 2013 in Kebumen was conducted. The data was collected using a questionnaire interviewing officers of government health care facilities which have implemented DOTS programs already. Data was analyzed by a logistic regression test with confidence interval (CI) of 95%.

Results: Eighty five pulmonary TB patients were studied consisting of 65 TB smear positive and 20 smear negative/roentgen positive patients. The median duration of delayed patients was 3.9 weeks; delayed health service was 2.6 weeks and delayed diagnosis was 7.6 weeks. Multivariate analysis showed that factors associated with delayed diagnosis were the type of health services which first visited (p value = 0.002 and OR = 6.87), level of knowledge about TB (p value = 0.002 and OR = 6.41), educational level (p value = 0.024 and OR = 5.68) and the number of visits to health service (p value = 0.021 and OR = 3.87).

Conclusion : The median duration of delayed diagnosis was 7.6 weeks. Delayed diagnosis among TB patients in Kebumen District was associated with type of health services which is non-DOTS, low level of knowledge about TB, low levels of education and the number of visits to health services e"3 times.

Keywords: delayed diagnosis, TB patients, Kebumen District

INTISARI

Pendahuluan: Tuberkulosis (TB) merupakan masalah kesehatan global yang utama. Diperkirakan 9 juta kasus baru pada tahun 2011 dan 1,4 juta meninggal karena TB. Diagnosis dini dan pengobatan yang efektif merupakan bagian penting dari program pengendalian TB. Keterlambatan diagnosis terkait dengan durasi penularan yang lebih lama, peningkatan keparahan penyakit dan menyebabkan komplikasi yang lebih berat dan bahkan kematian.

Tujuan: Untuk mengetahui faktor-faktor yang berhubungan dengan keterlambatan diagnosis antara pasien tuberkulosis di Kabupaten Kebumen.

Metode: Penelitian ini merupakan penelitian potomng lintang pada pasien TB yang didiagnosis baru pada bulan April sampai Juni 2013 di Kabupaten Kebumen. Data dikumpulkan dengan kuesioner dengan mewawancarai petugas di fasilitas kesehatan pemerintah yang telah melaksanakan program DOTS. Data dianalisis dengan uji regresi logistik dengan interval kepercayaan (IK) 95%.

Hasil: Delapan puluh lima pasien TB paru diteliti yang terdiri dari 65 pasien TB dengan BTA positif dan 20 pasien TB dengan BTA negatif/rontgen positif. Durasi rata-rata keterlambatan pasien adalah 3,9 minggu; keterlambatan pelayanan kesehatan adalah 2,6 minggu; dan keterlambatan diagnosis adalah 7,6 minggu. Analisis multivariabel menunjukkan bahwa faktor yang terkait dengan keterlambatan diagnosis adalah jenis pelayanan kesehatan yang pertama kali dikunjungi ($p = 0,002$ dan $OR = 6,87$), tingkat pengetahuan tentang TB ($p = 0,002$ dan $OR = 6,41$), tingkat pendidikan ($p = 0,024$ dan $OR = 5,68$) dan jumlah kunjungan ke pelayanan kesehatan ($p = 0,021$ dan $OR = 3,87$).

Simpulan: Durasi rata-rata keterlambatan diagnosis adalah 7,6 minggu. Keterlambatan diagnosis pasien TB di Kabupaten Kebumen terkait dengan jenis pelayanan kesehatan yang pertama kali dikunjungi adalah non-DOTS, rendahnya tingkat pengetahuan tentang TB, rendahnya tingkat pendidikan dan jumlah kunjungan ke pelayanan kesehatan e" 3 kali.

Kata kunci: keterlambatan diagnosis, pasien TB, Kabupaten Kebumen

INTRODUCTION

Tuberculosis is a major global health problem. It is estimated that almost 9 million of new cases were diagnosed in 2011 and 1.4 million were died because TB. It is the second leading cause of death from an infectious disease worldwide after HIV. In addition, an estimated of 95% of TB motality in the world occur in developing countries¹.

According to WHO (2006), a key element of the TB control program is early diagnosis and effective treatment. Delayed diagnosis can increase its ease transmission in the community, the severity of disease, and risk of death². Greenaway et al. suggested that there is a relationship between delayed diagnosis and treatment with mortality in patients with tuberculosis³.The other studies suggested that early diagnosis and treatment contribute to

decreasing mortality and transmission of TB in the community².

In Indonesia, the TB control program faces a major problem because of the high prevalence of TB which is 289 per 100,000 population, the incidence rate of 189 per 100,000 population and the mortality rate due to TB is 27 per 100,000 population by the number of deaths of 61,000 people per year⁴.

In Kebumen District, TB remains one of the priority program because of the high prevalence of TB cases which is 136.9 per 100,000 population in 2011 and 126.2 per 100,000 population in 2012, deaths due to TB increased from 18 cases in 2011 to 23 cases in 2012, yet the achievement of new case detection rate (CDR) is 60.4% in 2011 and 59.9% in 2012⁵. This might be caused by a delayed diagnosis and non-DOTS treatment.

MATERIALS AND METHODS

This study used a cross sectional design. The location of the study is Kebumen District which includes 35 units of Community Health Centers (PHC=Puskesmas). The study population was all new pulmonary TB patients who have been diagnosed and are being treated for tuberculosis at the Community Health Centers in Kebumen District with age e" 15 years who meet the inclusion and exclusion criteria. The sample size of this study is 85 individuals. Sampling was performed using a consecutive sampling technique .

The dependent variable in this study is delayed diagnosis, while the independent variables are demographic characteristics (age, gender, educational level, occupation, and socio-economic), access to health services (distance, travel time, and access to health care costs), knowledge about TB, stigma and treatment seeking patterns (action treatment was first performed, the type of health services which first visited and number of visits to health service).

The results were analyzed in univariate, bivariate and multivariate tests. For bivariate and multivariate analysis, a logistic regression with confidence level (á) of 95% was used.

RESULTS AND DISCUSSION

The median duration of delayed diagnosis was 7.6 weeks (range 2.4-31.0 weeks) and proportion of delayed diagnosis of 49.4%. The results of bivariate analysis showed that there are six variables which statistically significant with delayed diagnosis amongst tuberculosis patient in Kebumen District i.e. 1) the level of education (p = 0.008 ; OR = 5.09 and 95 % CI = 1.52-17.00), 2) socioeconomic status (p = 0.016 ; OR = 4.05 and 95% CI = 1.31- 12.59), 3) access to health care (p = 0.035 ; OR = 2.73 and 95 % CI = 1.07-6.93), 4) level of knowledge about TB (p = 0.000 ; OR = 6.49 and 95% CI = 2.52 - 16.73), 5) the type of health services which is first visited (p = 0,000 ; OR = 6.50 and 95% CI = 2.43 -7.38) and 6) the number of visit to health care (p = 0.001 ; OR = 4.62 and 95 % CI = 1.85 -11.52).

Table 1. Results of multivariate logistic regression analysis between risk factors with diagnosis delay in Kebumen District

No.	Variable	Diagnosis Delay		OR	CI 95%	p value
		≤ 7,6 weeks n (%)	> 7,6 weeks n (%)			
1	Type of health service					
	- Non DOTS	34 (66,7)	17 (33,3)	6,87	2,04-23,15	0,002**
	- DOTS	8 (23,5)	26 (76,5)	1		
2	Level of knowledge					
	- Low	29 (72,5)	11 (27,5)	6,41	1,99-20,60	0,002*
	- High	13 (28,9)	32 (71,1)	1		
3	Education level					
	- Low	38 (57,6)	28 (42,4)	5,68	1,26-25,53	0,024*
	- High	4 (21,1)	15 (78,9)			
4	The number of visits					
	- ≥ 3 times	29 (67,4)	14 (32,6)	3,87	1,23-12,21	0,021*
	- < 3 times	13 (31,0)	29 (69,0)			

Information : * = p value < 0,05 ** = p value < 0,01

The result of multivariate analysis (Table 1), showed that diagnosis delay was statistically associated with the type of health care which first visited ($p = 0.002$ and $OR = 6.87$), level of knowledge about tuberculosis ($p = 0.002$ and $OR = 6.41$), educational level (p value = 0.024 and $OR = 5.68$), and the number of visits to health service (p value = 0.021 and $OR = 3.87$).

The results showed that the median duration of delayed diagnosis in Kebumen District was 7.6 weeks, similar to the study in other studies which are 7 weeks (49 days) and 8 weeks^{6,7}. The duration of delayed diagnosis detected in this study is lower compared with studies in India (8.6 weeks) and Nepal (70.5 days)^{8,9}. The duration of diagnosis delay in Kebumen still longer than research in Yogyakarta (5.4 weeks)². This because of Yogyakarta is one of Indonesia's provinces with strong TB control programs and health systems are already well established², unlike in Kebumen District where TB control programs and health systems still need to be improved. This is evident from the high number of people with TB, case detection rate and the recruitment rate of tuberculosis suspects who are still below the target as well as the duration of the delayed diagnosis.

The education level is statistically associated with delayed diagnosis. Low level of education more at risk of delayed diagnosis compared to the high level of education. These results are consistent with research in India and Kenya^{10,11}. Educational level associated with a person's ability to absorb and receive information. People who have higher education levels are generally absorb and receive information about health issues easily compared with the less educated, thus affecting the decision in utilizing the

available health services. Delayed diagnosis is common among people who are illiterate and have low education level (did not finish primary/junior school). Level of education may determine the knowledge about the importance of early diagnosis and treatment seeking. In this case, people with low education levels tend to know the importance of seeking treatment after falling ill¹¹.

The level of knowledge about TB was statistically associated with delayed diagnosis. These results are consistent with research in Surakarta¹². Rosenstock cit. Notoatmodjo argued that knowledge of the disease is a factor that can influence a person's perception of a disease that ultimately can affect a person's behavior to reduce the threat of disease and knowledge is also needed to change the mindset and behavior¹³. According to Anderson cit. Samad, knowledge about illness and disease can be used as a basis in the use of health services, the higher the level of public knowledge about the disease, the higher to use health services¹⁰.

The type of health services which first visited statistically associated with delayed diagnosis and visiting non-DOTS health services more at risk of delayed diagnosis compared DOTS health services. These results are consistent with research conducted by Lönnroth et al. and Needhem et al. which suggests that patients visiting non DOTS-health service (private practice/personal) are more at risk of delayed diagnosis compared with who visited DOTS-health service. Delays in accessing DOTS-health services for TB diagnosis and treatment is a major challenge in Indonesia since it has a very wide geographic area. This causes most of the TB patients receive treatment from health

services outside the national TB program. The coverage DOTS-health services is not adequate and the number of health services (hospitals and private practices) that have not participated in the TB control program also causes TB patients experience delayed diagnosis and early treatment according to standard TB treatment². In Kebumen District, most of health professionals (doctors, nurses, laboratory personnel and TB programmers) in DOTS -health services (health centers/hospitals) have received training on the prevention and reduction of pulmonary TB. In addition, the number of health services and the availability of facilities such as DOTS diagnostic tool (binocular microscope) and supporting equipment diagnostics (reagents) are adequate¹⁴.

The number of visits to health care are statistically associated with delayed diagnosis. Visiting health services e" 3 times is more at risk of delayed diagnosis compared with visited health care <3 times. These results are consistent with research in India¹⁰. The more a person visiting health care, more likely to experience delayed diagnosis. Many TB patients have to visit several times to health care before being diagnosed. As many as two thirds of patients do visit more than 2 (two) times and nearly a quarter of patients do visit more than 6 (six) times before being diagnosed. This is because TB patients receive several different diagnosis before referred or deciding to visit higher health services¹⁵.

CONCLUSION

The median duration of delayed diagnosis amongst tuberculosis patients in Kebumen District was 7.6 weeks. Factors associated with

delayed diagnosis among tuberculosis patient in Kebumen are 1) the type of health services which first visited is Non DOTS, 2) low level of knowledge about TB, 3) low level of education and 4) the number of visits to health service e" 3 times.

RECOMMENDATION

Health Department of Kebumen District need to strengthen, improve and expand TB control programs to private health care (hospital/clinic/private practices) that have not participated in the implementation of the DOTS programs because long duration of delayed diagnosis and mostly occurring non-DOTS treatment. The health center is expected to provide information about TB to the community through health education, mass media and other medias (leaflets /brochures/posters) to increase public knowledge about TB in order to prevent delayed diagnosis. They are also expected to actively seeking information about TB in order to increase their knowledge.

REFERENCE

1. WHO. Global Tuberculosis Report 2012: WHO, Geneva; 2012.
2. Ahmad RA, Mahendradhata Y, Utarini A, de Vlas SJ. Diagnostic delay amongst tuberculosis patients in Jogjakarta Province, Indonesia is related to the quality of services in DOTS facilities. *Tropical Medicine & International Health*, 2011;16(4):412-23.
3. Pujiati S. Ketidaktepatan Waktu Pengambilan Obat Antituberkulosis pada Penderita Tuberkulosis Paru Baru Setelah Didiagnosis BTA Positif dan Faktor-Faktor Individu yang Mempengaruhinya di Puskesmas Kecamatan

- Ciracas, Jakarta Timur dan Puskesmas Satelitnya Periode 1 Januari 2007 – 23 Desember 2008: Program Pasca Sarjana, Fakultas Kesehatan Masyarakat, Universitas Indonesia, Depok, 2009.
4. Kemenkes RI. Strategi Nasional Pengendalian TB di Indonesia 2010-2014: Ditjen P2&PL, Jakarta, 2011.
 5. Dinkes Kabupaten Kebumen. Laporan Program Pencegahan dan Penanggulangan Tuberculosis: Bidang PMK, Kebumen, 2012.
 6. Leung EC, Leung C, Tam C. Delayed presentation and treatment of newly diagnosed pulmonary tuberculosis patients in Hong Kong. *Hong Kong Medical Journal*, 2007;13(3):221.
 7. Saqib M, Awan I, Rizvi S, Shahzad M, Mirza Z, Tahseen S, et al. Delay in diagnosis of tuberculosis in Rawalpindi, Pakistan. *BMC research notes*, 2011;4(1):165.
 8. Rajeswari R, Chandrasekaran V, Suhadev M, Sivasubramaniam S, Sudha G, Renu G. Factors associated with patient and health system delays in the diagnosis of tuberculosis in South India. *Int J Tuberc Lung Dis*, 2002;6: 789-95.
 9. Belay M, Bjune G, Ameni G, Abebe F. Diagnostic and treatment delay among Tuberculosis patients in Afar Region, Ethiopia: A cross-sectional study. *BMC Public Health*, 2012;12(1):369.
 10. Sumana M. Health Care Seeking Behaviour and Reasons for Delays in Diagnosis and Treatment of Tuberculosis Patients in Mysore City, 2010.
 11. Njau IW, Karanja SM, Wanzala P, Omolo JO. Factors associated with late presentation of suspected tuberculosis cases to tuberculosis management facilities: The case in Dagoretti district, Nairobi, Kenya. *Pan African Medical Journal*, 2012;12:93.
 12. Selleca C, Sarah. Hubungan Status Ekonomi dan Tingkat Pengetahuan TB dengan Keterlambatan Pasien dalam Diagnosis Kasus TB Paru. Surakarta: Universitas Sebelas Maret; 2012.
 13. Wulandari L. Peran Pengetahuan terhadap Perilaku Pencarian Pengobatan Penderita Suspek TB Paru di Indonesia (Analisis Data Survei Pengetahuan, Sikap dan Perilaku Tuberculosis Tahun 2010): Program Studi Ilmu Kesehatan Masyarakat, Fakultas Kesehatan Masyarakat, Universitas Indonesia. Depok, 2012.
 14. Efrizal. Evaluasi Program Penanggulangan Tuberculosis Di Kabupaten Kebumen Propinsi Jawa Tengah Tahun 2011, 2011.
 15. Yan F, Thomson R, Tang S, Squire SB, Wang W, Liu X, et al. Multiple perspectives on diagnosis delay for tuberculosis from key stakeholders in poor rural China: Case study in four provinces. *Health Policy*, 2007; 82(2): 186-99.

Tropical Medicine Journal

PAU Building
Jl. Teknik Utara, Berek, Yogyakarta 55281
0274-588483, email: tropmedjournal@gmail.com
Published by Faculty of Medicine, Universitas Gadjah Mada

Instructions to the Authors

Tropical Medicine Journal is a journal devoted to the publication of original articles in all field of basic, tropical medicine and tropical medical biotechnology.

This journal is a journal in tropical medical sciences that used as the media for dissemination of original research, innovative, ideas and new hypotheses in biomedicine, both for medical development, education and application. It also welcomes perspectives articles, biomedical history abridged articles, and reviews.

Statements and opinions expressed in the articles herein are those of author(s) responsibility and not necessary those of the Editor(s), the Faculty of Medicine, or Universitas Gadjah Mada.

Tropical Medicine Journal is published in June and December by the Faculty of Medicine, Universitas Gadjah Mada.

Submission of papers

Articles should be submitted in both hard copy and soft copy forms or in electronic form through e-mails as attachment to: The Editor-in-Chief, Tropical Medicine Journal, Faculty of Medicine, Universitas Gadjah Mada, Sekip Utara, Yogyakarta 55281, Phone: 0274-588483, Fax: 0274-588483
E-mail: tropmedjournal@gmail.com

Basic requirements for articles submitted to Tropical Medicine Journal are: a) original work; b) have not been previously published and not under consideration for publication elsewhere and if accepted will not be published elsewhere; c) should have obtained approval from the Ethics Committee; d) must have obtained signed informed

consent from subjects for articles involving human subjects.

Referee suggestions

Upon submission, the author should provide one cover letter. In the covering letter, authors should suggest names and addresses (including e-mail) of at least three experts in the field for evaluation of article. The choice of referees will however remain with the editorial board.

Language

Tropical Medicine Journal will publish the articles in English. Editors encourage authors to submit their articles in English. Even so, when a language barrier is encountered, editors allow authors to submit their article in Bahasa Indonesia and it will be translated in English by in-house translator.

Typescripts

Articles should be neatly typed in Times New Roman, 12 pt, double-spaced on A4 format with 3 cm on all margins. Receipt of papers will be acknowledged. Authors will be informed of the referee's comments.

Article types

Three types of articles may be submitted: a) Original research article (maximum: 25 pages, 35 references); b) Review article (maximum: 40 pages, 100 references); c) Case Report article (maximum: 10 pages, 20 references)

Proofs and Reprints

Proofs of manuscript will be sent to the author for approval prior to publication. Page proofs are considered to be the final version of the manuscript. With the exception of typographical or minor clerical errors, no changes will be made in the manuscript at the proof stage. Corrections should be returned to the Editor within one week. Authors of accepted article will receive 10 free off prints of their articles and can place order for additional off prints or hard copy of the journal after the acceptance of the articles.

Copyright

Submission of an article for publication implies to the transfer of the copyright from the author(s) to the publisher upon acceptance. Accepted articles become the permanent property of Tropical Medicine Journal and may not be reproduced by any means without the written consent of the Editor-in-Chief.

Manuscript preparation

The format of the typescript should be as follows:

- a. Title and authors:** The title should be a brief phrase describing the contents of the article. The title page should include the author's full names and affiliation that marked Arabic number. The name of the corresponding author should be indicated with postal adresse, phone, fax and e-mail information.
- b. Abstract:** The author should provide two abstract, in Indonesian and English language. All articles should be provided with an sbstract of between 200-300 words in one spacing. The abstract should be written in simple language with structured abstract style. Abstract should describe of the study using below headings: Introduction, Objectives, Methods, Results and Discussion, and Conclusion. Standard nomenclature should be used and abbreviations should be avoided.

- c. Keywords:** A maximum of 5 keywords must be given at the end of the abstract.
- d. Introduction:** The Introduction should provide the problem statement clearly, the relevant literature on the subject, and the proposed approach or solution.
- e. Materials and methods:** The materials and methods should be clear enough to allow experiments to be reproduced. Previously published research procedure should be cited, and important modifications of it should be mentioned briefly. If the conducted research involved the use of human subjects or animal laboratory, it should be stated that the clearance from the Research Ethics Committee was obtained. The Editor may request a copy of the clearance document or informed consent form for verification.
- f. Results and Discussion:** The Results should be presented with clarity and precision and explained without referring to the literature. The original and important findings should be stated. The Results should be illustrated with figures or tables where necessary but these should be kept to the minimum. The Discussion should interpret the findings in view of the results obtained against the background of existing knowledge. The Discussion should highlight what is new in the paper. Any assumption on which conclusions are made must be stated clearly
- g. Conclusions:** State the Conclusions in a few sentences at the end of the paper.
- h. Acknowledgments:** The Acknowledgments should be presented at the end of the text and before the references. Technical assistance, financial support and advice may be acknowledged.
- i. Tables:** The tables should be kept to a minimum and be designed to be as simple as possible. Each table should be numbered consecutively in Arabic numerals and supplied

with a heading and a legend. Tables should be self-explanatory without reference to the text.

j. Figure: The figures should be numbered consecutively with Arabic numerals. Graphics should be prepared using applications capable of generating high resolution GIF, TIFF, JPEG or Powerpoint before pasting in the Microsoft Word manuscript file. The figures should be constructed in such a manner that they can be understood without reading the text. Appropriate symbols should be used on graphs and explained in the legends. Graphs should not duplicate results presented in tables. Title and comments of the figures and photographs should be provided on separate page using MS Word.

k. References: References should be numbered consecutively in the order in which they are first mentioned in the text (Vancouver style). Identify references by Arabic number as superscript in order of appearance. A number must be used even if the author(s) is named in the text. The original number assigned to the reference is reused each time the reference is cited in the text, regardless of its previous position in the text. For example :

..... it has been reported¹

..... according to Sardjito²

..... Winstein & Swartz³ conducted

..... by Avon *et al.*⁴

Authors are responsible for the accuracy and the completeness of their references. References should be listed numerically (Vancouver style) at the end of the text and in the same order that they have been cited in the text. For citation references with six or less authors, all authors should be listed, when seven or more authors only first three authors should be listed followed by *et al.* Journal names are abbreviated according to Index Medicus and Index of Indonesia Learned Periodicals (PDIN 1974). References to journal articles, books, chapters in books, theses, etc. should be listed as given in Sample References.

Sample References

Scientific Journal

1. *Standard journal article*

You CH, Lee KY, Chey RY, Menguy R. Electro-gastro-graphic study of patients with unexplained nausea, bloating and vomiting. *Gastroenterology* 1980; 79(2):311-14.

Goate AM, Haynes AR, Owen MJ, Farral M, James LA, Lai LY, et al. Predisposing locus for Alzheimer's disease on chromosome 21. *Lancet* 1989;1:352-55.

2. *Organization as author*

The Royal Marsden Hospital Bone-marrow Transplantation. Team. Failure of syngeneic bone-marrow graft without preconditioning in post-hepatitis marrow aplasia. *Lancet* 1977;2:742-44.

3. *No author given*

Coffee drinking and cancer of the pancreas [editorial]. *BMJ* 1981;283-628.

4. *Article not in English*

Massone L, Borghi S, Pestarino A, Piccini R, Gambini C. Localisations palmaires purpuriques de la dermatite herpetiforme. *Ann Dermatol Venereol* 1987;114:1545-47.

5. *Volume with supplement*

Magni F, Rossoni G, Berti F, BN-52021 protects guinea-pig from heart anaphylaxis. *Pharmacol Res Commun* 1988;20 Suppl 5:75-78.

6. *Issue with supplement*

Gardos G, Cole JO, Haskell D, Marby D, Paine SS, Moore P. The natural history of tardive dyskinesia. *J Clin Psychopharmacol* 1988;8(4 Suppl):31S-37S.

7. *Volume with part*

Hanly C. Metaphysics and innateness: a psychoanalytic perspective. *Int J Psychoanal* 1988;69(Pt 3):389-99.

8. *Issue with part*

Edwards L, Meyskens F, Levine N. Effect of oral isotretinoin on dysplastic nevi. *J Am Acad Dermatol* 1989;20(2 Pt 1):257-60.

9. *Issue with no volume*
Baumeister AA. Origins and control of stereotyped movements. *Monogr Am Assoc Ment Defic* 1978; (3):353-84.
10. *No issue or volume*
Danoek K. Skiing in and through the history of medicine. *Nord Midicinhist Arsb* 1982;86-100.
11. *Pagination in roman numerals*
Ronne Y. Ansvarfall. Bloodtransfusion till fel patients. *Vard-facket* 1989;13:XXVI-XXVII.
12. *Type of article indicated as needed*
Spargo PM, Manners JM, DDAVP and open heart surgery [letter]. *Anaesthesia* 1989;44:363-64.
Fuhrman SA, Joiner KA. Binding of the third component of complement C3 by *Toxoplasma gondii* [abstract]. *Clin Res* 1987; 35:475A.
13. *Article containing retraction*
Shishido A. Retraction notice: Effect of platinum compounds on murine lymphocyte mitogenesis [Retraction of Alsabti EA, Ghalib ON, Salem MH. In: *Jpn J Med Sci Biol* 1979; 32:53-65). *Jpn J Med Sci Biol* 1980;33:235-37.
14. *Article retracted*
Alsabti EA, Ghalib ON, Salem Mh. Effect of platinum compounds on murine lymphocyte mitogenesis [Retracted by Shishido A. In: *Jpn J Med Sci Biol* 1980;33:235-7]. *Jpn J Med Sci Biol* 1979;32:53-65.
15. *Article containing comment*
Piccoli A, Bossatti A. Early steroid therapy in IgA neuropathy: still open question [comment]. *Nephron* 1989;51:289-91.
16. *Article in comment*
Kobayashi Y, Fujii K, Hiki Y, Tateno S, Kurokawa A, Kamiyama M. Steroid therapy in IgA nephropathy: a retrospective study in heavy proteinuric cases [see comments]. *Nephron* 1988;48:12-7. Comment in: *Nephron* 1989;51:289-91.
17. *Article with published erratum*
Schofield A. The CAGE questionnaire and psychological health [published erratum

appears in *Br J Addict* 1989;84:701]. *Br J Addict* 1988;83:761-64.

Books and Other Monographs

18. *Personal author(s)*
Colson JH, Armour WJ. Sports injuries and their treatment. 2nd rev. ed. London: S. Paul, 1986.
19. *Editor(s) as author*
Diener HC, Wilkinson M, editors. Drug-induced headache. New York: Springer-Verlag, 1988.
20. *Organization(s) as author*
Virginia Law Foundation. The medical and legal implications of AIDS. Charlottesville: The Foundation, 1987.
21. *Chapter in a book*
Winstein L, Swartz MN. Pathologic properties of invading microorganisms. In: Sodeman WA Jr, Sodeman WA, editors. *Pathologic Physiology, mechanisms of disease*. Philadelphia: Saunders, 1974:457-72.
22. *Conference proceedings*
Vivian VL, editor. Child abuse and neglect: a medical community response. Proceedings of the First AMA National Conference on Child Abuse and Neglect; 1984 Ma 30-31; Chicago. Chicago: American Medical Association, 1985.
23. *Conference paper*
Harley NH. Comparing radon daughter dosimetric and risk models. In: Gammage RB, Kaye SV, editors. *Indoor air and human health. Proceedings of the Seventh Life Sciences Symposium*; 1984 Oct 29-31; Knoxville (TN). Chelsea (MI):Lewis, 1985:69-78
24. *Scientific or technical report*
Akutsu T. Total heart replacement device. Bethesda (MD): National Institutes of Health. National Heart and Lung Institute; 1974 Apr. Report No.:NIH-NIHI-69-2185-4.
Disertasi Youssef NM. School adjustment of children with congenital heart disease [dissertation]. Pittsburg (PA): Univ. of Pittsburg, 1988.

25. *Dissertation*
Kay JG. Intracellular cytokine trafficking and phagocytosis in macrophages [Dissertation]. St Lucia, Qld: University of Queensland; 2007.

26. *Patent*
Harred JF, Knight AR, McIntyre JS, inventors. Dow Chemical Company, assignee. Epoxidation process. US patent 3,654,317, 1972 Apr 4.

Other Published Material

27. *Newspaper article*
Resberger B, Specter B. CFCs may be destroyed by natural process. The Washington Post 1989 Aug 7;Sect. A:2(col. 5).

28. *Audiovisual material*
AIDS epidemic: the physician's role [video-recording]. Cleveland (OH): Academy of Medicine of Cleveland, 1987.

29. *Computer program*
Renal system [computer program]. MS-DOS version. Edwardsville (KS): Medi-Sim, 1988.

30. *Legal material*
Toxic Substances Control Act: Hearing on S. 776 Before the Subcomm. on the Environment of the Senate Comm. on Commerce, 94th Cong., 1st Sess. 343(1975).

31. *Map*
Scotland [topographic map]. Washington: National Geographic Society (US), 1981.

32. *Dictionary or Encyclopaedia*
Ectasia. Dorland's illustrated medical dictionary. 27th ed. Philadelphia: Saunders, 1988: 527.

33. *Classic material*
The Winter's Tale: act 5, scene I, lines 13-16. The complete works of William Shakespeare. London: Rex, 1973.

34. *In press*
Lillywhite HB, Donald JA. Pulmonary blood flow regulation in an aquatic snake. Science. In press.

Electronic Material

35. *Journal article in the internet*
Morse SS. Factors in the emergence of infectious diseases. Emerg Infect Dis [serial online] 1995 Jan-Mar [cited 1996 Jun 5];1(1):[24 screens]. Available from: URL: <http://www.cdc.gov/ncidod/EID/eid.htm>

36. *Monograph in electronic format*
CDI, clinical dermatology illustrated [monograph on CD-ROM]. Reeves JRT, Maibach H. CMEA Multimedia Group, producers. 2nd ed. Version 2.0 San Diego: CMEA; 1995.

37. *Computer program*
Hemodynamics III: the ups and downs of hemodynamics [computer program]. Version 2.2. Orlando (FL): Computerized Educational System; 1993.

We thank to the reviewers of this edition:

dr. Abu Tholib, M.Sc., Ph.D., Sp.MK

dr. Ahmad Hamim Sadewa, Ph.D

dr. Arta Farmawati, Ph.D

dr. Elizabeth Henny Herningtyas, M.Si, Ph.D

Dr. Dra. Erna Kristin, Apt., M.Si.

dr. Hanggoro Tri Rinonce, Ph.D

dr. M. Lutfan Lazuardi, M.Kes., Ph.D

dr. Titik Nuryastuti, Ph.D, Sp.MK

dr. Tri Wibawa, Ph.D

Dra. Raden Ajeng Yayi Suryo Prabandari, M.Si., Ph.D