

# T-Cell Responses to the DBL $\alpha$ -Tag, a Short Semi-Conserved Region of the *Plasmodium falciparum* Membrane Erythrocyte Protein 1

Evelyn N. Gitau<sup>1,2</sup>, James Tuju<sup>1</sup>, Liz Stevenson<sup>1 $\ddagger$</sup> , Eva Kimani<sup>1</sup>, Henry Karanja<sup>1</sup>, Kevin Marsh<sup>1,3</sup>, Peter C. Bull<sup>1,3</sup>, Britta C. Urban<sup>1,2\*</sup>

**1** KEMRI-Wellcome Trust Collaborative Programme, Centre for Geographic Medicine Coast, Kilifi, Kenya, **2** Liverpool School of Tropical Medicine, Liverpool, United Kingdom, **3** Centre for Tropical Medicine, Nuffield Department of Internal Medicine, Oxford University, Oxford, United Kingdom

## Abstract

The *Plasmodium falciparum* erythrocyte membrane protein 1 (PfEMP1) is a variant surface antigen expressed on mature forms of infected erythrocytes. It is considered an important target of naturally acquired immunity. Despite its extreme sequence heterogeneity, variants of PfEMP1 can be stratified into distinct groups. Group A PfEMP1 have been independently associated with low host immunity and severe disease in several studies and are now of potential interest as vaccine candidates. Although antigen-specific antibodies are considered the main effector mechanism in immunity to malaria, the induction of efficient and long-lasting antibody responses requires CD4+ T-cell help. To date, very little is known about CD4+ T-cell responses to PfEMP1 expressed on clinical isolates. The DBL $\alpha$ -tag is a small region from the DBL $\alpha$ -domain of PfEMP1 that can be amplified with universal primers and is accessible in clinical parasite isolates. We identified the dominant expressed PfEMP1 in 41 individual clinical parasite isolates and expressed the corresponding DBL $\alpha$ -tag as recombinant antigen. Individual DBL $\alpha$ -tags were then used to activate CD4+ T-cells from acute and convalescent blood samples in children who were infected with the respective clinical parasite isolate. Here we show that CD4+ T-cell responses to the homologous DBL $\alpha$ -tag were induced in almost all children during acute malaria and maintained in some for 4 months. Children infected with parasites that dominantly expressed group A-like PfEMP1 were more likely to maintain antigen-specific IFN $\gamma$ -producing CD4+ T-cells than children infected with parasites dominantly expressing other PfEMP1. These results suggest that group A-like PfEMP1 may induce long-lasting effector memory T-cells that might be able to provide rapid help to variant-specific B cells. Furthermore, a number of children induced CD4+ T-cell responses to heterologous DBL $\alpha$ -tags, suggesting that CD4+ T-cells may recognise shared epitopes between several DBL $\alpha$ -tags.

**Citation:** Gitau EN, Tuju J, Stevenson L, Kimani E, Karanja H, et al. (2012) T-Cell Responses to the DBL $\alpha$ -Tag, a Short Semi-Conserved Region of the *Plasmodium falciparum* Membrane Erythrocyte Protein 1. PLoS ONE 7(1): e30095. doi:10.1371/journal.pone.0030095

**Editor:** Lars Hviid, University of Copenhagen, Denmark

**Received:** September 1, 2011; **Accepted:** December 9, 2011; **Published:** January 17, 2012

**Copyright:** © 2012 Gitau et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** The study was funded by Wellcome Trust Senior Fellowship Award to BCU (079082). KM and PCB are supported by Wellcome Trust Programme grants (077176, 084535) and JT is supported by a Wellcome Trust Strategic Award (084538). BCU and PCB are members of the EU FP6 and FP7 Networks of Excellence BioMalPar and EviMalaR. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

\* E-mail: [b.urban@liv.ac.uk](mailto:b.urban@liv.ac.uk)

$\ddagger$  Current address: Centre for Medical Parasitology, University of Copenhagen, Copenhagen, Denmark

## Introduction

Clinical immunity to malaria is achieved only after repeated infection with *Plasmodium falciparum* asexual bloodstage parasites. The *Plasmodium falciparum* erythrocyte membrane protein 1 (PfEMP1) mediates adhesion of mature forms of infected erythrocytes to endothelial cells and is central to pathogenesis and protective immune responses and also involved in immune evasion. (reviewed in [1]). Variants of PfEMP1 are encoded by approximately 60 var genes per haploid genome which undergo clonal antigenic variation and are extremely diverse both within and between parasite isolates [2]. In children living in malaria-endemic areas, repeated exposure to a wide range of different PfEMP1 expressed on parasite isolates results in the acquisition of a repertoire of antibodies against different variants that is associated with protection [3,4]. Furthermore, parasite isolates from children suffering from severe malaria or non-immune children were more likely to be recognised by sera from semi-immune children

suggesting that parasites from non-immune children and those with severe disease express antigenically restricted repertoires of PfEMP1 [5,6].

With the whole genome sequence of the laboratory isolate 3D7 it became evident that var genes encoding PfEMP1 can be grouped into three major types, Group A, B and C, suggesting their stratification into separate and potentially functionally distinct groups [7,8,9]. This basic concept has now been confirmed using sequence information from additional clinical and laboratory isolates [10]. We (Bull and colleagues) developed a sequence classification system based on a small semi-conserved area of the DBL $\alpha$ -domain of PfEMP1, the DBL $\alpha$ -tag, which allows classification of the entire var gene repertoire of clinical isolates. The amino acid sequence of amplified DBL $\alpha$ -tags can be grouped according to the number of cysteine (cys2 or cys4), the presence of sequence signatures at "Positions of Limited Variation" (PoLV) and through sharing of a limited number of sequence blocks within the hypervariable regions [11,12]. A schematic diagram of the

DBL $\alpha$ -tag is shown in the Figure S1. Using this system we showed differential expression of distinct subgroups of *var* genes in parasite isolates from children suffering from different syndromes of severe disease and in non-immune children [11,13]. Importantly, a subgroup of DBL $\alpha$ -tags that share blocks of diverse sequence overlap with group A *var* genes identified in the 3D7 genome [12] and are independently associated with young host age and severe malarial syndromes [3]. Although likely, whether serologically and genetically defined subgroups of PfEMP1 identify the same group of variants has not yet been determined.

T-cells play a critical role in protection, not only by providing help to B cells but also through the secretion of cytokines that activate macrophages and may have parasitocidal activity (reviewed in [14]). Rodent models of malaria have repeatedly shown that T-cells are essential for early control of parasitaemia and that elimination of parasites is dependent on both T and B-cells [15,16]. To date, only a few studies have analysed T-cell responses to PfEMP1 using either recombinant domains or peptides based on a very limited number of PfEMP1 molecules identified in laboratory isolates. These studies showed that a proportion of malaria-exposed children and adults responded to DBL $\alpha$  or exon2 by proliferation or cytokine secretion in an antigen-specific manner [17,18]. In both studies, proliferation of CD4+ T-cells in response to antigen was observed in a small number of unexposed European blood donors after seven days of culture. Under these conditions, priming of naive T-cells occurs in culture and it can not be determined whether the apparent CD4+ T-cell response of unexposed European blood donors was non-specific or an antigen-specific response of naive CD4+ T-cells [19]. By contrast, CD4+ T-cell responses to one CIDR $\alpha$  domain were observed in both malaria-exposed and unexposed individuals but only the response in malaria-exposed individuals was MHC class II restricted [17,20]. However, the use of recombinant domains representing PfEMP1 expressed on laboratory isolates as described in these studies may not capture T-cell responses specific for PfEMP1 expressed on parasite(s) circulating within a population and therefore miss a significant proportion of responses.

We were interested in whether there are differences in the type, magnitude or duration of CD4+ T-cell responses to PfEMP1 dominantly expressed at the time of acute disease in children suffering from severe malaria or infected with parasites expressing specific subgroups of PfEMP1. We expressed recombinant DBL $\alpha$ -tags from parasites isolated from clinical cases of malaria and determined CD4+ T-cell responses in those children who provided the clinical parasite isolate at the time of acute disease and during convalescence. Although the DBL $\alpha$ -tag is only a small part of the entire PfEMP1 molecule, it can serve as a model antigen to determine whether a specific subset of PfEMP1 is associated with a particular type of immune response.

Here we show that the magnitude, duration and type of DBL $\alpha$ -tag specific CD4+ T-cells responses were not different between children with severe or mild malarial symptoms. However, a small group of children infected with parasites expressing group A variants of PfEMP1 were more likely to induce antigen-specific IFN $\gamma$ -secreting CD4+ T-cells 16 weeks after follow-up indicating that the quality of T-cell responses to group A PfEMP1 differs from those induced against the more diverse non-GroupA PfEMP1. Surprisingly, we observed CD4+ T-cell responses to heterologous DBL $\alpha$ -tags (DBL $\alpha$ -tags from a clinical isolate a child was not infected with) in a sub-group of children tested despite the extreme sequence diversity of DBL $\alpha$ -tags. These data suggested that DBL $\alpha$ -tags contain a number of shared T-cell epitopes that may be recognised by cross-reactive CD4 T-cells.

## Results


### Phenotype of T-cell responses to PfEMP1 during acute disease and follow-up

The main aim of this study was to analyse the magnitude, duration and phenotype of CD4+ T-cell responses to the PfEMP1 variant expressed in parasites infecting individual children during acute disease and convalescence. PBMCs collected at presentation, 4 weeks and 16 weeks after the acute attack were stimulated with the DBL $\alpha$ -tag (EMBL accession nos. FR874861-FR874900, HE611335) representing the dominant PfEMP1 expressed on the parasites a given child was infected with at the time of acute disease and determined the proportion of DBL $\alpha$ -tag-specific CD4 T-cells producing either IFN $\gamma$ , IL10, IL2 or IL4 (Figure 1). We used the expression homologous DBL $\alpha$ -tags to identify CD4+ T-cell responses to the DBL $\alpha$ -tag derived from the clinical isolate a given child was infected with and heterologous DBL $\alpha$ -tags to identify CD4+ T-cell responses to DBL $\alpha$ -tags derived from clinical isolates of other children. Baseline haematological parameters are described in Table 1.

The majority of children (85%,  $n = 35$ ) induced CD4+ T-cells producing at least one of the tested cytokines IL2, IL4, IL10 and IFN $\gamma$  in response to activation with the homologous DBL $\alpha$ -tag during acute disease (Figure 2). Antigen-specific, cytokine producing CD4+ T-cells were detected in 58% ( $n = 24$ ) and 46% ( $n = 20$ ) of children 4 weeks and 16 weeks after the acute malaria attack, respectively. The proportion of antigen-specific, cytokine-producing CD4+ T-cells was comparable between different time points (Mann Whitney U test,  $p > 0.05$ ) although the number of children mounting a CD4+ T-cell response to the homologous DBL $\alpha$ -tag dropped significantly for CD4+IFN $\gamma$ +, CD4+IFN $\gamma$ +IL10+ and CD4+IL2+IL4+ T-cells over the 4 months following the acute episode (Pearson's  $\chi^2$  for trend,  $p < 0.05$ ; Table 2).

To describe changes in the type of CD4+ T-cell responses over time, we classified the CD4+ T-cell response to homologous DBL $\alpha$ -tags as a Th1 or Th2 response, CD4+IL2+ or CD4+IL10+ only response, a mixed response, no response. During acute disease, CD4+ T-cells from 11 children (27%) produced IFN $\gamma$  either alone or together with IL2 ( $n = 8$ ) indicative of a Th1 response. This profile was maintained in 3 children for 4 weeks and in only 1 child for 16 weeks whereas 6 children showed a Th1 profile during the follow-up period but not during acute disease (Figure 3A). We detected a Th2 profile (defined by secretion of IL4 either alone ( $n = 4$ ) or together with IL2 ( $n = 4$ ) and/or IL10 ( $n = 3$ ) in the absence of any CD4+IFN $\gamma$ + T-cells) in 4 children at acute disease but only one child maintained this profile over the entire period of 16 weeks. By contrast, 4 and 6 children gained CD4+IL4+ T-cell responses to the DBL $\alpha$ -tag at 4 weeks and 16 weeks, respectively, shifting from either a Th1 or mixed Th1/Th2 response at acute disease to a Th2 response in all but one child.

Antigen-specific, CD4+IL10+ T-cells were detected in 11 children either alone ( $n = 5$ ) or together with CD4+IFN $\gamma$ + T-cells ( $n = 5$ ). Four weeks after the acute episode, 6 children showed a sole CD4+IL10+ T-cell response, which was maintained in 3 children or switched from a CD4+IFN $\gamma$ + T-cell response either alone or mixed with CD4+IL10+ T-cells ( $n = 3$ ). We detected IFN $\gamma$ +IL10+ double producing CD4+ T-cells in seven children ( $n = 7$ ) but only in combination with either CD4+IL10+ or CD4+IFN $\gamma$ + T-cells. With the exception of one child, DBL $\alpha$ -tag specific CD4+IFN $\gamma$ +IL10+ T-cells were observed only at the time of acute disease. Although not significant, children who mounted DBL $\alpha$ -tag specific IFN $\gamma$ +IL10+ double producing CD4+ T-cells tended to have a higher parasitaemia at acute disease (median (25<sup>th</sup> and 75<sup>th</sup> interquartile) 255,680 parasites/ml blood (191,855–



**Figure 1. Gating Strategy used to identify T-cell responses.** PBMCs were gated to eliminate doublets and dead. CD3+lymphocytes were identified and gated for CD4+ and CD8+ T-cells. Within each T-cell subset, the proportion of IFN $\gamma$ , IL10, IL2, and IL4 producing cells were determined. IFN $\gamma$ +IL10+ and IL2+IL4+ double producers were determined by Boolean gating.  
doi:10.1371/journal.pone.0030095.g001

337,230) in children with IFN $\gamma$ +IL10+ double producing CD4+ T-cells and 148,240 parasites/ml blood (65,080–256,470) in those without; Mann Whitney  $p = 0.08$ ).

A small proportion of children displayed only CD4+IL2+ T-cells in response to the homologous DBL $\alpha$ -tag at acute disease ( $n = 6$ ) indicating that these T-cells were primed but not yet fully differentiated. IL2-producing CD4+ T-cell responses in the absence of any other cytokine-producing T-cells were also observed during follow-up but only in those children that either induced no CD4 T-cell response or a Th1 response during acute disease.

Together these results indicate that during acute malaria disease, CD4+ T-cell responses are dominated by a Th1 response sometimes in combination with IL10-producing CD4+ T-cells. By contrast, CD4+ T-cell responses 4 weeks and 16 weeks following the acute attack showed a more focused profile in those children that maintained a DBL $\alpha$ -tag specific response.

#### CD4+ T-cell responses to DBL $\alpha$ -tags in children with severe disease

We wondered whether children suffering from severe, life-threatening malaria differed with respect to the magnitude, duration or type of CD4+ T-cells responses specific for the homologous DBL $\alpha$ -tag from children suffering from moderate

malaria. Children with severe malaria were similar with respect to age, WBC and RBC count or Hb concentration to children with mild to moderate malaria. There were no differences in either the number of responders or the magnitude of cytokines produced by CD4+T-cells in response to homologous DBL $\alpha$ -tags between children suffering from severe or mild malaria either during acute disease or over the 4 months follow-up period. Likewise, there was no difference in the overall DBL $\alpha$ -tag specific T-cell profiles between children with severe disease and mild disease (Figure 3B).

#### T-cell responses to different subsets of DBL $\alpha$ -tag

Expression of *cys2* PfEMP1 and within these, the subgroup of group A-like PfEMP1, had been associated with severe disease and with low levels of existing immunity in young children [3]. We therefore determined the *var* gene classification of each recombinant DBL $\alpha$ -tag and compared CD4+ T-cell responses in children infected with parasites that expressed one dominant *cys2* PfEMP1 ( $n = 12$ ) or Group A-like PfEMP1 ( $n = 5$ , a subgroup of *cys2* sequences) with those that did not. As with severe disease, these children did not differ with respect to age, WBC, RBC counts or Hb. *Cys2* PfEMP1 were expressed on clinical isolates from three children suffering from severe malaria and nine children suffering from non-severe malaria. However, children infected with parasites that expressed one dominant *cys2* PfEMP1 showed a higher proportion of DBL $\alpha$ -tag specific CD4+IL10+ T-cells (median (25<sup>th</sup> and 75<sup>th</sup> percentile): 0.13 (0–0.28) for *cys2* DBL $\alpha$ -tags; 0 (0–0) for non-*cys2* DBL $\alpha$ -tags, Mann Whitney U test  $p = 0.015$ ) and a higher frequency of responders (6/9 responders for *cys2*; 5/27 responders for non *cys2*, Pearson's  $\chi^2 = 7.4$ ,  $df = 1$ ,  $p = 0.007$ ) during acute disease than children infected with parasites that expressed a dominant non-*cys2* PfEMP1 (Figure 4). Conversely, the phenotype of the CD4+ T-cell response in children infected with parasites that expressed a dominant *cys2* PfEMP1 at acute disease changed to a predominantly IFN $\gamma$ + response, both with respect to the number of children (3/6 responders for *cys2*; 2/18 responders for non *cys2*, Pearson's  $\chi^2 = 4.1$ ,  $df = 1$ ,  $p = 0.042$ ) and the magnitude of the response (median % of CD4+ T-cells (25<sup>th</sup> and 75<sup>th</sup> percentile): 0.03 (0–0.23) for *cys2* DBL $\alpha$ -tags; 0 (0–0) for non-*cys2* DBL $\alpha$ -tags, Mann Whitney U test  $p = 0.049$ ) 16 weeks after the acute episode. However, this trend was driven by a subgroup of children infected with parasites that expressed a dominant Group A-like PfEMP1 at acute disease (median % of CD4+ T-cells (25<sup>th</sup> and 75<sup>th</sup>

**Table 1. Baseline characteristics of study population.**

	Acute episode	4 weeks follow up	16 weeks follow up
age (months)	42.1 (25.6–57.4)	43.1 (26.6–58.4)	46.1 (29.6–61.4)
parasite density ( $10^3/\mu\text{l}$ )	190 (100–369)	0 <sup>a</sup>	0 <sup>b</sup>
WBC ( $10^6/\mu\text{l}$ )	8.4 (6.6–13.7)	8.2 (6.4–12.4)	9.1 (7–10.3)
RBC ( $10^9/\mu\text{l}$ )	4 (3.5–4.5)	4.8 (4.2–5.4)*	4.8 (4.2–5.1)*
Hb (g/dl)	9.2 (7.4–10.2)	10.6 (9.5–12)*	10.7 (9.4–11.9)*


Shown are median and in parenthesis 25<sup>th</sup> and 75<sup>th</sup> percentile.

\* $p \leq 0.05$  compared to acute disease, Wilcoxon signed rank test.

<sup>a</sup>one child had a parasite density of 514/ $\mu\text{l}$  blood 4 weeks after the acute event.

<sup>b</sup>one child had a parasite density of  $138 \times 10^3/\mu\text{l}$  blood 16 weeks after the acute event.

doi:10.1371/journal.pone.0030095.t001



**Figure 2. Percentage of CD4+ T-cells producing cytokines after stimulation with the homologous DBL $\alpha$ -tag.** PBMCs from children obtained during acute malaria, 4 weeks and 16 weeks after the acute event were stimulated with recombinant DBL $\alpha$ -tag representing the dominant expressed PfEMP1 a given child was infected with. CD4+ T-cells were stained for production of INF $\gamma$  and IL10 or IL2 or IL4 by intracellular cytokine staining. Shown are dot plots of percentages of CD4+ T-cells producing the indicated cytokine or cytokine combination during acute malaria or 4 and 16 weeks after the acute episode. Horizontal lines indicate the median.  
doi:10.1371/journal.pone.0030095.g002

percentile): 0.13 (0.05–0.3) for group A DBL $\alpha$ -tags; 0 (0–0) for non-group A DBL $\alpha$ -tags, Mann Whitney U test  $p=0.005$ ; 3/5 responders for group A; 2/20 responders for non group A, Pearson's  $\chi^2=8.5$ ,  $df=1$ ,  $p=0.003$ ). In summary, *cys2* DBL $\alpha$ -tags induced CD4+ T-cell responses that differed in phenotype and magnitude from CD4+ T-cell responses induced by non-*cys2* DBL $\alpha$ -tags suggesting that parasites that express a dominant *cys2* PfEMP1 induced a different CD4+ T-cell response compared to parasites that express a dominant *cys2* PfEMP1.


### T-cell responses to heterologous DBL $\alpha$ -tags

For some children we had sufficient number of PBMCs to evaluate CD4+ T-cell responses to another, heterologous DBL $\alpha$ -tag. The heterologous DBL $\alpha$ -tag tested was chosen randomly from a child tested at the same day. Overall, we analysed heterologous responses in 14 children during acute disease, in 16 children 4 weeks and in 10 children 16 weeks after the acute malaria episode. In total, CD4+ T-cell responses to 25 pairs of homologous and heterologous DBL $\alpha$ -tag were compared at one or more time points. The pairwise identity between homologous and heterologous DBL $\alpha$ -tags ranged from 33.6% to 56.7% (Table S1). For only four pairs, both the homologous and the heterologous DBL $\alpha$ -tag were classified as *cys2* PfEMP1. CD4+ T-cell responses to heterologous DBL $\alpha$ -tags varied between children, with some children inducing no CD4+ T-cell response, a similar CD4+ T-cell response profile or a different CD4+ T-cell response profile to the heterologous DBL $\alpha$ -tags compared to the homologous DBL $\alpha$ -tag (Figure 5A). There was no obvious tendency for either a similar or different type of CD4+ T-cell response to the heterologous DBL $\alpha$ -tag compared to the homologous DBL $\alpha$ -tag (Figure 5B). These data suggest that some children had been exposed to T-cell epitopes similar to those observed in the heterologous DBL $\alpha$ -tag before.

**Table 2. Magnitude of cytokine secretion by CD4+ T-cells after stimulation with DBL $\alpha$ -tags.**


		Acute episode	4 weeks follow up	16 weeks follow up
IFN $\gamma$ +	Median all	0.017 (0–0.18)	0 (0–0.02)	0 (0–0.025)
	N (%) responders*	15 (36.5%)	6 (14.6%)	5 (12.2%)
	Median responders	0.27 (0.08–0.42)	0.16 (0.09–0.54)	0.2 (0.06–0.34)
IL10+	Median all	0 (0–0.11)	0 (0–0.06)	0 (0–0.005)
	N (%) responders	11 (26.8%)	10 (24.3%)	5 (12.2%)
	Median responders	0.25 (0.14–0.42)	0.2 (0.08–0.61)	0.15 (0.08–0.33)
IFN $\gamma$ +IL10+	Median all	0 (0–0.007)	0 (0–0)	0 (0–0)
	N (%) responders*	7 (17%)	1 (2.4%)	1 (2.4%)
	Median responders	0.15 (0.08–0.61)	na	na
IL2+	Median all	0.06 (0–0.27)	0 (0–0.1)	0.008 (0–0.27)
	N (%) responders	15 (36.5%)	9 (22%)	9 (22%)
	Median responders	0.22 (0.14–0.5)	0.21 (0.1–0.69)	0.34 (0.07–0.55)
IL4+	Median all	0 (0–0.038)	0 (0–0.025)	0 (0–0.12)
	N (%) responders*	8 (19.5%)	5 (12.2%)	7 (17%)
	Median responders	0.18 (0.05–0.78)	0.13 (0.06–0.92)	0.19 (0.1–0.62)
IL2+IL4+	Median all	0.004 (0–0.008)	0 (0–0.008)	0.002 (0–0.011)
	N (%) responders	11 (26.8%)	3 (7.3%)	2 (4.8%)
	Median responders	0.1 (0.05–0.18)	0.06 (0.05–0.28)	0.04, 1.7

\* $p\leq 0.05$ , Pearson's  $\chi^2$  for reduction in the number of responders over time.  
doi:10.1371/journal.pone.0030095.t002




**Figure 3. Phenotype of CD4+ T-cells stimulated with the homologous DBL $\alpha$ -tag.** The phenotype of CD4+ T-cell in response to the homologous DBL $\alpha$ -tag was classified for each child and time point as follows: Th1, secretion of IFN $\gamma$  alone or together with IL2 and/or IFN $\gamma$ +IL10+; Th2, secretion of IL4 alone or in the presence of CD4+ T-cells secreting IL2 or IL10; IL2 alone: detection of only IL2-producing CD4+ T-cells; IL10 alone: detection of only IL10-producing CD4+ T-cells; mixed: detection of a mixed profile such as IFN $\gamma$ -producing CD4+ T-cells together with IL4- or IL10-producing CD4+ T-cells or IL2- and IL10-producing CD4+ T-cells; none: children who did not induce any antigen-specific CD4+ T-cells. Shown are pie charts of the proportion of all children (A) with a particular profile or (B) grouped by disease severity. doi:10.1371/journal.pone.0030095.g003

When we considered CD4+ T-cell responses to cys2 DBL $\alpha$ -tags and non-cys2 DBL $\alpha$ -tags for the homologous and heterologous DBL $\alpha$ -tags together, we observed a similar trend with respect to the differences in the type duration and magnitude of CD4+ T-cell responses. Overall, the magnitude of CD4+ T-cells responses and the proportion of responders was higher when PBMCs were stimulated with cys2 DBL $\alpha$ -tags or group A DBL $\alpha$ -tags compared to stimulation with non-cys2 DBL $\alpha$ -tags (Figure 6). Comparable to CD4+ T-cell responses to the homologous DBL $\alpha$ -tag, a higher percentage of CD4+IL10+ T-cells (median % of CD4+ T-cells (25<sup>th</sup> and 75<sup>th</sup> percentile): 0.13 (0–0.34) for cys2 DBL $\alpha$ -tags; 0 (0–0) for non-cys2 DBL $\alpha$ -tags, Mann Whitney U test  $p = 0.003$ ) and a higher proportion of responders (10/15 responders for cys2; 8/38 responders for non cys2, Pearson's  $\chi^2 = 9.9$ ,  $df = 1$ ,  $p = 0.002$ ) was detected during acute disease whereas CD4+IFN $\gamma$  positive T-cells dominated 16 weeks after the acute attack with respect to the number of responders (5/10 responders for cys2; 5/31 responders for non cys2, Pearson's  $\chi^2 = 4.7$ ,  $df = 1$ ,  $p = 0.03$ ). Likewise, the number of responders inducing CD4+IL10+ T-cells after stimulation with group A-like DBL $\alpha$ -tags was higher compared to other DBL $\alpha$ -tags during acute disease (5/7 responders for group A; 13/46 responders for non-group A, Pearson's  $\chi^2 = 5.05$ ,  $df = 1$ ,  $p = 0.025$ ) whereas induction of CD4+IFN $\gamma$ + T-cells in response to group A DBL $\alpha$ -tags dominated 16 weeks after the acute attack (4/6 responders for group A; 6/35 responders for non group-A, Pearson's  $\chi^2 = 6.8$ ,  $df = 1$ ,  $p = 0.009$ ). Together, these data indicate



**Figure 4. Cumulative percentage of cytokine producing CD4+ T-cells after stimulation with homologous DBL $\alpha$ -tag by PfEMP1 subgroup.** Shown are cumulative bar-graphs of CD4+ T-cells producing cytokines indicated for each child. Data are grouped by the PfEMP1 subclass of the homologous DBL $\alpha$ -tag at acute disease, week 4 and week 16 after the acute event. Children suffering from severe disease are indicated by a star. Note that the order of children is the same in each plot. doi:10.1371/journal.pone.0030095.g004



**Figure 5. CD4+ T-cells responses to homologous or heterologous DBL $\alpha$ -tags.** (A) Representative examples of CD4+ T-cell responses to homologous and heterologous DBL $\alpha$ -tags for 6 children. Pie charts indicate the IFN $\gamma$ + (red), IL10+ (blue) and IFN $\gamma$ +IL10+ double producing (green) CD4 T-cells as a proportion of the total cytokine producing CD4+ T-cells. White circles indicate the absence of a response. Within each cell, the sequence of circles indicates the proportion of CD4+ T-cells at acute disease (left), and 4 weeks (middle) and 16 weeks (right) after the acute episode. (B) Dot plots of the percentage of CD4+ T-cell responses to homologous (homo) and heterologous (hetero) DBL $\alpha$ -tags during acute malaria or 4 and 16 weeks after the acute attack. Horizontal lines indicate the median. doi:10.1371/journal.pone.0030095.g005

that cys2 PfEMP1, and particularly group A PfEMP1, may induce a different T-cell response than non-cys2 PfEMP1.

## Discussion


Here we report that DBL $\alpha$ -tag-specific T-cell responses are readily detected in children with acute malaria and can be identified 16 weeks after the acute episode in a proportion of children indicating that some of them may induce long-lasting effector T-cells. Importantly, the DBL $\alpha$ -tag used as antigen here was identified as the dominant PfEMP1 expressed on the clinical parasite isolate circulating in a given child and thus we measured T-cell responses to the individual PfEMP1 variant a child was actually exposed to.

Obviously, our approach has limitations. First, the entire PfEMP1 molecule will harbour many more T-cell epitopes and some of those lying outside the DBL $\alpha$ -tag may be more dominant. Second, we assumed that the T-cell phenotype(s) we detected using the DBL $\alpha$ -tag as an antigen will reflect the T-cell phenotype induced by the entire PfEMP1 molecule since activation of T-cell recognizing different epitopes of one PfEMP1 variant will occur in a similar environment with respect to co-stimulation and cytokines provided by antigen presenting cells. However, different parts of a given molecule may be processed at different rates and thus may affect epitope density or delays in T-cell priming by dendritic cells with consequences for T-cell function [21]. Bearing these assumptions in mind, we nevertheless made significant observations. First, we detected DBL $\alpha$ -tag specific IFN $\gamma$  and IL10+

double-producing CD4+ T-cells in a proportion of children. Second, CD4+ T-cell responses to cys2 DBL $\alpha$ -tag showed a distinct profile during acute disease but IFN $\gamma$  secretion 16 weeks after the acute event. Finally, we detected CD4+ T-cell responses to heterologous DBL $\alpha$ -tags despite considerable sequence diversity between DBL $\alpha$ -tags suggesting that at least some T-cell responses may be cross-reactive.

The majority of children responded to the homologous PfEMP1 by induction of CD4+IFN $\gamma$  and CD4+IL2+ producing T-cells, indicating a Th1 profile either alone or in combination with IL10-secreting CD4+ T-cells. In addition, we detected IFN $\gamma$  and IL10+ double-producing CD4+ T-cells, which have recently been described [22]. These T-cells reflect a normal differentiation process in Th1 responses providing a negative feedback loop in the presence of high antigen load or high concentrations of IL12 and are thought to regulate an otherwise dangerous inflammatory response. Most IL10-producing Th1 T-cells co-produce IFN $\gamma$  but sole production of IL10 after repeated stimulation has been reported [23,24,25,26,27]. Importantly, IL10-producing Th1 cells switch to IFN $\gamma$  production after a period of rest and some authors now suggest that at least some of the IL10-producing Tr1 cells may present late stages of Th1 [28]. In agreement with these studies, children with IL10-producing Th1 T-cells tended to have high parasitaemia during acute disease. IL10-producing Th1 T-cells had first been described over two decades ago but to our knowledge this is the first report of these cells in malaria [29].

When we analysed CD4+ T-cell responses by PfEMP1 subgroup, we observed that children infected with parasites



**Figure 6. CD4+ T-cells responses response to cys2 and non-cys2 DBL $\alpha$ -tags.** Dot plots of the percentage of CD4+ T-cell responses to cys2 and non-cys2 DBL $\alpha$ -tags whether or not they represent homologous or heterologous DBL $\alpha$ -tags during acute malaria or 4 and 16 weeks after the acute attack. Horizontal lines indicate the median.  
doi:10.1371/journal.pone.0030095.g006

expressing a cys2 PfEMP1 tended to mount IL10+ CD4+ T-cells during acute disease but displayed IFN $\gamma$ + CD4+ T-cell responses 16 weeks after the acute episode, the later was driven by group A-like DBL $\alpha$ -tags. When we considered all cys2 DBL $\alpha$ -tags tested independent of whether they reflected heterologous or homologous responses, we noted a similar trend to that observed with homologous cys2 DBL $\alpha$ -tag. The shift from a T-cell response dominated by IL10 at acute disease to one dominated by IFN $\gamma$  may be the result of a regulatory response to inflammation either due to normal differentiation of Th1 T-cells or due to induction of IL-10-producing regulatory T-cells [22,28]. During acute disease, high antigen-load and pro-inflammatory conditions may favour switching of Th1 cells to the production of IL10 [28]. Alternatively, antigen-specific Th1 T-cells may redistribute to the spleen during acute disease and re-enter the circulation only after the infection has been cleared [30]. To differentiate between these options, both IL10- and IFN $\gamma$ -producing DBL $\alpha$ -tag specific T-cells require further characterisation. Nevertheless, pending confirmation in a larger group of children, our results indicate that cys2 DBL $\alpha$ -tags and within this subset, group A-like DBL $\alpha$ -tags may induce a stable Th1 effector memory response that may contribute to the rapid acquisition of immunity to this particular subset of PfEMP1 variants. Although we can not exclude the possibility that antigen-specific T-cell responses detected 4 and 16 weeks after the acute episode were due to additional infections, it seems unlikely because all children were tested for presence of parasites by microscopy at the clinical examination during follow-up visits and they occurred mainly during the dry season when transmission is low. Stable populations of effector memory T-cells have been reported in humans in a rodent model of malaria [31,32]. Both cys2 and Group A PfEMP1 variants are frequently expressed in non-immune hosts such as young children but also in children suffering from severe malarial disease [3]. Thus, memory CD4+ T-cells specific for cys2 and Group A PfEMP1 may be induced during the first few infections and if they were expressed early in later infections but maybe not maintained due to negative immune-selection, adaptive immune responses against group A PfEMP1 variants might be boosted resulting in the maintenance of effector memory T-cells.

We were able to test CD4+ T-cell responses to heterologous DBL $\alpha$ -tags in a small subset of children. Overall, heterologous

responses varied both between antigens and between patients with no obvious trend towards reduced or increased antigen-specific CD4+ T-cell responses or the proportion of responders. CD4+ T-cell responses to heterologous DBL $\alpha$ -tags probably reflect previous or concurrent exposure of individual children to PfEMP1 variants with similar DBL $\alpha$ -domains either during the current or a previous infection. In all clinical isolates, more than one DBL $\alpha$ -tag sequence was detected probably representing expression of switched PfEMP1 or infection with additional parasite clones. Thus, all children in this study were exposed to more than one PfEMP1 variant over the course of acute infection and could have mounted an immune response to various DBL $\alpha$ -tag. However, over its entire length, different DBL $\alpha$ -tags and indeed DBL $\alpha$  domains show only limited identity and a large study on more than 14,000 DBL $\alpha$ -tag sequences in Kilifi, Kenya observed only a small number of identical sequences [13]. Therefore, chance exposure to the PfEMP1 represented by the heterologous DBL $\alpha$ -tag seems highly unlikely. By contrast, this observation suggests that the DBL $\alpha$ -tag region contains T-cell epitopes that are relatively conserved and shared between several DBL $\alpha$  domains. Indeed, a study by Sanni et al. suggested that targets of CD4+ T-cell responses to PfEMP1 are relatively conserved because they detected CD4+ T-cell responses to pools of peptides derived from semi-conserved blocks of 15 variants of PfEMP1 [18]. Our own preliminary analysis of predicted T-cell epitopes within the 41 DBL $\alpha$ -tags showed that predicted high affinity peptides often span semi-conserved areas. For instance the region between PoLV3 and PoLV4 shows distinct sequence signatures (also known as homology blocks) that differentiate between cys2 and cys4 DBL $\alpha$ -tags [10]. In our preliminary analysis we observed a higher proportion of peptides predicted to bind to MHC class II alleles in cys2 DBL $\alpha$ -tags than in cys4 DBL $\alpha$ -tags. Cross-recognition of peptides from a subset of antigenically restricted PfEMP1 within MHC class II supertypes would allow rapid activation of memory T-cells that can provide help to B-cells specific for the cognate antigen and may result in earlier and more efficient antibody responses. CD4+ T-cell epitopes may not be subjected to the same degree of immune selection as B-cell or CD8 T-cell epitopes because CD4+ T-cell effector function is often indirect through activation of other cell types, their frequency is generally lower than that of CD8+ T-cells and protection conferred by CD4+ T-

cells in other infections usually limits disease severity but does not induce sterile protection [33,34].

In summary, our data suggest that DBL $\alpha$ -tags from group A PfEMP1 may contain a limited set of T-cell epitopes that are shared between variants of group A DBL $\alpha$ -tags. If this were the case, subsequent encounter of group A PfEMP1 might activate cross-reactive CD4 T-cells that can provide rapid help to B cells. This hypothesis is testable using a combination of bioinformatic approaches and cellular assays. If it can be confirmed, the design of vaccines targeting PfEMP1 variants associated with severe, life threatening illness may become feasible.

## Methods

### Ethics Statement

The study was approved by the Kenyan National Ethics Review Committee (protocol no. 1131) and the Oxford Tropical Research Ethics Committee (protocol no. 30-06). Parents or Guardians of the children provided written informed consent.

### Study cohort

We analysed blood samples from 41 children with acute malaria attending Kilifi District Hospital between 2006 and 2009. After parents or guardians provided informed consent for participation of their children in the study, children donated a venous blood sample of 3 ml at presentation with acute malaria, and 4 weeks and 4 months after the acute episode. Children who were admitted with impaired consciousness (Blantyre Coma Score <5), severe anaemia (Hb <5 g/dl and at least 10,000 parasites/ $\mu$ l blood) or severe respiratory distress (deep breathing and chest recession) were classified as suffering from severe malaria (n = 9). During follow-up visits, children were clinically examined, tested for the presence of *Plasmodium* species by microscopy and received a differential blood count. Children who were acutely ill received medical treatment as required. Parents or Guardians of the children provided informed consent.

### Processing of blood samples

Heparinised blood was spun at 2000 rpm, plasma removed, aliquoted and stored at  $-80^{\circ}\text{C}$ . The remaining blood cells were diluted to a total volume of 5 ml in RPMI before separation of peripheral blood mononuclear cells (PBMC) with lymphoprep. PBMC were washed twice with RPMI, resuspended in 10% DMSO/FCS and stored in liquid nitrogen. The remaining red blood cells (RBCs) were washed twice in RPMI and RBCs and granulocytes were separated by plasmagel flotation. 100  $\mu$ l of packed RBCs were resuspended in 800  $\mu$ l Trizol and stored at  $-80^{\circ}\text{C}$  for extraction of RNA.

### Isolation and expression of dominant expressed DBL $\alpha$ tags

Dominant expressed PfEMP1 were identified in 41 clinical isolates by the method described in detail elsewhere [11] with the difference that we identified, classified and expressed the single transcript that was most abundant. In brief, RNA was extracted from RBCs, converted to cDNA and amplified using DBL $\alpha$ AF' (5'-GCACG(A/C)AGTTT(C\*/T)GC-3') and DBL $\alpha$ BR (5'-GCCATTC(G/C)TCGAACCA-3') primers targeting a semi-conserved region of the DBL $\alpha$ -domain. PCR-products were cloned into pCR2.1 cloning vector, plasmid extracted from 10–20 colonies and sequenced on an Applied Biosystems 3730 sequencer using Big Dye Terminator v.3.1 cycle sequencing. The dominant DBL $\alpha$ -tag sequence was identified (range 10%–85% of sequenced transcripts), amplified from a representative plasmid by PCR using

the DBL $\alpha$ AF' primer and a DBL $\alpha$ BR primer containing a 3'-stop codon (5'-TTAGCCCATTC(G/C)TCGAACCA-3'). PCR products were ligated into the pEXP5(NT) TOPO vector (Invitrogen) containing and N-terminal His-tag sequence, transformed into TOP10 cells and the insert confirmed by sequencing. The DBL $\alpha$ -tag-containing plasmids were transformed in BL21 DE3pLysS *E. coli* (Invitrogen). Individual colonies were grown to an OD<sub>600</sub> of 0.4, protein expression induced with 1 mM IPTG, and cells grown for another 4 hours. The cell pellet was harvested by centrifugation and lysed with Bugbuster NT (Novagen) in the presence of Benzonase (Novagen). All DBL $\alpha$ -tags formed inclusion bodies and were purified under denaturing condition on Probond Nickel-chelating resin (Invitrogen) according to manufacturers recommendation. Purified proteins were dialysed over 24 hours against 20 mM TrisHCl, 50 mM NaCl, 6 M Urea pH 4.5 with stepwise reduction of the Urea concentration. Final dialysis was against 20 mM TrisHCl, 50 mM NaCl pH 4.5 for 12 hours. Contaminating endotoxin was removed using Endotrap blue (Hyglos GmbH) according to manufacturers recommendation and the residual endotoxin concentration was determined using the Limulus amoebae-lysate assay (Cambrex Bioscience) according to manufacturers recommendation. The residual endotoxin concentrations ranged from 0.0004–0.003 U/ $\mu$ g protein. The final protein concentration was determined using the BCA assay (Pierce). Recombinant proteins were sterile-filtered with 0.2  $\mu$ m syringe filters and stored at  $-80^{\circ}\text{C}$ .

### Intracellular cytokine staining

PBMCs from acute and convalescent samples were thawed and  $0.5 \times 10^6$  cells seeded twice in triplicate into 96 well plates in medium (RPMI1698 supplemented with 5% pooled human AB serum, 5 mM glutamine, 10 mM HEPES, 50  $\mu$ M  $\beta$ -mercaptoethanol, 50  $\mu$ M kanamycin). Cells were rested overnight before activation with medium alone, 20  $\mu$ g/ml homologous recombinant DBL $\alpha$ -tag or anti-CD2/CD3/anti-CD28-coated MACS bead particles (Miltenyi Biotec) in the presence of 1  $\mu$ g/ml CD28 and CD49d for 2 hours. Cells were incubated for another 18 hours in the presence of brefeldinA. Cells were harvested and stained with ViViD Aqua (Invitrogen) before intracellular cytokine staining was performed as follows: Cells were fixed with Cytofix (Becton Dickinson) for 20 min at RT in the dark, washed twice with Cytoperm (Becton Dickinson) and subsequently stained in Cytoperm with CD3-ECD Beckman Coulter), CD4-PC7 (Beckman Coulter), CD8-APC H7 (Becton Dickinson) and either IFN $\gamma$ -FITC (R&D) and IL10-PE (Becton Dickinson) or IL2-FITC (R&D) and IL4-PE (Becton Dickinson), for 1 hour at  $4^{\circ}\text{C}$  in the dark. Cells were washed twice, resuspended in Sheath Fluid (Beckmann Coulter) and acquired on a Cyan Analyzer (Beckmann Coulter) within 24 hours. Live, CD3+CD4+ or CD3+CD8+ T-cells were identified and the proportion of IFN $\gamma$ +IL10-, IFN $\gamma$ -IL10+ and IFN $\gamma$ +IL10+ T-cells was determined using FlowJo Africa. Four samples with less than 1000 CD4+ T-cells (n = 2) or failed positive response (cytokine secretion in less than 1% of CD4+ T-cells stimulated with microbeads, n = 2) after staining for CD4+ T-cells secreting IL2, IL4 or both were excluded from further analysis. Values obtained from PBMCs incubated with medium alone were subtracted from values obtained after activation of PBMCs with DBL $\alpha$ -tags. When individual gates of cytokine-secreting CD4+ T-cells had less than 10 positive events or their percentage was below 0.03%, the response was recorded as zero.

### Statistical data analysis

All data were analysed using PAWStatistics version 18. Comparison of continuous variables between different groups was done using Mann-Whitney rank sign test and for comparison



of categorical variables the Pearson  $\chi^2$  or Fisher's exact test was used. P-values less than 0.05 were considered significant. For graphic representation of data PAWStatistics version 18.0 and Graphpad Prism version 5 was used.

## Supporting Information

**Figure S1 Schematic Overview of the DBL $\alpha$  domain organisation.** (A) Indicated is the relative position of homology blocks (HB) 1–5 common to all DBL domains. Universal primers amplifying the DBL $\alpha$ -tag target conserved sequences in HB3 and HB2 and are indicated by black arrows. The “Positions of Limited Variation” (PoLV) 1 to 4, which together with the number of cysteine's in each DBL $\alpha$ -tag form the basis of the DBL $\alpha$ -tag classification. (B) Sequence signature of cys2 and cys4 DBL $\alpha$ -tags. PoLV1-4 are indicated by blue brackets. (TIF)

## References

- Miller LH, Baruch DI, Marsh K, Doumbo OK (2002) The pathogenic basis of malaria. *Nature* 415: 673–679.
- Kyes S, Horrocks P, Newbold C (2001) Antigenic variation at the infected red cell surface in malaria. *Annu Rev Microbiol* 55: 673–707.
- Warimwe GM, Keane TM, Fegan G, Musyoki JN, Newton CR, et al. (2009) *Plasmodium falciparum* var gene expression is modified by host immunity. *Proc Natl Acad Sci U S A*.
- Bull PC, Lowe BS, Kortok M, Molyneux CS, Newbold CI, et al. (1998) Parasite antigens on the infected red cell surface are targets for naturally acquired immunity to malaria. *Nat Med* 4: 358–360.
- Bull PC, Lowe BS, Kortok M, Marsh K (1999) Antibody recognition of *Plasmodium falciparum* erythrocyte surface antigens in Kenya: evidence for rare and prevalent variants. *Infect Immun* 67: 733–739.
- Nielsen MA, Staalsoc T, Kurtzhals JA, Goka BQ, Dodoo D, et al. (2002) *Plasmodium falciparum* variant surface antigen expression varies between isolates causing severe and nonsevere malaria and is modified by acquired immunity. *J Immunol* 168: 3444–3450.
- Lavstsen T, Salanti A, Jensen A, Arnot D, Theander T (2003) Sub-grouping of *Plasmodium falciparum* 3D7 var genes based on sequence analysis of coding and non-coding regions. *Malaria Journal* 2: 27.
- Robinson BA, Welch TL, Smith JD (2003) Widespread functional specialization of *Plasmodium falciparum* erythrocyte membrane protein 1 family members to bind CD36 analysed across a parasite genome. *Mol Microbiol* 47: 1265–1278.
- Kraemer SM, Smith JD (2003) Evidence for the importance of genetic structuring to the structural and functional specialization of the *Plasmodium falciparum* var gene family. *Mol Microbiol* 50: 1527–1538.
- Rask TS, Hansen DA, Theander TG, Gorm Pedersen A, Lavstsen T (2010) *Plasmodium falciparum* erythrocyte membrane protein 1 diversity in seven genomes—divide and conquer. *PLoS Comput Biol* 6.
- Bull PC, Berriman M, Kyes S, Quail MA, Hall N, et al. (2005) *Plasmodium falciparum* variant surface antigen expression patterns during malaria. *PLoS Pathog* 1: e26.
- Bull PC, Buckee CO, Kyes S, Kortok MM, Thathy V, et al. (2008) *Plasmodium falciparum* antigenic variation. Mapping mosaic var gene sequences onto a network of shared, highly polymorphic sequence blocks. *Mol Microbiol* 68: 1519–1534.
- Warimwe GM, Keane TM, Fegan G, Musyoki JN, Newton CR, et al. (2009) *Plasmodium falciparum* var gene expression is modified by host immunity. *Proc Natl Acad Sci U S A* 106: 21801–21806.
- Langhorne J, Ndungu FM, Sponaas AM, Marsh K (2008) Immunity to malaria: more questions than answers. *Nat Immunol* 9: 725–732.
- Meding SJ, Langhorne J (1991) CD4+ T-cells and B cells are necessary for the transfer of protective immunity to *Plasmodium chabaudi chabaudi*. *Eur J Immunol* 21: 1433–1438.
- Stephens R, Albano FR, Quin S, Pascal BJ, Harrison V, et al. (2005) Malaria-specific transgenic CD4(+) T-cells protect immunodeficient mice from lethal infection and demonstrate requirement for a protective threshold of antibody production for parasite clearance. *Blood* 106: 1676–1684.
- Allsopp CE, Sanni LA, Reubsaet L, Ndungu F, Newbold C, et al. (2002) CD4 T-cell responses to a variant antigen of the malaria parasite *Plasmodium falciparum*, erythrocyte membrane protein-1, in individuals living in malaria-endemic areas. *J Infect Dis* 185: 812–819.
- Sanni LA, Allsopp CE, Reubsaet L, Sanni A, Newbold C, et al. (2002) Cellular responses to *Plasmodium falciparum* erythrocyte membrane protein-1: use of relatively conserved synthetic peptide pools to determine CD4 T-cell responses in malaria-exposed individuals in Benin, West Africa. *Malar J* 1: 7.
- Plebanski M, Saunders M, Burtles SS, Crowe S, Hooper DC (1992) Primary and secondary human in vitro T-cell responses to soluble antigens are mediated by subsets bearing different CD45 isoforms. *Immunology* 75: 86–91.
- Ndungu FM, Sanni L, Urban B, Stephens R, Newbold CI, et al. (2006) CD4 T-cells from malaria-nonexposed individuals respond to the CD36-Binding Domain of *Plasmodium falciparum* erythrocyte membrane protein-1 via an MHC class II-TCR-independent pathway. *J Immunol* 176: 5504–5512.
- Hensmann M, Li C, Moss C, Lindo V, Greer F, et al. (2004) Disulfide bonds in merozoite surface protein 1 of the malaria parasite impede efficient antigen processing and affect the in vivo antibody response. *Eur J Immunol* 34: 639–648.
- O'Garra A, Vieira P (2007) T(H)1 cells control themselves by producing interleukin-10. *Nat Rev Immunol* 7: 425–428.
- Jankovic D, Kullberg MC, Feng CG, Goldszmid RS, Collazo CM, et al. (2007) Conventional T-bet(+)Foxp3(−) Th1 cells are the major source of host-protective regulatory IL-10 during intracellular protozoan infection. *J Exp Med* 204: 273–283.
- Anderson CF, Oukka M, Kuchroo VJ, Sacks D (2007) CD4(+)CD25(−)Foxp3(−) Th1 cells are the source of IL-10-mediated immune suppression in chronic cutaneous leishmaniasis. *J Exp Med* 204: 285–297.
- Cardone J, Le Fric G, Vantourou P, Roberts A, Fuchs A, et al. (2010) Complement regulator CD46 temporally regulates cytokine production by conventional and unconventional T-cells. *Nat Immunol* 11: 862–871.
- Meiler F, Zumkehr J, Klunker S, Ruckert B, Akdis CA, et al. (2008) In vivo switch to IL-10-secreting T regulatory cells in high dose allergen exposure. *J Exp Med* 205: 2887–2898.
- Gabrysova L, Nicolson KS, Streeter HB, Verhagen J, Sabatos-Peyton CA, et al. (2009) Negative feedback control of the autoimmune response through antigen-induced differentiation of IL-10-secreting Th1 cells. *J Exp Med* 206: 1755–1767.
- Cope A, Le Fric G, Cardone J, Kemper C (2011) The Th1 life cycle: molecular control of IFN-gamma to IL-10 switching. *Trends Immunol* 32: 278–286.
- Del Prete G, De Carli M, Almerigogna F, Giudizi MG, Biagiotti R, et al. (1993) Human IL-10 is produced by both type 1 helper (Th1) and type 2 helper (Th2) T-cell clones and inhibits their antigen-specific proliferation and cytokine production. *J Immunol* 150: 353–360.
- Hviid L, Theander TG, Abu-Zeid YA, Abdulhadi NH, Jakobsen PH, et al. (1991) Loss of cellular immune reactivity during acute *Plasmodium falciparum* malaria. *FEMS Microbiol Immunol* 3: 219–227.
- Zhang HH, Song K, Rabin RL, Hill BJ, Perfetto SP, et al. (2010) CCR2 identifies a stable population of human effector memory CD4+ T-cells equipped for rapid recall response. *J Immunol* 185: 6646–6663.
- Stephens R, Langhorne J (2010) Effector memory Th1 CD4 T-cells are maintained in a mouse model of chronic malaria. *PLoS Pathog* 6: e1001208.
- Brown DM, Dilzer AM, Meents DL, Swain SL (2006) CD4 T-cell-mediated protection from lethal influenza: perforin and antibody-mediated mechanisms give a one-two punch. *J Immunol* 177: 2888–2898.
- MacLeod MK, Kappler JW, Marrack P (2010) Memory CD4 T-cells: generation, reactivation and re-assignment. *Immunology* 130: 10–15.

**Table S1 \*MUSCLE alignment, shown is the percentage and in parenthesis the number of identical amino acids. (DOC)**

## Acknowledgments

This study is published with the permission of the Director of KEMRI. We are particularly grateful to children and their guardians for participation in the study and to the clinical and field staff who are looking after severely ill children. We thank F. Menza and F. Kanyetta for sample collection, G. Warimwe, J. Musyoki and M. Opiyo for sample processing and M. Mackinnon for critical discussion.

## Author Contributions

Conceived and designed the experiments: BCU ENG. Performed the experiments: ENG JT LS EK HK. Analyzed the data: BCU ENG. Contributed reagents/materials/analysis tools: PCB KM. Wrote the paper: BCU ENG.