Parmar, D., Leone, T., Coast, E., Hukin, E., Murray, S. & Vwalika, B. (2014). Health system costs of safe abortion and post abortion care from unsafe induced abortions in Zambia. Paper presented at the Third International Conference of the African Health Economics and Policy Association (AfHEA), 11-03-2014-13-03-2014, Nairobi, Kenya.



City Research Online

Original citation: Parmar, D., Leone, T., Coast, E., Hukin, E., Murray, S. & Vwalika, B. (2014). Health system costs of safe abortion and post abortion care from unsafe induced abortions in Zambia. Paper presented at the Third International Conference of the African Health Economics and Policy Association (AfHEA), 11-03-2014-13-03-2014, Nairobi, Kenya.

Permanent City Research Online URL: http://openaccess.city.ac.uk/4132/

Copyright & reuse

City University London has developed City Research Online so that its users may access the research outputs of City University London's staff. Copyright © and Moral Rights for this paper are retained by the individual author(s) and/ or other copyright holders. All material in City Research Online is checked for eligibility for copyright before being made available in the live archive. URLs from City Research Online may be freely distributed and linked to from other web pages.

Versions of research

The version in City Research Online may differ from the final published version. Users are advised to check the Permanent City Research Online URL above for the status of the paper.

Enquiries

If you have any enquiries about any aspect of City Research Online, or if you wish to make contact with the author(s) of this paper, please email the team at publications@city.ac.uk.

Health System Costs of Safe Abortion and Post Abortion Care from Unsafe Induced Abortions in Zambia

Divya Parmar¹, Tiziana Leone¹, Ernestina Coast¹, Eleanor Hukin¹, Susan Fairley Murray², Bellington Vwalika³ ¹London School of Economics and Political Science, UK ²King's College London, UK ³University Teaching Hospital, Zambia

- Globally the number of unsafe abortions remain stubbornly high, especially in low-income countries.
- The global costs of treating complications from unsafe abortions are likely to be a significant burden on public health systems of these countries. However, there are little data and evidence that are of use to policymakers.
- In Zambia, despite induced abortion being legal since 1972, the proportion of maternal deaths attributable to unsafe abortion is estimated at 30% or more. The costs of safe and unsafe induced abortions to the Zambian health system have never been estimated.

AIM: To estimate the Zambia health system costs of induced abortions - PAC for unsafe induced abortions and safe induced abortions in health facilities

BOX 1. Definitions and terms

UNSAFE INDUCED ABORTION: "a procedure for terminating an unintended pregnancy, carried out either by persons lacking the necessary skills or in an environment that does not conform to minimal medical standards, or both" (WHO, 2012)1. Any abortions induced outside health facilities are regarded as unsafe abortions in this study.

SAFE INDUCED ABORTION (SA): Medical abortions (MA) for gestational age 5-9 weeks and manual vacuum aspiration (MVA) for gestation age 9-12 weeks conducted in health facilities are regarded as safe induced abortion for this study.

POST ABORTION CARE (PAC): Following an induced (safe or unsafe) or spontaneous abortion, women sometimes need PAC at health facilities, which ranges from treatment of abortion complication to counselling, and provision of contraception. In this study, PAC includes treatment for incomplete abortion, sepsis and shock. Time spend on counselling and contraception advice are included in personnel costs.

RESULTS FACILITY-LEVEL COSTS PER SERVICE IN ZAMBIA, 2013 **\$180** ■ Blood transfusion ■ Drugs and materials **\$160** ■ Lab tests & diagnostics ■ Personnel \$140 Hospitalisation \$120 **\$100** \$80 \$60 **\$40 \$20 \$0 MVA** MA Sepsis Incomplete Shock abortion Post abortion care (PAC) Safe induced abortions (SA)

| PAC POST- INDUCED UNSAFE ABORTION IN ZAMBIA, 2013 | |
|--|---------------------|
| | Costs (\$) |
| Facility-level | |
| Safe induced abortion (N=222) | |
| MA | 5,898 |
| MVA | 1,772 |
| Incomplete abortion | 856 |
| Total cost of SA | 8,525 |
| Average cost per SA | 38.29 |
| PAC post-induced unsafe abortion (N=2123) | |
| Incomplete abortion | 70,410 |
| Sepsis | 37,544 |
| Shock | 1,857 |
| Total cost for PAC post-induced unsafe abortion | 109,811 |
| Average cost per PAC post-induced unsafe abortion | 51.72 |
| National-level ^a | |
| Cost of SA (N=6,015 to 18,044) | 230,280 - 690,840 |
| Cost of PAC post-induced unsafe abortion (N= 45,471 to 11,368) | 588,005 – 2,351,966 |

Due to absence of nationally available data, these calculations are based on numbers obtained from other

require PAC for induced unsafe abortion were to receive an induced SA

studies and therefore presented as a range of costs as shown in Box 2. ^b Cost savings if all women who

FACILITY-LEVEL AND NATIONAL-LEVEL ANNUAL COSTS FOR SA AND

METHODS: Estimate the treatment costs and annual costs of providing SA and PAC services at University Teaching Hospital (UTH) in Lusaka and project these costs to generate indicative cost estimates for the Zambian public health system by using an adaptation of the WHO Mother-Baby Package Costing Spreadsheet.

DATA

- Key informant interviews (n=18) to collect information on treatment protocols and costs
- Medical case records review of women receiving SA and PAC (n=71) to validate the treatment protocols
- Review of facility aggregate records (logbooks) to estimate the number of women receiving treatment for abortions at UTH.

BOX 2. Number of safe and unsafe induced abortion in Zambia

Total number of induced abortions in Zambia (per year)

= Abortion Rate/1000 x Number of women 15-44 years

= 45,110 to 114,279

Total number of unsafe induced abortions in Zambia (per year)

= Unsafe abortion rate/1000 x Number of women 15-44 years = 27,066 to 108,264

Total number of unsafe induced abortions requiring PAC in Zambia (per year)

= Percentage of unsafe induced abortions requiring PAC x Total number of induced abortions = 11,368 to 45,471

Total number of safe induced abortions in Zambia

= Total number of induced abortions – Total number of unsafe induced abortions = 6,015 to 18,044

Value and sources of data

Unsafe Abortion Rate per 1000 women 15-44 years *: 15 to 38

Safe Abortion Rate per 1000 women 15-44 years *: 9 to 36

Percentage of unsafe induced abortions requiring PAC: 42% ²

Population of women 15-44 years: 3,007,336

*Abortion rates for Southern Africa are used as the lower values and abortion rates for Eastern Africa are used as higher values ³)

CONCLUSION

- The Zambian health system would save \$13.43 per case (i.e. \$152,774 to \$611,046 per year) if each woman treated for a complication of unsafe abortion had instead accessed these services for SA.
- The Zambian health system spends between \$588,005 and \$2.4 million per year on PAC due to unsafe induced abortions.
- Ultimately costs could be reduced further if unintended pregnancies were reduced through the uptake of family planning. Zambia has a high level of unmet need for contraception at 27% (DHS 2007).

REFERENCES

- 1. WHO 2012. Safe Abortion: Technical and Policy Guidance for Health Systems. Second ed. Geneva: World Health
- 2. DARROCH, J. & SINGH, S. 2011. Adding it up: The costs and benefits of investing in family planning and maternal and newborn health - Estimation methodology, New York, Guttmacher Institute.
- 3. SEDGH, G., SINGH, S., SHAH, I. H., ÅHMAN, E., HENSHAW, S. K. & BANKOLE, A. 2012. Induced abortion: incidence
- and trends worldwide from 1995 to 2008. The Lancet, 379, 625-632.



Cost savings ^b

This work was funded by the Economic & Social Research Council / DFID.



152,774 - 611,046