

Christensen, J.F., Gomila, A., Gaigg, S. B., Sivarajah, N. & Calvo-Merino, B. (2016). Dance Expertise modulates behavioural and psychophysiological responses to affective body movement. *Journal of Experimental Psychology*, doi: 10.1037/xhp0000176



**CITY UNIVERSITY
LONDON**

[City Research Online](#)

Original citation: Christensen, J.F., Gomila, A., Gaigg, S. B., Sivarajah, N. & Calvo-Merino, B. (2016). Dance Expertise modulates behavioural and psychophysiological responses to affective body movement. *Journal of Experimental Psychology*, doi: 10.1037/xhp0000176

Permanent City Research Online URL: <http://openaccess.city.ac.uk/13002/>

Copyright & reuse

City University London has developed City Research Online so that its users may access the research outputs of City University London's staff. Copyright © and Moral Rights for this paper are retained by the individual author(s) and/ or other copyright holders. All material in City Research Online is checked for eligibility for copyright before being made available in the live archive. URLs from City Research Online may be freely distributed and linked to from other web pages.

Versions of research

The version in City Research Online may differ from the final published version. Users are advised to check the Permanent City Research Online URL above for the status of the paper.

Enquiries

If you have any enquiries about any aspect of City Research Online, or if you wish to make contact with the author(s) of this paper, please email the team at publications@city.ac.uk.

**Dance Expertise modulates behavioural and psychophysiological responses to affective
body movement**

Julia F. Christensen*^{1,2,3}, Antoni Gomila², Sebastian B. Gaigg^{1,3}, Nithura Sivarajah¹, Beatriz Calvo-Merino^{1,4}

¹ Cognitive Neuroscience Research Unit, Department of Psychology, City University London, UK

² Department of Psychology, University of the Balearic Islands, Spain

³ Autism Research Group, Department of Psychology, City University London, UK

⁴ Department of Psychology, Universidad Complutense Madrid, Spain

Author note

Julia F. Christensen and Antoni Gomila, Human Evolution and Cognition (IFISC-CSIC) and Department of Psychology, University of the Balearic Islands, 07122 Palma, Spain; Sebastian Gaigg, Beatriz Calvo-Merino and Nithura Sivarajah, Department of Psychology, City University London, School of Arts and Social Science, St John Street, London EC1V 0HB, UK; Beatriz Calvo-Merino: Department of Psychology, Universidad Complutense Madrid, Campes de Somosaguas, Madrid, Spain.

Funded by a Newton International Fellowship of the British Academy (NF140935) FFI2010-20759; and AP2009-2889 by the Spanish Ministry of Education (JFC), RYC-2008-03090 and PSI2012-34558 (BCM). We are grateful to Danceworks London, Pineapple Studios, Rambert School of Ballet and Contemporary Dance, and the London School of Russian Ballet for letting us recruit participants on their premises. Special thank you to the dancers who participated in the study.

Correspondence: Julia F. Christensen, City University London; School of Social Sciences; Rhind Building; Northampton Square; London, EC1V 0HB, UK,

julia.christensen.1@city.ac.uk

Abstract

The present study shows how motor expertise increases sensitivity to affective body movement at the behavioural and physiological level. Nineteen affective movement experts (professional ballet dancers) and twenty-four controls watched 96 video clips of emotionally expressive body movements while they performed an affect rating task (subjective response) and their galvanic skin response was recorded (psychophysiological response). The movements in the clips were either sad or happy, and in half of the trials movements were played in the order in which they are learned (forward presentation), and in the other half, backwards (control condition). Results showed that motor expertise in affective body movement specifically modulated both behavioural and physiological sensitivity to others' affective body movement, and that this sensitivity is particularly strong when movements are shown in the way they are learnt (forward presentation). The evidence is discussed within current theories of proprioceptive arousal feedback and motor simulation accounts.

Keywords: affect; emotion; expertise; neuroaesthetics; galvanic response; motor simulation; empathy, dance.

Dance Expertise modulates behavioural and psychophysiological responses to affective body movement

1. Introduction

Cognitive neuroscience has begun to explore how expertise in the arts modulates behavioural, perceptual, and neurocognitive processes. Musicians process musical and auditory sounds more accurately than controls (Oechslin, Van De Ville, Lazeyras, Hauert, & James, 2013), they are more sensitive to musical dissonance (Dellacherie, Roy, Hugueville, Peretz, & Samson, 2010), and musical training results in changes in brain macro and microstructure, especially in regions implied in auditory processing and motor control, such as the temporal and frontal lobes (Bangert et al., 2006; Bengtsson et al., 2005; Gaser & Schlaug, 2003; Habib & Besson, 2009; Haslinger et al., 2005; Pantev et al., 1998; Schlaug, 2006). Dance expertise enhances perceptual sensitivity to familiar movements (Calvo-Merino, Ehrenberg, Leung, & Haggard, 2010), and modulates neural responses to familiar actions in the Action Observation Network (bilateral premotor and parietal cortices) (Calvo-Merino, Glaser, Grèzes, Passingham, & Haggard, 2005; Calvo-Merino, Grèzes, Glaser, Passingham, & Haggard, 2006; Cross, Hamilton, & Grafton, 2006; Fink, Graif, & Neubauer, 2009; Jang & Pollick, 2011; Orgs, Dombrowski, Heil, & Jansen-Osmann, 2008). Long-term dance training results in changes in brain structure, in particular, in premotor and sensorimotor regions (Hänggi, Koeneke, Bezzola, & Jäncke, 2010).

In contrast to the wealth of evidence describing expertise effects in the domains of action perception (see Bläsing et al., 2012 for a review of dance expertise effects in neurocognition), very little is known about how movement expertise modulates the processing of affective information in movement. Recent studies have shown that expert artists (an example of experts in emotional expression) have enhanced affective responses as compared to controls. For example, actors are more empathic than non-actors (Goldstein, 2009; Goldstein & Bloom, 2011; Goldstein & Winner, 2012), musicians are better at

recognizing vocal expressions of emotions than non-musicians (Lima & Castro, 2011), ballet dance ability is associated with trait emotional intelligence (Petrides, Niven, & Mouskounti, 2006), and participants with dance experience show a modulation of their aesthetic response to familiar movements (Kirsch, Drommelschmidt, Cross, 2013; Kirsch, Dawson, Cross, 2015; see Christensen and Calvo-Merino, 2013 for a review on dance expertise and aesthetic perception). This suggests that expertise in the arts facilitates the processing of emotional information. To what extent this influence operates at the level of perceptual processes (indexed by an ability to discriminate between expressions of emotion), or is deeper rooted in ‘hot affective’ processes (evidenced by changes in psychophysiological arousal), however, remains unclear.

Exactly what constitutes an emotion and what the role is of physiological arousal in the perception of emotions in others and the subjective experience of emotions in oneself has been the focus of debate for more than a century. James (1894) famously contended that the conscious experience of feeling an emotion is a consequence of physiological arousal responses such as changes in heart rate, breathing rate, muscle tension and galvanic skin responses. Although this view initially attracted criticism (Cannon, 1927), accumulating evidence lends support to James’ view and many contemporary theories of emotion continue to ascribe a central role to arousal in the elicitation of emotional experiences (Damasio, 1999; Lang, Bradley & Cuthbert, 1999; Scherer, 2009a,b; see also Laird & Lacasse, 2014). A robust finding in this context is that subjectively reported feelings are associated with particular changes in heart-rate, skin conductance and other physiological parameters (e.g., Lang et al., 1999). Interestingly, this association is often only moderate in general population (Mauss, Levenson, McCarter, Wilhelm, & Gross, 2005), but increased in expert dancers (Sze, Gyurak, Yuan, & Levenson, 2010). This suggests that expertise in the bodily expression of emotion can augment the extent to which arousal influences the subjective experience of feelings and there are reasons to believe that this could enhance sensitivity to the emotion expressed in the

movements of others. Specifically, theories of empathy suggest that emotional expressions directed toward us induce a form of embodied mimicry whereby our physiology instantiates the arousal and brain states that are suggested by the emotional expression of the other (e.g., Decety & Jackson, 2006). These states in turn give rise to subjective feelings in ourselves that serve as cues to allow us to perceive the emotion expressed by the other. Daily training in expressing affect through movement should enhance this embodied mimicry because of the repeated coupling between proprioceptive feedback from the dancers' own body and the exteroceptive sensory feedback due to self-observation and observation of colleagues in a dance studio mirror. Therefore, we expect dance expertise to enhance sensitivity to emotional body movements because of an enhanced embodiment of congruent arousal on the one hand, and greater influence of this arousal on subjective feelings on the other.

Based on the above arguments, the current study tested the hypothesis that expert dancers compared to non-dancers would be more accurate at discriminating the emotions expressed in dance at the level of subjective valence ratings, and that they will also be more responsive at the psychophysiological level to the emotions on display. Moreover, to establish whether these predicted effects are mediated by general expertise in affective body movement or more specific expertise with particular forms of movement, the responses in the two groups will be compared on displays of movements as they have been learnt (forward presentation), vs. movement displays that are less familiar (backwards presentation). Importantly, kinematic properties of the movements in these forward and backward presentation conditions (e.g., speed, degree of displacement, etc.) are matched. Therefore any difference in emotional responsiveness to forward as opposed to backward displays, particularly if observed only in expert dancers, would support the view that expertise with the specific type of movements the dancers have learned, rather than movement more generally, modulates affective processing. Finally, we explore whether expertise modulates the coupling between perceptual and psychophysiological emotional responses by examining correlations

between subjective ratings and physiological arousal. Based on the observations of Sze and colleagues (2010) noted above, the prediction here is that the subjective ratings of expert dancers will more closely reflect their physiological arousal than is the case for non-experts.

2. Method

2.1. Participants

Twenty-four female undergraduate students with no formal dance experience (age Controls: $M = 20.86$; $SD = 2.77$; range: 18-32 years) participated in exchange for course credits. Twenty female ballet dancers (in professional training or working professionally with Ballet as their main dance style) (age Dancers: $M = 24.85$; $SD = 4.22$; range: 20-36 years) participated in exchange for a small time reimbursement (£8/h). Further details about the dancers are provided in Table 1. One participant in the dance group felt very uncomfortable during the experimental task and was not included in the data analysis. Thus, 19 Ballet dancers were included in the analyses presented below.

Table 1

Participant characteristics. Shown are mean and (SD). “Other dance styles” include Step Dance, Jazz Dance, Jazz Ballet, Burlesque, Lyrical and Commercial Dance

GROUP	Age	Age range	DANCE STYLE					
			Ballet		Contemporary		Other Dance Styles	
			Years of experience	Hours training/ week	Years of experience	Hours training/ week	Years of experience	Hours training/ week
Dancers	24.85 (4.22)	20-36	17.90 (5.59)	20.50 (12.93)	9.46 (4.05)	6.54 (6.41)	3 (10.75)	3.67 (4.04)
Controls	20.86 (2.77)	18-32	0	0	0	0	0	0

2.2. Materials

Forty-eight ballet dance video clips were selected from an affective body movement library of ballet movements (Christensen, Nadal, Cela-Conde, & Gomila, 2014a). These movement stimuli are 5-6 seconds long, and show an extract of a solo dancer in a genuine live performance, in black and white, without soundtrack and with the dancer's face blurred. In the 48 videos selected for the current study there were a mean of 4.35 (SD =0.36) full academic ballet movements as established by the respective ballet syllabi (Vaganova method and Royal Academy of Dance). These ballet syllabi have a limited number of movements and the 48 clips contained a unique combination of these. See the supplementary material for sample video clips. Sample clip S1 is from Sleeping Beauty and sample clip S2 from Swan Lake. Table S1 contains information regarding the stimuli selection.

For this study we required stimuli with strong emotional expression. [However, ballet dancers do not always execute their movements in emotionally expressive manners, for example for training purposes or for abstract ballets without narrative or emotional content.](#) Therefore, importantly, in the stimulus library from which the stimuli were selected, each video had been coded in terms of its valence and arousal. We used these scores to select 24 movements depicting happiness and 24 movements depicting sadness, while ensuring that each category of clips (happy vs. sad) had, respectively, 12 of high arousal and 12 of low arousal. Paired t-tests confirmed a significant difference between happy and sad videos in valence ratings (Happy: $M = 4.75$; $SD = .84$; Sad: $M = 4.08$; $SD = .74$; $t(23) = -2.397$; $p = .025$) but no significant difference between the two video categories in arousal ratings (Happy: $M = 4.54$; $SD = 1.45$; Sad: $M = 3.75$; $SD = 1.31$; $t(23) = 1.588$; $p = .126$). Since the expression of happiness or sadness in a ballet movement is primarily dependant on the quality of the movement (i.e., how it is performed) rather than on any particular step, it was also possible to ensure that the happy and sad clips did not differ with respect to the number of pirouettes ($t(23) = 1.56$; $p = .127$), relevés ($t(23) = 0.00$; $p = 1.00$), large movements ($t(23) = .57$; $p = .57$) and high frequency movements ($t(23) = 1.17$; $p = .25$) comprising them.

To ensure that possible differences in the affective responses between happy and sad movements were not due to other differences in the stimuli, such as speed or amount of movement, we created a set of control stimuli. For this condition, the same 48 stimuli were played backwards (transformation was done by means of Adobe Premiere Elements 7.5), thereby ensuring that the speed and amount of movement were identical for both forward and backward stimuli. This resulted in a total of 96 stimuli (48 forwards, and 48 backwards with half of the stimuli being Happy movements and half Sad movements in each condition).

2.3. Procedure

Stimuli were randomly presented using the stimulus presentation programme E-prime (version E-Studio, v. 2.0.8.90; www.pstnet.com). Stimuli were displayed on a black background with each dancer occupying approximately 5.5 cm on the screen (head to heel). Viewing distance was ~40cm. A fixation cross was presented before (1500ms) and after (1000ms) each video clip, which lasted for 5-6 seconds ($M = 5.02$; $SD = .41$) and was faded in and out to minimize surprise. Participants performed a subjective affect rating task (self-paced) after each video clip in which they were asked to rate how sad or happy the movements made them feel. This procedure follows that used in the norming study from which the stimuli were selected (Christensen et al., 2014a) and also the procedures commonly used in studies that assess emotional responses at the level of subjective experience and psychophysiological arousal (e.g., Lang, et al., 1999).

Responses were collected using a continuous visual analogical scale (VAS) presented at the bottom of the screen ranging from 0 (very sad) to 100 (very happy); 50 was neutral. The labels “Sad” (left) and “Happy” (right) displayed on either side of the VAS, while the indication “Emotion?” was displayed in the centre of the screen. The cursor of the mouse appeared always in the centre of the screen to avoid response tendencies. After the mouse

click within the scale, the next trial was launched. Participants had a break after half the trials. Average experiment duration was 45 minutes. See figure 1 for the trial structure.




Figure 1. Trial structure. After a fixation cross (1500ms), the dance video was faded in (~6 seconds), faded out, and followed by a fixation cross (1000ms). Then the Visual Analogue Scale (ranging from Sad, 0; to Happy, 100) appeared below the word “Emotion?” written in the centre. The cursor of the mouse appeared always in the top centre of the screen rather than anywhere on the VAS scale to avoid the cursor position biasing the participants’ responses toward any of the extremes. Reproduced with permission (Christensen et al., 2014b).

Throughout the experiment, skin conductance was recorded at a frequency of 1kHz with an ADInstruments PowerLab System (ML845) including a GSR (ML 116) and Bioelectrical signal amplifier (ML408 with MLA2540 and MLA2505 5-lead shielded Bio Amp cables). Stainless steel bipolar GSR electrodes (MLT116F) were attached to the medial phalanges of the index and ring fingers of the participant’s non-dominant hand using fitted velcro tapes. A second computer running LabChart 7 (v.7.3.1. 1994-2004; www.adiinstruments.com) was connected with a parallel-to-serial port interface to the computer running the stimulus presentation programme. A trigger was sent from E-prime to the trace of the GSR online recording in LabChart marking each stimulus event.

2.4. Analyses

Repeated measures (RM) Analysis of variance (ANOVA) were conducted on both participants' VAS ratings and GSR data in order to examine the effects of Stimulus Presentation (Forward vs. Backward) and Dance Emotion (Happy vs. Sad). Given our specific predictions, any interactions indicated by the ANOVAs were followed up using planned comparisons (t-tests), without applying additional corrections for multiple comparisons (Rothman, 1990; Saville, 1990; McDonald, 2009). As effect sizes we report partial eta (η_p^2), where .01 is considered a small effect size, .06 a medium effect and .14 a large effect (Cohen, 1988).

Following standard procedures (e.g., Bradley, Codispoti, Cuthbert & Lang, 2001), the GSR data were quantified by first subtracting the maximum value within the six seconds of the video stimulus duration from the GSR value at the onset of the stimulus and then applying a log transformation ($\log[\text{GSR}+1]$) to normalize the distribution of the data (Bradley, Codispoti, Cuthbert, & Lang, 2001). Furthermore, all participants with 1.5 SD above or below the mean of their respective group were discarded. This left 18 Controls and 17 Dancers for the analyses of GSR data.

The final analysis was correlational and served to examine the extent to which subjective affective ratings reflected objectively measured arousal responses. For this purpose the VAS ratings and GSR responses were averaged for each stimulus across the participants in the two groups. These averages were then correlated with one another separately for forward and backward stimuli to quantify the association between the VAS ratings and GSR responses in the two groups (please refer to figure 4 for further details).

3. Results

3.1 Analysis of subjective affective ratings (VAS)

The analysis of participant's subjective VAS ratings confirmed that happy videos were rated as happier than sad videos and that forwards presentation resulted in more positive ratings than backwards ratings. The data also confirmed the prediction that dancers would be better able to differentiate happy and sad dance movements, particularly in the canonical forward presentation of the video clips. These conclusions were supported by a 2 x 2 x 2 RM ANOVA of the VAS ratings with the within group factors of Stimulus Presentation (Forwards, Backwards) and Dance Emotion (Happy, Sad), and the between group factor of Group (Controls, Dancers). This demonstrated a significant main effect of Dance Emotion (Happy: $M = 60.539$; $SE = 1.159$; Sad: $M = 39.679$; $SE = 1.159$; $F(1,41) = 175.794$, $p < .001$, $\eta_p^2 = .811$), confirming that videos in the Happy category received higher affective ratings than those in the Sad category. A significant main effect of Stimulus Presentation (Backwards: $m = 48.66$; $SE = .862$; Forwards: $m = 51.556$; $SE = .930$; $F(1,41) = 8.279$, $p = .006$, $\eta_p^2 = .168$), further showed that movements presented in their familiar forward direction were rated overall as of more positive valence than when played backwards. Although the main effect of group was not significant ($F(1,41) = .511$, $p = .479$, $\eta_p^2 = .012$), we observed an interaction between Group and Dance Emotion ($F(1,41) = 34.428$, $p < .001$, $\eta_p^2 = .456$), which is explained by a more pronounced differentiation in VAS ratings between the two displayed emotions (Happy, Sad) in the group of expert dancers (Happy: $M = 64.624$; $SE = 1.490$; Sad: $M = 34.533$; $SE = 1.732$) than in the control group (Happy: $M = 56.453$; $SE = 1.326$; Sad: $M = 44.825$; $SE = 1.541$), as set out in figure 2. The interaction between Group and Stimulus Presentation was also significant ($F(1,41) = 4.442$, $p = .042$, $\eta_p^2 = .098$) reflecting more pronounced differences in VAS ratings between forwards and backwards stimuli in the group of Dancers (Backwards: $M = 47.071$; $SE = 1.288$ vs. Forwards: $M = 52.086$; $SE = 1.389$) than in the Control group (Backwards: $M = 50.252$; $SE = 1.146$; Forwards: $M = 51.026$; $SE = 1.236$). Of most interest, however, is the fact that these 2-way

interactions were further characterised by a marginally significant three-way interaction between Stimulus, Dance Emotion and Group ($F(1,41) = 4.047$, $p = .051$, $\eta_p^2 = .090$).

To understand the source of the three-way interaction we performed two separate 2x2 ANOVAs (Stimulus Presentation x Dance Emotion); one for the Dancer group and one for the Control group. The ANOVA in the Dancer group showed a significant main effect of Stimulus Presentation ($F(1,18) = 6.05$, $p = .024$, $\eta_p^2 = .251$); Backward stimuli received lower ratings ($M = 47.07$, $SE = 1.48$) than the Forward stimuli ($M = 52.09$, $SE = 1.61$). There was also a main effect of Dance Emotion ($F(1,18) = 98.58$; $p < .001$, $\eta_p^2 = .846$); Happy stimuli were rated as more happy ($M = 64.62$; $SE = 1.60$) than Sad stimuli ($M = 24.53$; $SE = 2.22$). Importantly, there was an interaction of Stimulus Presentation and Dance Emotion ($F(1,18) = 7.87$; $p = .021$, $\eta_p^2 = .304$). We followed up this interaction with paired t-tests. For Happy movements, we found significant differences between the Forward ($M = 68.98$; $SE = 1.32$) and Backwards ($M = 60.27$; $SE = 2.08$) conditions ($t(18) = -5.471$; $p < .001$). No such difference was significant for Sad movements (Forward: $M = 35.19$; $SE = 3.22$; Backwards: $M = 33.87$; $SE = 2.02$), ($t(18) = -.424$; $p = .609$). The same ANOVA in the Control group only showed a significant main effect of Emotion, with Happy movements rated as more happy ($M = 56.45$; $SE = 1.28$) than Sad movements ($M = 44.83$; $SE = 1.09$; $F(1,23) = 64.15$; $p < .001$, $\eta_p^2 = .736$). Neither the main effect of Stimulus Presentation ($F(1,23) = 0.98$; $p = .332$, $\eta_p^2 = .041$) nor the interaction between Dance Emotion and Stimulus Presentation ($F(1,23) = 0.38$; $p = .544$, $\eta_p^2 = .016$) were significant in this group. These results suggest that Experts' subjective affect ratings are sensitive to the Stimulus Presentation (Forward or Backwards) when the movements express happiness, while no such effect of affective sensitivity was observed in the Control group; see figure 2.




Figure 2. Subjective affect ratings means in the 2x2x2 design. Bars reflect the different conditions: Stimulus Presentation (Forward, Backwards), Dance Emotion (Happy, Sad) and Group (Dancers, Controls). (** $p < .001$). Error bars reflect S.E.M. (VAS = Visual Analogical Scale for the affective ratings).

3.2. Analysis of physiological data (GSR)

To analyse participants' physiological responses two approaches were taken. First the data were analysed according to the same 'normative' (for ease of reference) principles as the VAS rating data above, comparing responses to happy and sad dance movements as defined by the norming study from which these stimuli were selected. In addition, however, it is also possible to examine physiological data on a 'subjective' (for ease of reference) subject-by-subject basis whereby each stimulus is classified as happy or sad according to each participant's own subjective rating (e.g., Cela-Conde et al., 2004; Salimpoor et al., 2009). The stimuli are sorted according to each participant's VAS ratings from lowest to highest (0 to 100) and the top half are classified as subjectively 'Happy' stimuli with the bottom half

considered to be the 'Sad' stimuli . This second procedure is particularly useful for examining physiological responses in relation to personally experienced feelings that may deviate from group averaged responses that make up normed stimulus libraries such as the one from which the materials for the current study were selected. Together, the analyses of normative and subjective GSR responses demonstrated overall higher GSR in Controls than Dancers but Dancers had a more differentiated response to happy and sad videos, which was particularly evident in an analysis of subjective GSR data that revealed significant differences between happy and sad videos only for Dancers and only when videos were played in their forward direction. These observations parallel the results of the subjective VAS data and were supported by the following analyses.

For the first analysis of normative GSR responses a 2 (Stimulus Presentation; Forward vs. Backwards) x 2 (Normative Dance Emotion; Happy vs. Sad) x 2 (Group; Controls vs. Dancers) mixed ANOVA was carried out, equivalent to the VAS analysis above. This revealed a main effect of Group ($F(1,33) = 19.90, p < .001, \eta_p^2 = .376$) with overall higher GSR responses in Controls ($m = .144, SE = .010$) than Dancers ($m = .08, SE = .010$) as well as a significant interaction between Dance Emotion and Group ($F(1,33) = 8.065, p = .008, \eta_p^2 = .196$). Follow-up comparisons showed that only for Dancers GSR responses differed significantly between the normatively Happy ($m = .086, SE = .009$) and Sad movements ($m = .077, SE = .008; F(1,16) = 5.251, p = .036, \eta_p^2 = .247$) whereas for controls this effect was weaker and fell short of conventional significance (Happy: $m = .151, SE = .012$; Sad: $m = .136, SE = .012; F(1,17) = 4.170, p = .057, \eta_p^2 = .197$). In this first analysis no other main effects or interactions were significant ($\eta_p^2 < .035$).

The second analysis of subjective GSR responses followed the same 2 (Stimulus Presentation; Forward vs. Backwards) x 2 (Subjective Dance Emotion; Happy vs. Sad) x 2 (Group; Controls vs. Dancers) as above and revealed a main effect of Emotion with more pronounced GSR for Happy movements ($M = .116, SE = .007$) than for Sad movements ($M =$

.108; SE = .007, $F(1,33) = 4.754$, $p = .036$, $\eta_p^2 = .126$). The main effect of Group was again also significant as in the first analysis above with Controls showing increased GSR ($M = .143$; SE = .010) compared to Dancers ($M = .081$, SE = .010; $F(1,33) = 19.680$, $p < .001$, $\eta_p^2 = .374$). No other main effects were significant but instead of the two-way interaction revealed by the normative analysis above, the current subjective analysis yielded a marginally significant three-way interaction between Stimulus Presentation, Dance Emotion and Group ($F(1,33) = 3.910$, $p = .056$, $\eta_p^2 = .106$) that parallels the 3-way interaction in the VAS analysis. Given this trend, our a priori hypothesis regarding group differences, and the significant between-group factor, we performed two additional RM ANOVAs separately for each group. A 2x2 RM ANOVA for the Dancer group showed a significant interaction between Stimulus Presentation and Subjective Dance Emotion ($F(1,33) = 5.634$, $p = .030$, $\eta_p^2 = .260$). Breaking down this interaction further with paired t-tests showed a significant difference in GSR in the two Dance Emotion categories as a function of Stimulus Presentation. Dancers' GSR was higher for Happy movements ($M = .0873$, SE = .009) than for Sad movements ($M = .0724$, SE = .008, $t(17) = -2.728$, $p = .015$) only in the Forward Condition. Conversely, there was no difference between the two Subjective Emotions for Backwards movements ($t(16) = .025$, $p = .980$). These data in the expert group show that physiological responses are sensitive to affective movement only when the movement is displayed in its familiar presentation (forward). By contrast a similar RM ANOVA for the Control group did not show any main effects (Dance Emotion: $F(1,17) = 2.204$; $p = .156$, $\eta_p^2 = .115$; Stimulus Presentation $F(1,17) = 0.015$; $p = .905$, $\eta_p^2 = .001$) or interaction between Dance Emotion and Stimulus Presentation ($F(1,17) = .716$; $p = .409$, $\eta_p^2 = .040$); their GSR during observation of self-rated Happy and Sad movements was similar, irrespective of Stimulus Presentation (figure 3).




Figure 3. Physiological responses measured through Galvanic Skin Response (GSR, Log transformed μs) of Dancers and Controls during observation of dance videos rated as Happy and Sad. Data are presented for the two stimuli presentations conditions (Forward, Backwards). (* $p < .05$). Error bars reflect S.E.M.

3.3. Correlations between subjective affective experience and physiological responses

The above analyses show that expertise in ballet dance does not only enhance the ability to discriminate the valence expressed in dance movements but it also sensitises a person's emotional responsiveness to emotional dance at the physiological level. The fact that this is observed only in relation to familiar forward presentations of relevant movements lends support to the idea that relevant expertise rather than spurious stimulus characteristics are mediating these effects. In a final analysis we examined whether dance expertise may also modulate the extent to which subjective experiential and psychophysiological facets of emotional responsiveness are coupled, which is thought to provide another indicator of affective sensitivity (Sze, et al., 2010). Thus, as explained in the analysis section, the

correlation between average VAS ratings and GSR responses were examined within each group for the forward and backward stimuli separately. The relevant scatter plots are illustrated in figure 4 and suggest that expert dancers were more sensitive affectively; their affective ratings correlated significantly with their physiological response ($p = .003$, $r = .419$) during observation of stimuli in their familiar presentation (forwards), while no such correlation was found when dancers observed the stimuli backwards ($p = .832$, $r = .031$). We did not find any significant correlation in the control group for neither the forward ($p = .229$, $r = -.177$) nor the backwards condition ($p = .554$, $r = .088$). These results suggest that when there is a strong degree of familiarity between the observer and the movement, (i.e. dancers observing forward dance movements) people reliably report their affective response in accordance with their bodily arousal.




Figure 4: Scatter plots illustrating the association between subjective VAS ratings and GSR responses for the Forward and Backward dance stimuli as a function of group (Dancers vs. Controls). Only the association for the forward stimuli in Dancers is significant. ** $p = .003$. Note that the data points represent the average values for the stimuli in the respective condition (forward vs. backward) across the participants in the respective groups (Dancers vs. Controls).

4. Discussion

Movement expertise modulates perceptual processes involved in the observation and appreciation of movements (Calvo-Merino et al., 2010, Kirsch, Drommelschmidt, & Cross, 2013), and an increased coupling between subjectively reported emotion and psychophysiological aspects of the emotional experience in expert dancers has been reported (Sze et al., 2010). The present study is the first to examine how movement expertise in the expression of affect through movement modulates sensitivity to such bodily expressed emotion at the level of subjective experiences and objective measures of physiological arousal. Expertise in this context was operationalised in two ways; by comparing expert dancers to control participants with no dance experience and by comparing responses to movements in their normal forward presentation (i.e., as they would be learned by experts) and in an unfamiliar backward presentation.

The principal finding was that expertise in affective body movements indeed augmented sensitivity to observed affective body movements. With their subjective ratings expert dancers discriminated more strongly between happy and sad dance clips played in the usual, forward direction than control participants did. Furthermore, controls had the same level of GSR to happy and sad movements, irrespective of movement presentation, while expert dancers had increased GSR to happy as compared to sad movements, specifically only for movements presented in their forward presentation. Moreover a correlation analysis

showed that only in the expert group subjective behavioural responses correlated with psychophysiological responses, and again this was specific when rating movements in their forward presentation. The observation that GSR responses were overall higher in controls rather than dancers. This is congruent with previous reported differences between experts and laypersons in other art domains (painting), where novel stimuli typically elicit greater physiological arousal responses than familiar responses (Pihko et al., 2011). This observation does, however, raise interesting questions for future studies in terms of how the effects of novelty/familiarity of stimuli interact with the sensitivity of the observer to affective dimensions.

Together, the findings suggest that dance training modulates intrapersonal as well as interpersonal emotional processes. Further research is needed into the involved mechanisms by examining the emotional sensitivity of dancers longitudinally as they become experts, and by looking at correlations between indices of emotional sensitivity (e.g., the correlation between subjective emotion ratings and physiological arousal) and years of dance experience. Additional individual difference variables such as trait emotional intelligence or Alexithymia (difficulty to identify and describe one's own emotion) are important to consider in future studies as possible mediators/moderators of the effects of expertise on emotional processes.

Beyond the role of expertise in the processing of emotion in movement, the present data raise questions about what precisely constitutes emotional movement. Dance naïve participants can identify with the emotion expressed in ballet dance (Christensen et al., 2014a; Christensen, Gaigg, Gomila, Oke & Calvo-Merino, 2014b), at least as far as their subjectively reported feelings are concerned. Given that the happy and sad dance clips selected for the present study did not differ in terms of the particular steps that comprised the dance sequences, the emotional salience of movements must be transmitted through the quality with which the movements are performed. This is not necessarily surprising. What is less expected is that some of the quality that renders movements emotionally expressive

appears to be preserved when the same movements are presented backwards. That is, both groups of participants reported differential happy and sad feelings in response to the dance clips irrespective of whether the clips were played in the normal forward or their unusual backward presentation. Dance expertise augmented the difference in the forward presentation but it did not diminish it in the backwards presentation. This finding could indicate that the emotional quality of movements is temporally relatively symmetrical such that temporal reversals do not result in a loss of emotional information. Or it could be that certain aspects of the temporal dynamics of movement are not critical for transmitting emotional information. It is worth noting, however, that dancers' GSR responses did not differentiate between happy and sad dance movements in the backward conditions, which suggests that the temporal dynamics of emotional movements impact differently on subjective and psychophysiological aspects of emotional responses. These issues warrant further investigation and future studies could seek to identify which properties of movement (e.g., angular velocity, jerk, etc.) predict subjective and/or psychophysiological responses. Such studies could lead to fruitful discoveries that may ultimately feed back into educational practices in ballet schools.

It will be important for the current observations to be replicated and extended to other expert groups such as actors, mimes and other performance artists who are experts in the bodily expression of emotion, and to use other types of stimuli materials; both artistic and everyday-type expressions of affect. The current study used a moderate sample size, given that some participants needed to be excluded from the analysis of GSR responses. Participant exclusions are unfortunately unavoidable in psychophysiological research and the recruitment of a specialist population (expert dancers) places certain constraints on achievable sample sizes.

In relation to the wider emotion literature, our results support the original conjecture made by James (1894) and reiterated in contemporary views (Laird & Lacasse, 2014; Niedenthal, 2007) that proprioceptive arousal feedback informs the conscious experience of

emotions. The data also speak to complementary motor simulation and embodiment accounts of social cognition (Keysers & Gazzola, 2006), which argue that a form of embodied simulation or mimicry of the behaviours and experiences of others is not only important for the understanding of others' actions (Jacob & Jeannerod, 2005; Jeannerod, 2001), but also for understanding and identifying with their affective experiences (Blackemore & Decety, 2001; Carr, Iacoboni, Dubeau, Mazziotta, & Lenzi, 2003; Chartrand & Bargh, 1999; Critchley, 2005; Dapretto et al., 2006; Di Dio & Gallese, 2009; Gallese, 2003; Goldman & Sripada, 2005; Molnar-Szakacs & Overy, 2006). This idea is intuitive when considering that all affective expression –be it facial or bodily– normally involves movement of our muscles. After all, emotion is also motion. Importantly, in the context of this wider literature, the current observations suggest that training in the bodily expression of emotions enhances an individual's sensitivity to the emotions expressed by others, with potentially important implications for the possible utility of dance and movement therapies for the management of disorders such as Autism Spectrum Disorders (ASD) that are characterised by impairments in social-emotional and wider social-cognitive processes (Chevallier, Kohls, Troiani, Brodtkin, & Schultz, 2012; Gaigg, 2012; see particularly Scharoun, Reinders, Bryden & Fletcher, 2014).

5. Author contributions

J.F.Christensen, Dr. B. Calvo-Merino, Dr. S.B. Gaigg and Dr. A. Gomila developed the study concept. J.F.Christensen, Dr. B. Calvo-Merino and S.B. Gaigg contributed to the study design. Testing and data collection were performed by J.F. Christensen and N. Sivarajah. J.F.Christensen, and Dr. B. Calvo-Merino performed the data analysis and interpretation under the supervision of Dr. S.B. Gaigg and A. Gomila. J.F. Christensen drafted the manuscript, and Dr. B. Calvo-Merino, Dr. S.B. Gaigg and Dr. A. Gomila provided critical revisions. All authors approved the final version of the manuscript for submission.

Table and Figure legends:

Table 1: Participant characteristics. Shown are mean and (SD). “Other dance styles” include Step Dance, Jazz Dance, Jazz Ballet, Burlesque, Lyrical and Commercial Dance

Figure 1. Trial structure. After a fixation cross (1500ms), the dance video was faded in (~6 seconds), faded out, and followed by a fixation cross (1000ms). Then the Visual Analogical Scale (ranging from Sad, 0; to Happy, 100) appeared below the word “Emotion?” written in the centre. The cursor of the mouse appeared always in the top centre of the screen rather than anywhere on the VAS scale to avoid the cursor position biasing the participants’ responses toward any of the extremes. Reproduced with permission (Christensen et al., 2014b).

Figure 2. Subjective affect ratings means in the 2x2x2 design. Bars reflect the different conditions: Stimulus Presentation (Forward, Backwards), Dance Emotion (Happy, Sad) and Group (Dancers, Controls). (** = $p < .001$). Error bars reflect S.E.M. (VAS = Visual Analogical Scale for the affective ratings).

Figure 3. Physiological responses measured through Galvanic Skin Response (GSR, Log transformed μs) of Dancers and Controls during observation of dance videos rated as Happy and Sad. Data are presented for the two stimuli presentations conditions (Forward, Backwards). (* = $p < .05$). Error bars reflect S.E.M.

Figure 4: Scatter plots illustrating the association between subjective VAS ratings and GSR responses for the Forward and Backward dance stimuli as a function of group (Dancers vs. Controls). Only the association for the forward stimuli in Dancers is significant. ** $p = .003$.

Note that the data points represent the average values for the stimuli in the respective condition (forward vs. backward) across the participants in the respective groups (Dancers vs. Controls).

6. References

- Bangert, M., Peschel, T., Schlaug, G., Rotte, M., Drescher, D., Hinrichs, H. (2006). Shared networks for auditory and motor processing in professional pianists: evidence from fMRI conjunction. *Neuroimage* 30, 917-926.
- Bastiaansen, J. A., Thioux, M., & Keysers, C. (2009). Evidence for mirror systems in emotions. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 364(1528), 2391-2404.
- Bengtsson, S. L., Nagy, Z., Skare, S., Forsman, L., Forssberg, H., & Ullen, F. (2005). Extensive piano practicing has regionally specific effects on white matter development. *Nature Neuroscience*, 8(9), 1148-1150. doi: 10.1038/nn1516
- Blackmore, S.L., & Decety, J. (2001). From the perception of action to the understanding of intention. *Nature Neuroscience*, 2, 561-567.
- Bläsing, B., Calvo-Merino, B., Cross, E., Jola, C., Honisch, J., Stevens, K. (2012). Neurocognitive control in dance perception and performance. *Acta Psychologica*, 139(2), 300-308.
- Calvo-Merino, B., Ehrenberg, S., Leung, D., & Haggard, P. (2010). Experts see it all: configural effects in action observation. *Psychological Research*, 74(4), 400-406.
- Calvo-Merino, B., Glaser, D. E., Grèzes, J., Passingham, R. E., & Haggard, P. (2005). Action observation and acquired motor skills: An fMRI study with expert dancers. *Cerebral Cortex*, 15(8), 1243-1249. doi: 10.1093/cercor/bhi007

- Calvo-Merino, B., Grèzes, J., Glaser, D. E., Passingham, R. E., & Haggard, P. (2006). Seeing or doing? Influence of visual and motor familiarity in action observation (vol 16, pg 1905, 2006). *Current Biology*, 16(22), 2277-2277. doi: 10.1016/j.cub.2006.10.065
- Cannon, W. B. (1927). The James-Lange theory of emotions: A critical examination and an alternative theory. *The American Journal of Psychology*, 39, 106-124.
- Carr, L., Iacoboni, M., Dubeau, M.C., Mazziotta, J.C., & Lenzi, G.L. (2003). Neural mechanisms of empathy in humans: a relay from neural systems for imitation to limbic areas. *Proceedings of the National Academy of Sciences of the United States of America*, 100, 5497-5502.
- Cela-Conde, C. J., Marty, G., Maestu, F., Ortiz, T., Munar, E., Fernandez, A. (2004). Activation of the prefrontal cortex in the human visual aesthetic perception. *Proceedings of the National Academy of Sciences of the United States of America*, 101(16), 6321-6325. doi: 10.1073/pnas.0401427101
- Chartrand, T., & Bargh, J. (1999). The chameleon effect: the perception-behavior link and social interaction. *Journal of personality and social psychology*, 76, 893-910.
- Chevallier, C., Kohls, G., Troiani, V., Brodtkin, E.S., & Schultz, R.T. (2012). The social motivation theory of autism. *Trends in Cognitive Science*, 16, 231-239.
- Christensen, J. F., & Calvo-Merino, B. (2013). Dance as a subject for empirical aesthetics. *Psychology of Aesthetics, Creativity and the Arts*, 7, 76-88.
- Christensen, J.F., Nadal, M., Cela-Conde, C.J., & Gomila, A. (2014a). A norming study and library of 203 dance movements. *Perception*, 43(2/3), 178-206. doi: 10.1068/p7581
- Christensen, J.F., Gaigg, S.B., Gomila, A., Oke, P., & Calvo-Merino, B. (2014b). Enhancing emotional experiences to dance through music: the role of valence and arousal in the cross-modal bias. *Frontiers in Human Neuroscience*, 8, 757.
- Cohen, J. (1988). *Statistical power analysis for the behavioral sciences* (2nd ed.). Hillsdale, NJ: Lawrence Erlbaum Associates Inc.

- Critchley, H.D. (2005). Neural mechanisms of autonomic, affective and cognitive integration. *The Journal of Comparative Neurology*, 493, 154-166. doi: 10.1002/cne.20749
- Cross, E.S., Hamilton, A.F. de C., & Grafton, S.T. (2006). Building a motor simulation de novo: Observation of dance by dancers. *Neuroimage*, 31(3), 1257-1267.
- Damasio, A. (1999). *The feeling of what happens: Body, emotion and the making of consciousness*. London, UK: Heinemann.
- Dapretto, M., Davies, M.S., Pfeifer, J.H., Scott, A.A., Sigman, M., Bookheimer, S.Y. (2006). Understanding emotions in others: mirror neuron dysfunction in children with autism spectrum disorders. *Nature Neuroscience*, 9, 28-30.
- Dellacherie, D., Roy, M., Hugueville, L., Peretz, I., & Samson, S. (2010). The effect of musical experience on emotional self-reports and psychophysiological responses to dissonance. *Psychophysiology*, 48(3), 337-349. doi: 10.1111/j.1469-8986.2010.01075.x
- Decety, J., & Jackson, P.L. (2006). A Social-Neuroscience Perspective on Empathy. *Current Directions in Psychological Science*, 15, 54-58.
- Di Dio, C., & Gallese, V. . (2009). Neuroaesthetics: a review. *Current Opinion in Neurobiology*, 19(6), 682-687.
- Effron, D. A., Niedenthal, P. M., Gil, S., & Droit-Volet, S. (2006). Embodied temporal perception of emotion. *Emotion*, 6(1).
- Evans, P., & Schubert, E. (2008). Relationships between expressed and felt emotions in music. *Musicae Scientiae*, 12(1), 75-99.
- Fink, A., Graif, B., & Neubauer, A. C. (2009). Brain correlates underlying creative thinking: EEG alpha activity in professional vs. novice dancers. *Neuroimage*, 46(3), 854-862. doi: 10.1016/j.neuroimage.2009.02.036

- Frischen, A., Eastwood, J. D., & Smilek, D. (2008). Visual search for faces with emotional expressions. *Psychological Bulletin*, 134(5).
- Gabrielsson, A. (2002). Emotion perceived and emotion felt: Same or different? *Musicae Scientiae*, Special Issue 2001-2002, 123-147
- Gaigg, S.B. (2012). The interplay between emotion and cognition in autism spectrum disorder: Implications for developmental theory. *Frontiers in Integrative Neuroscience* 4, 113.
- Gallese, V. (2003). The roots of empathy: the shared manifold hypothesis and the neural basis of intersubjectivity. *Psychopathology*, 36, 171-180.
- Gaser, C., & Schlaug, G. (2003). Brain structures differ between musicians and non-musicians. *The Journal of Neuroscience*, 16(6), 9240-9245.
- Goldman, A. I., & Sripada, C. S. (2005). Simulationist models of face-based emotion recognition. *Cognition* 94, 193-213. doi: doi:10.1016/j.cognition.2004.01.005
- Goldstein, T.R. (2009). Psychological Perspectives on Acting. *Psychology of Aesthetics, Creativity, and the Arts*, 3(1), 6-9.
- Goldstein, T.R., & Bloom, P. (2011). The Mind on stage: Why Cognitive Scientists Should Study Acting. *Trends in Cognitive Sciences*, 15(141-142).
- Goldstein, T.R., & Winner, E. (2012). Enhancing Empathy and Theory of Mind. *Journal of Cognition and Development*, 13(1), 19-37.
- Habib, M., & Besson, M. (2009). What do music training and musical experience teach us about brain plasticity? *Music Perception*, 26, 279-285.
- Hänggi, J., Koeneke, S., Bezzola, L., & Jäncke, L. (2010). Structural Neuroplasticity in the Sensorimotor Network of Professional Female Ballet Dancers. *Human Brain Mapping*, 31(8), 1196-1206. doi: 10.1002/hbm.20928

- Haslinger, B., Erhard, P., Altenmüller, E., Schroeder, U., Boecker, H., & Ceballos–Baumann, A.O. . (2005). Transmodal sensorimotor networks during action observation in professional pianists. *Journal of Cognitive Neuroscience*, 17, 282-293.
- Hunter, P. G., Schellenberg, E. G., & Schimmack, U. (2010). Feelings and Perceptions of Happiness and Sadness Induced by Music: Similarities, Differences, and Mixed Emotions. *Psychology of Aesthetics Creativity and the Arts*, 4(1), 47-56.
- Jacob, P., & Jeannerod, M. (2005). The motor theory of social cognition: a critique. *Trends in Cognitive Sciences*, 9, 21-25. doi: doi:10.1016/j.tics.2004.11.003
- James, W. (1894). Discussion: The physical basis of emotion. *Psychological Review*, 1, 516-529.
- Jang, S. H., & Pollick, F.E. (2011). Experience Influences Brain Mechanisms of Watching Dance. *Dance Research Journal*, 29(2), 352-377.
- Jeannerod, M. (2001). Neural Simulation of Action: A Unifying Mechanism for Motor Cognition. *NeuroImage*, 14, 103-109.
- Kallinen, K., & Ravaja, N. (2006). Emotion perceived and emotion felt: Same and different. *Musicae Scientiae*, 10(2), 191-213.
- Keysers, C., & Gazzola, V. (2006). Toward a unifying neural theory of social cognition. *Progress in Brain Research*, 156, 379-401. doi: 10.1016/S0079-6123(06)56021-2
- Kirsch, L. P., Drommelschmidt, K. A., Cross, E. S. (2013). The impact of sensorimotor experience on affective evaluation of dance. *Frontiers in Human Neuroscience*, 7(521), 1-10. doi: 10.3389/fnhum.2013.00521. eCollection 2013
- Kirsch, L.P., Dawson, K., Cross, E.S. (2015). Dance experience sculpts aesthetic perception and related brain circuits. *Annals of the New York Academy of Science*, 1337, 130-9. doi: 10.1111/nyas.12634

- Laird, J.D., & Lacasse, K. (2014). Bodily Influences on Emotional Feelings: Accumulating Evidence and Extensions of William James's Theory of Emotion. *Emotion Review*, 6(27). doi: 10.1177/1754073913494899
- Lang, P.J., Bradley, M.M., & Curthbert, B.N. (1998). Emotion, Motivation, and Anxiety: Brain Mechanisms and Psychophysiology. *Biological Psychiatry*, 44, 1248-63.
- Lang, P. J., Bradley, M. M., & Cuthbert, B. N. (2008). International affective picture system (IAPS): Affective ratings of pictures and instruction manual. Technical Report A-8, University of Florida, Gainesville.
- Lima, C.F., & Castro, S.L. (2011). Speaking to the trained ear: musical expertise enhances the recognition of emotions in speech prosody. *Emotion*, 11, 1021-1031.
- McDonald, J. H. (2009). *Handbook of Biological Statistics*. 2nd Edn. Baltimore, Maryland: Sparky House Publishing.
- Mauss, I.B., Levenson, R.W., McCarter, L., Wilhelm, F.H., & Gross, J.J. (2005). The tie that binds? Coherence among emotional experience, behavior, and autonomic physiology. *Emotion*, 5, 175-190.
- Molnar-Szakacs, I., & Overy, K. (2006). Music and mirror neurons: from motion to 'e'motion. *Social Cognitive and Affective Neuroscience*, 1, 235-241.
- Niedenthal, P.M. (2007). Embodying emotion. *Science*, 316(5827), 1002-1005. doi: 10.1126/science.1136930
- Oechslin, M. S., Van De Ville, D., Lazeyras, F., Hauert, C.A., & James, C.E. (2013). Degree of Musical Expertise Modulates Higher Order Brain Functioning. *Cerebral Cortex*, 23(9), 2213-2224. doi: 10.1093/cercor/bhs206
- Orgs, G., Dombrowski, J. H., Heil, M., & Jansen-Osmann, P. (2008). Expertise in dance modulates alpha/beta event-related desynchronization during action observation. *European Journal of Neuroscience*, 27(12), 3380-3384. doi: 10.1111/j.1460-9568.2008.06271.x

Pantev, C., Oostenveld, R., Engelien, A., Ross, B., Roberts, L.E., & M., Hoke. (1998).

Increased auditory cortical representation in musicians. *Nature*, 392(6678), 811-814.

Petrides, K. V., Niven, L., & Mouskounti, T. (2006). The trait emotional intelligence of ballet dancers and musicians. *Psicothema*, 18, 101-107.

Pihko, E., Virtanen, A., Saarinen, V.-M., Pannasch, S., Hirvenkari, L., Tossavainen, T., et al. (2011). Experiencing art: the influence of expertise and painting abstraction level. *Frontiers in Human Neuroscience*, 5(94).

Pollick, F. E., Paterson, H., Bruderlin, A., & Sanford, A. J. (2001). Perceiving affect from arm movement. *Cognition*, 82, B51–B61.

Rothman, K. J. (1990). No adjustments are needed for multiple comparisons. *Epidemiology* 1, 43–46. doi: 10.1097/00001648-199001000-00010

Salimpoor, V., Benovoy, M., Longo, G., Cooperstock, J.R., and Zatorre, R.J. (2009). The Rewarding Aspects of Music Listening are Related to Degree of Emotional Arousal, *PLoS ONE*, 4(10): e7487

Saville, D. J. (1990). Multiple comparison procedures: *Statistics*. 44, 174–180. doi: 10.2307/2684163

Scherer, K. (2009a). The dynamic architecture of emotion: evidence for the component process model. *Cognition & Emotion*, 23(7), 1307-1351.

Scherer, K. R. (2009b). Emotions are emergent processes: they require a dynamical computational architecture, *Philosophical Transactions of the Royal Society B*, 364/1535: doi: 10.1098/rstb.2009.0141.

Scherer, K. R. (2013). The evolutionary origin of multimodal synchronization in emotional expression. *Journal of Anthropological Sciences*, 91, 185–200. doi:10.4436/JASS.91005.

Scharoun, S.M., Reinders, N.J., Bryden, P.J., & Fletcher, P.C. (2014). Dance/Movement therapy as an intervention for children with Autism Spectrum Disorders. *American Journal of Dance Therapy*, 36, 209-228.

Schlaug, G. (2006). The Brain of Musicians: A Model for Functional and Structural Adaptation. *Annals of the New York Academy of Sciences*, 930, 281-299. doi: 10.1111/j.1749-6632.2001.tb05739.x

Sze, J. A., Gyurak, A., Yuan, J. W., & Levenson, R. W. (2010). Coherence Between Emotional Experience and Physiology: Does Body Awareness Training Have an Impact? *Emotion*, 10(6), 803-814. doi: 10.1037/a0020146