



City Research Online

City, University of London Institutional Repository

Citation: Murphy, J., Gray, K. & Cook, R. (2016). The Composite Face Illusion. *Psychonomic Bulletin & Review*, doi: 10.3758/s13423-016-1131-5

This is the accepted version of the paper.

This version of the publication may differ from the final published version.

Permanent repository link: <http://openaccess.city.ac.uk/15072/>

Link to published version: <http://dx.doi.org/10.3758/s13423-016-1131-5>

Copyright and reuse: City Research Online aims to make research outputs of City, University of London available to a wider audience. Copyright and Moral Rights remain with the author(s) and/or copyright holders. URLs from City Research Online may be freely distributed and linked to.

City Research Online:

<http://openaccess.city.ac.uk/>

publications@city.ac.uk

In press: *Psychonomic Bulletin & Review*

Format: *Review Article*

Running head: *The Composite Face Illusion*

The Composite Face Illusion

Jennifer Murphy^{1,2}, Katie Gray³, Richard Cook^{1,*}

¹Department of Psychology, City University London, London, U.K.

²MRC Social, Genetic and Developmental Psychiatry Centre, Institute of Psychiatry, Psychology, and Neuroscience, King's College London, London, U.K.

³Department of Psychology, University of Reading, Reading, U.K.

*Corresponding author: Richard.Cook.1@city.ac.uk

Department of Psychology,

City University London,

Whiskin Street

London, EC1R OJD

Tel: +44 20 7040 8644, Fax: +44 20 7040 8877

Abstract

Few findings in cognitive science have proved as influential as the composite face effect. When the top half of one face is aligned with the bottom half of another, and presented upright, the resulting composite arrangement induces a compelling percept of a novel facial configuration. Findings obtained using composite face procedures have contributed significantly to our understanding of holistic face processing, the detrimental effects of face inversion, the development of face perception, and aberrant face perception in clinical populations. Composite paradigms continue to advance our knowledge of face perception, as exemplified by their recent use for investigating the perceptual mechanisms underlying dynamic face processing. However, the paradigm has been the subject of intense scrutiny, particularly over the last decade, and there is a growing sense that the composite face illusion, whilst easy to illustrate, is deceptively difficult to measure and interpret. In this review we provide a focussed overview of the existing composite face literature, and identify six priorities for future research. Addressing these gaps in our knowledge will aid the evaluation and refinement of theoretical accounts of the illusion.

Key words:

Composite face effect; face perception; holistic representation; visual illusions

1. Introduction

In their seminal paper, Young, Hellawell and Hay (1987) presented a new ‘composite face paradigm’ for investigating the perception of facial configurations. The authors described how, when the top half of one face was aligned with the bottom half of another, the resulting composite arrangement induced the perception of a novel facial configuration; the presence of the bottom half altered the appearance of the top half, and *vice versa*. The illusory fusion of the two aligned halves hindered recognition – observers took longer to identify the source faces. Importantly, however, the illusion was observed only when the composite arrangements were aligned spatially and presented upright; free from illusion-induced interference, observers were able to identify the sources of upright, misaligned halves with comparative ease (Figure 1). When presented upside-down, observers’ recognition of the source faces was broadly similar in the aligned and misaligned conditions.

The composite face illusion has proved enormously influential. Crucially, the effect reveals a tendency to integrate feature information from disparate facial regions. Findings from studies employing composite face techniques have therefore informed the development of holistic theories of face perception, which posit that local facial features are integrated into a unified representation for the purposes of efficient analysis and interpretation (Farah, Wilson, Drain, & Tanaka, 1998; Maurer, Le Grand, & Mondloch, 2002; Piepers & Robbins, 2013). Moreover, the finding that greater illusory fusion is induced by upright arrangements suggests that the orientation-specific processes responsible for composite interference may also be disrupted in the classic face inversion effect, whereby orientation inversion disproportionately impairs the recognition of faces, relative to other classes of object (Yin, 1969).

The purpose of this review is to provide a focussed overview of the existing composite face literature with a view to identifying priorities for future research. We begin by discussing some of the practical issues associated with measurement (*Section 2*). We then consider how susceptibility to the illusion develops during ontogeny and the relative contribution of genetic and environmental factors (*Section 3*). In the following sections we consider the neural signatures associated with the illusion (*Section 4*), and discuss findings obtained from clinical populations (*Section 5*). Having reviewed the available empirical data on the composite face effect, we present and evaluate different theoretical accounts of the illusion (*Section 6*). We end the review with discussion of six priorities for future composite face research (*Section 7*).




Figure 1: When aligned with different lower halves, it is surprisingly difficult to recognise that the upper regions of the two composites are identical (left top). However, the illusion-induced interference is greatly diminished when the composites are misaligned (right top). When composite arrangements are shown upside-down, little illusion-induced interference is seen in either the aligned (left bottom) or misaligned (right bottom) conditions.

2. Measuring observers' susceptibility to the illusion

2.1 Paradigms

In the original composite study (A. W. Young et al., 1987), participants were presented with a single facial composite and the degree of illusion-induced interference was inferred from their response latencies. When asked to identify the source of a task-relevant half (e.g., upper half; hereafter 'target half'), whilst disregarding the remaining task-irrelevant half (e.g., lower half; hereafter 'distractor half'), observers were disproportionately slower in the upright-aligned condition. The application of this naming paradigm is limited, however, by the need to use familiar faces. To allow participants to name the source of the target half, composites must be constructed from personally familiar, celebrity, or otherwise learned faces. This is potentially problematic in light of known differences in the perceptual processing of familiar and unfamiliar faces (Jenkins & Burton, 2011; Murphy, Ipser, Gaigg, & Cook, 2015). For example, observers are better able to match familiar faces using their internal features, including the eyes, nose, and mouth (Osborne & Stevenage, 2008). In contrast, unfamiliar face matching is frequently based on external features, such as hairstyle and face shape. Familiar faces also place lower demands on visual working memory (Jackson & Raymond, 2008) and are easier to detect under conditions of reduced attention, compared with unfamiliar faces (Jackson & Raymond, 2006).

Since the initial description of the illusion, simultaneous and delayed matching procedures have been developed which overcome the need to use familiar faces (i.e., that allow investigation of holistic processing of unfamiliar faces). Crucially, these paradigms confirm that arrangements constructed from unfamiliar faces also produce composite face effects. In simultaneous-matching paradigms, two composite arrangements – either aligned or misaligned, upright or inverted – are presented simultaneously and observers asked whether two target regions (e.g., the upper halves) are identical or not (Hole, 1994). In delayed-matching paradigms, participants are again presented with two composite arrangements and asked to judge whether the target regions are identical or not. However, rather than appear simultaneously, the composites are presented sequentially (Goffaux & Rossion, 2006; Le Grand, Mondloch, Maurer, & Brent, 2004; Michel, Rossion, Han, Chung, & Caldara, 2006). In a popular variant, the first composite is presented briefly (e.g., 500 ms), followed by an inter-stimulus-interval, typically between 500 ms and 2 seconds. Thereafter, the second composite is presented until the participant responds (e.g., Michel et al., 2006). In both matching procedures, observers’ responses are disproportionately impaired (slower and/or less accurate) when composite arrangements are aligned and presented upright, than when misaligned or inverted.

2.2 Original and complete matching procedures

In simultaneous and sequential matching paradigms, the factorial combination of target-half (T_{same} , $T_{\text{different}}$) and distractor-half (D_{same} , $D_{\text{different}}$) yields four possible trial types: congruent-same (T_{same} , D_{same}), incongruent-same (T_{same} , $D_{\text{different}}$), congruent-different ($T_{\text{different}}$, $D_{\text{different}}$), incongruent-different ($T_{\text{different}}$, D_{same}). Note that on ‘congruent’ and ‘incongruent’ trials, susceptibility to the illusion therefore helps and hinders observers, respectively. Traditionally, researchers have employed a variant whereby distractor halves always differ, and therefore exert different illusory biases on the target regions (e.g., Cassia, Picozzi, Kuefner, Bricolo, & Turati, 2009; de Heering, Houthuys, & Rossion, 2007; Goffaux & Rossion, 2006; Konar, Bennett, & Sekuler, 2010; Le Grand et al., 2004; Robbins & McKone, 2007). The original matching procedure excludes congruent-same and incongruent-different trial types (Figure 2a). More recently, some groups have advocated the use of so-called ‘complete design’ matching procedures (Figure 2b), incorporating identical and different distractor halves and all four trial types (e.g., Cheung, Richler, Palmeri, & Gauthier, 2008; Richler, Cheung, & Gauthier, 2011a; Richler, Gauthier, Wenger, & Palmeri, 2008; Richler, Mack, Palmeri, & Gauthier, 2011). Importantly, upright face composite effects calculated using the complete

and original matching designs do not correlate (Richler & Gauthier, 2014), suggesting that the variants measure different phenomena, however the reason for the discrepancy remains contested (Meinhardt, Meinhardt-Injac, & Persike, 2014; Richler & Gauthier, 2013; Rossion, 2013).




Figure 2: Illustration of the (a) original and (b) complete designs used in simultaneous and sequential matching paradigms.

Proponents argue that, compared to the original matching procedure, the complete design estimates observers' susceptibility to the composite illusion more accurately. Unlike estimates inferred using the complete design, composite effects measured using the original matching task may be influenced by differences in response bias (Richler & Gauthier, 2014). For example, estimates of composite interference derived from original matching designs are disproportionately driven by responses on incongruent-same trials where the presence of the different distractors makes it harder to recognise that same target halves are identical. Where observers have a pre-existing bias to respond 'different', original matching designs may underestimate their susceptibility to the illusion. Consistent with this view, belief manipulations that affect response bias distort composite effect estimates measured using the original matching design, but not the complete design (Richler, Cheung et al., 2011a).

However, the complete design has also been criticised. Importantly, the effects of alignment seen in the additional congruent-same and incongruent-different trials may have a distinct origin to those seen in the incongruent-same and congruent-different trials included in the original matching design. Specifically, the facilitation seen on congruent-same trials, and the interference seen on incongruent-different trials, may reflect domain-general processes akin to response priming and response conflict, not perceptual integration (Rossion, 2013).

Accordingly, complete designs may be prone to strategic effects (see *Section 7.2*) and may be more likely to find effects with non-face composites (see *Section 6.1*); for example, composite effects have been observed for unfamiliar Chinese characters using this paradigm (Hsiao & Cottrell, 2009). A further objection relates to the predicted pattern of responses in the congruent-different condition. On these trials, different distractor regions are paired with different target regions. On some trials it is reasonable to assume that the additional different signal present in the distractors is fused with the physical differences between the targets, thereby making it easier for observers to respond ‘different’. However, where the physical difference between the targets is already large, it is conceivable that fusion with distractor regions with less obvious differences may actually dilute the difference signal, hindering correct responding (Robbins & McKone, 2007).

2.3 Control conditions

While different research groups employ different control conditions, the aim of these manipulations is consistent: to rule out simpler, generic accounts of the performance decrements seen when upright composites are aligned. The most commonly used control conditions, misalignment and inversion (see Figure 1), were introduced by Young and colleagues (1987). Interference is greatly reduced when composites are misaligned horizontally excluding the possibility that the interference is simply caused by the proximity of the distractor half. Transposing the position of upper and lower halves (Abbas & Duchaine, 2008) and presenting halves side-by-side (Richler, Tanaka, Brown, & Gauthier, 2008) also speak against this account. Similarly, the finding that composite interference is greatly diminished – although not eliminated entirely (e.g., Susilo, Rezlescu, & Duchaine, 2013) – when stimulus arrangements are shown upside-down confirms that performance decrements in the upright-aligned condition are not induced by the presence of the continuous boundary when the halves are aligned; whereas the boundary is disrupted by misalignment, it is preserved by inversion (Rossion, 2013). Where effects of alignment are seen for upright, but not inverted arrangements, authors can also exclude the possibility that the differential interference observed in these conditions reflects differences in the size of the spotlight of visuospatial attention (e.g., McKone et al., 2013). For these reasons, many consider the use of convergent inverted and misaligned control conditions necessary for clear interpretation of composite interference (McKone et al., 2013; Rossion, 2013). Finally, some investigations have included parallel tasks where observers judge composites constructed from non-face objects, including cars (Cassia et al., 2009) and dogs (Robbins & McKone, 2007). Where

observed, findings of face-specificity argue against the possibility that decrements in the upright-aligned condition reflect domain-general interference that may be produced by any object with a canonical orientation (e.g., Robbins & McKone, 2007).

2.4 Which judgments are affected?

In the original description of the illusion, observers were slower to name the identity of the faces from which target regions were sourced as a consequence of the composite interference (A. W. Young et al., 1987). It appears, however, that composite interference affects several other attributions, extending beyond facial identity. For example, viewers are slower to identify the emotion present in a target half, when paired with a distractor half sourced from the same face, but expressing a different emotion. Once again, disproportionate interference is seen when composite arrangements are presented upright and aligned, with little or no interference seen when the halves are misaligned or inverted (Calder, Young, Keane, & Dean, 2000; Palermo et al., 2011; Tanaka, Kaiser, Butler, & Le Grand, 2012; Wegrzyn, Bruckhaus, & Kissler, 2015). Composite interference also appears to bias the perception of facial gender and age. Observers were slower and made more errors in categorising the gender of target halves when composite faces were constructed using halves from different genders, relative to same-gender and misaligned control conditions (Baudouin & Humphreys, 2006). Similarly, when estimating the age of composite top halves, observers' estimates were biased towards the age of the bottom halves, a pattern of performance that was attenuated, but not eliminated, by inversion (Hole & George, 2011). Composite interference also distorts evaluative impressions. Top halves of upright composite faces were judged more attractive (Abbas & Duchaine, 2008) and trustworthy (Todorov, Loehr, & Oosterhof, 2010) when aligned with attractive and trustworthy bottom halves, respectively.

2.5 Low-level stimulus characteristics

Several studies have sought to investigate how observers' susceptibility to the composite face illusion is affected by low-level image manipulations. For example, results obtained using the original matching procedure suggest that filtered composites that preserve low-spatial frequencies, induce stronger illusion-induced interference than high-spatial frequency composites (Goffaux & Rossion, 2006). While this finding is consistent with the view that holistic processing is supported predominantly by low-spatial frequencies (Sergent, 1986), it has been contested; subsequent findings obtained using the complete design have found no effect of spatial frequency (Cheung et al., 2008). A related suggestion is that composite

effects vary as a function of face size, with arrangements resembling faces viewed at ~2m inducing the strongest illusions when measured using the complete design (Ross & Gauthier, 2015). Facial composites presented in grayscale have been found to induce greater composite interference in the original matching procedure, than composites shown in colour (Retter & Rossion, 2015). Interestingly, photographic negation has little effect on the strength of the illusion when measured using the original matching procedure; grayscale composites induce similar interference in positive and negative contrast polarity (Hole, George, & Dunsmore, 1999; Taubert & Alais, 2011).

3. Development

Susceptibility to the composite face illusion appears to be present early in development. For example, de Heering and colleagues (2007) found that four, five and six-year-olds displayed similar susceptibility to each other and to adults when tested using the original matching procedure (see also Carey & Diamond, 1994; Mondloch, Pathman, Maurer, Le Grand, & Schonen, 2007; Susilo, Crookes, McKone, & Turner, 2009). Comparable results have also been reported for expression composites (Durand, Gallay, Seigneureic, Robichon, & Baudouin, 2007). Susceptibility, however, may be present at an even younger age. Composite interference has been reported in 3-year-olds when tested on a simultaneous variant of the original matching procedure (Cassia et al., 2009). Remarkably, recent evidence from a visual preference paradigm indicates that infants as young as 3 months may also be susceptible to the illusion. When misaligned, 3-month-old infants exhibited a looking preference for familiar eye-regions over unfamiliar eye-regions. However, when aligned with an unfamiliar mouth, no systematic preference was observed (Turati, Di Giorgio, Bardi, & Simion, 2010).

Early susceptibility to the composite face illusion may partly reflect a genetic contribution. Having tested 173 twin pairs (102 monozygotic and 71 dizygotic, age 7–19 years) using the original matching procedure, the correlation between the composite effects estimated for the monozygotic twins exceeded that seen for the dizygotic twins (Zhu et al., 2010). The authors estimated that genetic heritability accounted for 31% of the variance in the twins' susceptibility to the illusion. As is often the case with heritable traits (e.g., Haworth et al., 2010), the genetic influence appeared to increase with age; the variance attributable to genetic factors was greater in the 13-19 year-olds, than in 7-12 year-olds. No concordance was observed for a non-face measure of global-to-local interference based on the Navon Task

(Navon, 1977), suggesting that the heritability of the composite face illusion may show a degree of face-specificity.

Complementary evidence suggests that visual experience also plays a critical role in shaping susceptibility to the composite face illusion. Twelve individuals (age 9-23 years) deprived of visual experience during the first 3-6 months of development due to bilateral congenital cataracts, showed little or no susceptibility to the composite face illusion when measured using the original matching design. Free from illusion-induced interference, these individuals outperformed controls when the target and distractor halves were aligned (Le Grand et al., 2004). When tested using the original matching design, Michel and colleagues (2006) found that observers' susceptibility to the composite illusion was also greater when participants viewed composite faces of their own race (but for conflicting results obtained with the complete design see Horry, Cheong, & Brewer, 2015; Zhao, Hayward, & Bühlhoff, 2014). For a review of this literature, see Hayward, Crookes, & Rhodes, (2013). Similarly, Susilo and colleagues (2009) found stronger effects using the original matching procedure when composites were constructed from own-age faces (but see Wiese, Kachel, & Schweinberger, 2013). However, preschool teachers (i.e. observers with extensive contact with child faces) failed to show an own-age effect when tested using the original matching task; they showed similar composite effects for arrangements constructed from own-age faces and the faces of children (de Heering & Rossion, 2008). Where observed, these biases may therefore be products of experience.

4. Neural basis

4.1 Neural markers revealed by fMRI and EEG

The neural representation of composite faces has been addressed using fMRI adaptation paradigms. These studies take advantage of the fact that repeated presentation of the same stimulus elicits a weaker change in the blood-oxygen-level-dependent (BOLD) signal, than presentation of two different stimuli. The same top half of a face elicits a larger response – a bigger release from adaptation – in the fusiform face area (FFA) when aligned with a lower half sourced from a different facial identity than when taken from the same face (Schiltz, Dricot, Goebel, & Rossion, 2010; Schiltz & Rossion, 2006). However, this difference is not found when the top and bottom face regions are spatially misaligned (Schiltz et al., 2010; Schiltz & Rossion, 2006; but see Harris & Aguirre, 2010) or when arrangements are inverted (Schiltz & Rossion, 2006)

The composite face illusion is also associated with a characteristic electroencephalography (EEG) signature. In comparison to aligned composite faces, consistent evidence indicates that misalignment elicits a delayed and enhanced N170 (Jacques & Rossion, 2009, 2010; Kuefner, Jacques, Prieto, & Rossion, 2010; Letourneau & Mitchell, 2008; Soria Bauser, Schriewer, & Suchan, 2014; Wiese et al., 2013), a response that bears striking similarity to the modulation elicited by face inversion (Eimer, 2000). Consistent with behavioural findings, the effect associated with misalignment is not observed when the composites are inverted (Jacques & Rossion, 2010). However, the response elicited may also depend on the stimulus attributes; a stronger effect of misalignment has been reported for young faces in comparison to older faces, regardless of participant age (Wiese et al., 2013). For aligned composite faces a larger N170 is also observed when sequentially presented top halves differ than when they are identical (Jacques & Rossion, 2009). Crucially, however, a similar increase in the N170 is observed when aligned distractor halves induce the illusion that two identical target halves differ (Jacques & Rossion, 2009; Kuefner et al., 2010). This effect is not seen with misaligned composites (Jacques & Rossion, 2009).

4.2 Brain stimulation and neuropsychology

Findings that illusory composite fusion can modulate FFA and N170 responses are of great interest. However, it is important to note that these observations are correlational. For example, the fMRI adaptation effects may reveal a causal role for the FFA in the emergence of the illusory percept, or may merely be a consequence of the illusion. The functional significance of these neural signatures thus remains unclear. In contrast, brain stimulation and neuropsychological paradigms potentially provide evidence that a particular neural substrate is causally involved in the experience of the illusory percept; where a region makes a necessary contribution to the illusion, interference in the form of stimulation or acquired lesion, should alter observers' susceptibility. To date, few investigations have employed brain stimulation to understand the neural basis of the composite face illusion. However, a recent study employing the complete design found that transcranial direct current stimulation (tDCS) applied to occipito-temporal cortex reduced observers' susceptibility to the composite face illusion, relative to a sham condition (Yang et al., 2014; but see Renzi et al., 2015).

In cases of Acquired Prosopagnosia (AP), individuals are left with face recognition difficulties following brain injury. To date, surprisingly few APs have been tested on composite face tasks. However, patients with damage to posterior (Busigny, Joubert, Felician,

Ceccaldi, & Rossion, 2010; Ramon, Busigny, & Rossion, 2010) or anterior (Busigny et al., 2014) regions of the right temporal lobe have been found to exhibit reduced composite face effects, measured using original matching procedures, relative to matched controls. Typical susceptibility to the original matching procedure was reported in patient Herschel, who developed AP following extensive lesions predominantly in the right occipitotemporal cortex (Rezlescu, Pitcher, & Duchaine, 2012). When tested using the complete design, patient LR, an individual who developed AP following focal damage to the anterior portion of the right temporal lobe, exhibited a typical composite effect (Bukach, Bub, Gauthier, & Tarr, 2006). However, this result is qualified by an abnormal pattern of responses; specifically, the time taken by LR to identify the top half of the face was influenced by the bottom half in both the aligned and misaligned conditions, indicative of an atypical composite effect (Busigny et al., 2014).

Patient CK, who exhibited severe integrative object agnosia, showed broadly typical recognition of upright aligned faces; for example, CK was able to identify celebrity faces as well as age-matched controls, even when the identity was obscured by a disguise. However, CK's recognition was drastically impaired relative to controls when upper and lower face halves were misaligned horizontally or when spatially aligned faces were inverted (Moscovitch, Winocur, & Behrmann, 1997). Interestingly, CK's recognition was largely unaffected when left and right facial halves were misaligned vertically, hinting at qualitative differences between the disruption induced by horizontal and vertical displacement. Because of the nature of CK's injury (MRI revealed no obvious circumscribed lesion), it is not possible to localize the sources of his impairments. Nevertheless, these results suggest that the neurocognitive mechanisms responsible for processing upright aligned faces dissociate from those mediating recognition of inverted or horizontally misaligned faces (see discussion of gated processing in *Section 6.1*).

5. The susceptibility of clinical populations to the illusion

5.1 Developmental Prosopagnosia

Individuals with Developmental Prosopagnosia (DP) experience life-long face recognition difficulties, despite normal levels of intelligence and typical low-level vision (Behrmann & Avidan, 2005; Cook & Biotti, 2016; Duchaine & Nakayama, 2006b; Susilo & Duchaine, 2013). DPs show reduced susceptibility to both the identity (Avidan, Tanzer, & Behrmann, 2011; Palermo et al., 2011; but see Susilo et al., 2010), and expression (Palermo et al., 2011)

variants of the composite face illusion, assessed using original matching procedures. However, DP is thought to be a heterogeneous condition (e.g., Susilo & Duchaine, 2013), and cases have been described who show broadly typical composite face effects on simultaneous variants of the original matching task (Le Grand et al., 2006). The contribution of task differences (e.g., simultaneous vs. sequential matching paradigms) to the discrepant findings remains unclear. We also note that several studies employing the original matching design fail to report participants' performance on different trials (e.g., Avidan et al., 2011; Palermo et al., 2011). While individual differences in illusion susceptibility may be most pronounced in the incongruent-same condition (e.g., Le Grand et al., 2004), failure to report observers' performance when targets differ (typically 50% of trials) makes it impossible to assess the potential contribution of response bias to group differences, where observed.

5.2 Autism Spectrum Disorder

It has been suggested that people with Autism Spectrum Disorders (ASD) may exhibit reduced susceptibility to the illusion. Observers with ASD often focus on local features and may therefore experience problems forming integrated global representations (Happé & Frith, 2006). Moreover, many individuals with ASD often exhibit reduced sensitivity to illusions induced by contextual influence (Behrmann, Thomas, & Humphreys, 2006; Simmons et al., 2009). Research examining composite face effects in ASD has yielded mixed results (Weigelt, Koldewyn, & Kanwisher, 2012). High-functioning adults with ASD showed broadly typical composite face effects to age- and IQ-matched controls on a sequential matching task (Nishimura, Rutherford, & Maurer, 2008). In contrast, a sample of adolescents with ASD failed to show the typical composite effect on a sequential matching task; their matching ability was very similar in the upright-aligned and upright-misaligned conditions (Teunisse & de Gelder, 2003). Whereas the foregoing studies employed the original matching design, Gauthier, Klaiman and Schultz (2009) found evidence of atypical composite face effects in a sample of adolescents with ASD using the complete design; the distractor halves induced similar interference in both aligned and misaligned conditions.

5.3 Schizophrenia

There is growing interest in the face recognition of observers with schizophrenia. In particular there has been suggestion that they may exhibit reduced holistic representation (Bortolon, Capdevielle, & Raffard, 2015). Schwartz, Marvel, Drapalski, Rosse and Deutsch (2002) utilized the composite face paradigm to examine holistic processing in a sample of 19

medicated observers with schizophrenia. Using the naming paradigm (A. W. Young et al., 1987), the authors found a comparable composite effect (i.e., greater composite interference in the aligned, than in the misaligned condition) in their clinical sample and matched controls, suggesting typical susceptibility to the illusion. However, in this particular investigation the authors utilized a misaligned control condition only; it is unclear whether the misalignment advantage is also seen for inverted composite arrangements (McKone et al., 2013).

6. Theoretical interpretations

Many explanations of the composite face effect appeal to the idea that upright aligned composites gain access to high-level face-specific processing, whereas misaligned and inverted composites do not. In contrast, some authors have challenged this view and have advanced domain-general accounts (*Section 6.1*). Domain-specific accounts have been heavily influenced by the concepts of configural and holistic representations (*Section 6.2*). However, while these ideas remain valuable theoretical heuristics, they are under-specified as theories in their own right (Burton, Schweinberger, Jenkins, & Kaufmann, 2015; Piepers & Robbins, 2013; Richler, Palmeri, & Gauthier, 2012). Drawing on the computer vision literature, attempts have therefore been made to provide instantiated image-processing accounts of the composite face illusion (*Section 6.3*).

6.1 Domain specific vs. domain general processing

Faces may recruit additional perceptual processing not engaged by other classes of object (Kanwisher, 2000; McKone, Kanwisher, & Duchaine, 2007; McKone & Robbins, 2011). However, employing resource-intensive processing indiscriminately is inefficient. Some stages of face processing may therefore be gated, whereby sophisticated domain-specific processing only commences once a face has been detected in the environment (Tsao & Livingstone, 2008). Detection of simple, proto-facial features, or ‘faciotopy’ (Henriksson, Mur, & Kriegeskorte, 2015), possibly mediated by subcortical visual processing, may be sufficient to engage more complex cortical processing (Johnson, 2005; Shah, Gaule, Bird, & Cook, 2013). When composite faces are aligned and presented upright, the presence of the intact facial arrangement may therefore permit access to face-specific processing, responsible for the composite illusion. Inverted or misaligned composites may lack the basic faciotopy necessary to gain access to the highest levels of face-specific processing, and therefore do not induce composite interference (Tsao & Livingstone, 2008). It is striking that these manipulations also disrupt the residual face recognition ability seen in patient CK (see

Section 4.2). Potential accounts of this high-level face processing are described in the following sections (see *Sections 6.2* and *6.3*).

In contrast, some authors reject the idea that the composite-face illusion is a product of face-specific representation mechanisms; rather, they argue that composite interference reflects a form of automatic processing recruited by ‘objects of expertise’. Objects of expertise are categories of objects i) with which the observer has extensive visual experience, and ii) comprising exemplars that share a common prototypical feature arrangement. Thus, birds (Gauthier, Skudlarski, Gore, & Anderson, 2000) and dogs (Diamond & Carey, 1986) may be objects of expertise for ornithologists and dog show judges, respectively. Following extensive experience individuating exemplars, observers are thought to process the separate features of objects of expertise as a unified whole (Diamond & Carey, 1986; Richler et al., 2012; Richler, Wong, & Gauthier, 2011; A. C. Wong, Palmeri, & Gauthier, 2009).

Consistent with this domain-general account, composite effects obtained using the complete design have been reported for non-face objects of expertise encountered naturally, including cars (Bukach, Phillips, & Gauthier, 2010; Gauthier, Curran, Curby, & Collins, 2003), words (A. C. Wong et al., 2011), Chinese characters (A. C. Wong et al., 2012), and chess boards (Bogdan, Bartlett, & Krawczyk, 2012). Similar composite effects have been reported with synthetic non-face objects, including ‘Greebles’ (Gauthier & Tarr, 2002; Gauthier, Williams, Tarr, & Tanaka, 1998) and ‘Ziggerins’ (A. C. Wong et al., 2009), where expertise is acquired through lab-based training. Crucially, the type of training administered appears to modulate illusion susceptibility. Participants who received individuation training – naming particular exemplars – exhibited greater composite effects than those who received categorisation training – judging which group a given exemplar belonged to (A. C. Wong et al., 2009). There have been relatively few reports of composite effects for non-face objects obtained using the original matching procedure. For example, Robbins and McKone (2007) failed to find a composite effect when dog experts were presented with arrangements constructed from dog stimuli. Similarly, Greeble experts failed to show composite effects for Greebles (Gauthier & Tarr, 2002; Gauthier et al., 1998). Recently, however, a study found evidence of composite effects for body postures using the original procedure (Willems, Vrancken, Germeys, & Verfaillie, 2014), possibly reflecting similarities in the way that faces and bodies are processed (Minnebusch & Daum, 2009; see *Section 7.4*).

6.2 Configural and holistic processing

It has been widely suggested that upright faces may recruit a rapid parallel analysis of the whole face. The terms ‘holistic’ and ‘configural’ are often used interchangeably to describe this whole-face processing, leading to some confusion in the literature (e.g., Piepers & Robbins, 2013). However, two broadly distinct accounts may be delineated.

Faces are defined by the presence and prototypical arrangement of certain basic features; i.e., two eyes above a nose and mouth, so-called first-order relations. Because all faces share this common arrangement, the spatial distances between internal features, so-called second-order relations, may be particularly important for the recognition of individual faces (Diamond & Carey, 1986). The processing of these second-order relations has been commonly termed ‘configural’ processing. It is possible that the composite face illusion alters configural processing; for example, when composite faces are presented upright and aligned, observers may perceive a novel configuration that hinders recognition of the constituent parts (Hancock, Burton, & Bruce, 1996; A. W. Young et al., 1987). Consistent with this interpretation, observers’ ability to discriminate faces that share common features and differ only in their spatial arrangement (e.g., eyes close together or far apart), is greatly diminished when stimuli are inverted (Barton, Keenan, & Bass, 2001; Freire, Lee, & Symons, 2000; Goffaux & Rossion, 2006; Haig, 1984; Leder & Bruce, 2000; Leder, Candrian, Huber, & Bruce, 2001; Rhodes, Brake, & Atkinson, 1993; Searcy & Bartlett, 1996).

Alternatively, variation in feature shape and the spatial relations may be described within a single non-decomposable holistic representation – a ‘Gestalt’ that cannot be broken down into its constituent parts or the inter-relations between them (Farah et al., 1998). Because different distractor regions induce the modelling of a new face Gestalt, they alter the appearance of the target half. This view accords with recent findings that the composite face illusion alters observers’ perception of feature shape as well as feature configurations (Hayward, Crookes, Chu, Favelle, & Rhodes, 2016). This holistic account is also consistent with the part-whole effect, whereby individual features are judged more accurately in the context of an upright face than when shown in isolation, despite the context remaining uninformative (Tanaka & Farah, 1993). Interestingly, this contextual advantage disappears when stimulus arrangements are shown upside-down. Similarly, individuals’ ability to discriminate exemplars that differ only in terms of i) their features or ii) their inter-feature

spacing, are highly-correlated for upright faces, but not for inverted faces or non-face objects (Yovel & Kanwisher, 2008).

In an attempt to reconcile these views, it has been proposed that holistic, configural, and piecemeal ('parts-based') processing, are points on a processing continuum defined by degree of feature integration (Reed, Stone, Grubb, & McGoldrick, 2006). However, the distinction between holistic and configural processing remains contentious. Changing the spatial relations between features necessarily changes the Gestalt representation hypothesised by the holistic account. The distinction between the accounts therefore relies on the assumption that feature shape may be changed independently of the spatial relations between the features. While this view is accepted by some authors (Farah et al., 1998; Yovel & Kanwisher, 2008), it is disputed by others on the ground that changing local contour and shading cues also changes the configural relations between features (Hancock et al., 1996; McKone & Yovel, 2009; Piepers & Robbins, 2013).

6.3 Image processing models

According to the Gabor jet model (Biederman & Kalocsais, 1997; Wiskott, Fellous, Kuiger, & Von Der Malsburg, 1997), faces are described by columns of multi-scale, multi-orientation filters (Figure 3a), with their receptive fields centered on facial landmarks defined relative to the first-order features (Figure 3b). The model retains biological plausibility, insofar as a Gabor jet approximates the multiscale, multiorientation tuning properties of cells in a V1 hypercolumn (De Valois & De Valois, 1988). For a given face, a Gabor jet centered on the left pupil, with different filters for eight orientations and five scales would yield 40-item vectors describing the contrast variation around the left eye (Figure 3c). Face recognition is achieved by comparing the combined readout from all of the Gabor jets, with stored templates for known faces. Importantly, variation in one part of the face will affect kernels with medium and large receptive fields centered on other parts of the face (Xu, Biederman, & Shah, 2014). Where aligned composite faces are shown upright, image variation present in the lower face half will therefore alter the readout of Gabor Jets centered on features in the upper half, biasing observers' perception of its identity (Herald, Shah, Xu, Biederman, & Juarez, 2015).




Figure 3: (a) Illustration of a Gabor Jet, comprising different multi-scale, multi-orientation filters. (b) The receptive fields of the filters in a particular Gabor Jet are centered on a particular facial landmark. The receptive fields of the intermediate and larger scale filters overlap and cover wide areas of the face. (c) Gabor jets comprising n filters yield a vector of n items describing the contrast variation at the particular facial landmark sampled by the Gabor Jet.

There has also been interest in the utility of unsupervised data reduction algorithms as models of human face perception. Following an initial pre-processing stage – input images are typically cropped to a common aspect ratio and faces aligned using sets of fiducial points – algorithms are able to extract dimensions that describe the variation within a set of input images. To date, many authors have used Principal Components Analysis (PCA) to illustrate the value of this approach (Calder, 2011; Calder, Burton, Miller, Young, & Akamatsu, 2001; Calder & Young, 2005; Hancock et al., 1996). When applied to faces, PCA returns the n Eigenface dimensions (*whole face* components that may be combined linearly to reconstruct a given facial exemplar) that most effectively describe the variation present within the set of sampled images (Turk & Pentland, 1991). Thereafter, facial exemplars are represented as vectors in the facespace defined by the n Eigenface dimensions. Importantly, where PCA is applied only to intact, naturalistic faces, components will reflect the natural covariation present in different face halves; for example, happy and angry mouth regions will have been encountered in the presence of accompanying happy and angry eye-regions. Unless the algorithm encounters faces with different emotion signals in their top and bottom halves, the resulting PCA space will lack the dimensionality with which to represent mixed-emotion composites (Cottrell, Branson, & Calder, 2002). The illusion-induced interference first reported by Calder et al., (2000) may therefore reflect the process of modelling emotion-

incongruent composites within an emotion-congruent face-space. By appealing to the same logic, PCA models can explain other types of composite binding such as that seen for age and gender (see *Section 2.4*). Many of the principles and predictions derived from PCA models generalize well to other data reduction algorithms, notably Architecture II Independent Components Analysis (ICA; Bartlett, Movellan, & Sejnowski, 2002; Calder & Young, 2005; Nestor, Plaut, & Behrmann, 2013).

7. Future directions

In the foregoing sections, we have reviewed the body of empirical findings reported with composite paradigms and existing theoretical accounts of the illusion. In this section, we look forward, highlighting priorities for future research. In particular, several critical questions remain unresolved. These gaps in our knowledge undermine attempts to evaluate and refine theoretical accounts of the composite face illusion.

7.1 The functional significance of the composite face illusion

The view that holistic representation, as measured by the composite face illusion, is causally related to face recognition ability remains extremely popular; for example, holistic processing may permit accurate and efficient representation (Maurer et al., 2002; Piepers & Robbins, 2013). However, the functional significance of the composite face illusion remains uncertain. First, the literature is inconsistent with respect to the relationship between individuals' susceptibility to the composite face illusion and other markers of holistic representation, including the part-whole (Tanaka & Farah, 1993) and face-inversion (Yin, 1969) effects. Whilst some authors have found associations between these measures using the complete design (DeGutis, Wilmer, Mercado, & Cohan, 2013), others using the original matching procedure have not (Durand et al., 2007; Wang, Li, Fang, Tian, & Liu, 2012). For example, Wang and colleagues (2012) found susceptibility to the composite and part-whole effects were unrelated. Second, studies comparing observers' susceptibility to the composite face illusion and face recognition ability have also yielded equivocal results; whilst some studies have found little or no correlation using the original matching procedure (de Heering & Maurer, 2014; Konar et al., 2010; Wang et al., 2012), other authors have observed associations with ability using the complete (DeGutis et al., 2013) and original procedures (Avidan et al., 2011). It is likely that differences between the composite tasks used (e.g., Richler, Cheung, & Gauthier, 2011b), their analyses (e.g., DeGutis et al., 2013), the sensitivity of face recognition measures (e.g., Duchaine & Nakayama, 2006a), and sample

composition (e.g., Avidan et al., 2011), all contribute to the discrepant findings. It is imperative that future research teases apart these sources of variation to understand the functional significance of the illusion.

7.2 Strategic and contextual influences

The composite face illusion is frequently attributed to automatic holistic processing – either mediated by face-specific mechanisms or a product of perceptual expertise (see *Section 6*). Importantly, however, several results suggest that strategic and contextual factors influence susceptibility to the illusion. For example, Navon priming manipulations that bias observers' attention towards global form increase susceptibility to the composite-face illusion measured using the complete design (Gao, Flevaris, Robertson, & Bentin, 2011). Conversely, induction of negative mood (Curby, Johnson, & Tyson, 2012) and the presence of unrelated observers (Garcia-Marques, Fernandes, Fonseca, & Prada, 2015), were found to decrease illusion susceptibility when measured using the original matching procedure. In a study employing the complete design, greater distractor interference was observed for composites made from unfamiliar non-face objects when trials were preceded by an aligned composite face (Richler, Bukach, & Gauthier, 2009). Aligned composites also induce more interference when preceded by misaligned composites – possibly reflecting the creation of a larger 'attentional window' that biases subsequent perceptual processing – than when preceded by aligned composites (Richler et al., 2009). Where observed, composite effects for unfamiliar non-face objects (e.g., Hsiao & Cottrell, 2009), may be a product of contextual and strategic influences, whereas composite effects observed with objects-of-expertise, including faces, may be the product of increasingly automatic holistic processing (Richler, Wong et al., 2011). Understanding how strategic and contextual factors influence performance on composite paradigms will inform investigation into the perceptual origins of the illusion (Richler, Cheung et al., 2011a; Rossion, 2013).

7.3 Top-down or bottom up?

There is growing interest in how flexible strategic factors, including attention and response base-rates, can affect the size of composite face effects (*Section 7.2*). A related question is the extent to which the illusion itself is a product of 'bottom-up' feed-forward processing of upright aligned composite arrangements, or a 'top-down' interpretation imposed on a sensory description (see Gregory, 1997). Several theoretical approaches, including the Gabor Jet (Xu et al., 2014) and PCA accounts (Cottrell et al., 2002), model composite interference as an

emergent property of feed-forward descriptive processes. Alternatively, however, the composite illusion may be a product of a top-down modelling process. The goal of visual perception is to infer the causes of sensory input; percepts can be thought of as inferences from sensory data and knowledge derived from the past (Friston, 2005; Gregory, 1997; Kersten, Mamassian, & Yuille, 2004). The generative models we have acquired for faces will likely reflect the whole-face covariation we experience during the course of our lives. These whole-face models will aid perception of ambiguous forms or expressions under typical conditions, but will tend to generate holistic illusions when inputs come from different faces.




Figure 4: Compelling demonstrations of the composite illusion can be provided using highly abstract cartoon-face stimuli. In the upright-aligned condition (left), the presence of the difference mouths makes it harder to recognise that the eyes are the same. The illusion is diminished in the misaligned (middle) and inverted (right) conditions.

While little attempt has been made to distinguish these possibilities, several lines of evidence suggest that a top-down account is plausible. For example, should the composite illusion be an emergent property of feed-forward processing, some degree of illusory bias might be expected where distractor regions are masked and fail to reach conscious awareness. However, studies employing continuous flash suppression suggest that distractor face regions do not induce illusory bias when presented outside of conscious awareness (Axelrod & Rees, 2015). Conversely, when observers are consciously aware of stimulus arrangements, a surprising array of stimuli successfully elicit the composite illusion. For example, photographic negation drastically distorts the appearance of faces (Galper, 1970); in particular, the inversion of shape-from-shading cues, whereby patterns of shading are used to infer three dimensional form, give negated faces a grotesque appearance (Kemp, Pike, White, & Musselman, 1996). While this manipulation might be expected to severely hamper feed-forward processing, studies employing the original matching paradigm suggest that

arrangements shown in typical polarity and photographic negative induce broadly similar composite effects (Hole et al., 1999; Taubert & Alais, 2011). Similarly, compelling demonstrations of the composite illusion can be achieved using abstract cartoon faces (Figure 4). While cartoon and naturalistic faces may share conceptual characteristics, they bear little physical resemblance. Knowledge about faces therefore appears to modulate the composite face illusion in the absence of typical appearance.

7.4 Not all facial composites are created equal

While considerable research has addressed differences in individuals' susceptibility to the composite face illusion (Avidan et al., 2011; DeGutis et al., 2013; Richler, Cheung et al., 2011b; Wang et al., 2012), there is a paucity of research examining inter-stimulus differences. While it is recognised that some facial composites induce stronger illusion-induced interference than others (e.g., Richler & Gauthier, 2014; Ross, Richler, & Gauthier, 2015), little explanation has been offered for these differences. A better understanding of these differences might inform theoretical accounts; for example, it will be interesting to see whether image processing approaches (e.g. Gabor Jet and PCA accounts; see *Section 6.3*) can model this inter-stimulus variability. Moreover, an appreciation of inter-stimulus-variability may help disambiguate some of the equivocal aspects of the literature (e.g., the functional significance of the illusion, findings from clinical populations). Some of the variability may reflect low-level image differences including image scale, spatial frequency and colour information (see *Section 2.5*). Differences in shape and texture variation may also be important. For example, in original matching paradigms, distractors with incongruent shape but congruent texture exert more influence on a target region, than distractors with incongruent texture but congruent shape (Jiang, Blanz, & Rossion, 2011).

A further possibility is that observers are detecting emotion cues in some of the ostensibly 'neutral' faces used to construct facial composites (Figure 5). When posing for photos, actors seeking to appear neutral may appear anxious or bored. Moreover, it is not always easy to distinguish a stranger's permanent facial shape from their transient facial expressions; e.g., whether an unfamiliar actor is scowling or simply has narrow eyes (e.g., Todorov, Said, Engell, & Oosterhof, 2008). Crucially, observers may therefore perceive emotion where actors do not intend to convey emotion. It is well established that facial emotions induce strong composite effects (Calder et al., 2000; Palermo et al., 2011; Tanaka et al., 2012). Facial composites rich in perceived emotion cues may therefore induce stronger composite

effects when observers are asked to judge whether target regions are, or are not identical (e.g., in original or complete matching paradigms). Interestingly, gaze direction – known to modulate the perception of facial emotion; for example direct gaze makes expressions appear angrier (Adams & Kleck, 2003, 2005) – has been found to influence the magnitude of composite effects measured using the original matching design (S. G. Young, Slepian, Wilson, & Hugenberg, 2014). We also note that a study employing the original matching procedure recently found composite effects with expressive body postures (Willems et al., 2014), suggestive of holistic coding of body posture. In contrast, authors have failed to observe composite effects with neutral bodies (e.g., Soria Bauser, Suchan, & Daum, 2011).



Figure 5: Examples of facial composites taken from a commonly used stimulus set developed by Le Grand et al. (2004). While the composites are constructed with ostensibly neutral faces, subtle emotion cues are present in many of the faces.

7.5 The nature of holistic information representation

The composite face illusion is frequently attributed to whole-face processing (see *Section 6.1*). However, surprisingly little is known about the nature of the integrated representations derived from this analysis. At least two lines of evidence suggest that current ‘holistic’ and ‘configural’ accounts may be under-specified. First, observers’ face recognition is surprisingly insensitive to vertical and horizontal stretching; for example observers can easily

recognise faces stretched to twice their true height (Hole, George, Eaves, & Rasek, 2002). Whole-face processing is thought to mediate accurate and efficient description of the spatial relationships between features (see Rossion, 2008). However, such findings suggest that spatial relations may be coded relatively – for example, the height of one feature may be described through reference to the height of another – not through absolute metrics – where horizontal and vertical distances are measured in units akin to degrees of visual angle (Burton et al., 2015). Second, recent findings suggest that, in comparison to vertical information, the horizontal information structure of faces contributes disproportionately to face recognition (Dakin & Watt, 2009) and is disproportionately sensitive to orientation inversion (Goffaux & Dakin, 2010). These findings raise the possibility that holistic representations may be dominated by horizontal information. Composite face paradigms may offer researchers a way to test these hypotheses.

7.6 Dynamic Face Processing

While the overwhelming majority of existing research has examined the composite face illusion using static faces, the faces we typically encounter outside of the lab are moving (O'Toole, Roark, & Abdi, 2002). A recent study employing the naming paradigm found that the presence of an aligned dynamic distractor impairs identification of dynamic target halves, learned beforehand (Favelle, Tobin, Piepers, Burke, & Robbins, 2015). Importantly, composite interference is diminished when the dynamic halves are misaligned or presented upside-down. Similarly, when asked to judge the speed of eye opening and closing movements, the presence of task-irrelevant mouth opening and closing induces illusory slowing of the eye changes (Cook, Aichelburg, & Johnston, 2015). Interestingly, this illusion-induced interference is only seen in upright dynamic faces, at particular relative-phase relationships. Whereas static composite effects are disrupted by spatial misalignment, phase-specificity suggests that dynamic integration may be sensitive to temporospatial alignment.

It is important that future research addresses the relationship between composite interference seen with dynamic faces and the recognition of moving faces. The presence of motion cues often improves facial recognition (Knight & Johnston, 1997; Lander, Christie, & Bruce, 1999). However, observers' ability to use motion cues is sensitive to orientation, suggestive of holistic or configural representation (Hill & Johnston, 2001; Knight & Johnston, 1997). Interestingly, observers with ASD are relatively poor at recognising individuals' facial motion signature (O'Brien, Spencer, Girges, Johnston, & Hill, 2014) and show little or no

susceptibility to the illusory slowing induced by dynamic cross-feature interactions (Shah, Bird, & Cook, 2016). Moreover, the relationship between the dynamic and static composites remains unclear; while both illusions exhibit sensitivity to alignment and orientation, it is uncertain whether they are products of common processes or representations.

8. Conclusion

Results obtained from composite face procedures have contributed significantly to our understanding of holistic face processing, the detrimental effects of inversion, and aberrant face perception in clinical populations. The ongoing value of the paradigm is illustrated by its recent application to investigate dynamic face processing. However, composite procedures have been the subject of intense scrutiny, particularly over the last decade, and there is a growing sense that the composite face illusion, whilst easy to illustrate, is deceptively difficult to measure and interpret. Considerable debate has focussed on the use of original and complete designs (Richler & Gauthier, 2014; Richler et al., 2012; Rossion, 2013), control conditions (McKone et al., 2013), the analyses employed (DeGutis et al., 2013), and the effects of strategic and contextual factors (Richler, Wong et al., 2011). We have suggested that inter-stimulus variability may also affect the susceptibility estimates obtained by different authors and warrants greater consideration. Despite overarching disagreements on how to refine the paradigm, the composite face effect remains a compelling and fascinating visual illusion. As a tool, it still holds much potential in our investigation of how faces and other objects of expertise are processed.

References

- Abbas, Z. A., & Duchaine, B. (2008). The role of holistic processing in judgments of facial attractiveness. *Perception, 37*(8), 1187-1196.
- Adams, R. B., & Kleck, R. E. (2003). Perceived gaze direction and the processing of facial displays of emotion. *Psychological Science, 14*(6), 644-647.
- Adams, R. B., & Kleck, R. E. (2005). Effects of direct and averted gaze on the perception of facially communicated emotion. *Emotion, 5*(1), 3-11.
- Avidan, G., Tanzer, M., & Behrmann, M. (2011). Impaired holistic processing in congenital prosopagnosia. *Neuropsychologia, 49*(9), 2541-2552.
- Axelrod, V., & Rees, G. (2015). Conscious awareness is required for holistic face processing. *Consciousness and Cognition, 27*, 233-245.
- Bartlett, M. S., Movellan, J. R., & Sejnowski, T. J. (2002). Face recognition by independent component analysis. *IEEE Transactions on Neural Networks, 13*(6), 1450-1464.
- Barton, J. J., Keenan, J. P., & Bass, T. (2001). Discrimination of spatial relations and features in faces: Effects of inversion and viewing duration. *British Journal of Psychology, 92*(3), 527-549.
- Baudouin, J. Y., & Humphreys, G. W. (2006). Configural information in gender categorisation. *Perception, 35*(4), 531-540.
- Behrmann, M., & Avidan, G. (2005). Congenital prosopagnosia: face-blind from birth. *Trends in Cognitive Sciences, 9*(4), 180-187.
- Behrmann, M., Thomas, C., & Humphreys, K. (2006). Seeing it differently: visual processing in autism. *Trends in Cognitive Sciences, 10*(6), 258-264.
- Biederman, I., & Kalocsais, P. (1997). Neurocomputational bases of object and face recognition. *Philosophical Transactions of the Royal Society B- Biological Sciences, 352*(1358), 1203-1219.

- Boggan, A. L., Bartlett, J. C., & Krawczyk, D. C. (2012). Chess masters show a hallmark of face processing with chess. *Journal of Experimental Psychology: General*, *141*(1), 37-42.
- Bortolon, C., Capdevielle, D., & Raffard, S. (2015). Face recognition in schizophrenia disorder: A comprehensive review of behavioral, neuroimaging and neurophysiological studies. *Neuroscience and Biobehavioral Reviews*, *53*, 79-107.
- Bukach, C. M., Bub, D. N., Gauthier, I., & Tarr, M. J. (2006). Perceptual expertise effects are not all or none: Spatially limited perceptual expertise for faces in a case of prosopagnosia. *Journal of Cognitive Neuroscience*, *18*(1), 48-63.
- Bukach, C. M., Phillips, W. S., & Gauthier, I. (2010). Limits of generalization between categories and implications for theories of category specificity. *Attention, Perception, & Psychophysics*, *72*(7), 1865-1874.
- Burton, A. M., Schweinberger, S. R., Jenkins, R., & Kaufmann, J. M. (2015). Arguments against a configural processing account of familiar face recognition. *Perspectives on Psychological Science*, *10*(4), 482-496.
- Busigny, T., Joubert, S., Felician, O., Ceccaldi, M., & Rossion, B. (2010). Holistic perception of the individual face is specific and necessary: evidence from an extensive case study of acquired prosopagnosia. *Neuropsychologia*, *48*(14), 4057-4092.
- Busigny, T., Van Belle, G., Jemel, B., Hosein, A., Joubert, S., & Rossion, B. (2014). Face-specific impairment in holistic perception following focal lesion of the right anterior temporal lobe. *Neuropsychologia*, *56*, 312-333.
- Calder, A. J. (2011). Does facial identity and facial expression recognition involve separate visual routes? In A. J. Calder, G. Rhodes, J. V. Haxby & M. H. Johnson (Eds.), *The Oxford Handbook of Face Perception* Oxford, UK: Oxford University Press.
- Calder, A. J., Burton, A. M., Miller, P., Young, A. W., & Akamatsu, S. (2001). A principal component analysis of facial expressions. *Vision Research*, *41*(9), 1179-1208.
- Calder, A. J., & Young, A. W. (2005). Understanding the recognition of facial identity and facial expression. *Nature Reviews Neuroscience*, *6*(8), 641-651.

- Calder, A. J., Young, A. W., Keane, J., & Dean, M. (2000). Configural information in facial expression perception. *Journal of Experimental Psychology: Human Perception and Performance*, 26(2), 527-551.
- Carey, S., & Diamond, R. (1994). Are faces perceived as configurations more by adults than by children? *Visual Cognition*, 1, 253-274.
- Cassia, V. M., Picozzi, M., Kuefner, D., Bricolo, E., & Turati, C. (2009). Holistic processing for faces and cars in preschool-aged children and adults: evidence from the composite effect. *Developmental Science*, 12(2), 236-248.
- Cheung, O. S., Richler, J. J., Palmeri, T. J., & Gauthier, I. (2008). Revisiting the role of spatial frequencies in the holistic processing of faces. *Journal of Experimental Psychology: Human Perception and Performance*, 34(6), 1327-1336.
- Cook, R., Aichelburg, C., & Johnston, A. (2015). Illusory feature slowing: evidence for perceptual models of global facial change. *Psychological Science*, 26(4), 512-517.
- Cook, R., & Biotti, F. (2016). Developmental prosopagnosia. *Current Biology*, 26(8), R312-R313.
- Cottrell, G. W., Branson, K. M., & Calder, A. J. (2002). *Do expression and identity need separate representations*. Paper presented at the Proceedings of the 24th Annual Conference of the Cognitive Science Society.
- Curby, K. M., Johnson, K. J., & Tyson, A. (2012). Face to face with emotion: holistic face processing is modulated by emotional state. *Cognition & Emotion*, 26(1), 93-102.
- Dakin, S. C., & Watt, R. J. (2009). Biological "bar codes" in human faces. *Journal of Vision*, 9(4), 2.
- de Heering, A., Houthuys, S., & Rossion, B. (2007). Holistic face processing is mature at 4 years of age: Evidence from the composite face effect. *Journal of Experimental Child Psychology*, 96(1), 57-70.
- de Heering, A., & Maurer, D. (2014). Face memory deficits in patients deprived of early visual input by bilateral congenital cataracts. *Developmental Psychobiology*, 56(1), 96-108.

- de Heering, A., & Rossion, B. (2008). Prolonged visual experience in adulthood modulates holistic face perception. *PLoS One*, 3(5), e2317.
- De Valois, R. L., & De Valois, K. K. (1988). *Spatial Vision*. Oxford, UK: Oxford University Press.
- DeGutis, J., Wilmer, J., Mercado, R. J., & Cohan, S. (2013). Using regression to measure holistic face processing reveals a strong link with face recognition ability. *Cognition*, 126(1), 87-100.
- Diamond, R., & Carey, S. (1986). Why faces are and are not special: an effect of expertise. *Journal of Experimental Psychology: General*, 115(2), 107-117.
- Duchaine, B., & Nakayama, K. (2006a). The Cambridge Face Memory Test: results for neurologically intact individuals and an investigation of its validity using inverted face stimuli and prosopagnosic participants. *Neuropsychologia*, 44(4), 576-585.
- Duchaine, B., & Nakayama, K. (2006b). Developmental prosopagnosia: a window to content-specific face processing. *Current Opinion in Neurobiology*, 16, 166-173.
- Durand, K., Gallay, M., Seigneuric, A., Robichon, F., & Baudouin, J. Y. (2007). The development of facial emotion recognition: The role of configural information. *Journal of Experimental Child Psychology*, 97(1), 14-27.
- Eimer, M. (2000). Effects of face inversion on the structural encoding and recognition of faces. Evidence from event-related brain potentials. *Cognitive Brain Research*, 10(1-2), 145-158.
- Farah, M. J., Wilson, K. D., Drain, M., & Tanaka, J. N. (1998). What is "special" about face perception? *Psychological Review*, 105(3), 482-498.
- Favelle, S., Tobin, A., Piepers, D., Burke, D., & Robbins, R. A. (2015). Dynamic composite faces are processed holistically. *Vision Research*, 112, 26-32.
- Freire, A., Lee, K., & Symons, L. A. (2000). The face-inversion effect as a deficit in the encoding of configural information: direct evidence. *Perception*, 29(2), 159-170.
- Friston, K. (2005). A theory of cortical responses. *Philosophical Transactions of the Royal Society B-Biological Sciences*, 360(1456), 815-836.

- Galper, R. E. (1970). Recognition of faces in photographic negative. *Psychonomic Science*, 19(4), 207-208.
- Gao, Z., Flevaris, A. V., Robertson, L. C., & Bentin, S. (2011). Priming global and local processing of composite faces: revisiting the processing-bias effect on face perception. *Attention, Perception, & Psychophysics*, 73(5), 1477-1486.
- Garcia-Marques, T., Fernandes, A., Fonseca, R., & Prada, M. (2015). Social presence and the composite face effect. *Acta Psychologica*, 158, 61-66.
- Gauthier, I., Curran, T., Curby, K. M., & Collins, D. (2003). Perceptual interference supports a non-modular account of face processing. *Nature Neuroscience*, 6(4), 428-432.
- Gauthier, I., Klaiman, C., & Schultz, R. T. (2009). Face composite effects reveal abnormal face processing in Autism spectrum disorders. *Vision Research*, 49(4), 470-478.
- Gauthier, I., Skudlarski, P., Gore, J. C., & Anderson, A. W. (2000). Expertise for cars and birds recruits brain areas involved in face recognition. *Nature Neuroscience*, 3(2), 191-197.
- Gauthier, I., & Tarr, M. J. (2002). Unraveling mechanisms for expert object recognition: bridging brain activity and behavior. *Journal of Experimental Psychology: Human Perception and Performance*, 28(2), 431-446.
- Gauthier, I., Williams, P., Tarr, M. J., & Tanaka, J. W. (1998). Training 'greeble' experts: a framework for studying expert object recognition processes. *Vision Research*, 38(15), 2401-2428.
- Goffaux, V., & Dakin, S. C. (2010). Horizontal information drives the behavioral signatures of face processing. *Frontiers in Psychology*, 1, 143.
- Goffaux, V., & Rossion, B. (2006). Faces are "spatial"--holistic face perception is supported by low spatial frequencies. *Journal of Experimental Psychology: Human Perception and Performance*, 32(4), 1023-1039.
- Gregory, R. L. (1997). Knowledge in perception and illusion. *Philosophical Transactions of the Royal Society B- Biological Sciences*, 352(1358), 1121-1127.

- Haig, N. D. (1984). The effect of feature displacement on face recognition. *Perception, 13*(5), 505-512.
- Hancock, P. J., Burton, A. M., & Bruce, V. (1996). Face processing: human perception and principal components analysis. *Memory and Cognition, 24*(1), 21-40.
- Happe, F., & Frith, U. (2006). The weak coherence account: detail-focused cognitive style in autism spectrum disorders. *Journal of Autism and Developmental Disorders, 36*(1), 5-25.
- Harris, A., & Aguirre, G. K. (2010). Neural tuning for face wholes and parts in human fusiform gyrus revealed by fMRI adaptation. *Journal of Neurophysiology, 104*(1), 336-345.
- Haworth, C. M., Wright, M. J., Luciano, M., Martin, N. G., de Geus, E. J., van Beijsterveldt, C. E., et al. (2010). The heritability of general cognitive ability increases linearly from childhood to young adulthood. *Molecular Psychiatry, 15*(11), 1112-1120.
- Hayward, W. G., Crookes, K., Chu, M. H., Favelle, S. K., & Rhodes, G. (2016). Holistic processing of face configurations and components. *Journal of Experimental Psychology: Human Perception and Performance.*
- Hayward, W. G., Crookes, K., & Rhodes, G. (2013). The other-race effect: Holistic coding differences and beyond. *Visual Cognition, 21*(9-10), 1224-1247.
- Henriksson, L., Mur, M., & Kriegeskorte, N. (2015). Faciotopy-A face-feature map with face-like topology in the human occipital face area. *Cortex, 72*, 156-167.
- Herald, S. B., Shah, M. P., Xu, X., Biederman, I., & Juarez, J. J. (2015). A neurocomputational account of the magnitude of the face offset effect. *Journal of Vision, 15*(12), 419-419.
- Hill, H., & Johnston, A. (2001). Categorizing sex and identity from the biological motion of faces. *Current Biology, 11*(11), 880-885.
- Hole, G. (1994). Configurational factors in the perception of unfamiliar faces. *Perception, 23*(1), 65-74.
- Hole, G., & George, P. (2011). Evidence for holistic processing of facial age. *Visual Cognition, 19*(5), 585-615.

- Hole, G., George, P. A., & Dunsmore, V. (1999). Evidence for holistic processing of faces viewed as photographic negatives. *Perception*, 28(3), 341-359.
- Hole, G., George, P. A., Eaves, K., & Rasek, A. (2002). Effects of geometric distortions on face-recognition performance. *Perception*, 31(10), 1221-1240.
- Horry, R., Cheong, W., & Brewer, N. (2015). The other-race effect in perception and recognition: Insights from the complete composite task. *Journal of Experimental Psychology: Human Perception and Performance*, 41(2), 508-524.
- Hsiao, J. H., & Cottrell, G. W. (2009). Not all visual expertise is holistic, but it may be leftist the case of Chinese character recognition. *Psychological Science*, 20(4), 455-463.
- Jackson, M. C., & Raymond, J. E. (2006). The role of attention and familiarity in face identification. *Perception and Psychophysics*, 68, 543-557.
- Jackson, M. C., & Raymond, J. E. (2008). Familiarity enhances visual working memory for faces. *Journal of Experimental Psychology: Human Perception and Performance*, 34, 556-568.
- Jacques, C., & Rossion, B. (2009). The initial representation of individual faces in the right occipito-temporal cortex is holistic: electrophysiological evidence from the composite face illusion. *Journal of Vision*, 9(6), 1-16.
- Jacques, C., & Rossion, B. (2010). Misaligning face halves increases and delays the N170 specifically for upright faces: implications for the nature of early face representations. *Brain Research*, 1318, 96-109.
- Jenkins, R., & Burton, A. M. (2011). Stable face representations. *Philosophical Transactions of the Royal Society B- Biological Sciences*, 366(1571), 1671-1683.
- Jiang, F., Blanz, V., & Rossion, B. (2011). Holistic processing of shape cues in face identification: Evidence from face inversion, composite faces, and acquired prosopagnosia. *Visual Cognition*, 19(8), 1003-1034.
- Johnson, M. H. (2005). Subcortical face processing. *Nature Reviews Neuroscience*, 6(10), 766-774.
- Kanwisher, N. (2000). Domain specificity in face perception. *Nature Neuroscience*, 3(8), 759-763.

- Kemp, R., Pike, G., White, P., & Musselman, A. (1996). Perception and recognition of normal and negative faces: the role of shape from shading and pigmentation cues. *Perception*, 25(1), 37-52.
- Kersten, D., Mamassian, P., & Yuille, A. (2004). Object perception as Bayesian inference. *Annual Review of Psychology*, 55, 271-304.
- Knight, B., & Johnston, A. (1997). The role of movement in face recognition. *Visual Cognition*, 4(3), 265-273.
- Konar, Y., Bennett, P. J., & Sekuler, A. B. (2010). Holistic processing is not correlated with face-identification accuracy. *Psychological Science*, 21(1), 38-43.
- Kuefner, D., Jacques, C., Prieto, E. A., & Rossion, B. (2010). Electrophysiological correlates of the composite face illusion: Disentangling perceptual and decisional components of holistic face processing in the human brain. *Brain and Cognition*, 74(3), 225-238.
- Lander, K., Christie, F., & Bruce, V. (1999). The role of movement in the recognition of famous faces. *Memory and Cognition* 27, 974-985
- Le Grand, R., Cooper, P. A., Mondloch, C. J., Lewis, T. L., Sagiv, N., de Gelder, B., et al. (2006). What aspects of face processing are impaired in developmental prosopagnosia? *Brain and Cognition*, 61(2), 139-158.
- Le Grand, R., Mondloch, C. J., Maurer, D., & Brent, H. P. (2004). Impairment in holistic face processing following early visual deprivation. *Psychological Science*, 15(11), 762-768.
- Leder, H., & Bruce, V. (2000). When inverted faces are recognized: The role of configural information in face recognition. *Quarterly Journal of Experimental Psychology*, 53(2), 513-536.
- Leder, H., Candrian, G., Huber, O., & Bruce, V. (2001). Configural features in the context of upright and inverted faces. *Perception*, 30(1), 73-83.
- Letourneau, S. M., & Mitchell, T. V. (2008). Behavioral and ERP measures of holistic face processing in a composite task. *Brain and Cognition*, 67(2), 234-245.

- Maurer, D., Le Grand, R., & Mondloch, C. J. (2002). The many faces of configural processing. *Trends in Cognitive Sciences*, 6(6), 255-260.
- McKone, E., Davies, A. A., Darke, H., Crookes, K., Wickramariyaratne, T., Zappia, S., et al. (2013). Importance of the inverted control in measuring holistic face processing with the composite effect and part-whole effect. *Frontiers in Psychology*, 4(33), 1-21.
- McKone, E., Kanwisher, N., & Duchaine, B. C. (2007). Can generic expertise explain special processing for faces? *Trends in Cognitive Sciences*, 11(1), 8-15.
- McKone, E., & Robbins, R. (2011). Are faces special? In A. J. Calder, G. Rhodes, J. V. Haxby & M. H. Johnson (Eds.), *The Oxford Handbook of Face Perception* Oxford, UK: Oxford University Press.
- McKone, E., & Yovel, G. (2009). Why does picture-plane inversion sometimes dissociate the perception of features and spacing in faces, and sometimes not? Toward a new theory of holistic processing. *Psychonomic Bulletin & Review*, 16(5), 778-797.
- Meinhardt, G., Meinhardt-Injac, B., & Persike, M. (2014). The complete design in the composite face paradigm: role of response bias, target certainty, and feedback. *Frontiers in Human Neuroscience*, 8(885), 1-14.
- Michel, C., Rossion, B., Han, J., Chung, C. S., & Caldara, R. (2006). Holistic processing is finely tuned for faces of one's own race. *Psychological Science*, 17(7), 608-615.
- Minnebusch, D. A., & Daum, I. (2009). Neuropsychological mechanisms of visual face and body perception. *Neuroscience and Biobehavioral Reviews*, 33(7), 1133-1144.
- Mondloch, C. J., Pathman, T., Maurer, D., Le Grand, R., & de Schonen, S. (2007). The composite face effect in six-year-old children: Evidence of adult-like holistic face processing. *Visual Cognition*, 15(5), 564-577.
- Moscovitch, M., Winocur, G., & Behrmann, M. (1997). What is special about face recognition? Nineteen experiments on a person with visual object agnosia and dyslexia but normal face recognition. *Journal of Cognitive Neuroscience*, 9(5), 555-604.

- Murphy, J., Ipser, A., Gaigg, S. B., & Cook, R. (2015). Exemplar variance supports robust learning of facial identity. *Journal of Experimental Psychology: Human Perception and Performance*, *41*(3), 577-581.
- Navon, D. (1977). Forest before trees: The precedence of global features in visual perception. *Cognitive Psychology*, *9*(3), 353-383.
- Nestor, A., Plaut, D. C., & Behrmann, M. (2013). Face space architectures: Evidence for the use of independent color-based features. *Psychological Science*, *24*(7), 1294-1300.
- Nishimura, M., Rutherford, M. D., & Maurer, D. (2008). Converging evidence of configural processing of faces in high-functioning adults with autism spectrum disorders. *Visual Cognition*, *16*, 859-891.
- O'Brien, J., Spencer, J., Girges, C., Johnston, A., & Hill, H. (2014). Impaired perception of facial motion in autism spectrum disorder. *PLoS One*, *9*(7), e102173.
- O'Toole, A. J., Roark, D. A., & Abdi, H. (2002). Recognizing moving faces: a psychological and neural synthesis. *Trends in Cognitive Sciences*, *6*(6), 261-266.
- Osborne, C. D., & Stevenage, S. V. (2008). Internal feature saliency as a marker of familiarity and configural processing. *Visual Cognition*, *16*, 23-43.
- Palermo, R., Willis, M. L., Rivolta, D., McKone, E., Wilson, C. E., & Calder, A. J. (2011). Impaired holistic coding of facial expression and facial identity in congenital prosopagnosia. *Neuropsychologia*, *49*(5), 1226-1235.
- Piepers, D. W., & Robbins, R. A. (2013). A Review and Clarification of the Terms "holistic," "configural," and "relational" in the Face Perception Literature. *Frontiers in Psychology*, *3*(559), 1-11.
- Ramon, M., Busigny, T., & Rossion, B. (2010). Impaired holistic processing of unfamiliar individual faces in acquired prosopagnosia. *Neuropsychologia*, *48*(4), 933-944.
- Reed, C. L., Stone, V. E., Grubb, J. D., & McGoldrick, J. E. (2006). Turning configural processing upside down: part and whole body postures. *Journal of Experimental Psychology: Human Perception and Performance*, *32*(1), 73-87.

- Renzi, C., Ferrari, C., Schiavi, S., Pisoni, A., Papagno, C., Vecchi, T., et al. (2015). The role of the occipital face area in holistic processing involved in face detection and discrimination: A tDCS study. *Neuropsychology*, *29*(3), 409-416.
- Retter, T. L., & Rossion, B. (2015). Global shape information increases but color information decreases the composite face effect. *Perception*, *44*(5), 511-528.
- Rezlescu, C., Pitcher, D., & Duchaine, B. (2012). Acquired prosopagnosia with spared within-class object recognition but impaired recognition of degraded basic-level objects. *Cognitive Neuropsychology*, *29*(4), 325-347.
- Rhodes, G., Brake, S., & Atkinson, A. P. (1993). What's lost in inverted faces? *Cognition*, *47*(1), 25-57.
- Richler, J. J., Bukach, C. M., & Gauthier, I. (2009). Context influences holistic processing of nonface objects in the composite task. *Attention, Perception, & Psychophysics*, *71*(3), 530-540.
- Richler, J. J., Cheung, O. S., & Gauthier, I. (2011a). Beliefs alter holistic face processing ... if response bias is not taken into account. *Journal of Vision*, *11*(13), 1-13.
- Richler, J. J., Cheung, O. S., & Gauthier, I. (2011b). Holistic processing predicts face recognition. *Psychological Science*, *22*(4), 464-471.
- Richler, J. J., & Gauthier, I. (2013). When intuition fails to align with data: A reply to Rossion (2013). *Visual Cognition*, *21*(2), 254-276.
- Richler, J. J., & Gauthier, I. (2014). A meta-analysis and review of holistic face processing. *Psychological Bulletin*, *140*(5), 1281-1302.
- Richler, J. J., Gauthier, I., Wenger, M. J., & Palmeri, T. J. (2008). Holistic processing of faces: perceptual and decisional components. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, *34*(2), 328-342.
- Richler, J. J., Mack, M. L., Palmeri, T. J., & Gauthier, I. (2011). Inverted faces are (eventually) processed holistically. *Vision Research*, *51*(3), 333-342.

- Richler, J. J., Palmeri, T. J., & Gauthier, I. (2012). Meanings, mechanisms, and measures of holistic processing. *Frontiers in Psychology, 3*(553), 1-6.
- Richler, J. J., Tanaka, J. W., Brown, D. D., & Gauthier, I. (2008). Why does selective attention to parts fail in face processing? *Journal of Experimental Psychology: Learning, Memory, and Cognition, 34*(6), 1356-1368.
- Richler, J. J., Wong, Y. K., & Gauthier, I. (2011). Perceptual expertise as a shift from strategic interference to automatic holistic processing. *Current Directions in Psychological Science, 20*(2), 129-134.
- Robbins, R., & McKone, E. (2007). No face-like processing for objects-of-expertise in three behavioural tasks. *Cognition, 103*(1), 34-79.
- Ross, D. A., & Gauthier, I. (2015). Holistic processing in the composite task depends on face size. *Visual Cognition, 23*(5), 533-545.
- Ross, D. A., Richler, J. J., & Gauthier, I. (2015). Reliability of composite-task measurements of holistic face processing. *Behavior Research Methods, 47*(3), 736-743.
- Rossion, B. (2008). Picture-plane inversion leads to qualitative changes of face perception. *Acta Psychologica, 128*(2), 274-289.
- Rossion, B. (2013). The composite face illusion: A whole window into our understanding of holistic face perception. *Visual Cognition, 21*(2), 139-253.
- Schiltz, C., Dricot, L., Goebel, R., & Rossion, B. (2010). Holistic perception of individual faces in the right middle fusiform gyrus as evidenced by the composite face illusion. *Journal of Vision, 10*(2), 1-16.
- Schiltz, C., & Rossion, B. (2006). Faces are represented holistically in the human occipito-temporal cortex. *Neuroimage, 32*(3), 1385-1394.
- Schwartz, B. L., Marvel, C. L., Drapalski, A., Rosse, R. B., & Deutsch, S. I. (2002). Configural processing in face recognition in schizophrenia. *Cognitive Neuropsychiatry, 7*(1), 15-39.

- Searcy, J. H., & Bartlett, J. C. (1996). Inversion and processing of component and spatial-relational information in faces. *Journal of Experimental Psychology: Human Perception and Performance*, 22(4), 904-915.
- Sergent, J. (1986). Microgenesis of face perception. In H. D. Ellis, M. A. Jeeves, F. Newcombe & A. M. Young (Eds.), *Aspects of Face Processing* (pp. 17-33). Dordrecht: Martinus Nijhoff.
- Shah, P., Bird, G., & Cook, R. (2016). Face processing in autism: Reduced integration of cross-feature dynamics. *Cortex*, 75, 113-119.
- Shah, P., Gaule, A., Bird, G., & Cook, R. (2013). Robust orienting to protofacial stimuli in autism. *Current Biology*, 23(24), R1087-R1088.
- Simmons, D. R., Robertson, A. E., McKay, L. S., Toal, E., McAleer, P., & Pollick, F. E. (2009). Vision in autism spectrum disorders. *Vision Research*, 49(22), 2705-2739.
- Soria Bauser, D. A., Schriewer, E., & Suchan, B. (2014). Dissociation between the behavioural and electrophysiological effects of the face and body composite illusions. *British Journal of Psychology*, 106(3), 414-432.
- Soria Bauser, D. A., Suchan, B., & Daum, I. (2011). Differences between perception of human faces and body shapes: Evidence from the composite illusion. *Vision Research*, 51(1), 195-202.
- Susilo, T., Crookes, K., McKone, E., & Turner, H. (2009). The composite task reveals stronger holistic processing in children than adults for child faces. *PLoS One*, 4(7), e6460.
- Susilo, T., & Duchaine, B. (2013). Advances in developmental prosopagnosia research. *Current Opinion in Neurobiology*, 23, 423-429.
- Susilo, T., McKone, E., Dennett, H., Darke, H., Palermo, R., Hall, A., et al. (2010). Face recognition impairments despite normal holistic processing and face space coding: Evidence from a case of developmental prosopagnosia. *Cognitive Neuropsychology*, 27(8), 636-664.
- Susilo, T., Rezlescu, C., & Duchaine, B. (2013). The composite effect for inverted faces is reliable at large sample sizes and requires the basic face configuration. *Journal of Vision*, 13(13), 1-9.

- Tanaka, J. W., & Farah, M. J. (1993). Parts and wholes in face recognition. *Quarterly Journal of Experimental Psychology*, *46*(2), 225-245.
- Tanaka, J. W., Kaiser, M. D., Butler, S., & Le Grand, R. (2012). Mixed emotions: holistic and analytic perception of facial expressions. *Cognition & Emotion*, *26*(6), 961-977.
- Taubert, J., & Alais, D. (2011). Identity aftereffects, but not composite effects, are contingent on contrast polarity. *Perception*, *40*(4), 422-436.
- Teunisse, J. P., & de Gelder, B. (2003). Face processing in adolescents with autistic disorder: the inversion and composite effects. *Brain and Cognition*, *52*(3), 285-294.
- Todorov, A., Loehr, V. G., & Oosterhof, N. (2010). The obligatory nature of holistic processing of faces in social judgments. *Perception*, *39*, 514-532
- Todorov, A., Said, C. P., Engell, A. D., & Oosterhof, N. N. (2008). Understanding evaluation of faces on social dimensions. *Trends in Cognitive Sciences*, *12*(12), 455-460.
- Tsao, D. Y., & Livingstone, M. S. (2008). Mechanisms of face perception. *Annual Review of Neuroscience*, *31*, 411-437.
- Turati, C., Di Giorgio, E., Bardi, L., & Simion, F. (2010). Holistic face processing in newborns, 3-month-old infants, and adults: evidence from the composite face effect. *Child Development*, *81*(6), 1894-1905.
- Turk, M., & Pentland, A. (1991). Eigenfaces for recognition. *Journal of Cognitive Neuroscience*, *3*(1), 71-86.
- Wang, R., Li, J., Fang, H., Tian, M., & Liu, J. (2012). Individual differences in holistic processing predict face recognition ability. *Psychological Science*, *23*(2), 169-177.
- Wegrzyn, M., Bruckhaus, I., & Kissler, J. (2015). Categorical perception of fear and anger expressions in whole, masked and composite faces. *PLoS One*, *10*(8), e0134790.
- Weigelt, S., Koldewyn, K., & Kanwisher, N. (2012). Face identity recognition in autism spectrum disorders: A review of behavioral studies. *Neuroscience and Biobehavioral Reviews*, *36*(3), 1060-1084.

- Wiese, H., Kachel, U., & Schweinberger, S. R. (2013). Holistic face processing of own-and other-age faces in young and older adults: ERP evidence from the composite face task. *Neuroimage*, *74*, 306-317.
- Willems, S., Vrancken, L., Germeys, F., & Verfaillie, K. (2014). Holistic processing of human body postures: evidence from the composite effect. *Frontiers in Psychology*, *5*(618), 1-9.
- Wiskott, L., Fellous, J. M., Kuiger, N., & Von Der Malsburg, C. (1997). Face recognition by elastic bunch graph matching. *Pattern Analysis and Machine Intelligence, IEEE Transactions on*, *19*(7), 775-779.
- Wong, A. C., Bukach, C. M., Hsiao, J., Greenspon, E., Ahern, E., Duan, Y., et al. (2012). Holistic processing as a hallmark of perceptual expertise for nonface categories including Chinese characters. *Journal of Vision*, *12*(13:7), 1-15.
- Wong, A. C., Bukach, C. M., Yuen, C., Yang, L., Leung, S., & Greenspon, E. (2011). Holistic processing of words modulated by reading experience. *PLoS One*, *6*(6), e20753.
- Wong, A. C., Palmeri, T. J., & Gauthier, I. (2009). Conditions for facelike expertise with objects: becoming a Ziggerin expert--but which type? *Psychological Science*, *20*(9), 1108-1117.
- Xu, X., Biederman, I., & Shah, M. P. (2014). A neurocomputational account of the face configural effect. *Journal of Vision*, *14*(8:9), 1-9.
- Yang, L. Z., Zhang, W., Shi, B., Yang, Z., Wei, Z., Gu, F., et al. (2014). Electrical stimulation over bilateral occipito-temporal regions reduces N170 in the right hemisphere and the composite face effect. *PLoS One*, *9*(12), e115772.
- Yin, R. K. (1969). Looking at upside-down faces. *Journal of Experimental Psychology*, *81*(1), 141-145.
- Young, A. W., Hellawell, D., & Hay, D. C. (1987). Configurational information in face perception. *Perception*, *16*(6), 747-759.
- Young, S. G., Slepian, M. L., Wilson, J. P., & Hugenberg, K. (2014). Averted eye-gaze disrupts configural face encoding. *Journal of Experimental Social Psychology*, *53*, 94-99.

Yovel, G., & Kanwisher, N. (2008). The representations of spacing and part-based information are associated for upright faces but dissociated for objects: evidence from individual differences. *Psychonomic Bulletin & Review*, *15*(5), 933-939.

Zhao, M., Hayward, W. G., & Bülthoff, I. (2014). Holistic processing, contact, and the other-race effect in face recognition. *Vision Research*, *105*, 61-69.

Zhu, Q., Song, Y., Hu, S., Li, X., Tian, M., Zhen, Z., et al. (2010). Heritability of the specific cognitive ability of face perception. *Current Biology*, *20*(2), 137-142.