

Inactivation of TRPM2 Channels by Extracellular Divalent Copper



Wenyue Yu¹, Lin-Hua Jiang^{2,3}, Yang Zheng¹, Xupang Hu¹, Jianhong Luo^{1*}, Wei Yang^{1*}

1 Key Laboratory of Medical Neurobiology of the Ministry of Health of China, Zhejiang Province Key Laboratory of Neurobiology, Department of Neurobiology, School of Medicine, Zhejiang University, Hangzhou, Zhejiang Province, China, **2** Department of Physiology and Neurobiology, Xinxiang Medical University, Xinxiang, Henan Province, China, **3** School of Biomedical Sciences, Faculty of Biological Sciences, University of Leeds, Leeds, United Kingdom

Abstract

Cu²⁺ is an essential metal ion that plays a critical role in the regulation of a number of ion channels and receptors in addition to acting as a cofactor in a variety of enzymes. Here, we showed that human melastatin transient receptor potential 2 (hTRPM2) channel is sensitive to inhibition by extracellular Cu²⁺. Cu²⁺ at concentrations as low as 3 μM inhibited the hTRPM2 channel completely and irreversibly upon washing or using Cu²⁺ chelators, suggesting channel inactivation. The Cu²⁺-induced inactivation was similar when the channels conducted inward or outward currents, indicating the permeating ions had little effect on Cu²⁺-induced inactivation. Furthermore, Cu²⁺ had no effect on single channel conductance. Alanine substitution by site-directed mutagenesis of His995 in the pore-forming region strongly attenuated Cu²⁺-induced channel inactivation, and mutation of several other pore residues to alanine altered the kinetics of channel inactivation by Cu²⁺. In addition, while introduction of the P1018L mutation is known to result in channel inactivation, exposure to Cu²⁺ accelerated the inactivation of this mutant channel. In contrast with the hTRPM2, the mouse TRPM2 (mTRPM2) channel, which contains glutamine at the position equivalent to His995, was insensitive to Cu²⁺. Replacement of His995 with glutamine in the hTRPM2 conferred loss of Cu²⁺-induced channel inactivation. Taken together, these results suggest that Cu²⁺ inactivates the hTRPM2 channel by interacting with the outer pore region. Our results also indicate that the amino acid residue difference in this region gives rise to species-dependent effect by Cu²⁺ on the human and mouse TRPM2 channels.

Citation: Yu W, Jiang L-H, Zheng Y, Hu X, Luo J, et al. (2014) Inactivation of TRPM2 Channels by Extracellular Divalent Copper. *PLoS ONE* 9(11): e112071. doi:10.1371/journal.pone.0112071

Editor: Shang-Zhong Xu, University of Hull, United Kingdom

Received: September 9, 2014; **Accepted:** October 11, 2014; **Published:** November 11, 2014

Copyright: © 2014 Yu et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability: The authors confirm that all data underlying the findings are fully available without restriction. All relevant data are within the paper.

Funding: This work was supported by National Basic Research Program of China (2013CB910204), Natural Science Foundation of China (81371302), Zhejiang Province Qianjiang Talent Project (2012R10037), and Zhejiang University basic research fund (2012FZA7014) to WY; University of Leeds-Zhejiang University Strategic Collaboration Award and Natural Science Foundation of China (31471118) to LHH; and National Basic Research Program of China (2014CB910300 and 2010CB912002) to JHL. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* Email: luojianhong@zju.edu.cn (JHL); yangwei@zju.edu.cn (WY)

Introduction

The TRPM2 channel belongs to the melastatin subfamily of the mammalian transient receptor potential (TRP) channels, which share several conserved domains with other TRPM channels, such as the TRPM homology domains (MHD domains) in the N-terminus and the TRP box and coiled-coil domain in the C-terminus [1–4]. The TRPM2 channel is a homo-tetramer and each subunit contains six transmembrane segments with a pore-forming region between the fifth and sixth segments and intracellular N- and C-termini [5]. The TRPM2 channel is a non-selective cation channel and permeates calcium ion, and is activated by intracellular ADP-ribose (ADPR) [1,6] or intracellular calcium [7–10]. Accumulating evidence indicates that the TRPM2 channel plays an important role in a number of physiological and pathophysiological processes, including neurodegeneration, immunological functions, insulin release [11–15]. Previous studies showed that the TRPM2 channel can undergo rapid inactivation upon exposure to extracellular proton and Zn²⁺ that interact selective residues in the pore region [9,16,17]. Mutation of the residues in the pore region can strongly alter the channel inactivation. Thus, the disease-associated P1018L mutation

conferred rapid inactivation of the hTRPM2 channel, whereas manipulation of the pore region by site-directed mutagenesis resulted in a TRPM2-LDE mutant channel that exhibited no inactivation, suggesting alterations in the conformation and structure of the pore region represent an important molecular mechanisms of the TRPM2 channel inactivation [18,19].

Cu²⁺ is the third abundant trace metal in the human body, and plays a critical role in a variety of physiological and pathological conditions. Cu²⁺ is a cofactor for a variety of enzymes, and relates to the formation of reactive oxygen species. Like zinc, excessive Cu²⁺ is toxic for neurons [20,21]. Cu²⁺ is involved in several human diseases [22–24], and the Cu²⁺ chelators have been intensively used as therapeutic treatments Cu²⁺ related diseases, such as Wilson's disease and cancer [25]. Several studies suggest Cu²⁺ and Zn²⁺ regulate cell functions via altering the activity of a variety of ion channels [26,27]. For example, Cu²⁺ reduces the tonic inhibition of neurons by blocking the GABA_A receptors [28]. Therefore, elucidating the mechanisms regulating ion channels by Cu²⁺ is critical for a better understanding of its physiological and pathological roles in humans.

It is well known that Cu²⁺ can activate, modulate or inhibit ion channels. For example, Cu²⁺ activates the TRPV1 and TRPA1




Figure 1. TRPM2 open channels inactivated by extracellular Cu²⁺. (A) Representative recordings of the inward currents evoked by 500 μM ADPR at -80 mV, using a 500 ms voltage ramp of -100 mV to $+100$ mV applied every 5 s, before and after exposure to the indicated Cu²⁺ concentrations. The dotted lines indicate zero currents. (B–C) Summary of the percentage inhibition (B) and time required for inward current amplitude reached 90% inhibition after Cu²⁺ exposure (C). (D) Left panel, the ADPR-induced inward currents when fully inhibited by 100 μM Cu²⁺ were not reversed after treating with 20 μM 2-ME; Right panel, summary of the current recovery during exposure to 2-ME. (E) Left panel, the ADPR-induced inward currents when fully inhibited by 100 μM Cu²⁺ were not reversed after treating with 5 mM EDTA; Right panel, summary of the current recovery during exposure to EDTA. Residual current expressed as the percentage of the currents immediately before exposure to Cu²⁺ is $3.3 \pm 1.7\%$ after inactivation by Cu²⁺, which returned to $2.6 \pm 0.9\%$ after washing with EDTA. In 2-ME group, residual current changed from $1.8 \pm 0.5\%$ to $2.0 \pm 0.7\%$. The number of cells examined in each case is 4–6. doi:10.1371/journal.pone.0112071.g001

channels [29,30] and, by contrast, Cu²⁺ inhibits endothelial Na⁺ channels [31], BK and Shaker K⁺ channels [32]. A recent study has reported that extracellular Cu²⁺ induces the hTRPM2 channel inactivation [33], but the underlying molecular or structural basis still remains unclear. Here, using site-directed mutagenesis and patch-clamp recording, we identified His995 in the pore region to be crucial in Cu²⁺-induced hTRPM2 channel inactivation. In addition, the mTRPM2 channel is insensitive to Cu²⁺ and such a species-dependent effect by extracellular Cu²⁺ arises from replacement of His995 in the hTRPM2 channel with glutamine at the equivalent position in the mTRPM2 channel.

Materials and Methods

Clones, cells and molecular biology

The cDNAs encoding the hTRPM2 and mTRPM2 were kindly provided by Dr AM Scharenberg (Washington University, USA) and Dr Y Mori (Kyoto University, Japan), respectively. Tetracycline-inducible HEK293 cells stably expressing hTRPM2 was

kindly provided by Dr AM Scharenberg. The point mutations were introduced by site-directed mutagenesis [34] and confirmed by sequencing. Human embryonic kidney (HEK) 293 cells were used to transiently express wild-type (WT) and mutant channels. HEK293 cells were maintained in DMEM/F-12 medium supplemented with 50 units/ml penicillin, 50 μg/ml streptomycin (Gibco, USA) and 10% fetal bovine serum (Gibco, USA). Cell transfection was described previously [4]. Tetracycline-inducible hTRPM2-expressing HEK293 cells were used in some experiments and the expression of the hTRPM2 channel was induced by adding 1 μg/ml tetracycline in the culture medium for 12–24 h before use.

Electrophysiology

Whole-cell current recordings were performed using an Axonpatch 200B amplifier at room temperature as described previously [4]. The extracellular solution contained (in mM): 147 NaCl, 2 KCl, 1 MgCl₂, 2 CaCl₂, 10 HEPES, and 13 glucose,




Figure 2. Voltage-independent effects by extracellular Cu²⁺ on inward and outward TRPM2 channel currents. (A) ADPR-induced currents mediated by the hTRPM2 channels at a holding membrane potentials (HP) of +40 mV (outward currents) or -40 mV (inward currents) and the effect of 100 μM Cu²⁺. The dotted lines indicate the baseline. (B–C) Summary of the outward or inward current amplitude before and after exposure to Cu²⁺ and upon washout, as shown in (A). The number of cells examined in each case is 4. ***, $p < 0.005$ compared with the currents before and after exposure to the indicated Cu²⁺. (D) Summary of the time for 90% inhibition at both 40 mV and -40 mV, there is no significant difference between these groups. doi:10.1371/journal.pone.0112071.g002

pH 7.4. The intracellular solution contained (in mM): 147 NaCl, 0.05 EGTA, 1 MgCl₂, 10 HEPES, and 0.5 ADPR, pH 7.3. Thus, the currents were mainly carried by Na⁺. The cell membrane potential was held at 0 mV. To record ADPR-induced currents, voltage ramps with 500 ms duration from -100 mV to 100 mV were applied every 5 s. The glass microelectrodes with resistance of 3–5 MΩ were used. Single channel activity recordings were performed using a HEKA EPC10 amplifier controlled with PatchMaster software (HEKA), and carried out in the outside-out configuration as previous described [17]. Data were acquired at 10 kHz and filtered offline at 50 Hz. CuSO₄ stock solution was prepared by dissolving in the extracellular solutions. Changes of extracellular solutions and applications of Cu²⁺, clotrimazole (CLT), 2-ME and EDTA were achieved using an RSC-160 system (Biologic Science Instruments) with a solution change time of ~300 ms [4].

Data analysis

The data are presented in the text and figures, where appropriate, as mean ± S.E. values. The inhibition was expressed by the currents in the indicated solutions as the percentage of the currents before the solution changes, and the kinetics were estimated by determining the time required for 90% inhibition ($t_{90\%}$). Single-channel conductance was estimated from all-point histograms constructed from the current events recorded at

-80 mV. A double-Gaussian function was fitted to the histograms as previously described [35]. Statistical analysis was performed using Student's t-test with $p < 0.05$ designated as significant difference.

Results

Extracellular Cu²⁺ induces hTRPM2 channel inactivation

We first investigated the effect of extracellular Cu²⁺ on the open hTRPM2 channels. The activity of the hTRPM2 channels was induced by application of ADPR (500 μM) in the pipette solution, as illustrated by the inward currents at -80 mV (Fig. 1A). Perfusion with 3 μM Cu²⁺ abolished the hTRPM2 channel currents in several minutes, as reported by a recent study [33]. While the hTRPM2 channels were completely inhibited by Cu²⁺ independently of concentrations from 3 μM to 1 mM, the time required for 90% inhibition ($t_{90\%}$) was concentration-dependent from 98.4 ± 16.7 s ($n = 4$) at 3 μM to 7.1 ± 1.5 s ($n = 5$) at 1 mM (Fig. 1B). The inhibitory effect of Cu²⁺ was irreversible on washout, which suggests that Cu²⁺ induced channel inactivation like proton and zinc [16,17,36]. Such inactivation could result from either channel inactivation or tight binding of Cu²⁺ to the hTRPM2 channels. To clarify this issue, we applied 2-ME or EDTA to chelate Cu²⁺, after Cu²⁺ induced complete inhibition of ADP-induced currents. The inhibition was still not rescued by treating with the 2-ME or EDTA for up to two min (Fig. 1C, D),




Figure 3. Effects of external Cu^{2+} on human TRPM2 single channel conductance. (A) Representative recordings in the outside-out configuration of the effects of $30 \mu\text{M Cu}^{2+}$ on ADPR-induced TRPM2 currents (in red). Single channel events are clearly visible in the expanded traces illustrated below. (B) The grey and red histograms of single channel events indicated the exposure in ECS and $30 \mu\text{M Cu}^{2+}$, respectively. The superimposed curve represents a fit of a double-Gaussian function.
doi:10.1371/journal.pone.0112071.g003

which suggests that Cu^{2+} may not bind with the TRPM2 after TRPM2 inactivation, or alternatively Cu^{2+} tightly binds with the hTRPM2 channels in the small pocket and both EDTA and 2-ME cannot access.

Our previous study showed that Zn^{2+} inactivates the hTRPM2 channel and such inactivation is strongly affected by the permeant ions [17]. It was thus interesting to know whether Cu^{2+} -induced hTRPM2 inactivation was similar as that by Zn^{2+} . Unlike Zn^{2+} , $100 \mu\text{M Cu}^{2+}$ still inhibited irreversibly the hTRPM2 channel currents at 40 mV (Fig. 2A), which suggests the hTRPM2 inactivation induced by Cu^{2+} may be different from that by Zn^{2+} . Moreover, we investigated whether Cu^{2+} affects the single channel conductance using the patch-clamp recording in outside-out configuration. As shown in Fig. 3, the single channel conductance of the hTRPM2 channels was not changed by Cu^{2+} .

His995 is a key residue in the outer pore in Cu^{2+} -induced inactivation

Metal ions can interact with polar amino acid like histidine, glutamate, aspartate, and lysine [9,16,36,37]. Several studies have identified histidine, cysteine, aspartic acid, arginine and glutamine determine Cu^{2+} -induced channel inhibition or inactivation, such as the P2X7 receptor [38], GABA_A receptor [28], ENaC channel [31], BK channel [32]. In addition, a previous study of olfactory CNG channel gating suggests that the extracellular pore region is

importantly involved in the channel gating [39]. Our previous studies have identified several amino acids which are responsible for channel inactivation induced by proton and Zn^{2+} (Fig. 4A). We hypothesized that Cu^{2+} may interact with similar amino acid residues in the pore region which Zn^{2+} binds to (Fig. 3). We introduced alanine substitution into the candidate amino acid residues as described in previous studies [16,36], and determined the effects of $100 \mu\text{M Cu}^{2+}$ on the functional mutant channels. All of these mutants did not strongly result in strong channel inactivation as illustrated by several examples in Fig. 4. Representative current traces of the hTRPM2 mutant channels are illustrated in Fig. 4A, and the mutational effects on the Cu^{2+} -induced inhibition and inactivation are summarized in Fig. 4B. Several mutants, albeit still be inactivated by Cu^{2+} , exhibited significantly slower inactivation kinetics, such as K952A ($88.8 \pm 6.7 \text{ s}$), R961A ($92.3 \pm 11.7 \text{ s}$), H973A ($126.9 \pm 10.3 \text{ s}$), D994A ($88.7 \pm 6.1 \text{ s}$), D1002A ($88.8 \pm 4.6 \text{ s}$), E1010A ($90.6 \pm 9.8 \text{ s}$) and R1017A ($90.4 \pm 7.8 \text{ s}$) as compared to the WT channel ($48.5 \pm 2.5 \text{ s}$) (Fig. 4A, B). On the contrary, some other mutations accelerated inactivation kinetics, including K1005A ($16.7 \pm 1.9 \text{ s}$), K1007A ($32.9 \pm 7.6 \text{ s}$), E1022A ($38.2 \pm 3.4 \text{ s}$) (Fig. 4B). Strikingly, the H995A mutant channel was insensitive to inactivation by $100 \mu\text{M Cu}^{2+}$ (Fig. 4A, B). These results suggest that Cu^{2+} inactivates the hTRPM2 channel via engaging the outer pore region.




Figure 4. Alanine substitution of Cu²⁺-binding candidate residues in the outer pore of hTRPM2 channel. (A) The amino acid sequences of the pore region between the S5 and S6 of the hTRPM2 channel; the residues examined in this study are numbered and highlighted in bold. Residues in the extracellular ends of S5 and S6 are indicated in the left and right shading boxes, respectively. Alanine substitutions leading to loss of function are indicated by asterisks. (B) Representative recordings of the ADPR-induced currents in HEK cells expressing WT or the indicated mutant channel before and after exposure to 100 μM Cu²⁺ (denoted by the black bars). The currents for the H995A mutant channels show incomplete inhibition by Cu²⁺ and complete inhibition by subsequent application of 20 μM CLT (denoted by the grey bars). The dotted lines indicate zero currents. (C) Summary of the time for 90% inhibition (left) and the residual currents upon exposure to Cu²⁺ (right). The dotted lines indicate the time or residual currents for the WT channel. The number of cells examined in each case is 3–22. The mutant channels showing significant difference from the WT channels are indicated in parentheses, *, p<0.05, **, p<0.01, ***, p<0.005. doi:10.1371/journal.pone.0112071.g004




Figure 5. P1018L mutation facilitates Cu²⁺-induced hTRPM2 channel inactivation. (A) The inward current of cell transfected with human TRPM2 is blocked by Cu²⁺. (B) P1018L mutant exhibited strong channel inactivation in extracellular solution (ECS). (C) Cu²⁺ accelerated the P1018L mutant channel inactivation. (D) Summary of the time for 90% inhibition after exposure to Cu²⁺. The number of cells examined in each case is 3–13. The significant difference is indicated in parentheses, ***, $p < 0.005$. doi:10.1371/journal.pone.0112071.g005

Cu²⁺ accelerates the inactivation of hTRPM2 P1018L mutant channel

A previous study showed that the P1018L mutation in hTRPM2 resulted in channel inactivation [18], it is interesting to know the effect of extracellular Cu²⁺ on this mutant channel. Consistent with the previous study [18], the hTRPM2 P1018L mutant channel exhibited strong channel inactivation (121.6 ± 14.8 s) (Fig. 5A, B). Interestingly, 100 μM Cu²⁺ significantly accelerated the inactivation kinetics of the hTRPM2 P1018L mutant channel (15.3 ± 0.7 s) (Fig. 5C, D).

Mouse TRPM2 channel shows insensitivity to Cu²⁺-induced inactivation

Species difference exists between the human and mouse TRPM2 channels in terms of inactivation of the TRPM2 channels by protons and zinc, as a result of difference in the amino acid sequences of the pore region [17,36]. By comparing with the sequences, three residues in the pore region of the hTRPM2 channel are different with those of the mTRPM2. Arg961, His995 and Arg1017 residues in the hTRPM2 are replaced with Ser958, Gln992, and Ala1014 in the mTRPM2, respectively (Fig. 4A). The results that His995 plays a crucial role in Cu²⁺-induced hTRPM2 channel inactivation suggest that Cu²⁺ may not strongly interact with the pore region of the mTRPM2 channel to induce inactivation of the mTRPM2 channel. To address this hypothesis,

we expressed the mTRPM2 channels in HEK293 cells and determined the effect of Cu²⁺ on the mTRPM2 channels. As shown in Fig. 6A, 100 μM Cu²⁺ failed to induce significant inhibition/inactivation of the mTRPM2 channels. We further generated the construct expressing the H995Q hTRPM2 mutant channel. The H995Q mutation almost completely abolished the hTRPM2 channel inactivation induced by Cu²⁺, even better than the H995A mutation (Fig. 6B). These results provide further evidence to indicate that His995 is critical in determining hTRPM2 channel inactivation by Cu²⁺.

Discussion

In this study, we found that extracellular Cu²⁺ inactivates the human but not the mouse TRPM2 channel, and revealed a striking species-dependent effect. Moreover, we identified several residues are involved in this process, particularly His995 which is the key residue in determining the hTRPM2 channel inactivation induced by Cu²⁺.

Cu²⁺-induced TRPM2 inactivation is independent on electrostatic repulsion and channel inhibition

Unlike the actions of extracellular Cu²⁺ on other ion channels, the present study shows Cu²⁺ induced an irreversible and concentration-independent inhibition of the hTRPM2 channel. Moreover, we found that Cu²⁺ induced similar TRPM2 inactivation




Figure 6. His⁹⁹⁵ is essential in Cu²⁺ interacting with human TRPM2 channel. (A) Recordings of the inward currents in cells expressing the human, mouse wild-type or mutant TRPM2 channels. (B) Summary of the effects of Cu²⁺ on these three channels. The number of cells examined in each case is 3–5. The significant difference is indicated in parentheses. ***, $p < 0.005$. doi:10.1371/journal.pone.0112071.g006

tion at the holding potential of -40 mV driving inward currents, as at 40 mV, driving outward currents. This is different with the membrane potential-dependent action of zinc shown in our previous study [17]. Thus, the action of Cu²⁺ may mechanistically differ from the action of zinc [17]. Remarkably, our data indicate that Cu²⁺-induced hTRPM2 inactivation is independent on electrostatic repulsion, suggesting that Cu²⁺ binds with the residues outside the electrical field and induces conformational changes that lead to hTRPM2 channel inactivation. Both EDTA and 2-ME failed to reverse the Cu²⁺-induced channel inactivation. One simple explanation is that Cu²⁺ induces substantial conformational changes that prevent accessibility of such Cu²⁺ chelators.

His995 is the key residue determining Cu²⁺-induced inactivation of human TRPM2 channel

As we mentioned before, many studies have identified several residues interact with Cu²⁺, such as histidine, cysteine, glutamate, aspartic acid and arginine. By alanine scanning, we have shown several residues in of pore region that are involved in the Cu²⁺-induced inactivation of the hTRPM2 channel (Fig. 4). Particularly, the H995A mutation that is located at the large turret of the pore region strongly attenuated the hTRPM2 channel inactivation by Cu²⁺ (Fig. 4). Although several other residues replaced by alanine did not prevent the hTRPM2 channel inactivation by Cu²⁺, the kinetics were substantially changed at these mutant channels (Fig. 4). Specifically, K952A, R961A, H973A, D994A, D1002A, E1010A and R1017A mutations slowed down the inactivation kinetics (Fig. 4A, B), whereas K1005A, K1007A and E1022A resulted in faster inactivation kinetics (Fig. 4B). These findings support that Cu²⁺

induces the hTRPM2 channel inactivation by acting on the outer pore region. In future, it is important to uncover the conformational changes that give rise to Cu²⁺-induced hTRPM2 channel inactivation.

Species specific effects of Cu²⁺ on human and mouse TRPM2 channels

The TRPM2 channels are highly conserved in different species including the human and mouse. Our previous studies found three different residues in the pore region, Arg⁹⁶¹/Ser⁹⁵⁸, His⁹⁹⁵/Gln⁹⁹², Arg¹⁰¹⁷/Ala¹⁰¹⁴, between the human and mouse TRPM2 channels and showed that His⁹⁹⁵/Gln⁹⁹² is important in determining the different kinetics of channel inactivation induced by proton or Zn²⁺ between the human and mouse TRPM2 channels [17,36]. Here, we have shown that all of these three sites are critical for the hTRPM2 channel inactivation induced by Cu²⁺, which is confirmed by the observation that there was no mTRPM2 channel inactivation by Cu²⁺ (Fig. 6). Taken together, the previous and present studies suggest that the hTRPM2 channel is more sensitive to Cu²⁺, Zn²⁺ and proton than the mTRPM2 channel. Many lines of evidence have indicated that Cu²⁺ is involved in the pathogenesis of neurodegenerative disorders such as Alzheimer's disease, Parkinson's disease and prion disease [40,41]. On the other hand, accumulating evidence implicates that the TRPM2 channels as an oxidative stress sensor is also related to these mental disorders [5]. In future, it is interesting to investigate whether the role of Cu²⁺ in the neurodegenerative disorders is related to the hTRPM2 channel. However, according to our results, the striking species difference between the human and mouse TRPM2 channels

suggest unsuitable to use mice as animal models to investigate the regulation of TRPM2 channels by Cu^{2+} in relation to the human diseases.

Comparing the effects on TRPM2 channels induced by Cu^{2+} and Zn^{2+}

Both Zn^{2+} and Cu^{2+} are metal ions, but the Cu^{2+} -induced TRPM2 channel inactivation is different with that by Zn^{2+} . Firstly, the key residues that determine the TRPM2 inactivation by Cu^{2+} and Zn^{2+} are different. Secondly, Cu^{2+} -induced TRPM2 channel inactivation is independent on the membrane potential and/or the direction of Na^+ permeation (Fig. 2). Finally, the Cu^{2+} -induced inactivation was accelerated by the K1005A, K1007A, E1022A mutations, however, the Zn^{2+} -induced inactivation was accelerated by the D964A, H973A, K1005A, K1007A, R1017A, and E1022A mutations. The Cu^{2+} -induced inactivation was slowed by the K952A, R961A, H973A, D994A, D1002A, E1010A and R1017A, whereas the Zn^{2+} -induced inactivation was slowed by H958A, E994A, H995A, E1010A, and D1012A. Although there are some mutations present the similar effect on the TRPM2 channel inactivation by Cu^{2+} and Zn^{2+} , many mutations resulted in the opposite effects between Cu^{2+} -induced inactivation and Zn^{2+} -induced inactivation. These differences suggest the critical site of Cu^{2+} -binding overlap with but differ from the Zn^{2+} -binding site and some differences in the molecular mechanisms in the TRPM2 inactivation induced by Cu^{2+} and Zn^{2+} . However, there was no effect of Cu^{2+} on the single channel conductance as previously reported for Zn^{2+} [33], which suggests that both Cu^{2+} and Zn^{2+} induce TRPM2 inactivation without affecting the ion conducting pathway. Future study is required to know how the TRPM2 channels undergo conformational changes during the inactivation during Cu^{2+} or Zn^{2+} treatment.

Our data have shown that the P1018L mutation located at the

pore region of hTRPM2 resulted in strong channel inactivation (Fig. 5) as reported previously [18]. We investigated whether hTRPM2 channel inactivation induced by Cu^{2+} and P1018L mutation occurred through similar mechanisms. Our result showed that Cu^{2+} significantly accelerated the inactivation kinetics of the hTRPM2 P1018L mutant channel (Fig. 5), indicating that Cu^{2+} can induce strong channel inactivation of this mutant channel. Taken together, our results suggest that TRPM2 inactivation is complex, it will be interesting to know how to induce channel inactivation by these different factors in the future.

In summary, we have shown that extracellular Cu^{2+} inactivates the hTRPM2 channel independently of the ion conducting pathway. By introducing point mutation, we identified multiple residues, especially His995, in the outer pore region that are involved in Cu^{2+} -induced hTRPM2 channel inactivation. We also discovered that the mTRPM2 channel is insensitive to Cu^{2+} . These findings open a new way for us to understand the roles of TRPM2 channel in Cu^{2+} related physiological and pathological processes.

Acknowledgments

We thank Dr AM Scharenberg (Washington University, USA) for kindly providing the hTRPM2 cDNA and the tetracycline-inducible hTRPM2 stable cells, and Dr Y Mori (Kyoto University, Japan) for the mTRPM2 cDNA, and Dr J Zheng (UC Davis, USA) for kindly supporting single channel recording.

Author Contributions

Conceived and designed the experiments: WY LHJ WYY. Performed the experiments: WYY YZ XPH WY. Analyzed the data: WYY WY. Contributed reagents/materials/analysis tools: JHL LHJ WY. Wrote the paper: WYY WY LHJ. Revised this manuscript: JHL LHJ.

References

- Perraud AL, Fleig A, Dunn CA, Bagley LA, Launay P, et al. (2001) ADP-ribose gating of the calcium-permeable LTRPC2 channel revealed by Nudix motif homology. *Nature* 411: 595–599.
- Nilius B (2007) Transient receptor potential (TRP) cation channels: rewarding unique proteins. *Bull Mem Acad R Med Belg* 162: 244–253.
- Wu IJ, Sweet TB, Clapham DE (2010) International Union of Basic and Clinical Pharmacology. LXXVI. Current progress in the mammalian TRP ion channel family. *Pharmacol Rev* 62: 381–404.
- Mei ZZ, Mao HJ, Jiang LH (2006) Conserved cysteine residues in the pore region are obligatory for human TRPM2 channel function. *Am J Physiol Cell Physiol* 291: C1022–1028.
- Jiang LH, Yang W, Zou J, Beech DJ (2010) TRPM2 channel properties, functions and therapeutic potentials. *Expert Opin Ther Targets* 14: 973–988.
- Sano Y, Inamura K, Miyake A, Mochizuki S, Yokoi H, et al. (2001) Immuncyte Ca^{2+} influx system mediated by LTRPC2. *Science* 293: 1327–1330.
- McHugh D, Flemming R, Xu SZ, Perraud AL, Beech DJ (2003) Critical intracellular Ca^{2+} dependence of transient receptor potential melastatin 2 (TRPM2) cation channel activation. *J Biol Chem* 278: 11002–11006.
- Csanady L, Torocsik B (2009) Four Ca^{2+} ions activate TRPM2 channels by binding in deep crevices near the pore but intracellularly of the gate. *J Gen Physiol* 133: 189–203.
- Du J, Xie J, Yue L (2009) Modulation of TRPM2 by acidic pH and the underlying mechanisms for pH sensitivity. *J Gen Physiol* 134: 471–488.
- Sumoza-Toledo A, Penner R (2011) TRPM2: a multifunctional ion channel for calcium signalling. *J Physiol* 589: 1515–1525.
- Di A, Gao XP, Qian F, Kawamura T, Han J, et al. (2012) The redox-sensitive cation channel TRPM2 modulates phagocyte ROS production and inflammation. *Nat Immunol* 13: 29–34.
- Haraguchi K, Kawamoto A, Isami K, Maeda S, Kusano A, et al. (2012) TRPM2 contributes to inflammatory and neuropathic pain through the aggravation of pronociceptive inflammatory responses in mice. *J Neurosci* 32: 3931–3941.
- Massullo P, Sumoza-Toledo A, Bhagat H, Partida-Sanchez S (2006) TRPM channels, calcium and redox sensors during innate immune responses. *Semin Cell Dev Biol* 17: 654–666.
- Uchida K, Tominaga M (2011) TRPM2 modulates insulin secretion in pancreatic beta-cells. *Islets* 3: 209–211.
- Orfanelli U, Wenke AK, Dogliani C, Russo V, Bosserhoff AK, et al. (2008) Identification of novel sense and antisense transcription at the TRPM2 locus in cancer. *Cell Res* 18: 1128–1140.
- Yang W, Zou J, Xia R, Vaal ML, Seymour VA, et al. (2010) State-dependent inhibition of TRPM2 channel by acidic pH. *J Biol Chem* 285: 30411–30418.
- Yang W, Manna PT, Zou J, Luo J, Beech DJ, et al. (2011) Zinc inactivates melastatin transient receptor potential 2 channels via the outer pore. *J Biol Chem* 286: 23789–23798.
- Hermosura MC, Cui AM, Go RC, Davenport B, Shetler CM, et al. (2008) Altered functional properties of a TRPM2 variant in Guamanian ALS and PD. *Proc Natl Acad Sci U S A* 105: 18029–18034.
- Toth B, Csanady L (2012) Pore collapse underlies irreversible inactivation of TRPM2 cation channel currents. *Proc Natl Acad Sci U S A* 109: 13440–13445.
- Ahuja A, Dev K, Tanwar RS, Selwal KK, Tyagi PK (2014) Copper mediated neurological disorder: Visions into amyotrophic lateral sclerosis, Alzheimer and Menkes disease. *J Trace Elem Med Biol*.
- Kocer G, Naziroglu M, Celik O, Onal L, Ozcelik D, et al. (2013) Basic fibroblast growth factor attenuates bisphosphonate-induced oxidative injury but decreases zinc and copper levels in oral epithelium of rat. *Biol Trace Elem Res* 153: 251–256.
- Chelly J, Monaco AP (1993) Cloning the Wilson disease gene. *Nat Genet* 5: 317–318.
- Ellingsen DG, Chashchin V, Haug E, Chashchin M, Tkachenko V, et al. (2007) An epidemiological study of reproductive function biomarkers in male welders. *Biomarkers* 12: 497–509.
- Tisato F, Marzano C, Porchia M, Pelli M, Santini C (2010) Copper in diseases and treatments, and copper-based anticancer strategies. *Med Res Rev* 30: 708–749.
- Ding XQ, Fitzgerald JB, Quiambao AB, Harry CS, Malykhina AP (2010) Molecular pathogenesis of achromatopsia associated with mutations in the cone cyclic nucleotide-gated channel CNGA3 subunit. *Adv Exp Med Biol* 664: 245–253.
- Kiss T, Osipenko ON (1994) Toxic effects of heavy metals on ionic channels. *Pharmacol Rev* 46: 245–267.

27. Restrepo-Angulo I, De Vizcaya-Ruiz A, Camacho J (2010) Ion channels in toxicology. *J Appl Toxicol* 30: 497–512.
28. McGee TP, Houston CM, Brickley SG (2013) Copper block of extrasynaptic GABAA receptors in the mature cerebellum and striatum. *J Neurosci* 33: 13431–13435.
29. Gu Q, Lin RL (2010) Heavy metals zinc, cadmium, and copper stimulate pulmonary sensory neurons via direct activation of TRPA1. *J Appl Physiol* (1985) 108: 891–897.
30. Riera CE, Vogel H, Simon SA, le Coutre J (2007) Artificial sweeteners and salts producing a metallic taste sensation activate TRPV1 receptors. *Am J Physiol Regul Integr Comp Physiol* 293: R626–634.
31. Chen J, Myerburg MM, Passero CJ, Winarski KL, Sheng S (2011) External Cu²⁺ inhibits human epithelial Na⁺ channels by binding at a subunit interface of extracellular domains. *J Biol Chem* 286: 27436–27446.
32. Ma Z, Wong KY, Horrigan FT (2008) An extracellular Cu²⁺ binding site in the voltage sensor of BK and Shaker potassium channels. *J Gen Physiol* 131: 483–502.
33. Zeng B, Chen GL, Xu SZ (2012) Divalent copper is a potent extracellular blocker for TRPM2 channel. *Biochem Biophys Res Commun* 424: 279–284.
34. Yang W, Jiang LH (2013) Site-directed mutagenesis to study the structure-function relationships of ion channels. *Methods Mol Biol* 998: 257–266.
35. Cao X, Ma L, Yang F, Wang K, Zheng J (2014) Divalent cations potentiate TRPV1 channel by lowering the heat activation threshold. *J Gen Physiol* 143: 75–90.
36. Zou J, Yang W, Beech DJ, Jiang LH (2011) A residue in the TRPM2 channel outer pore is crucial in determining species-dependent sensitivity to extracellular acidic pH. *Pflügers Arch* 462: 293–302.
37. Coddou C, Acuna-Castillo C, Bull P, Huidobro-Toro JP (2007) Dissecting the facilitator and inhibitor allosteric metal sites of the P2X4 receptor channel: critical roles of CYS132 for zinc potentiation and ASP138 for copper inhibition. *J Biol Chem* 282: 36879–36886.
38. Liu X, Surprenant A, Mao HJ, Roger S, Xia R, et al. (2008) Identification of key residues coordinating functional inhibition of P2X7 receptors by zinc and copper. *Mol Pharmacol* 73: 252–259.
39. Gavazzo P, Picco C, Eismann E, Kaupp UB, Menini A (2000) A point mutation in the pore region alters gating, Ca²⁺ blockage, and permeation of olfactory cyclic nucleotide-gated channels. *J Gen Physiol* 116: 311–326.
40. Squitti R (2012) Copper dysfunction in Alzheimer's disease: from meta-analysis of biochemical studies to new insight into genetics. *J Trace Elem Med Biol* 26: 93–96.
41. Desai V, Kaler SG (2008) Role of copper in human neurological disorders. *Am J Clin Nutr* 88: 855S–858S.