

1 **Title:** Neuromuscular responses to mild-muscle damaging eccentric exercise in a low glycogen state

2

3 **Authors:** James P. Gavin, Stephen D. Myers, and Mark E.T. Willems

4

5 **Affiliation:** Department of Sport & Exercise Sciences, University of Chichester, Chichester, UK

6

7 **Keywords:** Eccentric contractions; Muscle glycogen; Neuromuscular fatigue; Muscle damage

8

9

10

11

12

13

14

15

16

17 **Correspondence address:**

18 Prof Mark Willems

19 Department of Sport & Exercise Sciences

20 University of Chichester

21 College Lane

22 Chichester

23 West Sussex

24 PO19 6PE

25 UK.

26 Email: [m.willems@chi.ac.uk](mailto:m.willems@chi.ac.uk)

27 Phone: +44 (0) 1243 816468

28

29

30

## 1 **Abstract**

2 The aim of this study was to examine the effect of low muscle glycogen on the neuromuscular responses to  
3 maximal eccentric contractions. Fourteen healthy men ( $22\pm 3$  yr) performed single-leg cycling (20 min at  $\sim 75\%$   
4 maximal oxygen uptake ( $\text{VO}_{2\text{max}}$ ); eight 90 s sprints at a 1:1 work-to-rest ratio (5% decrements from 90 to 55%  
5  $\text{VO}_{2\text{max}}$ ); and  $\sim 85\%$   $\text{VO}_{2\text{max}}$  until exhaustion) the evening before 100 eccentric knee extensions ( $1.57 \text{ rad}\cdot\text{s}^{-1}$ ) with  
6 reduced (RED) and normal glycogen (NORM). Neuromuscular responses were measured during and up to 48 h  
7 after with maximal voluntary and involuntary (twitch, 20 Hz and 50 Hz) isometric contractions. During eccentric  
8 contractions, peak torque decreased (RED:  $-16.1\pm 2.5\%$ ; NORM:  $-6.2\pm 5.1\%$ ) and EMG frequency increased  
9 according to muscle length. EMG activity decreased for RED only. After eccentric contractions, maximal  
10 isometric force was reduced up to 24 h for NORM ( $-13.5\pm 5.8\%$ ) and 48 h for RED ( $-7.4\pm 10.9\%$ ). Twelve hours  
11 after eccentric contractions, twitch force and the 20:50 Hz ratio were decreased for RED but not for NORM.  
12 Immediate involuntary with prolonged voluntary force loss suggests that reduced glycogen is associated with  
13 increased susceptibility to mild muscle-damaging eccentric exercise with contributions of peripheral and central  
14 mechanisms to be different during recovery.

15

## 16 **Introduction**

17 Eccentric contractions are common for the knee extensors in daily activities, whether unaccustomed  
18 (such as downhill running) or habitual (such as stair descent). Vigorous eccentric contractions induce muscle  
19 damage, characterised by muscle soreness (Clarkson and Sayers, 1999), force loss (Behrens et al., 2012) and  
20 altered muscle glycogen metabolism (Asp et al., 1998; Widrick et al., 1992). Impairments in force-producing  
21 capacity have been attributed to central and peripheral neuromuscular origins (Edwards, 1988). Central  
22 mechanisms occur between the motor cortex and proximal to the neuromuscular junction and refer to the failure  
23 of central activation in muscle recruitment during contraction. Peripheral mechanisms reside below the  
24 neuromuscular junction and are associated with excitation-contraction (E-C) coupling failure due to: i) disturbed  
25  $\text{Ca}^{2+}$  homeostasis (Warren et al., 2001; Verburg et al., 2006), and ii) sarcomeric disruption (Morgan et al., 2004).  
26 Furthermore, unique neuromuscular characteristics are exhibited for eccentric exercise, including reduced  
27 activation (Tesch et al., 1990) and early type II fibre recruitment (Nardone et al., 1989), with preserved  
28 recruitment order (Beltman et al., 2004). Following eccentric exercise, low-frequency fatigue (LFF) suggests  
29 disturbance in E-C coupling (Skurvydas et al., 2006) and intramuscular  $\text{Ca}^{2+}$  homeostasis (Verburg et al., 2006).  
30 LFF is characterised by force loss at low-, as opposed to high stimulation frequencies (Verburg et al., 2006).

1 Previously, decreased sarcoplasmic reticulum  $\text{Ca}^{2+}$  release, and thus reduced tetanic  $\text{Ca}^{2+}$  concentration, have  
2 been attributed to the emergence of LFF (Westerblad et al., 1993). Therefore, with force production associated  
3 with intramuscular  $\text{Ca}^{2+}$  release/uptake, muscle glycogen reduction is likely to present a metabolic stress,  
4 impairing sarcoplasmic reticulum function during subsequent exercise.

5 Neuromuscular origins contributing to fatigue and muscle damage have been studied widely. Some  
6 have reported little evidence of central mechanisms (Bigland-Ritchie et al., 1986; Kent-Braun, 1999), whereas  
7 others have demonstrated significant central involvement in response to knee extensor fatigue (Stewart et al.,  
8 2008). As maximal eccentric contractions produce forces exceeding isometric and concentric actions (Tesch et  
9 al., 1990), accompanying reduced neural drive indicates an inhibitory response, preventing further muscle-  
10 tendon unit damage (Clarkson and Sayers, 1999). Furthermore, metabolic consequences of eccentric exercise  
11 include disturbed  $\text{Ca}^{2+}$  homeostasis, preferential type II fibre damage (Asp et al., 1998; McHugh et al., 2000) and  
12 impaired glycogen recovery (Widrick et al., 1992). Maximal eccentric knee extensor exercise appears to  
13 compromise subsequent concentric exercise performance by increasing relative workload, and therefore utilising  
14 already reduced muscle glycogen stores (Asp et al., 1998). Mechanical and metabolic responses to knee extensor  
15 muscle damage are well characterised, however the neuromuscular responses during eccentric exercise with  
16 lowered muscle glycogen are unclear. Low muscle glycogen availability following prolonged exercise may be  
17 associated with susceptibility to, and recovery from muscle damage; and therefore relevant to those undertaking  
18 new training regimen or athletic cohorts performing heavy and/or repeat training bouts. Although neuromuscular  
19 responses to eccentric knee extensor exercise are well documented, it remains unresolved as whether an additive  
20 effect exists for exercise-induced mechanical and metabolic stresses on the i) recovery of neuromuscular  
21 function, and ii) the contributory mechanisms. This study protocol aimed to examine the effect of reduced  
22 muscle glycogen availability on the neuromuscular responses to maximal eccentric contractions.

23

## 24 **Methods**

### 25 *Participants*

26 Fourteen healthy males (age,  $22\pm 3$  years; height,  $179\pm 6$  cm; weight,  $76\pm 15$  kg; body fat,  $9\pm 3\%$ ) volunteered and  
27 provided written informed consent for participation. Participants were recreationally active, free from  
28 musculoskeletal injury and had not been involved in, or were presently engaged in resistance training. Sample  
29 size estimation was based on data for maximal voluntary contraction (MVC) force loss ( $-19.8\%$ ,  $\alpha$  level = 0.05,

1 power = 0.80 (1 -  $\beta$ ) (Faul et al., 2007). The research protocol was approved by the University of Chichester  
2 Research Ethics Committee.

3

#### 4 *Study protocol*

5 Participants visited the laboratory on seven occasions within 10 days. Visits one and two were familiarisation  
6 sessions to accustom participants with single-leg cycling, electrical stimulation procedures, and to establish  
7 EMG sensor placement. Visit three was a pre-eccentric, evening session involving an exercise-diet intervention  
8 to manipulate glycogen state. With dietary control, this involved single-leg cycling to reduce muscle glycogen in  
9 the right leg; the resting left leg served as control. The protocol was validated in a similar cohort by directly  
10 measuring muscle glycogen (muscle glycogen reduced by ~44.7%) (Pilegaard et al., 2002). Visit four followed  
11 early the next morning, with participants performing 100 maximal, eccentric knee extensions with the reduced  
12 glycogen availability (RED) right leg, and then the normal glycogen availability (NORM) left leg. Visits five to  
13 seven involved assessment of neuromuscular function and muscle soreness at 12 h intervals, up to 48 h.  
14 Neuromuscular assessment also occurred immediately after single-leg cycling, for the RED leg, and then the  
15 NORM leg. Aside from visits three and four, where diet was manipulated to limit glycogen resynthesis,  
16 volunteers were instructed to arrive 3 h after their final meal and caffeine intake. Water was provided *ad libitum*.  
17 Throughout the experimental period, participants were requested to avoid strenuous physical activity.

18

#### 19 *Familiarisation*

20 Familiarisation one commenced with a maximal oxygen uptake ( $\dot{V}O_{2max}$ ) test on a two-leg cycle ergometer  
21 (Excalibur Sport 925900, Lode, Groningen, the Netherlands) to determine single-leg cycling workload. After a  
22 10 min warm up, participants cycled at 80 W for 60 s (~80 rpm), thereafter workload increased by 30 W every  
23 60 s until volitional exhaustion. A 15 min rest was then followed by submaximal, single-leg cycling to adjust the  
24 cycle ergometer settings to the individual, and to calculate participant-workload based upon two-leg  $\dot{V}O_{2max}$   
25 values. The right leg cycled (70 to 80 rpm) unloaded for 10 min, then 3 min load increments were introduced  
26 until individual workload corresponding to 75% two-leg  $\dot{V}O_{2max}$  was maintained. This ~30 min procedure was  
27 followed by a 15 min rest, and then the electrical stimulation protocol to determine maximal (right leg, 657±98  
28 mA; left leg, 670±110 mA) and submaximal (5% MVC) currents (right leg, 111±24 mA; left leg, 114±17 mA) to  
29 be used in subsequent assessments. Forty-eight hours later, familiarisation two involved further electrical  
30 stimulation and single-leg cycling practice, and then isokinetic dynamometer and EMG configuration.

1

2 *Glycogen reduction exercise*

3 Participants arrived at the laboratory between 19:00 and 20:00 h in a 3 h fasted state, to perform exhaustive,  
4 single-leg cycling. The left crank was removed from a cycle ergometer (Monark Ergonomic 824E, Varberg,  
5 Sweden) and replaced with a platform to support the resting left leg. The right foot was strapped to the pedal of  
6 the right crank, and the left leg was secured in a relaxed position on the support stand. Maximal single-leg  
7 cycling workload was calculated as 74% of maximal two-leg workload (Pernow and Saltin, 1971) achieved in  
8 the familiarisation cycling trial. Subsequent workload calculations for single-leg cycling were derivatives of  
9 single-leg maximal workload, and referred to henceforth. Respiratory responses (Table 1) were measured using a  
10 portable metabolic system (Cosmed K4b<sup>2</sup>, Rome, Italy). Single-leg cycling involved: 20 min at 75%  $\dot{V}O_{2max}$   
11 power ( $2.3 \pm 0.3$  kg;  $178 \pm 27$  W); eight 90 s sprints at a 1:1 work-to-rest ratio (5% decrements from 90 to 55%  
12  $\dot{V}O_{2max}$ ); and exhaustive cycling at 85%  $\dot{V}O_{2max}$  for the right limb (Pilegaard et al., 2002). Cadence was  
13 maintained between 70 and 80 rpm; exhaustion was confirmed with an inability to sustain 50 rpm. To limit  
14 localised glycogen resynthesis in the RED leg, 30 min of arm cranking exercise (Monark 864, Varberg, Sweden)  
15 followed single-leg cycling (Camera et al., 2012; Pilegaard et al., 2002). Finally, a low carbohydrate diet was  
16 provided ( $\sim 3932$  kJ: protein  $\sim 960$  kJ, carbohydrate  $\sim 70$  kJ, fat 2907 kJ) consisting of three eggs and six bacon  
17 slices, and for breakfast the next morning, two eggs (Steensberg et al., 2002).

18

19 <<< INSERT TABLE 1 HERE >>>

20

21 *Eccentric exercise*

22 The next morning ( $\sim 06:30$  h), maximal, eccentric knee extensions for the RED leg, and then the NORM leg were  
23 completed. Contractions were performed in a non-randomised order to limit the time-course for glycogen  
24 resynthesis in the RED leg, as resynthesis rate is elevated after exhaustive exercise (Casey et al., 1995). Ten sets  
25 of ten eccentric contractions were performed ( $1.57 \text{ rad}\cdot\text{s}^{-1}$ ), separated by 60 s rests (Byrne et al., 2001) on an  
26 isokinetic dynamometer (Humac Norm, Cybex International Inc., NY, USA). Seated and secured upright (with  
27 the hip joint at 1.57 rad) in the dynamometer chair; the lateral femoral epicondyle of the contracting leg was  
28 aligned with the rotational axis of the gravity corrected dynamometer and the lever arm was fastened to the ankle  
29 joint. Instruction was given to resist the lever arm moving through a 0.09 to 1.74 rad range of motion (ROM)  
30 (full knee extension = 0 rad). Standardised verbal encouragement, and instantaneous torque-time curve feedback

1 from the dynamometer monitor, were provided to ensure maximal effort throughout entire ROM. Peak torque  
2 and negative work done per eccentric set were normalised to mean of eccentric set one.

3

#### 4 *Surface electromyography recording and processing*

5 *Vastus lateralis*, *vastus medialis* and *rectus femoris* EMG were recorded via bipolar sensors, from a Delsys  
6 Bagnoli-8 system (Delsys Inc., Boston, USA) during eccentric exercise. To minimise skin-to-electrode  
7 impedance, the skin was shaved, cleansed and abraded. Sensors were fixed on the mid-aspect of the muscle belly  
8 according to recommendations (Hermens et al., 2000). The reference electrode was placed approximately 4 cm  
9 proximal to the patella apex of the assessed leg. Signals were sampled at 1000 Hz by amplifiers embedded in the  
10 EMG sensors (10 x 1 mm sensor contacts, 10 mm inter-electrode distance; bandwidth = 20-240 Hz; common  
11 mode rejection ratio = 92 dB; input impedance = >1015  $\Omega$ ). Raw EMG data were then processed with a 2<sup>nd</sup> order  
12 Butterworth filter (bandwidth of 10-350 Hz), and the root mean square (RMS) accounting for time-phase shifts  
13 (EMGworks® Data Acquisition and Analysis software, Delsys Inc., Boston, USA). Filtered data were also  
14 treated with a fast Fourier transformation to quantify median frequency (MDF). To ensure isovelocity and  
15 examine EMG activity at different stages of the contraction mid-range, peak torque, RMS and MDF were  
16 calculated for three, 0.26 rad epochs. The first and final 0.48 rad ranges were excluded from analysis due to: i)  
17 some participants being unable to maintain a constant contraction velocity at range of motion limits, and ii) time  
18 for the lever arm to reach constant angular velocity (Fig. 1). Epochs corresponded to estimates of short (first;  
19 0.48 to 0.74 rad), medium (second; 0.74 to 1.0 rad) and long muscle lengths (third; 1.0 to 1.26).

20

21 <<< INSERT FIG. 1. HERE >>>

22

#### 23 *Electrical stimulation and muscle soreness*

24 Knee extensor neuromuscular function was measured on a custom-built chair, with a DS7A electrical stimulator  
25 and NeuroLog pulse generator (Digitimer Limited, Welwyn Garden City, UK). Participants were seated and  
26 secured, (hip and knee at 1.57 rad), with the ankle joint attached via a steel chain leading to a mechanically  
27 calibrated load-cell (RS 250 kg, Tedeo Huntleigh, Cardiff, UK) at the chair base. Electrical stimulation was  
28 received through two saline treated electrodes (9 x 18 cm) positioned over the proximal and distal part of the  
29 thigh. Muscle soreness was reported at rest using a visual analogue scale (0 = not at all sore; 10 = extremely  
30 sore), with the muscle-belly palpated until enough pressure was exerted to blanch the fingernail. Warming up

1 involved three 5 s isometric, submaximal knee extensions and standardised stretching. Maximal twitch was  
2 ascribed, progressively applying 50 mA pulses until the twitch force reached a plateau. Unlike nerve stimulation,  
3 muscle stimulation with large electrodes may involve antagonist co-activation and incomplete agonist activation,  
4 yet it presents less discomfort and can be considered a valid alternative to nerve stimulation (Place et al., 2010).  
5 Submaximal twitch was then calculated (equivalent to 5% isometric MVC force), and measured by delivering a  
6 single, 1 ms stimuli (100 mA, 400 V) to the relaxed knee extensors. Then, participants produced three separate, 3  
7 to 5 s isometric MVC (2 min rests), with verbal encouragement and instantaneous force-time feedback from a  
8 computer monitor. After a 2 min rest, 20 and 50 Hz stimulations (0.5 s duration) were applied using the  
9 submaximal current. Ratio between 20 and 50 Hz forces was calculated; a decrease of which indicates LFF.

10

### 11 *Statistical analysis*

12 Two-way repeated measure ANOVA was used to examine condition-set interaction (RED and NORM) for  
13 neuromuscular responses during and after eccentric exercise. A post-hoc Bonferroni correction was used to  
14 control for multiple comparisons. Eccentric exercise responses (peak torque, negative work done, RMS and  
15 MDF) were compared between contractions one and one-hundred for conditions (RED and NORM), separately  
16 for each three epochs. Effect sizes (Cohen's  $d$ ) were calculated to detect meaningful difference; values are  
17 interpreted as 0.2 for small, 0.5 for moderate, and 0.8 for large differences. Data are expressed as means $\pm$ SD,  
18 with statistical significance set as  $P < 0.05$ .

19

## 20 **Results**

### 21 *Peak torque and negative work*

22 Baseline peak torque was 195.8 $\pm$ 32.9 N·m for RED, and 178.2 $\pm$ 26.8 N·m for NORM ( $P=0.01$ ), respectively.  
23 Peak torque showed a condition-set interaction ( $F_{(1,13)}=11.9$ ,  $P=0.0001$ ,  $d=0.57$ ), decreasing by -16.1 $\pm$ 2.5%  
24 ( $P=0.0001$ ) for RED, and -6.2 $\pm$ 5.1% ( $P=0.004$ ) for NORM during eccentric contractions (Fig. 2). Peak torque  
25 decreased at short muscle length for both legs (RED,  $P=0.02$ , -19.3 $\pm$ 2.8 %,  $d=0.6$ ; NORM,  $P=0.01$ , -18.4 $\pm$ 2.2%,  
26  $d=0.6$ ), and at long muscle length ( $P=0.02$ , -21.7 $\pm$ 1.9%,  $d=0.63$ ) for RED. Negative work decreased during  
27 eccentric contractions ( $F_{(1,13)}=4.9$ ,  $P=0.05$ ,  $d=0.27$ ) by -11.2 $\pm$ 1.4% ( $P=0.002$ ,  $d=0.4$ ) for RED, but not for  
28 NORM ( $P=0.6$ , 3.1 $\pm$ 0.6%,  $d=-0.09$ ) by set ten.

29

30 <<< INSERT FIG. 2. HERE >>>

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30

*Electromyography responses*

*Rectus femoris* RMS was unchanged during eccentric contractions for RED at short ( $P=0.1$ ,  $-2.6\pm 1.7\%$ , Fig. 3) and medium muscle lengths ( $P=0.07$ ,  $-9.6\pm 2.8\%$ ), but increased at long muscle length ( $P=0.03$ ,  $16.2\pm 4.1\%$ ,  $d=0.22$ ). *Rectus femoris* RMS was unchanged during eccentric contractions for NORM at all muscle lengths. *Vastus lateralis* ( $F_{(1,13)}=6.0$ ,  $P=0.08$ ,  $d=0.35$ ) and *vastus medialis* ( $F_{(1,13)}=1.79$ ,  $P=0.07$ ,  $d=0.13$ ) RMS were unchanged for both conditions.

<<< INSERT FIG. 3. HERE >>>

*Rectus femoris* MDF increased at short muscle length for RED ( $P=0.01$ ,  $13.2\pm 4.0\%$ ,  $d=0.56$ ) and at long muscle length for both conditions (RED:  $P=0.02$ ,  $17.9\pm 3.6\%$ ,  $d=0.85$ ; NORM:  $P=0.02$ ,  $15.5\pm 3.2\%$ ,  $d=0.81$ ). *Vastus medialis* MDF differed between conditions ( $F_{(1,13)}=1.98$ ,  $P=0.04$ ,  $d=0.18$ ), unchanged for RED, but decreased for NORM ( $P=0.006$ ,  $-16.3\pm 4.5\%$ ,  $d=0.83$ ) during eccentric exercise. *Vastus lateralis* MDF decreased ( $F_{(1,13)}=2.47$ ,  $P=0.01$ ,  $d=0.25$ ) for both conditions during eccentric exercise (RED:  $P=0.01$ ,  $-20.7\pm 6.2\%$ ,  $d=0.8$ ; NORM:  $P=0.003$ ,  $-15.8\pm 5.2\%$ ,  $d=0.55$ ).

*Neuromuscular recovery*

Single-leg cycling decreased MVC ( $F_{(1,13)}=13.9$ ,  $P=0.0001$ ,  $d=0.50$ ) for RED ( $P=0.0001$ ,  $-23.2\pm 9.2\%$ ,  $d=1.20$ ) and NORM ( $-5.4\pm 5.3\%$ ,  $P=0.002$ ,  $d=0.30$ ) from pre-condition. Eccentric contractions decreased MVC in both conditions at 0 h (RED:  $P=0.0001$ ,  $-23.9\pm 12.0\%$ ,  $d=1.42$ ; NORM:  $P=0.0001$ ,  $-19.4\pm 5.6\%$ ,  $d=1.15$ ), 12 h (RED:  $P=0.0001$ ,  $-10.6\pm 12.1\%$ ,  $d=0.58$ ; NORM:  $P=0.01$ ,  $-9.7\pm 6.6\%$ ,  $d=0.57$ ) and 24 h (RED:  $P=0.0001$ ,  $-15.8\pm 10.3\%$ ,  $d=0.84$ ; NORM:  $P=0.0001$ ,  $-13.5\pm 5.8\%$ ,  $d=0.78$ ). At 48 h MVC was decreased for RED only ( $P=0.003$ ,  $-7.4\pm 10.9\%$ ,  $d=0.36$ ; Fig. 4a). After single-leg cycling, twitch force decreased ( $F_{(1,13)}=7.7$ ,  $P=0.001$ ,  $d=0.39$ ) for RED ( $P=0.001$ ,  $-46.7\pm 14.5\%$ ,  $d=1.47$ ) and increased for NORM ( $P=0.05$ ,  $20.9\pm 14.4\%$ ,  $d=-0.64$ ). Eccentric contractions decreased twitch for RED ( $P=0.0001$ ,  $-41.3\pm 9.1\%$ ,  $d=1.95$ ), but not NORM ( $P=0.2$ ,  $-13.7\pm 8.7\%$ ,  $d=0.44$ ; Fig. 4b). At 12 h, twitch force had recovered for both conditions.

<<< INSERT FIG. 4. HERE >>>



1 Single-leg cycling decreased 20:50 Hz force ratio ( $F_{(1,13)}=14.0$ ,  $P=0.0001$ ,  $d=0.52$ ) for RED only ( $P=0.0001$ ,  
2  $d=1.30$ ). Eccentric contractions decreased 20:50 Hz ratio for both conditions (RED:  $P=0.003$ ,  $0.75\pm0.09$  to  
3  $0.64\pm0.11$ ,  $d=0.93$ ; NORM:  $P=0.01$ ,  $0.76\pm0.12$  to  $0.66\pm0.09$ ,  $d=0.79$ ). The 20:50 Hz ratio was decreased at 12 h  
4 for RED only ( $P=0.03$ ,  $0.70\pm0.09$ ,  $d=0.38$ ; Fig. 4c). At 24 h the 20:50 Hz ratio had recovered in both conditions.  
5 Muscle soreness increased ( $F_{(1,13)}=0.64$ ,  $P=0.03$ ,  $d=0.17$ ) after single-leg cycling for both conditions (RED:  
6  $P=0.006$ ,  $0.7\pm0.3$ ,  $d=0.96$ ; NORM:  $P=0.01$ ,  $0.5\pm0.4$ ,  $d=0.72$ ), and immediately after eccentric contractions  
7 (RED:  $P=0.001$ ,  $2.2\pm0.5$ ,  $d=0.85$ ; NORM:  $P=0.03$ ,  $1.5\pm0.6$ ,  $d=0.68$ ). Soreness remained up to 48 h in both  
8 conditions ( $P>0.05$ , RED:  $2.0\pm1.1$ ,  $2.1\pm0.9$ ,  $1.8\pm1.0$ ; NORM:  $1.8\pm1.0$ ,  $2.2\pm1.4$ ,  $2.0\pm0.8$ ; Fig. 4d).

9

## 10 Discussion

11 This study examined the effect of muscle glycogen reduction on the neuromuscular responses to, and recovery  
12 from mild exercise-induced muscle damage. Performance decline during eccentric contractions with RED was  
13 accompanied by decreased *rectus femoris* and *vastus lateralis* EMG activity, and increased *rectus femoris* EMG  
14 frequency. In recovery, voluntary force loss persisted for longer (up to 48 h), and immediate involuntary force  
15 loss was greater for RED. Twitch force was recovered after 12 h, but 20:50 Hz ratio remained decreased for  
16 RED. Persistent voluntary force loss, with short-term involuntary force loss following eccentric contractions  
17 suggests peripheral and central mechanisms in early recovery, and then central-mediated mechanisms during  
18 prolonged recovery with lowered glycogen.

19 Hody et al., (2013) found negative work during maximal eccentric knee extensions to be associated  
20 with the magnitude of several indirect muscle damage markers. Therefore, the preservation of negative work by  
21 the NORM leg during eccentric contractions is expected to induce greater neuromuscular damage. Conversely,  
22 the greater immediate peripheral fatigue shown for RED may be partially explained by single-leg cycling  
23 imposing metabolic stress, including disrupted  $Ca^{2+}$  homeostasis (Kent-Braun, 1999). That negative work done  
24 did not account for induced damage can also be attributed to the high variability of symptoms reported by Hody  
25 and colleagues (2013). These authors measured plasma creatine kinase activity, muscle soreness and stiffness,  
26 whereas we assessed voluntary force, involuntary force and muscle soreness. We used single-leg cycling  
27 previously shown to reduce muscle glycogen to  $337\pm33$  mmol $\cdot$ kg $^{-1}$  dry weight in the exercising leg (compared to  
28  $609\pm47$  mmol $\cdot$ kg $^{-1}$  dry weight in the resting, control) (Pilegaard et al., 2002). Altered metabolic acidosis  
29 following single-leg cycling, particularly lowered pH, would be likely to disturb muscle membrane excitability  
30 and E-C coupling, subsequently incurring greater peripheral fatigue. Stewart et al., (2008) found cycling to

1 fatigue on three consecutive days decreased 10 Hz force after the first (-40%), and third bouts (-34%). For the  
2 same time-points, MVC force was decreased by -14% and -11%, and 100 Hz force unchanged and decreased by  
3 -16%. After eccentric contractions we found peripheral fatigue transient for RED, which did not account for  
4 persistent neuromuscular impairment. Involuntary force losses were recovered 12 h after eccentric exercise, yet  
5 more severe muscle damage would cause greater neuromuscular impairment with which to study glycogen-  
6 mediated differences with greater sensitivity.

7         We found little change in muscle performance during eccentric contractions under normal conditions;  
8 this concurs with Tesch et al., (1990) who showed torque loss between -34 and -47% for maximal, concentric  
9 contractions, but minimal change for maximal, eccentric contractions. Therefore, the greater forces produced  
10 during eccentric contractions are better preserved. Whereas muscle activity increased, and frequency decreased  
11 with concentric actions, both were unchanged with eccentric (Tesch et al., 1990). Similarly, RMS was  
12 unchanged for NORM, which was supported by maintained muscular performance. In the RED condition,  
13 decreased RMS indicates reduced motor neurone firing, which may reflect an inhibitory response to prevent  
14 further neuromuscular damage. In addition, range of motion data allowed analysis at constant angular velocity.  
15 Decreased peak torque was observed from repetition one to one-hundred at short (-19.3%) and long muscle  
16 lengths (-21.7%) for RED, but only at short length for NORM (-18.4%). Greater force loss at short lengths may  
17 indicate a right-ward shift in muscle length-tension characteristics (Byrne et al., 2001) for NORM, but similar  
18 loss at long length for RED, suggests single-leg cycling also altered contractile structures. Furthermore, peak  
19 torque decreased at short and long muscle lengths for RED indicating advancement, or early onset of contractile  
20 damage. As motor unit activation is incomplete for eccentric contractions (McHugh et al., 2000), NORM would  
21 be expected to have a greater motor unit reserve, than RED, such that it could activate ‘fresh’ units during the  
22 repeated contractions, thus postponing fatigue. Increased MDF during eccentric exercise indicates fatigue, and  
23 may reflect additional recruitment of type II fibres with greater conduction velocity (Kamen and Caldwell,  
24 1996), more so if single-leg cycling depleted type I fibres. We speculate that the higher *rectus femoris* MDF in  
25 RED, than in NORM, during eccentric exercise would indicate higher conduction velocity and greater type II  
26 fibre recruitment, as single-leg cycling preferentially utilised type I muscle fibres and their glycogen stores.  
27 However, because eccentric contractions with RED resulted in unchanged *vastus medialis* MDF, and decreased  
28 *vastus lateralis* MDF, we cannot find a robust association between low glycogen availability and muscle fibre  
29 recruitment during eccentric exercise.

1           Following eccentric contractions, voluntary force production was decreased up to 24 h for NORM and  
2 48 h for RED. Single-leg cycling did not incur lasting neuromuscular function, yet disturbed intramuscular  
3 *milieu* may have activated group III and IV afferents in RED, more than NORM. Heightened perceptual  
4 sensitivity could explain the larger effect sizes of soreness shown for RED after single-leg cycling and eccentric  
5 contractions. Furthermore, afferent feedback is known to facilitate central fatigue (Amann et al., 2011), which is  
6 consistent with our findings of decreased voluntary force (up to 48 h) and limited LFF (up to 24 h) shown for  
7 RED. Low glycogen availability leads to a decreased ATP resynthesis rate, and therefore reduced intracellular  
8 ATP. This compromises energy demand to satisfy E-C coupling, and results in an inability to convert neural  
9 drive into force output (Green, 1991). In this study LFF remained up to 12 h post-eccentric in both conditions,  
10 suggesting glycogen availability had little effect on E-C coupling disruption. However, the active limb  
11 experienced decreased 20 (-47.9%) and 50 Hz forces (-32.8%) after single-leg cycling.

12           A limitation to the current study is the lack of biopsy/blood sampling, making it difficult to: i) confirm  
13 muscle glycogen reduction, and ii) discern whether single-leg cycling resulted in structural and metabolic  
14 changes in the RED leg. We are confident the pre-validated, single-leg cycling protocol reduced glycogen in the  
15 RED leg, however biopsies would have provided direct assessment of glycogen reduction, and recovery.  
16 Furthermore, it is likely that the neuromuscular differences between conditions were not solely due to muscle  
17 glycogen concentration, but other structural and metabolic alterations caused by single-leg cycling. In terms of  
18 measurement tools, EMG, muscle stimulation and indirect calorimetry are easily administered, non-invasive  
19 methods to quantify neuromuscular behaviour.

20

## 21 **Conclusions**

22 During eccentric exercise, reduced muscle glycogen resulted in a failure to maintain muscular performance, and  
23 a decreased muscle activity. In recovery, eccentric exercise with reduced glycogen caused greater, early evoked  
24 force loss, delayed voluntary force recovery, independent of muscle soreness. Immediate involuntary force loss,  
25 with prolonged voluntary force loss, indicates reduced glycogen is associated with both central and peripheral  
26 mechanisms in early recovery and then central mechanisms in the later recovery. These results suggest that  
27 reduced muscle glycogen impairs muscular work capacity and alters neuromuscular fatigue profiles following  
28 mild exercise-induced muscle damage of the knee extensors. These findings have application to individuals  
29 performing repeat training bouts and mechanistically, understanding the impact of performing damaging exercise  
30 in a pre-fatigued state.

1

2 **Acknowledgements:** No external funding sources were used to prepare this article.

3

4 **Conflicts of Interest:** There are no conflicts of interests concerning the preparation of this article.

5

6 **References**

7 Amann M, Blain GM, Proctor LT, Sebranek JJ, Pegelow DF, Dempsey JA. Implications of group III and IV  
8 muscle afferents for high-intensity endurance exercise performance in humans. *J Physiol* 2011;589(21):5299-  
9 309.

10 Asp S, Daugaard JR, Kristiansen S, Kiens B, Richter EA. Exercise metabolism in human skeletal muscle  
11 exposed to prior eccentric exercise. *J Physiol* 1998;509(1):305-13.

12 Behrens M, Mau-Moeller A, Bruhn S. Effect of exercise-induced muscle damage on neuromuscular function of  
13 the quadriceps muscle. *Int J Sports Med* 2012;33(8):600-6.

14 Beltman JG, Sargeant AJ, van Mechelen W, de Haan A. Voluntary activation level and muscle fiber recruitment  
15 of human quadriceps during lengthening contractions. *J Appl Physiol* 2004;97(2):619-26.

16 Bigland-Ritchie B, Furbush F, Woods JJ. Fatigue of intermittent submaximal voluntary contractions: central and  
17 peripheral factors. *J Appl Physiol* 1986;61(2):421-29.

18 Byrne C, Eston RG, Edwards RH. Characteristics of isometric and dynamic strength loss following eccentric  
19 exercise-induced muscle damage. *Scand J Med Sci Sports* 2001;11(3):134-40.

20 Camera DM, West DW, Burd NA, Phillips SM, Garnham AP, Hawley JA, et al. Low muscle glycogen  
21 concentration does not suppress the anabolic response to resistance exercise. *J Appl Physiol* 2012;113(2):206-14.

22 Casey A, Short AH, Hultman E, Greenhaff PL. Glycogen resynthesis in human muscle fibre types following  
23 exercise-induced glycogen depletion. *J Physiol* 1995;483(1):265-71.

24 Clarkson PM, Sayers SP. Etiology of exercise-induced muscle damage. *Can J Appl Physiol* 1999;24(3):234-48.

25 Edwards RH. Hypotheses of peripheral and central mechanisms underlying occupational muscle pain and injury.  
26 *Eur J Appl Physiol Occup Physiol* 1988;57(3):275-81.

27 Faul F, Erdfelder E, Lang AG, Buchner A. G\*Power 3: a flexible statistical power analysis program for the  
28 social, behavioral, and biomedical sciences. *Behav Res Methods* 2007;39(2):175-91.

29 Green H J. How important is endogenous muscle glycogen to fatigue in prolonged exercise? *Can J Physiol*  
30 *Pharmacol* 1991;69(2):290-97.

1 Hermens HJ, Freriks B, Disselhorst-Klug C, Rau G. Development of recommendations for SEMG sensors and  
2 sensor placement procedures. *J Electromyogr Kines* 2000;10(5):361-74.

3 Hody S, Rogister B, Leprince P, Wang F, Croisier JL. Muscle fatigue experienced during maximal eccentric  
4 exercise is predictive of the plasma creatine kinase (CK) response. *Scand J Med Sci Sports* 2013;23(4):501-7.

5 Kamen G, Caldwell GE. Physiology and interpretation of the electromyogram. *J Clin Neurophysiol*  
6 1996;13(5):366-84.

7 Kent-Braun JA. Central and peripheral contributions to muscle fatigue in humans during sustained maximal  
8 effort. *Eur J Appl Physiol Occup Physiol* 1999;80(1):57-63.

9 McHugh MP, Connolly DA, Eston RG, Gleim GW. Electromyographic analysis of exercise resulting in  
10 symptoms of muscle damage. *J Sports Sci* 2000;18(3):163-72.

11 Morgan DL, Gregory JE, Proske U. The influence of fatigue on damage from eccentric contractions in the  
12 gastrocnemius muscle of the cat. *J Physiol* 2004;561(3):841-50.

13 Nardone A, Romano C, Schieppati M. Selective recruitment of high-threshold human motor units during  
14 voluntary isotonic lengthening of active muscles. *J Physiol* 1989;409:451-71.

15 Pernow B, Saltin B. Availability of substrates and capacity for prolonged heavy exercise in man. *J Appl Physiol*  
16 1971;31(3):416-22.

17 Pilegaard H, Keller C, Steensberg A, Helge JW, Pedersen BK, Saltin B, et al. Influence of pre-exercise muscle  
18 glycogen content on exercise-induced transcriptional regulation of metabolic genes. *J Physiol* 2002;541(1):261-  
19 71.

20 Place N, Casartelli N, Glatthorn JF, Maffiuletti NA. Comparison of quadriceps inactivation between nerve and  
21 muscle stimulation. *Muscle Nerve* 2010;42(6):894-900.

22 Skurvydas A, Sipaviciene S, Krutulyte G, Gailiuniene A, Stasiulis A, Mamkus G, et al. Dynamics of indirect  
23 symptoms of skeletal muscle damage after stretch-shortening exercise. *J Electromyogr Kines* 2006;16(6):629-36.

24 Steensberg A, van Hall G, Keller C, Osada T, Schjerling P, Pedersen BK, et al. Muscle glycogen content and  
25 glucose uptake during exercise in humans: influence of prior exercise and dietary manipulation. *J Physiol*  
26 2002;541(1):273-81.

27 Stewart RD, Duhamel TA, Rich S, Tupling AR, Green HJ. Effects of consecutive days of exercise and recovery  
28 on muscle mechanical function. *Med Sci Sports Exerc* 2008;40(2):316-25.

29 Tesch PA, Dudley GA, Duvoisin MR, Hather BM, Harris RT. Force and EMG signal patterns during repeated  
30 bouts of concentric or eccentric muscle actions. *Acta Physiol Scand* 1990;138(3):263-71.

1 Verburg E, Dutka TL, Lamb GD. Long-lasting muscle fatigue: partial disruption of excitation-contraction  
2 coupling by elevated cytosolic Ca<sup>2+</sup> concentration during contractions. *Am J Physiol Cell Physiol*  
3 2006;290(4):C1199-208.

4 Warren GL, Ingalls CP, Lowe DA, Armstrong RB. Excitation-contraction uncoupling: major role in contraction-  
5 induced muscle injury. *Exerc Sport Sci Rev* 2001;29(2):82-7.

6 Westerblad H, Duty S, Allen DG. Intracellular calcium concentration during low-frequency fatigue in isolated  
7 single fibers of mouse skeletal muscle. *J Appl Physiol* 1993; 75(1): 382-388.

8 Widrick JJ, Costill DL, McConnell GK, Anderson DE, Pearson DR, Zachwieja JJ. Time course of glycogen  
9 accumulation after eccentric exercise. *J Appl Physiol* 1992;72(5):1999-2004.

10  
11  
12  
13  
14  
15  
16  
17  
18  
19

1 **Tables**

2 **Table 1** Mean physiological responses to single-leg muscle glycogen reduction cycling.

	75% VO <sub>2max</sub>				90%	85%	80%	75%	70%	65%	60%	55%	85% VO <sub>2max</sub>
	5 min	10 min	15 min	20 min	90 s sprints (1:1 work-to-rest ratio)								Exhaustive
Workload (W)	124±30	126±29	126±32	138±29	162 ±33	152±34	147±34	140±36	130±37	125±34	121±29	121±25	134±35
Cadence (rpm)	71.9±5.2	70.4±3.0	69.7±3.7	68.6±7.4	74.8±6.2	74.8±5.1	76.6±4.7	76.6±5.8	77.8±7.4	78.9±6.0	80.8±6.6	81.0±7.6	66.1±7.7
V <sub>E</sub> (L·min <sup>-1</sup> )	76.2±14.3	87.7±21.6	93.2±18.7	93.7±21.8	92.6±16.6	98.5±19.8	92.3±20.7	87.4±19.5	86.8±20.3	80.1±19.2	72.9±20.7	72.4±20.1	102.2±23.0
VO <sub>2</sub> (L·min <sup>-1</sup> ) 1)	2.28±0.55	2.30±0.54	2.46±0.63	2.49±0.53	2.59±0.44	2.6±0.47	2.55±0.4	2.5±0.5	2.46±0.53	2.44±0.43	2.33±0.56	2.19±0.48	2.45±0.69
VCO <sub>2</sub> (L·min <sup>-1</sup> )	2.39±0.65	2.30±0.59	2.35±0.53	2.44±0.54	2.57±0.44	2.46±0.43	2.33±0.32	2.21±0.38	2.20±0.46	2.09±0.26	1.95±0.47	1.89±0.41	2.33±0.74
VO <sub>2</sub> (mL·kg <sup>-1</sup> ·min <sup>-1</sup> ) 1)	29.9±8.9	30.0±8.4	32.0±9.1	32.4±7.7	33.8±7.2	34.0±7.5	33.3±6.6	32.5±7.2	32.2±7.7	32.1±7.3	30.3±7.5	28.7±6.8	31.6±9.0
RER	1.04±0.13	1.00±0.14	0.95±0.15	0.98±0.12	1.00±0.11	0.95±0.12	0.92±0.11	0.89±0.1	0.90±0.1	0.87±0.09	0.84±0.1	0.87±0.11	0.93±0.2

---

HR (b·min <sup>-1</sup> )	170±14	165±17	159±21	153±20	171±15	172±14	170±16	170±12	166±12	162±12	160±13	157±12	177±8
---------------------------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	-------

---


1 Values are mean±SD; V<sub>E</sub>, minute ventilation; VO<sub>2</sub>, oxygen uptake; VCO<sub>2</sub>, carbon dioxide production; RER, respiratory exchange ratio; HR, heart rate.

2


3




1 Figures




2



3




1



2


3 **Fig. 1.** A schematic of the knee angles during maximal, eccentric knee extensions performed on an isokinetic  
 4 dynamometer at  $1.57 \text{ rad}\cdot\text{s}^{-1}$  (a). To ensure isovelocity and examine the influence of muscle length, peak torque  
 5 (b), angular velocity (c), EMG root mean square (RMS) (d) and median frequency (MDF) recorded during  
 6 eccentric contractions were averaged across three,  $0.26 \text{ rad}$  epochs. With the first and final  $0.48 \text{ rad}$  excluded,  
 7 epochs corresponded to short (**S**) (first;  $0.48$  to  $0.74 \text{ rad}$ ), medium (**M**) (second;  $0.74$  to  $1.0 \text{ rad}$ ) and long (**L**)  
 8 muscle lengths (third;  $1.0$  to  $1.26$ ).



1


2 **Fig. 2.** Peak torque (normalised to set one) during maximal eccentric knee extensions with reduced (RED) and  
 3 normal (NORM) glycogen availability legs. Data presented are mean±SD. \* Significant difference between  
 4 conditions, # significant difference between sets one and ten,  $P < 0.05$ .

5




6


**a**



1



2



**d**

1


2 **Fig. 3.** *Rectus femoris* root mean square (RMS) (a) and median frequency (MDF) (b), and *vastus medialis* MDF

3 (c) and *vastus lateralis* MDF (d) during maximal eccentric knee extensions with reduced (RED) and normal

4 (NORM) glycogen availability legs. Data presented are mean±SD. \* Significant difference between conditions, #


5 significant difference between sets one and ten,  $P < 0.05$ . MDF data refer to total epoch mean, not individual

6 epoch mean.




**a**


7



1



2



1

2 **Fig. 4.** Isometric maximal voluntary contraction (MVC) force (a), twitch force (b), 20:50 Hz force ratio (c) and  
 3 muscle soreness (d) following maximal eccentric knee extensions with reduced (RED) and normal (NORM)  
 4 glycogen availability legs. Data presented are mean±SD. \* Significant difference between conditions, P < 0.05.

5