

ON COHERENT SYSTEMS OF TYPE  $(n, d, n + 1)$   
ON PETRI CURVES

*U. N. Bhosle, L. Brambila-Paz and P. E. Newstead*

Comunicación Técnica No I-08-05/22-01-2008  
(MB/CIMAT)



# ON COHERENT SYSTEMS OF TYPE $(n, d, n + 1)$ ON PETRI CURVES

U. N. BHOSLE, L. BRAMBILA-PAZ AND P. E. NEWSTEAD

ABSTRACT. We study coherent systems of type  $(n, d, n + 1)$  on a Petri curve  $X$  of genus  $g \geq 2$ . We describe the geometry of the moduli space of such coherent systems for large values of the parameter  $\alpha$ . We determine the top critical value of  $\alpha$  and show that the corresponding “flip” has positive codimension. We investigate also the non-emptiness of the moduli space for smaller values of  $\alpha$ , proving in many cases that the condition for non-emptiness is the same as for large  $\alpha$ . We give some detailed results for  $g \leq 5$  and applications to higher rank Brill-Noether theory and the stability of kernels of evaluation maps, thus proving Butler’s conjecture in some cases in which it was not previously known.

## 1. INTRODUCTION

Let  $X$  be a smooth irreducible projective curve. A coherent system of type  $(n, d, k)$  on  $X$  is a pair  $(E, V)$  where  $E$  is a vector bundle on  $X$  of rank  $n$  and degree  $d$  and  $V$  is a linear subspace of  $H^0(E)$  with  $\dim V = k$ . A notion of stability for coherent systems, dependent on a real variable  $\alpha$ , can be defined and leads to the construction of moduli spaces  $G(\alpha; n, d, k)$  for  $\alpha$ -stable coherent systems (see [16], [19], [26]). There is a natural compactification  $\tilde{G}(\alpha; n, d, k)$  obtained by considering equivalence classes of  $\alpha$ -semistable coherent systems. For  $k = 0$ ,  $G(\alpha; n, d, 0)$  is independent of  $\alpha$  and coincides with the moduli space  $M(n, d)$  of stable bundles of rank  $n$  and degree  $d$  on  $X$ , while  $\tilde{G}(\alpha; n, d, 0)$  coincides with the corresponding moduli space  $\tilde{M}(n, d)$  of S-equivalence classes of semistable bundles. If  $k \geq 1$ , a necessary condition for non-emptiness of  $G(\alpha; n, d, k)$  (resp.  $\tilde{G}(\alpha; n, d, k)$ ) is  $\alpha > 0$  (resp.  $\alpha \geq 0$ ). For  $n = 1$ , all coherent systems are  $\alpha$ -stable for all  $\alpha > 0$  and  $G(\alpha; 1, d, k)$  coincides with the classical variety of linear systems  $G_d^{k-1}$ .

A systematic study of coherent systems on curves of genus  $g \geq 2$  defined over the complex numbers was begun in [5] (see also [4]) and continued in [6] and [7]. In particular, precise conditions for non-emptiness of  $G(\alpha; n, d, k)$  are known when  $k \leq n$  [6, Theorem 3.3]. For  $k > n$ , much less is known. There are general results due to E. Ballico [2] and

---

*Date:* December 13, 2007.

*2000 Mathematics Subject Classification.* 14H60.

*Key words and phrases.* coherent systems, stability, Brill-Noether, Petri curve.

The authors are members of the research group VBAC (Vector Bundles on Algebraic Curves). The first two authors were supported by EPSRC grant GR/T22988/01 for a visit to the University of Liverpool. The second author acknowledges the support of CONACYT grant 48263-F. The third author thanks CIMAT, Guanajuato, México and California State University Channel Islands, where a part of this paper was completed, and acknowledges support from the Academia Mexicana de Ciencias, under its exchange agreement with the Royal Society of London.

M. Teixidor i Bigas [30]; Teixidor's results are much the stronger, but are certainly not best possible. Some more detailed results have been obtained in [8, 9]. It is known that the  $\alpha$ -stability condition stabilises for  $\alpha > d(n-1)$ ; we denote the corresponding "large  $\alpha$ " moduli space  $G(\alpha; n, d, k)$  by  $G_L(n, d, k)$  (see section 2 for more details).

Our object in this paper is to study the case  $k = n + 1$  when the curve  $X$  is a *Petri curve*, in other words, for every line bundle  $\mathcal{L}$  on  $X$ , the multiplication map

$$H^0(\mathcal{L}) \otimes H^0(\mathcal{L}^* \otimes K) \rightarrow H^0(K)$$

is injective. In this case  $G_L := G_L(\alpha; n, d, n+1)$  is non-empty if and only if the *Brill-Noether number*

$$\beta := \beta(n, d, n+1) = g - (n+1)(n-d+g)$$

is non-negative [5, Theorem 5.11]. When in addition  $d \leq g+n$ ,  $G(\alpha) := G(\alpha; n, d, n+1)$  is independent of  $\alpha > 0$  and its structure has been determined [8, Theorem 2]. Our first main theorem (Theorem 3.1) generalises these results and gives a significant improvement of the estimate  $\alpha > d(n-1)$  for  $G(\alpha)$  to coincide with  $G_L$ . The detailed statement, which includes additional information on the structure of  $G_L$ , is as follows (here  $E'$  denotes the subsheaf image of the evaluation map  $V \otimes \mathcal{O} \rightarrow E$ ; for the definitions of *generated* and *generically generated*, see section 2).

**Theorem 3.1.** *Suppose that  $X$  is a Petri curve of genus  $g \geq 2$  and  $\alpha > \max\{0, \alpha_l\}$ , where*

$$\alpha_l := d(n-1) - n \left( n-1 + g - \left\lfloor \frac{g}{n} \right\rfloor \right).$$

*Then*

- (1)  $G(\alpha) \neq \emptyset$  if and only if  $\beta \geq 0$ ;
- (2)  $G(\alpha) = G_L$ ;
- (3)  $(E, V) \in G(\alpha)$  if and only if  $(E, V)$  is generically generated and  $H^0(E'^*) = 0$ ;
- (4) if  $\beta > 0$ ,  $G(\alpha)$  is smooth and irreducible of dimension  $\beta$ ; moreover the generic element of  $G(\alpha)$  is generated;
- (5) if  $\beta = 0$ ,  $G(\alpha)$  is a finite set of cardinality

$$g! \prod_{i=0}^n \frac{i!}{(g-d+n+i)!};$$

*moreover every element of  $G(\alpha)$  is generated.*

It follows in particular that, if  $(E, V) \in G_L$ , then the cokernel  $E/E'$  of the evaluation map  $V \otimes \mathcal{O} \rightarrow E$  is a torsion sheaf. In section 4, we define a stratification of  $G_L$  in terms of the length of  $E/E'$ . More precisely, for every integer  $t \geq 0$ , we write

$$\Sigma_t = \{(E, V) \in G_L : E/E' \text{ has length } t\} \quad \text{and} \quad S_t = \bigcup_{i \geq t} \Sigma_i.$$

Then

**Theorem 4.2.** *Suppose  $\beta \geq 0$  and that the subsets  $S_t$  of  $G_L$  are defined as above. Then*

- (1)  $S_t$  is closed in  $G_L$  and is non-empty if and only if  $0 \leq t \leq t_1 := \lfloor \frac{\beta}{n+1} \rfloor$ ;
- (2) for  $1 \leq t \leq t_1$ ,  $S_t \subset \overline{S_{t-1}} \setminus S_t$ ;
- (3) for  $1 \leq t \leq t_1$ ,  $\dim S_t = \beta - t$ ;
- (4)  $S_t$  is irreducible for  $t < \frac{\beta}{n+1}$ ;
- (5) if  $\frac{\beta}{n+1}$  is an integer, then all irreducible components of  $S_{t_1}$  have the same dimension.

In section 5, we show that there exists  $(E, V) \in G_L$  such that  $(E, V)$  is not  $\alpha_l$ -stable, in other words  $\alpha_l$  is an (actual) critical value in the sense of [5, Definition 2.4]. In view of Theorem 3.1,  $\alpha_l$  is in fact the top critical value of  $\alpha$ .

Sections 6 – 8 are concerned with the moduli space  $G(\alpha)$  for arbitrary  $\alpha$ . It was proved in [8] that, if  $G(\alpha) \neq \emptyset$ , then  $\beta \geq 0$ . Several results on the non-emptiness of  $G(\alpha)$  when  $\beta \geq 0$  were also proved in [8]. In section 6, we extend these results using the techniques of elementary transformations and extensions of coherent systems. In particular for  $n = 2, 3, 4$ , we show in section 7 that  $G(\alpha) \neq \emptyset$  if and only if  $\beta \geq 0$  (see Theorems 7.1, 7.2 and 7.3 for details). We then consider in section 8 the case  $g \leq 5$  (including  $g = 0$  and  $g = 1$ , which have been excluded from our general discussion). For  $g \leq 2$ , the results are complete, while for  $g = 3, 4, 5$ , there are a few cases still to be solved.

In section 9, we give some applications to higher rank Brill-Noether theory (see section 2 for definitions). We first obtain some irreducibility and smoothness results for Brill-Noether loci using the programme envisaged in [5, section 11]. For the second application, suppose that  $\mathcal{L}$  is a generated line bundle of degree  $d > 0$  and let  $V$  be a linear subspace of  $H^0(\mathcal{L})$  of dimension  $n + 1$  which generates  $\mathcal{L}$  (in other words,  $(\mathcal{L}, V)$  is a generated coherent system of type  $(1, d, n + 1)$ ). We have an evaluation sequence

$$0 \longrightarrow M_{V, \mathcal{L}} \longrightarrow V \otimes \mathcal{O} \longrightarrow \mathcal{L} \longrightarrow 0.$$

The bundles  $M_{V, \mathcal{L}}$  arise in several contexts and have been used in the study of Picard bundles [13], normal generation of vector bundles [25, 11], syzygies and projective embeddings [14], higher rank Brill-Noether loci [20], theta-divisors [3, 23] and coherent systems [12, 5, 8].

A particular point of interest is to determine whether or not  $M_{V, \mathcal{L}}$  is stable. In fact, in [12], Butler conjectured that  $M_{V, \mathcal{L}}$  is stable for general choices of  $X$ ,  $\mathcal{L}$  and  $V$ . His conjecture [12, Conjecture 2] is concerned more generally with generated coherent systems of any type  $(n, d, k)$ . We shall be concerned only with the case  $n = 1$ ; Butler's conjecture can then be stated as follows.

**Conjecture 9.5.** *Let  $X$  be a Petri curve of genus  $g \geq 3$ . Suppose that  $\beta := \beta(1, d, n + 1) \geq 0$  and that  $\mathcal{L}$  is a general element of  $B(1, d, n + 1)$  (when  $\beta = 0$ ,  $\mathcal{L}$  can be any element*

of the finite set  $B(1, d, n + 1)$ ) and let  $V$  be a general subspace of  $H^0(\mathcal{L})$  of dimension  $n + 1$ . Then  $M_{V, \mathcal{L}}$  is stable.

In most of the above references,  $V$  is taken to be  $H^0(\mathcal{L})$ , which implies by Riemann-Roch that  $d \leq g + n$  and the stability problem has been solved in this case [12, 8]. However the case where  $V$  is a proper subspace of  $H^0(\mathcal{L})$  seems equally interesting; this is mentioned but not used in [12], used in a minor way in [5] and studied for low values of the codimension in [23]. However, the restriction placed on  $d$  in [23] implies that  $d \leq 2n$ , so this case (although not the remaining results of [23]) is also covered in [20, 22]. In the present paper, we do not use the stability of  $M_{V, \mathcal{L}}$  except through citations from earlier papers. We are therefore able to use our methods to prove the stability of  $M_{V, \mathcal{L}}$  in some cases where it is not (to our knowledge) already known. These new examples for which  $M_{V, \mathcal{L}}$  is stable depend essentially on the use of extensions of coherent systems (more specifically on Propositions 6.9, 6.10, 6.12, 7.5 and 7.6).

We assume throughout that  $X$  is a Petri curve of genus  $g$ , where, except in section 8,  $g \geq 2$ . We assume also that  $X$  is defined over the complex numbers. We denote the canonical line bundle on  $X$  by  $K$ .

## 2. PRELIMINARIES

In this section, we recall some facts about coherent systems, most of which can be found in [5] and [15].

For  $\alpha \in \mathbb{R}$ , we define the  $\alpha$ -slope of the coherent system  $(E, V)$  of type  $(n, d, k)$  by

$$\mu_\alpha(E, V) := \frac{d}{n} + \alpha \frac{k}{n}.$$

A *coherent subsystem* of  $(E, V)$  is a pair  $(F, W)$ , where  $F$  is a subbundle of  $E$  and  $W \subset V \cap H^0(F)$ .

**Definition 2.1.** For any  $\alpha \in \mathbb{R}$ , a coherent system  $(E, V)$  on  $X$  is  $\alpha$ -stable (respectively  $\alpha$ -semistable) if, for every proper coherent subsystem  $(F, W)$ ,

$$\mu_\alpha(F, W) < \mu_\alpha(E, V) \quad (\text{respectively } \leq).$$

We denote by  $G(\alpha; n, d, k)$  the moduli space of  $\alpha$ -stable coherent systems of type  $(n, d, k)$  ([16], [19], [26]) and by  $\tilde{G}(\alpha; n, d, k)$  the moduli space of S-equivalence classes of  $\alpha$ -semistable coherent systems (see [5, section 2]). It follows from the definition of  $\alpha$ -stability that, if  $k \geq 1$  and  $G(\alpha; n, d, k) \neq \emptyset$ , then  $\alpha > 0$  and  $d > 0$  [5, section 2 and Lemmas 4.1 and 4.3].

**Remark 2.2.** Given a coherent system  $(E, V)$  and an effective line bundle  $\mathcal{L}$ , let  $\tilde{E} = E \otimes \mathcal{L}$ . Choose a non-zero section  $s$  of  $\mathcal{L}$  and let  $\tilde{V}$  be the image of  $V$  in  $H^0(\tilde{E})$  under the induced inclusion  $H^0(E) \hookrightarrow H^0(\tilde{E}) : v \mapsto v \otimes s$ . Then

- (1)  $E$  is (semi)stable if and only if  $\tilde{E}$  is (semi)stable.

(2)  $(E, V)$  is  $\alpha$ -(semi)stable if and only if  $(\tilde{E}, \tilde{V})$  is  $\alpha$ -(semi)stable [26, Lemma 1.5].

**Remark 2.3.** It follows from Remark 2.2 that, if  $G(\alpha; n, d, k) \neq \emptyset$  for all integers  $d \in [a, b]$  with  $a, b \in \mathbb{Z}$  and  $b - a \geq n - 1$ , then  $G(\alpha; n, d, k) \neq \emptyset$  for all  $d \geq a$ .

For any triple  $(n, d, k)$ , we define the *Brill-Noether number*  $\beta(n, d, k)$  by

$$\beta(n, d, k) = n^2(g - 1) + 1 - k(k - d + n(g - 1)).$$

For a coherent system  $(E, V)$ , the *Petri map* at  $(E, V)$  is the map

$$(2.1) \quad V \otimes H^0(E^* \otimes K) \rightarrow H^0(E \otimes E^* \otimes K)$$

given by multiplication of sections. We have the following fundamental result (see [15, Corollaire 3.14], [5, Corollary 3.6 and Proposition 3.10]).

**Proposition 2.4.** *Every irreducible component of  $G(\alpha; n, d, k)$  has dimension  $\geq \beta(n, d, k)$ . Moreover, if  $(E, V) \in G(\alpha; n, d, k)$ , then  $G(\alpha; n, d, k)$  is smooth of dimension  $\beta(n, d, k)$  at  $(E, V)$  if and only if (2.1) is injective.*

For a line bundle  $\mathcal{L}$  with  $V = H^0(\mathcal{L})$ , the Petri map (2.1) takes the form

$$(2.2) \quad H^0(\mathcal{L}) \otimes H^0(\mathcal{L}^* \otimes K) \rightarrow H^0(K)$$

**Definition 2.5.** The curve  $X$  is a *Petri curve* if (2.2) is injective for every line bundle  $\mathcal{L}$  on  $X$ .

It is a classical fact (see [1]) that the general curve of any given genus  $g$  is a Petri curve. It should however be emphasised that, except for certain low values of the genus, there exist  $\alpha$ -stable coherent systems  $(E, V)$  on the general curve for which (2.1) is not injective (see, for example, [29, §5]).

The  $\alpha$ -range is divided into a finite set of intervals by a set of *critical values*  $\{\alpha_i\}$ , where, for  $k \geq n$ ,

$$0 = \alpha_0 < \alpha_1 < \cdots < \alpha_L < \infty$$

[5, Proposition 4.6]. For  $\alpha, \alpha' \in (\alpha_i, \alpha_{i+1})$ , we have  $G(\alpha; n, d, k) = G(\alpha'; n, d, k)$  and we denote this moduli space by  $G_i := G_i(n, d, k)$ . In particular, for  $\alpha > \alpha_L$ , we have the “large  $\alpha$ ” moduli space  $G_L := G_L(n, d, k)$ .

The relation between two consecutive moduli spaces  $G_{i-1}$  and  $G_i$  is given by the so called “flips” (see [5] for a more complete description). For any critical value  $\alpha_i$ , we denote by  $\alpha_i^-$ ,  $\alpha_i^+$  values of  $\alpha$  in the intervals respectively immediately before and after  $\alpha_i$  and let

$$G_i^+ := \{(E, V) \in G_i \mid (E, V) \text{ is not } \alpha_i^- \text{-stable}\}$$

and

$$G_i^- := \{(E, V) \in G_{i-1} \mid (E, V) \text{ is not } \alpha_i^+ \text{-stable}\}.$$

These are called *flip loci* and

$$(2.3) \quad G_i - G_i^+ = G_{i-1} - G_i^-.$$

For any critical value  $\alpha_i$ , the flip locus  $G_i^+$  consists of the coherent systems  $(E, V) \in G_i$  for which there exists an exact sequence

$$(2.4) \quad 0 \rightarrow (E_1, V_1) \rightarrow (E, V) \rightarrow (E_2, V_2) \rightarrow 0,$$

with  $(E_j, V_j)$  of type  $(n_j, d_j, k_j)$ ,  $\alpha_i$ -semistable and  $\alpha_i^+$ -stable for  $j = 1, 2$  and

$$(2.5) \quad \mu_{\alpha_i}(E_1, V_1) = \mu_{\alpha_i}(E_2, V_2), \quad k_1/n_1 < k/n$$

(see [5, Lemma 6.5] for more details). Similarly, the flip locus  $G_i^-$  consists of the coherent systems  $(E, V) \in G_{i-1}$  for which there exists an exact sequence

$$0 \rightarrow (E_2, V_2) \rightarrow (E, V) \rightarrow (E_1, V_1) \rightarrow 0,$$

with  $(E_j, V_j)$   $\alpha_i$ -semistable and  $\alpha_i^-$ -stable for  $j = 1, 2$  and satisfying (2.5).

In [5], numerical criteria were obtained to help determine whether the flip loci have positive codimension. More generally, these criteria can be used to estimate the number of parameters on which the coherent systems  $(E, V)$  given by extensions (2.4) depend. Define, for  $\{j, l\} = \{1, 2\}$ ,

$$(2.6) \quad \begin{aligned} C_{jl} &= n_j n_l (g-1) - n_j d_l + n_l d_j + k_j d_l - k_j n_l (g-1) - k_j k_l \\ &= (k_j - n_j)(d_l - n_l (g-1)) + n_l d_j - k_j k_l \end{aligned}$$

and

$$(2.7) \quad \mathbb{H}_{jl}^0 = \text{Hom}((E_j, V_j), (E_l, V_l)), \quad \mathbb{H}_{jl}^2 = H^0(E_l^* \otimes N_j \otimes K)^*,$$

$N_j$  being the kernel of the evaluation map  $V_j \otimes \mathcal{O} \rightarrow E_j$ . We have, by [5, equations (8) and (11)],

$$(2.8) \quad \dim \text{Ext}^1((E_j, V_j), (E_l, V_l)) = C_{jl} + \dim \mathbb{H}_{jl}^0 + \dim \mathbb{H}_{jl}^2.$$

The following lemma can be regarded as a simplified version of [5, Lemma 6.8].

**Lemma 2.6.** *Suppose that, for  $j = 1, 2$ ,  $(E_j, V_j)$  has type  $(n_j, d_j, k_j)$  and varies in a family depending on at most  $\beta(n_j, d_j, k_j)$  parameters. Suppose further that, for some  $h_0, h_2$ ,*

$$\dim \mathbb{H}_{21}^0 \leq h_0, \quad \dim \mathbb{H}_{21}^2 \leq h_2$$

for all  $(E_j, V_j)$  occurring in these families and that

$$C_{12} - h_0 - h_2 > 0.$$

Then the coherent systems  $(E, V)$  arising as non-trivial extensions of the form (2.4) depend on at most  $\beta(n, d, k) - 1$  parameters.

*Proof.* By (2.8), for fixed  $(E_1, V_1), (E_2, V_2)$ , the coherent systems  $(E, V)$  depend on at most

$$C_{21} + h_0 + h_2 - 1$$

parameters. The result follows from [5, Corollary 3.7].  $\square$

**Remark 2.7.** Note that, if we assume in addition that  $(E, V)$  is  $\alpha$ -stable for some  $\alpha$ , then we can take  $h_0 = 0$ , since a non-zero homomorphism  $(E_2, V_2) \rightarrow (E_1, V_1)$  would contradict [5, Proposition 2.2(ii)].

The “small  $\alpha$ ” moduli spaces  $G_0(n, d, k)$  and  $\tilde{G}_0(n, d, k)$  are closely related to the Brill-Noether locus  $B(n, d, k)$  of stable bundles, which is defined by

$$B(n, d, k) := \{E \in M(n, d) \mid h^0(E) \geq k\}.$$

Similarly one defines the Brill-Noether locus  $\tilde{B}(n, d, k)$  for semistable bundles by

$$\tilde{B}(n, d, k) := \{[E] \in \tilde{M}(n, d) \mid h^0(\text{gr}(E)) \geq k\},$$

where  $\tilde{M}(n, d)$  is the moduli space of S-equivalence classes of semistable bundles,  $[E]$  is the S-equivalence class of  $E$  and  $\text{gr}(E)$  is the graded object associated to a semistable bundle  $E$ . The formula  $(E, V) \mapsto [E]$  defines a morphism

$$\psi : G_0(n, d, k) \rightarrow \tilde{B}(n, d, k),$$

whose image contains  $B(n, d, k)$ . We shall use this morphism  $\psi$  in section 9.

We finish this section with a useful definition and some notation.

**Definition 2.8.** A coherent system  $(E, V)$  is

*generated* if the evaluation map  $V \otimes \mathcal{O} \rightarrow E$  is surjective;

*generically generated* if the cokernel of the evaluation map is a torsion sheaf.

**Notation.** We shall write  $\beta$ ,  $G(\alpha)$ ,  $\tilde{G}(\alpha)$ ,  $G_L$  for  $\beta(n, d, n+1)$ ,  $G(\alpha; n, d, n+1)$ ,  $\tilde{G}(\alpha; n, d, n+1)$ ,  $G_L(n, d, n+1)$  respectively. For any coherent system  $(E, V)$ , we shall consistently denote by  $E'$  the subsheaf image of the evaluation map. We shall also denote by  $(n_i, d_i, k_i)$  the type of a coherent system  $(E_i, V_i)$ .

### 3. THE MODULI SPACE FOR LARGE $\alpha$

In this section we assume that  $X$  is a Petri curve and obtain a strengthening of [5, Theorem 5.11]. In particular we obtain a much better lower bound on the parameter  $\alpha$  which ensures that  $G(\alpha) = G_L$ . In later sections we shall prove that this bound is best possible and describe a natural stratification of  $G_L$ . For  $d \leq g+n$ , Theorem 3.1 has been proved in [8, Theorem 2]. We recall that, for any coherent system  $(E, V)$ ,  $E'$  denotes the subsheaf image of  $V \otimes \mathcal{O}$  in  $E$ .

**Theorem 3.1.** *Suppose that  $X$  is a Petri curve and  $\alpha > \max\{0, \alpha_l\}$ , where*

$$(3.1) \quad \alpha_l := d(n-1) - n \left( n-1 + g - \left\lfloor \frac{g}{n} \right\rfloor \right).$$

*Then*

- (1)  $G(\alpha) \neq \emptyset$  if and only if  $\beta \geq 0$ ;
- (2)  $G(\alpha) = G_L$ ;



- (3)  $(E, V) \in G(\alpha)$  if and only if  $(E, V)$  is generically generated and  $H^0(E'^*) = 0$ ;
- (4) if  $\beta > 0$ ,  $G(\alpha)$  is smooth and irreducible of dimension  $\beta$ ; moreover the generic element of  $G(\alpha)$  is generated;
- (5) if  $\beta = 0$ ,  $G(\alpha)$  is a finite set of cardinality

$$g! \prod_{i=0}^n \frac{i!}{(g-d+n+i)!};$$

moreover every element of  $G(\alpha)$  is generated.

We shall prove Theorem 3.1 by means of a sequence of propositions. We begin with two lemmas, the first of which is a variant of [8, Lemma 3.1]. Since the hypotheses are not exactly the same as those of [8, Lemma 3.1], we include a proof.

**Lemma 3.2.** *Let  $X$  be a Petri curve and  $(E, V)$  a coherent system of type  $(n, d, k)$ . If  $(E, V)$  is generically generated and  $H^0(E'^*) = 0$ , then  $k \geq n + 1$  and  $d \geq g + n - \lfloor \frac{g}{n+1} \rfloor$ . Moreover, if  $(E_2, V_2)$  is a quotient coherent system of  $(E, V)$ , then  $(E_2, V_2)$  is generically generated and  $H^0(E_2'^*) = 0$ .*

*Proof.* Certainly  $k \geq n$ . If  $k = n$ , then  $E' \cong \mathcal{O}^n$ , contradicting the hypothesis  $H^0(E'^*) = 0$ . So  $k \geq n + 1$ .

Replacing  $V$ , if necessary, by a subspace of dimension  $n + 1$  which generates  $E'$ , we have an exact sequence

$$(3.2) \quad 0 \rightarrow \mathcal{L}^* \rightarrow V \otimes \mathcal{O} \rightarrow E' \rightarrow 0,$$

where  $\mathcal{L} = \det E'$ . From the dual of (3.2) and the hypothesis  $H^0(E'^*) = 0$ , we see that  $h^0(\mathcal{L}) \geq n + 1$ . By classical Brill-Noether theory, this implies that

$$\deg E' = \deg \mathcal{L} \geq \frac{ng}{n+1} + n = g + n - \frac{g}{n+1}.$$

Hence  $d \geq \deg E' \geq g + n - \lfloor \frac{g}{n+1} \rfloor$  as required.

For the last part, note that the image of  $E'$  in  $E_2$  is precisely  $E_2'$ . Hence  $E_2'$  is a quotient of  $E'$  and the result follows.  $\square$

**Remark 3.3.** Note that

$$(3.3) \quad \alpha_l = (n-1)(d-g-n) - \left( g - n \left\lfloor \frac{g}{n} \right\rfloor \right) = (n-1)(d-n) - n \left( g - \left\lfloor \frac{g}{n} \right\rfloor \right).$$

and that

$$d \geq g + n - \left\lfloor \frac{g}{n+1} \right\rfloor \Leftrightarrow d \geq \frac{ng}{n+1} + n \Leftrightarrow \beta \geq 0.$$

Note in particular that, by (3.3),

$$\alpha_l \geq 0 \Rightarrow d \geq g + n \Rightarrow \beta \geq 0.$$

**Lemma 3.4.** Let  $f : \mathbb{Z}_{>0} \rightarrow \mathbb{Q}$  be defined by

$$f(r) := \frac{1}{r} \left( g - \left[ \frac{g}{r+1} \right] \right).$$

Then  $f$  is a decreasing function of  $r$ .

*Proof.* If  $g \geq r+1$ , we have

$$f(r) \geq \frac{1}{r} \left( g - \frac{g}{r+1} \right) = \frac{g}{r+1}$$

and

$$f(r+1) \leq \frac{1}{r+1} \left( g - \frac{g-r-1}{r+2} \right) = \frac{g+1}{r+2} \leq \frac{g}{r+1}.$$

On the other hand, if  $g < r+1$ , then

$$f(r) = \frac{g}{r} > \frac{g}{r+1} = f(r+1).$$

□

**Proposition 3.5.** Suppose that  $(E, V)$  is a generically generated coherent system of type  $(n, d, n+1)$  and  $H^0(E^*) = 0$ . Then  $(E, V)$  is  $\alpha$ -stable for  $\alpha > \max\{0, \alpha_l\}$ .

*Proof.* Let  $(E_2, V_2)$  be a proper quotient coherent system of  $(E, V)$  of type  $(n_2, d_2, k_2)$ . It follows from Lemma 3.2 that  $k_2 \geq n_2 + 1$  and  $d_2 \geq g + n_2 - \left[ \frac{g}{n_2+1} \right]$ . Hence

$$(3.4) \quad \mu_\alpha(E_2, V_2) \geq 1 + \frac{1}{n_2} \left( g - \left[ \frac{g}{n_2+1} \right] \right) + \alpha \left( \frac{n_2+1}{n_2} \right).$$

If  $\alpha > \max\{0, \alpha_l\}$  then, since  $0 < n_2 < n$ ,

$$(3.5) \quad \alpha \left( \frac{1}{n_2} - \frac{1}{n} \right) = \alpha \left( \frac{n-n_2}{nn_2} \right) \geq \frac{\alpha}{n(n-1)} > \frac{d}{n} - 1 - \frac{1}{n-1} \left( g - \left[ \frac{g}{n} \right] \right).$$

Hence, from (3.4) and Lemma 3.4,

$$\mu_\alpha(E_2, V_2) - \mu_\alpha(E, V) > \frac{1}{n_2} \left( g - \left[ \frac{g}{n_2+1} \right] \right) - \frac{1}{n-1} \left( g - \left[ \frac{g}{n} \right] \right) \geq 0.$$

Since this holds for all  $(E_2, V_2)$ , it follows that  $(E, V)$  is  $\alpha$ -stable. □

**Remark 3.6.** Suppose  $(E_2, V_2)$  is a coherent system of type  $(n_2, d_2, k_2)$  with

$$0 < n_2 < n, \quad k_2 \geq n_2 + 1, \quad d_2 \geq g + n_2 - \left[ \frac{g}{n_2+1} \right].$$

If  $\alpha \geq \alpha_l > 0$ , then (3.4) still holds as does the first inequality in (3.5), while the second inequality in (3.5) becomes  $\geq$ . So

$$\mu_\alpha(E_2, V_2) \geq \mu_\alpha(E, V)$$

with equality if and only if  $\alpha = \alpha_l$  and

$$n_2 = n - 1, \quad k_2 = n, \quad d_2 = g + n - 1 - \left\lfloor \frac{g}{n} \right\rfloor.$$

**Proposition 3.7.** *For given  $n$  and  $d$ , the following three conditions are equivalent:*

- (a) *there exists a generated coherent system  $(E, V)$  of type  $(n, d, n + 1)$  with  $H^0(E^*) = 0$ ;*
- (b) *there exists a generically generated coherent system  $(E, V)$  of type  $(n, d, n + 1)$  with  $H^0(E'^*) = 0$ ;*
- (c)  $\beta \geq 0$ .

*Proof.* Clearly (a) implies (b) and, by Lemma 3.2 and Remark 3.3, (b) implies (c).

Now suppose (c) holds. By classical Brill-Noether theory,  $G(1, d, n + 1) \neq \emptyset$  and its general element  $(\mathcal{L}, W)$  is generated (in the case  $\beta = 0$ ,  $G(1, d, n + 1)$  is finite and all elements are generated). If we define  $E$  by the exact sequence

$$0 \rightarrow E^* \rightarrow W \otimes \mathcal{O} \rightarrow \mathcal{L} \rightarrow 0,$$

then  $(E, W^*)$  satisfies (a). □

**Proposition 3.8.** *Suppose that  $\alpha > \max\{0, \alpha_l\}$  and  $(E, V)$  is an  $\alpha$ -semistable coherent system of type  $(n, d, n + 1)$ . Then  $(E, V)$  is generically generated and  $H^0(E'^*) = 0$ .*

*Proof.* Since  $(E', V)$  is a generated coherent system, we can write  $(E', V) \cong (\mathcal{O}^s, H^0(\mathcal{O}^s)) \oplus (G, W)$  where  $H^0(G^*) = 0$ ,  $W = H^0(G) \cap V$  and  $(G, W)$  is generated. Let  $r$  denote the rank of  $G$ . Note that, since  $h^0(E') \geq n + 1$ , we must have  $r \geq 1$ . We require to show that  $r = n$ .

Suppose to the contrary that  $r \leq n - 1$ . Since the coherent system  $(G, W)$  is generated, we have, by Lemma 3.2,  $\deg G \geq g + r - \left\lfloor \frac{g}{r+1} \right\rfloor$ . Hence

$$\frac{1}{r} \left( g - \left\lfloor \frac{g}{r+1} \right\rfloor \right) + 1 + \alpha \frac{n+1-s}{r} \leq \mu_\alpha(G, W).$$

Since  $(E, V)$  is  $\alpha$ -semistable, it follows that

$$\frac{1}{r} \left( g - \left\lfloor \frac{g}{r+1} \right\rfloor \right) + 1 + \alpha \frac{n+1-s}{r} \leq \frac{d}{n} + \alpha \frac{n+1}{n}.$$

Now  $s \leq n - r$ ; so, for any fixed  $r$ , the minimum value for the left-hand side of this inequality is given by  $s = n - r$ . By Lemma 3.4, this minimum value is then a decreasing function of  $r$ . Hence

$$\frac{1}{n-1} \left( g - \left\lfloor \frac{g}{n} \right\rfloor \right) + 1 + \alpha \frac{n}{n-1} \leq \frac{d}{n} + \alpha \frac{n+1}{n},$$

i. e.

$$\frac{\alpha}{n(n-1)} \leq \frac{d-n}{n} - \frac{1}{n-1} \left( g - \left\lfloor \frac{g}{n} \right\rfloor \right),$$

contradicting the hypothesis that  $\alpha > \alpha_l$ . □

**Remark 3.9.** Under the hypotheses of Proposition 3.8, we have an exact sequence

$$(3.6) \quad 0 \rightarrow E' \rightarrow E \rightarrow \tau \rightarrow 0,$$

where  $\tau$  is a torsion sheaf. If  $t$  is the length of  $\tau$ , then  $\deg E' = d - t$ . Since  $(E', V)$  is generated and  $H^0(E'^*) = 0$ , Lemma 3.2 gives  $d - t \geq g + n - \left\lfloor \frac{g}{n+1} \right\rfloor$ , or equivalently

$$(3.7) \quad t \leq t_1 := d - g - n + \left\lfloor \frac{g}{n+1} \right\rfloor = \left\lfloor \frac{\beta}{n+1} \right\rfloor.$$

We shall see later (Theorem 4.2) that this bound is best possible. In particular, if we write

$$d_0 = g + n - \left\lfloor \frac{g}{n+1} \right\rfloor,$$

then, for  $d > d_0$ , we have  $t_1 \geq 1$ , so there exists a non-generated coherent system  $(E, V)$  in  $G_L$ .

*Proof of Theorem 3.1.* Parts (2) and (3) follow from Propositions 3.5 and 3.8, and (1) then follows from Proposition 3.7.

(4) If  $\beta > 0$ , it follows from [8, Lemma 4.2] and [5, Theorem 5.11] that  $G(\alpha)$  is smooth and irreducible of dimension  $\beta$ . The fact that the generic element is generated then follows from Proposition 3.7.

(5) If  $\beta = 0$ , it follows from [8, Lemma 4.2] that  $G(\alpha)$  is finite and that, as a scheme, it is reduced. By (3.6) and (3.7), every element is generated. The formula for the cardinality of  $G(\alpha)$  now follows from [1, Chapter V, formula (1.2)].  $\square$

#### 4. A STRATIFICATION OF $G_L$

Let

$$(4.1) \quad \Sigma_0 = \{(E, V) \in G_L \mid (E, V) \text{ is generated}\}.$$

Clearly  $\Sigma_0$  is open in  $G_L$ . If  $\beta \geq 0$ , we know from Theorem 3.1 that  $\Sigma_0 \neq \emptyset$ . Moreover, by Remark 3.9, the complement of  $\Sigma_0$  in  $G_L$  is a disjoint union of locally closed subsets  $\Sigma_t$ , defined for  $1 \leq t \leq t_1$  by

$$(4.2) \quad \Sigma_t = \{(E, V) \in G_L \mid \exists \text{ an exact sequence (3.6) with } \tau \text{ of length } t\}.$$

We now define

$$S_t = \bigcup_{i \geq t} \Sigma_i,$$

where the  $\Sigma_i$  are the locally closed subsets of  $G_L$  defined in (4.1) and (4.2). Clearly  $G_L = S_0 \supset S_1 \supset \cdots \supset S_t \supset \cdots$ . We would like to show that the subsets  $S_t$  define a well-behaved stratification of  $G_L$ .

We begin with a lemma, which will be needed again later

**Lemma 4.1.** *Suppose that we have an exact sequence*

$$0 \longrightarrow F \longrightarrow E \longrightarrow \tau \longrightarrow 0,$$

where  $\tau$  is a torsion sheaf of length  $t$ , and that  $V$  is a subspace of  $H^0(F)$  of dimension  $n + 1$ . Then

$$(E, V) \in G_L(n, d, n + 1) \Leftrightarrow (F, V) \in G_L(n, d - t, n + 1).$$

*Proof.* It is clear that  $(E, V)$  is generically generated if and only if  $(F, V)$  is generically generated and that  $E' = F'$ . The result follows at once from Theorem 3.1(3).  $\square$

**Theorem 4.2.** *Suppose  $\beta \geq 0$  and that the subsets  $S_t$  of  $G_L$  are defined as above. Then*

- (1)  $S_t$  is closed in  $G_L$  and is non-empty if and only if  $0 \leq t \leq t_1 := \lfloor \frac{\beta}{n+1} \rfloor$ ;
- (2) for  $1 \leq t \leq t_1$ ,  $S_t \subset \overline{S_{t-1}} \setminus S_t$ ;
- (3) for  $1 \leq t \leq t_1$ ,  $\dim S_t = \beta - t$ ;
- (4)  $S_t$  is irreducible for  $t < \frac{\beta}{n+1}$ ;
- (5) if  $\frac{\beta}{n+1}$  is an integer, then all irreducible components of  $S_{t_1}$  have the same dimension.

*Proof.* The fact that  $S_t$  is empty if  $t > t_1 = \lfloor \frac{\beta}{n+1} \rfloor$  has already been proved in Remark 3.9. We prove the rest of the theorem by induction on  $t_1$ , the result being an immediate consequence of Theorem 3.1 if  $t_1 = 0$ .

Suppose therefore that  $t_1 \geq 1$ . We consider the moduli space

$$G_{L,d-1} := G_L(n, d - 1, n + 1)$$

and denote by  $S_{t,d-1}$  the subset of  $G_{L,d-1}$  given by

$$S_{t,d-1} := \{(F, V) \in G_{L,d-1} \mid \exists \text{ an exact sequence (3.6) with } \tau \text{ of length } \geq t\}.$$

The maximum value of  $t$  on  $G_{L,d-1}$  is

$$\left\lfloor \frac{\beta(1, d - 1, n + 1)}{n + 1} \right\rfloor = t_1 - 1,$$

so we can assume inductively that the theorem holds for  $G_{L,d-1}$ .

Note next that, if  $(F, V) \in G_{L,d-1}$  and  $E$  is defined by an elementary transformation

$$(4.3) \quad 0 \rightarrow F \rightarrow E \rightarrow \tau \rightarrow 0,$$

with  $\tau$  a torsion sheaf of length 1, then  $(E, V) \in G_L$  by Lemma 4.1. In fact it is easy to see that the  $(E, V)$  obtained in this way are precisely the elements of  $S_1$  and, more generally, for  $1 \leq t \leq t_1$ ,

$$(4.4) \quad (E, V) \in S_t \Leftrightarrow (F, V) \in S_{t-1,d-1}.$$

The next step is to carry out this construction for families of coherent systems. Since  $(n, d - 1, n + 1)$  are coprime there is a universal family  $(\mathcal{U}, \mathcal{V})$  parametrised by  $G_{L,d-1}$

[6, Proposition A.8]. Denote by  $p : \mathbb{P}\mathcal{U} \rightarrow X \times G_{L,d-1}$  the natural projection. As in the Hecke correspondence of [24],  $\mathbb{P}\mathcal{U}$  parametrises the triples

$$(F, V, 0 \rightarrow F \rightarrow E \rightarrow \tau \rightarrow 0)$$

for which  $(F, V) \in G_{L,d-1}$  and  $\tau$  has length 1. The universal property of  $G_L$  now gives us a diagram

$$\begin{array}{ccc} \mathbb{P}\mathcal{U} & \xrightarrow{\Psi} & G_L \\ p \downarrow & & \\ X \times G_{L,d-1} & & \end{array}$$

By (4.4), we have

$$(4.5) \quad S_t = \Psi(p^{-1}(X \times S_{t-1,d-1})), \quad \Psi^{-1}(S_{t-1} \setminus S_t) = p^{-1}(X \times (S_{t-2,d-1} \setminus S_{t-1,d-1})).$$

The fact that  $S_t \neq \emptyset$  for  $t \leq t_1$  follows at once. Moreover  $G_{L,d-1}$  is a projective variety and, by inductive hypothesis,  $S_{t-1,d-1}$  is closed and, provided  $t-1 < \frac{\beta}{n+1} - 1$ , also irreducible; hence  $S_t$  is closed in  $G_L$ , completing the proof of (1). Properties (2) and (4) follow immediately from (4.5).

For (3), note that, by the inductive hypothesis,

$$(4.6) \quad \dim(p^{-1}(X \times S_{t-1,d-1})) = \beta(n, d-1, n+1) - (t-1) + 1 + (n-1) = \beta - t.$$

Moreover, if  $(E, V) \in \Sigma_t$  and the torsion sheaf  $\tau$  of (4.2) has support consisting of  $t$  distinct points, then  $\Psi^{-1}(E, V)$  consists of precisely  $t$  points. Hence  $\Psi$  is generically finite on  $(p^{-1}(X \times S_{t-1,d-1}))$ , so (3) follows from (4.6).

Finally, for (5), suppose  $\frac{\beta}{n+1}$  is an integer and let  $S'$  be any irreducible component of  $S_{t_1-1,d-1}$ ; by inductive hypothesis,  $\dim S' = \beta(n, d-1, n+1) - (t_1-1)$ . As in (4.6), we have

$$\dim(\Psi(p^{-1}(X \times S'))) = \beta - t_1.$$

The result follows. □

## 5. THE TOP CRITICAL VALUE

In the previous sections we gave a description of  $G_L(n, d, n+1)$ . We shall show now that the bound of Theorem 3.1 is best possible if  $\alpha_l > 0$  and analyse what happens at this value of the parameter. Note that the condition  $\alpha_l > 0$  implies that  $n \geq 2$ .

**Theorem 5.1.** *Suppose  $\alpha_l > 0$ . Then there exists a coherent system  $(E, V)$  which is  $\alpha_l^+$ -stable and  $\alpha_l$ -semistable, but not  $\alpha_l$ -stable.*

*Proof.* We shall construct  $(E, V)$  as an extension

$$(5.1) \quad 0 \rightarrow (E_1, V_1) \rightarrow (E, V) \rightarrow (E_2, V_2) \rightarrow 0,$$

where

$$(5.1a) \quad (E_2, V_2) \in G_L(n-1, d_2, n) \text{ with } d_2 = g + n - 1 - \left\lfloor \frac{g}{n} \right\rfloor;$$

$$(5.1b) \quad (E_1, V_1) \text{ is of type } (1, d - d_2, 1).$$

Note that  $d > d_2$  by (3.3), so  $(E_1, V_1)$  exists. Moreover  $\beta(n-1, d_2, n) \geq 0$ ; so, by Theorem 3.1,  $(E_2, V_2)$  also exists and indeed is  $\alpha$ -stable for all  $\alpha > 0$  and in particular for  $\alpha = \alpha_l$ . It is easy to check from the definition (3.1) that

$$(5.2) \quad \mu_{\alpha_l}(E_1, V_1) = \mu_{\alpha_l}(E_2, V_2),$$

so  $(E, V)$  is  $\alpha_l$ -semistable but not  $\alpha_l$ -stable. Moreover, since  $(E_1, V_1)$  and  $(E_2, V_2)$  are both  $\alpha_l$ -stable but not isomorphic, it follows from (5.2) that

$$(5.3) \quad \text{Hom}((E_1, V_1), (E_2, V_2)) = 0 = \text{Hom}((E_2, V_2), (E_1, V_1)).$$

Now any subsystem of  $(E, V)$  which contradicts  $\alpha_l^+$ -stability must also contradict  $\alpha_l$ -stability. If the extension (5.1) is non-trivial, the only subsystem which contradicts  $\alpha_l$ -stability is  $(E_1, V_1)$  and clearly this does not contradict  $\alpha_l^+$ -stability. It remains only to prove that there exists a non-trivial extension (5.1), or equivalently to prove that

$$\text{Ext}^1((E_2, V_2), (E_1, V_1)) \neq 0.$$

Now, by (2.8) and (2.6),

$$\dim \text{Ext}^1((E_2, V_2), (E_1, V_1)) \geq C_{21} = (k_2 - n_2)(d_1 - n_1(g - 1)) + n_1 d_2 - k_1 k_2.$$

Here we have  $(n_1, d_1, k_1) = (1, d - d_2, 1)$ ,  $(n_2, d_2, k_2) = (n - 1, d_2, n)$ , so

$$C_{21} = (d - d_2 - g + 1) + d_2 - n = d - g - n + 1.$$

Since  $\alpha_l > 0$ , it follows from (3.3) that  $d - g - n > 0$  and so  $C_{21} > 0$  as required.  $\square$

**Corollary 5.2.** *If  $\alpha_l > 0$ , then it is equal to the top critical value  $\alpha_L$ . Moreover the flip locus  $G_L^+$  is given precisely by the non-trivial extensions (5.1) which satisfy (5.1a) and (5.1b) and has dimension  $\leq \beta - 1$ .*

*Proof.* The fact that  $\alpha_L = \alpha_l$  follows at once from Theorems 3.1 and 5.1. If  $(E, V) \in G_L^+$ , we have a sequence (2.4) for which  $(E_2, V_2)$  is  $\alpha_l^+$ -stable and (2.5) holds with  $\alpha_i = \alpha_l$ . By Lemma 3.2, we must have  $k_2 \geq n_2 + 1$  and  $d_2 \geq g + n_2 - \left\lfloor \frac{g}{n_2 + 1} \right\rfloor$ . By Remark 3.6, it follows that

$$n_2 = n - 1, \quad k_2 = n, \quad d_2 = g + n - 1 - \left\lfloor \frac{g}{n} \right\rfloor.$$

Hence all the conditions of (5.1) hold.

According to Lemma 2.6 and Remark 2.7, it remains to prove that

$$C_{12} - h^0(E_1^* \otimes N_2 \otimes K) > 0.$$

Putting in values from (5.1), we have, since  $\alpha_l > 0$ ,

$$C_{12} = (n - 1) \left( d - g - n + 1 + \left\lfloor \frac{g}{n} \right\rfloor \right) - n > g - \left\lfloor \frac{g}{n} \right\rfloor - 1 \geq 0.$$

On the other hand,  $E_1^* \otimes N_2 \otimes K$  is a line bundle of degree  $2g - 2 - d$ . If  $d > 2g - 2$ , we are finished. If  $d \leq 2g - 2$ , then, by Clifford's Theorem,

$$h^0(E_1^* \otimes N_2 \otimes K) \leq g - \frac{d}{2} < g - \frac{g + n}{2}.$$

It is therefore sufficient to prove that

$$\frac{g+n}{2} \geq \left\lfloor \frac{g}{n} \right\rfloor + 1.$$

Since  $n \geq 2$ , this is obvious.  $\square$

**Remark 5.3.** The estimate for the dimension of  $G_L^+$  in the proof of Corollary 5.2 is sufficient for our purposes, but is quite crude and can certainly be improved.

We now turn to the determination of the flip locus  $G_L^-$ .

**Proposition 5.4.** *If  $\alpha_l > 0$ , then the flip locus  $G_L^-$  consists of the non-trivial extensions*

$$(5.4) \quad 0 \rightarrow (E_2, V_2) \rightarrow (E, V) \rightarrow (E_1, V_1) \rightarrow 0,$$

where  $(E_1, V_1)$  and  $(E_2, V_2)$  satisfy the same properties as in (5.1), and has dimension  $\leq \beta - 1$ .

*Proof.* If  $(E, V) \in G_L^-$ , then there certainly exists a non-trivial extension (5.4) with  $(E_2, V_2)$   $\alpha_l^-$ -stable and

$$\mu_{\alpha_l}(E_2, V_2) = \mu_{\alpha_l}(E, V), \quad k_2 \geq n_2 + 1$$

(see (2.5)). By [8, Theorem 1(1)], we must have  $\beta(n_2, d_2, n_2 + 1) \geq 0$  and so, by Remark 3.3,  $d_2 \geq g + n_2 - \left\lfloor \frac{g}{n_2+1} \right\rfloor$ . By Remark 3.6, it follows that

$$n_2 = n - 1, \quad k_2 = n, \quad d_2 = g + n - 1 - \left\lfloor \frac{g}{n} \right\rfloor.$$

Hence all the conditions of (5.1) hold. Now note that  $N_1 = 0$  and  $C_{21} > 0$  as shown in the proof of Theorem 5.1. The proposition follows from Remark 2.7.  $\square$

**Remark 5.5.** Taking  $\alpha = \alpha_l$  in the proof of Proposition 3.8 gives a slightly different description of  $G_L^-$ , namely

$$G_L^- = \{(E, V) \mid (E, V) \text{ generically generated, } E' \cong \mathcal{O} \oplus G, H^0(G^*) = 0, G \text{ saturated in } E\}.$$

It is easy to see that these two descriptions are equivalent.

**Theorem 5.6.** *Suppose  $\alpha_l > 0$ . Then  $G_{L-1}$  is non-empty and irreducible, and is birational to  $G_L$ .*

*Proof.* This follows from Corollary 5.2, Proposition 5.4 and (2.3).  $\square$

## 6. MODULI SPACES FOR ANY $\alpha$

As we have seen (see Theorems 3.1 and 5.6), for  $\beta(n, d, n + 1) \geq 0$  and  $\alpha > \alpha_{L-1}$ , the moduli space  $G(\alpha; n, d, n + 1)$  is non-empty and the non-emptiness is related with the existence of coherent systems  $(E, V)$  such that  $E$  is generically generated and  $H^0(E^*) = 0$ . Our object in this section is to try to generalise these results to arbitrary  $\alpha > 0$ . For  $d \leq g + n$ , these results are largely contained in the unpublished [12] (see also [11]) and in [8].



We begin by recalling the results of [8] which we require.

**Proposition 6.1.** [8, Theorem 1(1)] *Let  $X$  be a Petri curve and  $\beta < 0$ . Then  $G(\alpha) = \emptyset$  for all  $\alpha > 0$ .*

Before proceeding further, we define

$$U(n, d, n+1) := \{(E, V) \in G_L : E \text{ is stable}\}$$

and

$$U^s(n, d, n+1) := \{(E, V) : (E, V) \text{ is } \alpha\text{-stable for all } \alpha > 0\}.$$

Note that  $U(n, d, n+1)$  can be defined alternatively as

$$U(n, d, n+1) := \{(E, V) : E \text{ is stable and } (E, V) \text{ is } \alpha\text{-stable for all } \alpha > 0\}$$

and in particular  $U(n, d, n+1) \subset U^s(n, d, n+1)$ . In the converse direction, note that, if  $(E, V) \in U^s(n, d, n+1)$ , then  $E$  is semistable. However it is not generally true that  $U(n, d, n+1) = U^s(n, d, n+1)$  and we can have  $U^s(n, d, n+1) \neq \emptyset$ ,  $U(n, d, n+1) = \emptyset$ . Our main object in the remainder of the paper is to determine when these sets are non-empty.

**Remark 6.2.** By openness of  $\alpha$ -stability,  $U(n, d, n+1)$  and  $U^s(n, d, n+1)$  are open subsets of  $G_L$ , thus inheriting natural structures of smooth variety, and with these same structures they are also embedded as open subsets of every  $G(\alpha)$ . If either  $U(n, d, n+1)$  or  $U^s(n, d, n+1)$  is non-empty, then, by Theorem 3.1, it is irreducible of dimension  $\beta$  (or finite when  $\beta = 0$ ) and its generic element  $(E, V)$  is generated with  $H^0(E^*) = 0$ .

**Proposition 6.3.** [8, Proposition 2.5(4)] *Let  $(E, V)$  be a generated coherent system of type  $(n, d, n+1)$  such that  $E$  is semistable. Then  $(E, V) \in U^s(n, d, n+1)$ .*

**Proposition 6.4.** [8, Proposition 4.1(2)] *Let  $X$  be a Petri curve and suppose that  $g + n - \lfloor \frac{g}{n+1} \rfloor \leq d \leq g + n$  and that  $g$  and  $n$  are not both equal to 2. Then  $U(n, d, n+1)$  is non-empty.*

**Proposition 6.5.** [8, Proposition 4.6] *Let  $X$  be a Petri curve and  $\beta \geq 0$ . If  $g \geq n^2 - 1$ , then  $U(n, d, n+1) \neq \emptyset$ .*

In the remainder of this section, we shall introduce two further techniques for constructing coherent systems. The first is that of elementary transformations, which we shall use in two distinct ways.

Since any stable bundle of degree  $\geq n(2g - 1)$  is generated by its sections, Proposition 6.3 implies that  $U(n, d, n+1) \neq \emptyset$  for  $d \geq n(2g - 1)$  (see also [8, Proposition 2.6]). The next proposition provides a significant improvement on this.

**Proposition 6.6.** *Let  $X$  be a Petri curve. If*

$$d_0 = \begin{cases} \frac{n(g+3)}{2} & \text{if } g \text{ is odd} \\ \frac{n(g+2)}{2} & \text{if } g \text{ is even,} \end{cases}$$

*then  $U^s(n, d_0, n+1) \neq \emptyset$ .*

If  $d \geq d_1$ , where

$$d_1 = \begin{cases} \frac{n(g+3)}{2} + 1 & \text{if } g \text{ is odd} \\ \frac{n(g+2)}{2} + 1 & \text{if } g \text{ is even and } n \leq \frac{g!}{(\frac{g}{2})!(\frac{g}{2}+1)!} \\ \frac{n(g+4)}{2} + 1 & \text{if } g \text{ is even and } n > \frac{g!}{(\frac{g}{2})!(\frac{g}{2}+1)!}, \end{cases}$$

then  $U(n, d, n+1) \neq \emptyset$ .

*Proof.* It is easy to check that, with the above definition of  $d_0$ ,  $\beta(1, \frac{d_0}{n}, 2) \geq 0$  (in fact,  $\frac{d_0}{n}$  is the smallest integer for which this is true). Hence, by classical Brill-Noether theory, there exists a line bundle  $\mathcal{L}$  of degree  $\frac{d_0}{n}$  such that  $h^0(\mathcal{L}) \geq 2$  and  $\mathcal{L}$  is generated by its sections. Now let  $\mathcal{L}_1, \dots, \mathcal{L}_n$  be any such line bundles and let  $V$  be a subspace of  $H^0(\mathcal{L}_1 \oplus \dots \oplus \mathcal{L}_n)$  of dimension  $n+1$  such that  $(\mathcal{L}_1 \oplus \dots \oplus \mathcal{L}_n, V)$  is generated. Hence  $(\mathcal{L}_1 \oplus \dots \oplus \mathcal{L}_n, V) \in U^s(n, d_0, n+1)$  by Proposition 6.3.

Again by classical Brill-Noether theory, one can find pairwise non-isomorphic line bundles  $\mathcal{L}_1, \dots, \mathcal{L}_n$  of degree  $\frac{d_1-1}{n}$  such that, for all  $i$ ,  $h^0(\mathcal{L}_i) \geq 2$  and  $\mathcal{L}_i$  is generated by its sections (in the case  $g$  even and  $d_1 = \frac{n(g+2)}{2} + 1$ , the number of distinct line bundles of degree  $\frac{d_1-1}{n}$  with  $h^0 \geq 2$  is  $\frac{g!}{(\frac{g}{2})!(\frac{g}{2}+1)!}$  [1, Chapter V, formula (1.2)]). Now consider extensions

$$0 \rightarrow \mathcal{L}_1 \oplus \dots \oplus \mathcal{L}_n \rightarrow E \rightarrow \tau \rightarrow 0,$$

where  $\tau$  is a torsion sheaf of length  $t \geq 1$ . These extensions are classified by  $n$ -tuples  $(e_1, \dots, e_n)$  with  $e_i \in \text{Ext}^1(\tau, \mathcal{L}_i)$ . It can be shown (see [21, Théorème A.5]) that, for any  $t$ , there exists an extension of this type for which  $E$  is stable. Moreover  $V$  can be regarded as a subspace of  $H^0(E)$ , making  $(E, V)$  a coherent system. If  $(E_1, V_1)$  is a proper subsystem of  $(E, V)$  with  $E_1 \neq E$ , then  $V_1 \subset V \cap H^0(E_1 \cap \mathcal{L}_1 \oplus \dots \oplus \mathcal{L}_n)$ . It follows from the  $\alpha$ -stability of  $(\mathcal{L}_1 \oplus \dots \oplus \mathcal{L}_n, V)$  for large  $\alpha$  that  $\frac{k_1}{n_1} \leq \frac{k}{n}$ . Since  $E$  is stable, we have also  $\frac{d_1}{n_1} < \frac{d}{n}$ . It follows that  $(E, V) \in U(n, d, n+1)$ .  $\square$

**Remark 6.7.** For a general curve  $X$ , the second part of Proposition 6.6 is valid with

$$d_1 = \begin{cases} \frac{n(g+1)}{2} + 1 & \text{if } g \text{ is odd} \\ \frac{n(g+2)}{2} + 1 & \text{if } g \text{ is even} \end{cases}$$

by [30]. However, this does not imply the result for an arbitrary Petri curve.

Our second use of elementary transformations is to prove

**Proposition 6.8.** *Suppose that  $U(n, na, n+1) \neq \emptyset$  for some integer  $a$ . Then  $U(n, d, n+1) \neq \emptyset$  for all  $d$  with  $d > na$  and  $d \equiv \pm 1 \pmod{n}$ .*

*Proof.* In view of Remark 2.2, it is sufficient to prove this for  $d = na + 1$  and for  $d = na + n - 1$ .

Suppose first that  $d = na + 1$ . Let  $(F, V) \in U(n, na, n+1)$  and define  $E$  as an elementary transformation (4.3). Then  $(E, V) \in G_L(n, na + 1, n+1)$  by Lemma 4.1. The stability of  $E$  follows easily from the stability of  $F$ , so  $(E, V) \in U(n, d, n+1)$ .

Now suppose  $d = na + n - 1$ . Again let  $(F, V) \in G_L(n, na, n + 1)$  and let  $x \in X$ . Let  $\tau$  be the torsion sheaf of length 1 supported at  $x$  and define  $E$  as an elementary transformation

$$0 \rightarrow E \rightarrow F(x) \rightarrow \tau \rightarrow 0.$$

Then  $F$  can be regarded as a subsheaf of  $E$  and  $V$  as a subspace of  $H^0(E)$ . By Lemma 4.1, the coherent system  $(E, V) \in G_L(n, na + n - 1, n + 1)$ . The stability of  $E$  follows from the stability of  $F(x)$ .  $\square$

The second technique is the use of extensions of coherent systems. The idea is to take a generic element  $(E, V)$  of  $G_L$  and try to prove that  $E$  is stable. If this is not the case, there exists a quotient  $E_2$  of  $E$  with  $\mu(E_2) \leq \mu(E)$  and we can choose  $E_2$  to be stable. We have therefore an extension

$$0 \rightarrow E_1 \rightarrow E \rightarrow E_2 \rightarrow 0,$$

and, taking  $V_1 = V \cap H^0(E_1)$  and  $V_2 = V/V_1$ , we obtain an extension of coherent systems

$$(6.1) \quad 0 \rightarrow (E_1, V_1) \rightarrow (E, V) \rightarrow (E_2, V_2) \rightarrow 0.$$

We are assuming that  $(E, V)$  is a generic element of  $G_L$ , so  $(E, V)$  is generated and  $H^0(E^*) = 0$ . Using Lemma 3.2, we see that (6.1) is subject to the following conditions:

- $\mu(E_2) \leq \mu(E)$ ;
- $E_2$  is stable,  $(E_2, V_2)$  is generated and  $k_2 \geq n_2 + 1$ ;
- $\mu(E_2) \geq 1 + \frac{1}{n_2} \left( g - \left[ \frac{g}{n_2 + 1} \right] \right)$ .

**Proposition 6.9.** *Suppose that  $X$  is a Petri curve,  $n \geq 3$ ,  $d < g + n + \frac{g}{n-1}$  and  $n_2 \leq n - 2$ . Then no extension (6.1) exists satisfying the stated conditions.*

*Proof.* Suppose we have such an extension. Then

$$1 + \frac{1}{n_2} \left( g - \left[ \frac{g}{n_2 + 1} \right] \right) \leq \mu(E_2) \leq \frac{d}{n}.$$

By Lemma 3.4, the left hand side of this inequality is a decreasing function of  $n_2$ ; so we have

$$1 + \frac{1}{n-2} \left( g - \left[ \frac{g}{n-1} \right] \right) \leq \frac{d}{n},$$

i.e.

$$\begin{aligned} d &\geq g + n + \frac{2g}{n-2} - \frac{n}{n-2} \left[ \frac{g}{n-1} \right] \\ &\geq g + n + \frac{2g}{n-2} - \frac{ng}{(n-2)(n-1)} \\ &= g + n + \frac{g}{n-1}. \end{aligned}$$

This gives the required contradiction.  $\square$

It remains to consider the extensions (6.1) for which  $n_2 = n - 1$ . We have two cases:

$$(6.2) \quad 0 \rightarrow (E_1, V_1) \rightarrow (E, V) \rightarrow (E_2, V_2) \rightarrow 0, \quad n_1 = k_1 = 1$$

and

$$(6.3) \quad 0 \rightarrow (E_1, 0) \rightarrow (E, V) \rightarrow (E_2, V_2) \rightarrow 0, \quad n_1 = 1.$$

**Proposition 6.10.** *Suppose that  $X$  is a Petri curve,  $n \geq 2$  and  $d > g + n$ . Then the extensions (6.2) which satisfy the conditions stated above depend on at most  $\beta - 1$  parameters.*

*Proof.* Since  $E_2$  is stable and  $(E_2, V_2)$  is generated,  $(E_2, V_2) \in G_L(n_2, d_2, n_2 + 1)$  by Proposition 6.3. Hence  $(E_2, V_2)$  depends on  $\beta(n_2, d_2, n_2 + 1)$  parameters, while  $(E_1, V_1)$  depends on  $d_1 = \beta(1, d_1, 1)$  parameters. By Remark 2.7,

$$\mathbb{H}_{21}^0 = \text{Hom}((E_2, V_2), (E_1, V_1)) = 0.$$

By Lemma 2.6, it remains to prove that

$$(6.4) \quad C_{12} > \dim \mathbb{H}_{21}^2.$$

Now, by (2.6),

$$C_{12} = (n - 1)d_1 - n,$$

while

$$\dim \mathbb{H}_{21}^2 = h^0(E_1^* \otimes N_2 \otimes K),$$

where  $N_2$  is the kernel of the evaluation map  $V_2 \otimes \mathcal{O} \rightarrow E_2$ . Now  $E_1^* \otimes N_2 \otimes K$  is a line bundle of degree  $2g - 2 - d$ . If  $d \leq 2g - 2$ , then, by Clifford's Theorem,

$$h^0(E_1^* \otimes N_2 \otimes K) \leq g - 1 - \frac{d}{2} + 1 = g - \frac{d}{2}.$$

So (6.4) holds if

$$(n - 1)d_1 - n > g - \frac{d}{2}.$$

Since  $d_1 \geq \frac{d}{n}$ , this will be true if

$$\frac{(n - 1)d}{n} - n > g - \frac{d}{2},$$

i.e. if

$$\frac{3n - 2}{2n}d > g + n.$$

This is certainly true since  $d > g + n$ .

If  $d > 2g - 2$ , then  $h^0(E_1^* \otimes N_2 \otimes K) = 0$  and we require to prove only that  $C_{12} > 0$ . In fact

$$C_{12} = (n - 1)d_1 - n \geq \frac{n - 1}{n}d - n > \frac{n - 1}{n}(g + n) - n = \frac{n - 1}{n}g - 1 \geq 0.$$

□

**Remark 6.11.** Propositions 6.9 and 6.10 are directed towards proving that  $U(n, d, n + 1) \neq \emptyset$ . If we wish only to prove that  $U^s(n, d, n + 1) \neq \emptyset$ , we are not concerned with the stability of  $E$  and we need to consider extensions (6.2) under the usual conditions of [5, section 6.2] for the flip loci  $G_i^+$ . We can still assume that  $(E, V)$  is generated with  $H^0(E^*) = 0$ , so  $(E_2, V_2)$  is also generated with  $H^0(E_2^*) = 0$ , hence  $d_2 \geq g + n_2 - \left\lfloor \frac{d}{n_2+1} \right\rfloor$ , and now  $\mu(E_2) < \mu(E)$ . So the result of Proposition 6.9 holds under the assumption  $d \leq g + n + \frac{g}{n-1}$ . In Proposition 6.10, note that  $(E_2, V_2) \in G_L(n_2, d_2, n_2 + 1)$  by Theorem 3.1(3); so  $(E_2, V_2)$  depends on precisely  $\beta(n_2, d_2, n_2 + 1)$  parameters and the rest of the proof goes through.

We turn now to the consideration of the extensions (6.3).

**Proposition 6.12.** *Let  $X$  be a Petri curve and  $n \geq 3$ . Suppose that  $d < g + n + \frac{g}{n-1}$ . Then there exist no extensions (6.3) satisfying the conditions of (6.1) with*

$$(6.5) \quad \frac{d}{n} < \frac{2g}{2n-1} + 2.$$

*Proof.* Since  $(E_2, V_2)$  is generated, we can write as usual

$$0 \rightarrow N_2 \rightarrow V_2 \otimes \mathcal{O} \rightarrow E_2 \rightarrow 0.$$

Note that  $H^0(N_2) = 0$  and that  $(N_2^*, V_2^*)$  is generated. Moreover  $N_2^*$  has rank 2 and, since  $h^0(E_2^*) = 0$ ,  $h^0(N_2^*) \geq n + 1$ . Suppose we prove that, for any line subbundle  $\mathcal{L}_1$  of  $N_2^*$ ,

$$(6.6) \quad h^0(\mathcal{L}_1) \leq 1.$$

Then, by [25, Lemma 3.9],

$$h^0(\det N_2^*) \geq 2n - 1.$$

Hence, by classical Brill-Noether theory and the assumption  $\mu(E_2) \leq \mu(E)$ ,

$$\frac{(n-1)d}{n} \geq d_2 = \deg N_2^* \geq \frac{(2n-2)g}{2n-1} + 2n - 2,$$

which contradicts (6.5).

It remains to prove (6.6). Consider an exact sequence

$$0 \rightarrow \mathcal{L}_1 \rightarrow N_2^* \rightarrow \mathcal{L}_2 \rightarrow 0.$$

Since  $N_2^*$  is generated, so is  $\mathcal{L}_2$ . But  $\mathcal{L}_2$  is certainly not trivial since  $h^0(N_2) = 0$ , so  $h^0(\mathcal{L}_2) = s \geq 2$  and

$$\deg \mathcal{L}_2 \geq \frac{(s-1)g}{s} + s - 1.$$

If  $s < n$ , then  $h^0(\mathcal{L}_1) \geq n + 1 - s \geq 2$  and

$$\deg \mathcal{L}_1 \geq \frac{(n-s)g}{n-s+1} + n - s.$$

So

$$\begin{aligned} d_2 = \deg N_2^* &\geq \frac{(s-1)g}{s} + s - 1 + \frac{(n-s)g}{n-s+1} + n - s \\ &= 2g - \frac{(n+1)g}{s(n-s+1)} + n - 1. \end{aligned}$$

Since  $2 \leq s \leq n-1$ , this gives

$$(6.7) \quad d_2 \geq 2g - \frac{(n+1)g}{2(n-1)} + n - 1 \geq g + n - 1;$$

since  $\frac{(n-1)d}{n} \geq d_2$ , this contradicts the assumption that  $d < g + n + \frac{g}{n-1}$ . It follows that  $s \geq n$ , so

$$\deg \mathcal{L}_2 \geq \frac{(n-1)g}{n} + n - 1$$

and

$$(6.8) \quad \deg \mathcal{L}_1 = d_2 - \deg \mathcal{L}_2 < g + n - 1 - \frac{(n-1)g}{n} - n + 1 = \frac{g}{n}.$$

The inequality (6.6) now follows from classical Brill-Noether theory. This completes the proof.  $\square$

**Remark 6.13.** The non-strict inequality

$$(6.9) \quad d \leq g + n + \frac{g}{n-1}$$

is sufficient except when  $n = 3$ , when (6.7) fails to give a contradiction. The other place where the inequality  $d < g + n + \frac{g}{n-1}$  is used is (6.8). In this case (6.9) gives  $\deg \mathcal{L}_1 \leq \frac{g}{n}$ , which is sufficient for (6.6). In particular, if  $n \geq 4$ , (6.9) and (6.5) are sufficient for the validity of Proposition 6.12.

## 7. THE CASES $n = 2$ , $n = 3$ AND $n = 4$

In this section we shall assume that  $g \geq 3$ .

**Theorem 7.1.** *Let  $X$  be a Petri curve of genus  $g \geq 3$ . Then  $U(2, d, 3) \neq \emptyset$  if and only if  $\beta(2, d, 3) \geq 0$ .*

*Proof.* This follows at once from Propositions 6.1 and 6.5.  $\square$

**Theorem 7.2.** *Let  $X$  be a Petri curve of genus  $g \geq 3$ . Then  $U(3, d, 4) \neq \emptyset$  if and only if  $\beta(3, d, 4) \geq 0$ .*

*Proof.* According to Proposition 6.5, the result holds for  $g \geq 8$ . For lower values of  $g$ , the result holds by Proposition 6.4 in the following cases

- $g = 3, d = 6$ ;
- $g = 4, d = 6, 7$ ;
- $g = 5, d = 7, 8$ ;
- $g = 6, d = 8, 9$ ;

- $g = 7, d = 9, 10$ .

For  $g \neq 5$ , Proposition 6.8 and Remark 2.3 give the result for all  $d \geq g + 3 - \lfloor \frac{g}{4} \rfloor$ , i.e. for all  $\beta \geq 0$ .

When  $g = 5$ , Remark 2.2 gives the result for  $d = 10, 11$  and Proposition 6.6 for  $d \geq 13$ , leaving only  $d = 9, 12$  open. For  $g = 5, d = 9$ , the inequalities  $d < g + n + \frac{g}{n-1}$ ,  $d > g + n$  and  $\frac{d}{n} < \frac{2g}{2n-1} + 2$  are all satisfied and the result follows from Propositions 6.9, 6.10 and 6.12. Finally, the case  $d = 12$  now follows using Remark 2.2.  $\square$

**Theorem 7.3.** *Let  $X$  be a Petri curve of genus  $g \geq 3$ . Then  $U(4, d, 5) \neq \emptyset$  if and only if  $\beta(4, d, 5) \geq 0$ .*

*Proof.* Proposition 6.5 gives  $U(4, d, 5) \neq \emptyset$  for  $g \geq 15$ . Now Proposition 6.4 covers the following cases

- $g = 3, d = 7$ ;
- $g = 4, d = 8$ ;
- $g = 5, d = 8, 9$ ;
- $g = 6, d = 9, 10$ ;
- $g = 7, d = 10, 11$ ;
- $g = 8, d = 11, 12$ ;
- $g = 9, d = 12, 13$ ;
- $g = 10, d = 12, 13, 14$ ;
- $g = 11, d = 13, 14, 15$ ;
- $g = 12, d = 14, 15, 16$ ;
- $g = 13, d = 15, 16, 17$ ;
- $g = 14, d = 16, 17, 18$ .

Proposition 6.8 now gives the following additional cases

- $g = 4, d = 9, 11$ ;
- $g = 5, d = 11$ ;
- $g = 8, d = 13$ ;
- $g = 9, d = 15$ ;
- $g = 10, d = 15$ ;
- $g = 12, d = 17$ ;
- $g = 14, d = 19$ .

Remark 2.3 now completes the argument for  $g = 10, 12, 14$ .

For other  $g$ , we try using extensions of coherent systems. Propositions 6.9, 6.10 and 6.12, together with Proposition 6.8, give the following additional cases

- $g = 5, d = 10$ ;
- $g = 6, d = 11$ ;
- $g = 7, d = 12, 13$ ;

- $g = 8, d = 14$ ;
- $g = 9, d = 14$ ;
- $g = 11, d = 16$ ;
- $g = 13, d = 18$ .

Again using Remark 2.3, this completes the argument for  $g = 5, 7, 8, 9, 11, 13$ . Moreover, in view of Proposition 6.6, the only outstanding cases are  $g = 3, d = 8, 9, 10, 12, g = 4, d = 10, 14$  and  $g = 6, d = 12, 16$ .

**Proposition 7.4.** *Suppose that  $X$  is a Petri curve of genus 3 and  $d = 8, 9$  or 12. Then  $U(4, d, 5) \neq \emptyset$ .*

*Proof.* Suppose first that  $d = 8$ . Since  $d = 2n$ , the result then follows from [7, Theorem 5.4]. For  $d = 9$ , we now use Proposition 6.8 and, for  $d = 12$ , we apply Remark 2.2.  $\square$

**Proposition 7.5.** *Suppose that  $X$  is a Petri curve of genus 6 and  $d = 12$  or 16. Then  $U(4, d, 5) \neq \emptyset$ .*

*Proof.* In view of Remark 2.2, it is sufficient to prove that  $U(4, 12, 5) \neq \emptyset$ . Note that in this case we have

$$12 = d = g + n + \frac{g}{n-1} \text{ and } \frac{d}{n} = 3 < \frac{2g}{2n-1} + 2 = \frac{12}{7} + 2.$$

Let  $(E, V)$  be a generic element of  $G_L(4, 12, 5)$  and suppose that  $E$  is not stable. By Remark 6.13 and Proposition 6.10, the only possible form for a destabilising sequence is

$$(7.1) \quad 0 \rightarrow (E_1, V_1) \rightarrow (E, V) \rightarrow (E_2, V_2) \rightarrow 0, \quad E_2 \text{ stable}, n_2 \leq 2.$$

Moreover, all the inequalities in the proof of Proposition 6.9 must be equalities, which is the case if and only if

$$n_1 = n_2 = 2 \text{ and } d_1 = d_2 = 6.$$

Since (7.1) is the only possible form for a destabilising sequence with  $E_2$  stable, it follows that  $E$  is semistable. If  $k_2 > 3$ , then [25, Lemma 3.9] applies to give  $h^0(\det E_2) \geq 5$ , which would require  $d_2 \geq 9$  by classical Brill-Noether theory, a contradiction. So  $k_2 = 3$  and  $k_1 = 2$ .

Since  $(E_2, V_2)$  is generated and  $h^0(E_2^*) = 0$ , we have  $(E_2, V_2) \in U(2, 6, 3)$ , which has dimension  $\beta(2, 6, 3) = 0$ . Since  $E$  is semistable and  $\mu(E_1) = \mu(E)$ ,  $E_1$  is also semistable. Moreover,  $(E_1, V_1)$  must be generically generated, otherwise it would have a subsystem  $(\mathcal{L}, V_1)$  with  $\mathcal{L}$  a line bundle, contradicting the  $\alpha$ -stability of  $(E, V)$ . It follows that any subsystem  $(\mathcal{L}_1, W_1)$  of  $(E_1, V_1)$  with  $\mathcal{L}_1$  of rank 1 has  $\deg \mathcal{L}_1 \leq 3$  and  $\dim W_1 \leq 1$ , so  $(E_1, V_1)$  is  $\alpha$ -semistable for all  $\alpha > 0$ . Now, by [5, Theorem 5.6],

$$\dim G_L(2, 6, 2) = \beta(2, 6, 2) = 9.$$

On the other hand, if  $(E_1, V_1) \notin G_L(2, 6, 2)$ , we have

$$(7.2) \quad 0 \rightarrow (\mathcal{L}_1, W_1) \rightarrow (E_1, V_1) \rightarrow (\mathcal{L}_2, W_2) \rightarrow 0$$



with

$$\deg \mathcal{L}_1 = \deg \mathcal{L}_2 = 3 \text{ and } \dim W_1 = \dim W_2 = 1.$$

Moreover, for the extensions (7.2), we have, by (2.6),

- $C_{21} = 3 - 1 = 2$ ;
- $\dim \mathbb{H}_{21}^0 = \dim \text{Hom}((\mathcal{L}_2, W_2), (\mathcal{L}_1, W_1)) \leq 1$ ;
- $\dim \mathbb{H}_{21}^2 = 0$  by (2.7),

so

$$\dim \text{Ext}^1((\mathcal{L}_2, W_2), (\mathcal{L}_1, W_1)) \leq C_{21} + 1 = 3.$$

Since  $(\mathcal{L}_1, W_1)$  and  $(\mathcal{L}_2, W_2)$  each depend on 3 parameters, the extensions (7.2) depend on at most

$$3 + 3 + 3 - 1 = 8 < \beta(2, 6, 2)$$

parameters.

We now consider the extensions (7.1) with  $(E_1, V_1)$ ,  $(E_2, V_2)$  as above. We have, by (2.6) and (2.7),

- $C_{12} = 12 - 6 = 6$ ;
- $\dim \mathbb{H}_{21}^2 = h^0(E_1^* \otimes N_2 \otimes K) \leq 3$  by [10, Theorem 2.1] since  $E_1^* \otimes N_2 \otimes K$  is semistable of rank 2 and slope

$$-\frac{d_1}{2} + \deg N_2 + \deg K = -3 - 6 + 10 = 1;$$

- $\mathbb{H}_{21}^0 = 0$  by Remark 2.7.

So, by Lemma 2.6, the general  $(E, V) \in G_L(4, 12, 5)$  does not admit an extension (7.1) and we are done. □

**Proposition 7.6.** *Suppose that  $X$  is a Petri curve of genus 3 or 4 and  $d = 10$  or 14. Then  $U(4, d, 5) \neq \emptyset$ .*

*Proof.* In view of Remark 2.2, it is sufficient to prove that  $U(4, 10, 5) \neq \emptyset$ . Let  $(E, V)$  be a generic element of  $G_L(4, 10, 5)$  and suppose that  $E$  is not stable. Then we have a destabilising sequence

$$(7.3) \quad 0 \rightarrow (E_1, V_1) \rightarrow (E, V) \rightarrow (E_2, V_2) \rightarrow 0$$

satisfying the conditions of (6.1). We have the following possibilities.

- $n_2 = 1$ :  $3 \leq \mu(E_2) \leq \frac{5}{2}$ , which is a contradiction.
- $n_2 = 2$ :  $\frac{1}{2}(g+1) \leq \mu(E_2) \leq \frac{5}{2}$ , so  $d_2 = 4$  or  $5$  if  $g = 3$ ,  $d_2 = 5$  if  $g = 4$ ; moreover  $k_2 \geq 3$  and, by [27],  $h^0(E_2) \leq \frac{7}{2}$ , so  $k_2 = 3$ .
- $n_2 = 3$ :  $2 \leq \mu(E_2) \leq \frac{5}{2}$ , so  $d_2 = 6$  or  $7$ ; moreover  $k_2 \geq 4$  and, by [27],  $h^0(E_2) \leq \frac{d_2+3}{2}$ , giving the possibilities  $(d_2, k_2) = (6, 4), (7, 4), (7, 5)$ .

We consider first the case  $n_2 = 3$ . If  $k_2 = 4$ , we are in the situation of (6.2) and Proposition 6.10 applies. In the remaining case  $d_2 = 7$ ,  $k_2 = 5$ , we have  $h^0(\det E_2) = 8 - g \leq 5$ . So, by [25, Lemma 3.9],  $E_2$  possesses either a line subbundle  $\mathcal{L}$  with  $h^0(\mathcal{L}) \geq 2$  or a subbundle  $F$  of rank 2 with  $h^0(F) \geq 3$ . In the first case, since  $E_2$  is stable, we have  $\deg \mathcal{L} \leq 2$ , a contradiction. In the second case  $d_F := \deg F \leq 4$  and any line subbundle of  $F$  has  $\deg \mathcal{L} \leq 2$ , hence  $h^0(\mathcal{L}) \leq 1$ . It follows that, for any subspace  $W$  of  $H^0(F)$  of dimension 3,  $(F, W) \in G_L(2, d_F, 3)$ . Hence, by Theorem 3.1(1),  $\beta(2, d_F, 3) \geq 0$ . Since  $d_F \leq 4$ , this holds only when  $g = 3$ ,  $d_F = 4$ . It follows that  $F$  is semistable and, by [27],  $h^0(F) \leq 3$  and hence  $h^0(F) = 3$ . Note further that  $F$  is not strictly semistable, for otherwise we would have a sequence  $0 \rightarrow \mathcal{L}_1 \rightarrow F \rightarrow \mathcal{L}_2 \rightarrow 0$  with  $\deg \mathcal{L}_1 = \deg \mathcal{L}_2 = 2$ , so that  $h^0(F) \leq 2$ . Hence  $F$  is stable and  $(F, W) \in U(2, 4, 3)$ . Now let  $W_1 := H^0(F) \cap V_2$  and consider the exact sequence

$$(7.4) \quad 0 \rightarrow (F, W_1) \rightarrow (E_2, V_2) \rightarrow (\mathcal{L}, W_2) \rightarrow 0,$$

where  $\dim W_1 \leq 3$ . If  $\dim W_1 < 3$ , then  $\dim W_2 \geq 3$ , contradicting the fact that  $\deg \mathcal{L} = 3$ . So  $\dim W_2 = 2$ ,  $\dim W_1 = 3$  and

$$(F, W_1) \in U(2, 4, 3), \quad (\mathcal{L}, W_2) \in U(1, 3, 2).$$

For the extensions (7.4), we have, by (2.6) and (2.7),

- $C_{21} = 4 - 4 + 6 - 6 = 0$ ;
- $\mathbb{H}_{21}^0 = 0$  by Remark 2.7;
- $\dim \mathbb{H}_{21}^2 = h^0(F^* \otimes \mathcal{L}^* \otimes K)^* = 0$  since  $F^* \otimes \mathcal{L}^* \otimes K$  is stable of degree  $-2$ .

So, by (2.8), the extension (7.4) splits, which contradicts the stability of  $E_2$ . We have therefore proved that the only possible destabilising sequences for a general  $(E, V)$  of type (7.3) with  $E_2$  stable are those with  $n_2 = 2$ .

Suppose then that  $n_2 = 2$ . We have  $k_2 = h^0(E_2) = 3$  and we know that  $(E_2, V_2)$  is generated and  $h^0(E_2^*) = 0$ , so  $(E_2, V_2) \in U(2, d_2, 3)$ . Suppose now that  $E$  is semistable, so that  $d_2 = 5$ . Then also  $E_1$  is semistable and in fact stable since  $\gcd(n_1, d_1) = 1$ . It follows that any line subbundle  $\mathcal{L}$  of  $E_1$  has  $\deg \mathcal{L} \leq 2$  and hence  $h^0(\mathcal{L}) \leq 1$ . So  $(E_1, V_1) \in U(2, 5, 2)$ . For the extensions (7.3), we have, by (2.6) and (2.7),

- $C_{12} = 10 - 6 = 4$ ;
- $\dim \mathbb{H}_{21}^2 = h^0(E_1^* \otimes N_2 \otimes K) = 0$  since  $E_1^* \otimes N_2 \otimes K$  is stable with slope  $< 0$ ;
- $\mathbb{H}_{21}^0 = 0$  by Remark 2.7.

So, by Lemma 2.6, the general  $(E, V)$  does not admit an extension of this type.

It remains to consider the possibility that  $E$  is not semistable. From the above, this can happen only when  $g = 3$  and we have an extension (7.3) with

$$n_1 = n_2 = 2, \quad d_1 = 6, \quad d_2 = 4, \quad k_1 = 2, \quad k_2 = 3.$$

We certainly have  $(E_2, V_2) \in U(2, 4, 3)$ , but we can no longer guarantee that  $E_1$  is semistable. However the maximal degree of a line subbundle of  $E_1$  is 4, for otherwise

$E$  would have a quotient bundle of rank 3 and degree  $\leq 5$ ; this cannot be stable since  $E$  has no stable quotient bundles of rank 3 contradicting the stability of  $E$ . It follows that  $E$  would have either a quotient line bundle of degree  $\leq 1$  or a stable quotient bundle of rank 2 of degree  $\leq 3$ ; both of these are impossible (see the itemized list following (7.3)). Moreover, we can still argue as in the proof of Proposition 7.5 to show that  $(E_1, V_1)$  depends on at most  $\beta(2, 6, 2)$  parameters. Now for the extensions (7.3), we have, by (2.6) and (2.7),

- $C_{12} = 12 - 6 = 6$ ;
- $\dim \mathbb{H}_{21}^2 = h^0(E_1^* \otimes N_2 \otimes K) = 0$  since  $\deg N_2 \otimes K = 0$  and the maximal degree of a line subbundle of  $E_1^*$  is  $-2$ ;
- $\mathbb{H}_{21}^0 = 0$  by Remark 2.7.

The result now follows from another application of Lemma 2.6. □

This completes the proof of Theorem 7.3. □

**Remark 7.7.** In the course of proving Proposition 7.6, we have shown that there is no coherent system  $(E_2, V_2)$  of type  $(3, 7, 5)$  on a Petri curve of genus 3 or 4 with  $E_2$  stable. A slight modification of the proof shows that  $G(\alpha; 3, 7, 5) = \emptyset$  for all  $\alpha > 0$  and all  $g \geq 3$  (we have to prove that  $E_2$  is stable for all  $(E_2, V_2) \in G(\alpha; 3, 7, 5)$ ). Since  $\beta(3, 7, 5) = 17 - 6g < 0$  for  $g \geq 3$ , this is to be expected, but, so far as we are aware, it has previously been proved only for  $g \geq 6$  (see [8, Theorem 3.9], where it is shown that, for  $k > n$ ,  $G(\alpha; n, d, k) = \emptyset$  if  $\beta(n, d, n+1) < 0$ ; in this case  $\beta(3, 7, 4) = 16 - 3g < 0$  if and only if  $g \geq 6$ ).

## 8. LOW GENUS

The cases  $g = 0$  and  $g = 1$  have been excluded from the earlier part of this paper since they present special features and have been handled elsewhere [17, 18].

For  $g = 0$ , there are no stable bundles of rank  $\geq 2$ , so  $U(n, d, n+1)$  is always empty if  $n \geq 2$ . Moreover, if  $d$  is not divisible by  $n$ , there exist no semistable bundles; hence  $U^s(n, d, n+1) = \emptyset$ . For the remaining case, when  $d$  is divisible by  $n$ ,  $U^s(n, d, n+1) \neq \emptyset$  (see [17, Proposition 6.4]). One may note that in this case  $\beta \geq 0$  is equivalent to  $d \geq n$ .

For  $g = 1$ , the moduli spaces  $G(\alpha)$  are well understood (see [18]). The results for  $U(n, d, n+1)$  and  $U^s(n, d, n+1)$  are summarised in the following theorem.

**Theorem 8.1.** *Let  $X$  be a curve of genus 1 and  $n \geq 2$ . Then*

- $U^s(n, d, n+1) \neq \emptyset$  if and only if  $d \geq n+1$ ;
- $U(n, d, n+1) \neq \emptyset$  if and only if  $d \geq n+1$  and  $\gcd(n, d) = 1$ .

*Proof.* The first part follows from the main theorem of [18] and [18, Remark 6.3]. For the second part, recall that, on an elliptic curve, stable bundles exist if and only if  $(n, d) = 1$ , and, in this case, all semistable bundles are stable.  $\square$

The condition  $d \geq n + 1$  here is precisely equivalent to  $\beta \geq 0$ .

For  $g = 2$ , note first that the case  $g = n = 2$ ,  $d = 4$  is a genuine exception in Proposition 6.4 (see [8, Lemma 6.6(1)]). More generally, if  $E$  is any bundle of rank  $n \geq 2$  and degree  $2n$  with  $h^0(E) \geq n + 1$  on a curve of genus 2, then  $E$  cannot be stable. In fact, by Riemann-Roch, we have  $h^1(E) \geq 1$ , so there exists a non-zero homomorphism  $E \rightarrow K$ , which immediately contradicts stability. There do exist semistable bundles with  $h^0(E) \geq n + 1$ , which can be constructed as in the proof of Proposition 6.6 or by using sequences

$$0 \rightarrow E^* \rightarrow V \otimes \mathcal{O} \rightarrow \mathcal{L} \rightarrow 0$$

with  $\deg \mathcal{L} = 2n$  and  $V$  a subspace of  $H^0(\mathcal{L})$  of dimension  $n + 1$  which generates  $\mathcal{L}$ ; the coherent system  $(E, V^*)$  is then  $\alpha$ -stable for all  $\alpha > 0$ . We deduce

**Theorem 8.2.** *Let  $X$  be a curve of genus 2 and  $n \geq 2$ . Then*

- $U^s(n, d, n + 1) \neq \emptyset$  if and only if  $d \geq n + 2$  (or equivalently  $\beta \geq 0$ );
- $U(n, d, n + 1) \neq \emptyset$  if and only if  $d \geq n + 2$ ,  $d \neq 2n$ .

*Proof.* We have  $U(n, d, n + 1) \neq \emptyset$  in the following cases:

- $d \geq 3n$  by [8, Proposition 2.6];
- $d = n + 2, \dots, 2n - 1$  by [7, Theorem 5.5];
- $d = 2n + 2, \dots, 3n - 1$  by Remark 2.2.

Moreover  $U^s(n, 2n, n + 1) \neq \emptyset$  by Proposition 6.6. It remains to prove

- (i)  $U(n, 2n, n + 1) = \emptyset$ ;
- (ii)  $U(n, 2n + 1, n + 1) \neq \emptyset$ .

For (i), we have already remarked that a vector bundle  $E$  of rank  $n$  and degree  $2n$  with  $h^0(E) \geq n + 1$  cannot be stable (see also [22, Théorème 2]).

For (ii), every stable bundle  $E$  of rank  $n$  and degree  $2n + 1$  has  $h^0(E) \geq n + 1$ . If we can prove that there exists such a bundle which is generated, we can choose a subspace  $V$  of  $H^0(E)$  of dimension  $n + 1$  such that  $(E, V)$  is generated. Then  $(E, V) \in U(n, d, n + 1)$  by Proposition 6.3.

To show that  $E$  is generated, we need to prove that  $h^1(E(-x)) = 0$  for all  $x \in X$ . Now  $E(-x)$  is stable of degree  $n + 1$  and  $E(-x)^* \otimes K$  is stable of degree  $n - 1$ . We consider the Brill-Noether locus  $B(n, n - 1, 1)$ . By [28] or [10], this locus has dimension  $\beta(n, n - 1, 1)$  and hence codimension

$$1 - (n - 1) + n(g - 1) = 2$$

in  $M(n, n-1)$ . It follows that the generic  $E \in M(n, 2n+1)$  has

$$h^1(E(-x)) = h^0(E(-x)^* \otimes K) = 0$$

for all  $x \in X$  as required.

This completes the proof of the theorem.  $\square$

**Theorem 8.3.** *Let  $X$  be a Petri curve of genus 3 and  $n \geq 2$ . Then  $U(n, d, n+1) \neq \emptyset$  if  $\beta \geq 0$ , except possibly when  $n \geq 5$ ,  $d = 2n + 2$ .*

*Proof.* For  $n = 2, 3, 4$ , this has already been proved. For  $n \geq 5$ , we have  $U(n, d, n+1) \neq \emptyset$  in the following cases:

- $d \geq 3n + 1$  by Proposition 6.6;
- $d = n + 3, \dots, 2n$  by [7, Theorem 5.4];
- $d = 2n + 1$  by Proposition 6.8;
- $d = 2n + 3, \dots, 3n$  by Remark 2.2.

$\square$

**Remark 8.4.** For general  $X$  (but not necessarily for all Petri  $X$ ), the exception can be removed using Teixidor's degeneration methods [30].

**Remark 8.5.** For  $g = 4, 5$  and  $n \geq 5$ , a similar argument works with the following possible exceptions

- $g = 4$ ,  $d = 2n + 2, 2n + 3, 3n + 2, 3n + 3$ ;
- $g = 5$ ,  $n = 5$ ,  $d = 12, 13, 17, 18$ ;
- $g = 5$ ,  $n \geq 6$ ,  $d = 2n + 2, 2n + 3, 2n + 4, 3n + 2, 3n + 3, 3n + 4$ .

For general  $X$ , one can use Teixidor's result to rule out some of the exceptions.

## 9. APPLICATIONS TO BRILL-NOETHER THEORY AND DUAL SPANS

We recall from section 2 that the *Brill-Noether locus*  $B(n, d, k)$  and  $\tilde{B}(n, d, k)$  are defined by

$$B(n, d, k) = \{E \in M(n, d) \mid h^0(E) \geq k\}$$

and

$$\tilde{B}(n, d, k) = \{[E] \in \tilde{M}(n, d) \mid h^0(\text{gr}(E)) \geq k\},$$

It follows that the formula  $(E, V) \mapsto [E]$  defines a morphism

$$\psi : G_0(n, d, k) \rightarrow \tilde{B}(n, d, k),$$

whose image contains  $B(n, d, k)$ .

The following theorem, which is essentially a restatement of [5, Theorem 11.4 and Corollary 11.5] for the case  $k = n + 1$ , is true for any smooth curve; we state it in a very general and formal way to make it applicable in a wide variety of situations.

**Theorem 9.1.** *Suppose that  $B(n, d, n + 1) \neq \emptyset$ . Then*

- (1)  $\psi$  is one-to-one over  $B(n, d, n + 1) - B(n, d, n + 2)$ ;
- (2) if  $G_0(n, d, n + 1)$  is irreducible, then  $B(n, d, n + 1)$  is irreducible;
- (3) if  $\beta(n, d, n + 1) \leq n^2(g - 1)$  and  $G_0(n, d, n + 1)$  is smooth and irreducible, then

$$\text{Sing}B(n, d, n + 1) = B(n, d, n + 2)$$

and  $G_0(n, d, n + 1)$  is a desingularisation of the closure  $\overline{B}(n, d, n + 1)$  of  $B(n, d, n + 1)$  in  $\widetilde{M}(n, d)$ .

*Proof.* (1) is obvious.

(2) follows from (1) and the fact that  $B(n, d, n + 1)$  is a Zariski-open subset of  $\psi(G_0(n, d, n + 1))$ . [Note that the hypothesis  $\beta(n, d, n + 1) \leq n^2(g - 1)$  of [5, Conditions 11.3] is not needed here.]

(3) follows from [5, Corollary 11.5]. □

Of course, if  $U(n, d, n + 1) \neq \emptyset$ , then  $B(n, d, n + 1) \neq \emptyset$ . Thus we have many instances in this paper for which  $B(n, d, n + 1) \neq \emptyset$ . We shall not list all of them as we shall be stating a more specific result later. For the time being, we note the following two corollaries. The first is a slightly extended version of [8, Corollary 4.5], the second is new.

**Corollary 9.2.** *Suppose that  $X$  is a Petri curve,  $g + n - \lfloor \frac{g}{n+1} \rfloor \leq d \leq g + n$  and  $(g, n) \neq (2, 2)$ . Then*

- (1)  $B(n, d, n + 1)$  is irreducible of dimension  $\beta(n, d, n + 1)$  and smooth outside  $B(n, d, n + 2)$ ;
- (2)  $G_L(n, d, n + 1)$  is a desingularisation of  $\overline{B}(n, d, n + 1)$ ;
- (3) if either  $d < g + n$  or  $d = g + n$  and  $n \nmid g$ ,  $B(n, d, n + 1)$  is projective and  $G_L(n, d, n + 1)$  is a desingularisation of  $B(n, d, n + 1)$ .

*Proof.* The condition on  $d$  implies that  $\alpha_l \leq 0$ . Hence, by Theorem 3.1,  $G_0(n, d, n + 1) = G_L(n, d, n + 1)$  and is smooth and irreducible of dimension  $\beta(n, d, n + 1)$ . Moreover  $U(n, d, n + 1) \neq \emptyset$  by Proposition 6.4. (1) and (2) now follow from Theorem 9.1. For (3), we note that, under the stated conditions on  $d$ ,  $E$  is stable for every  $(E, V) \in G_L(n, d, n + 1)$  [8, Proposition 3.5]; hence  $\psi(G_L(n, d, n + 1)) = B(n, d, n + 1)$ . □

**Remark 9.3.** When  $g = n = 2$  and  $d = 4$ ,  $B(2, 4, 3) = \emptyset$  by [8, Lemma 6.6], but  $G_L(2, 4, 3) \neq \emptyset$ . In this case, the image of  $\psi$  is contained in  $\widetilde{M}(2, 4) \setminus M(2, 4)$ .

**Corollary 9.4.** *Suppose that  $X$  is a Petri curve and that all the flip loci for coherent systems of type  $(n, d, n + 1)$  have dimension  $\leq \beta(n, d, n + 1) - 1$ . If  $B(n, d, n + 1) \neq \emptyset$ , then*

- $B(n, d, n + 1)$  is irreducible;

- $B(n, d, n+1)$  is smooth of dimension  $\beta(n, d, n+1)$  at  $E$  whenever  $E$  is generically generated and  $h^0(E) = n+1$ .

*Proof.* The hypotheses imply that  $G_0(n, d, n+1)$  is birational to  $G_L(n, d, n+1)$  and is therefore irreducible. Irreducibility of  $B(n, d, n+1)$  follows from Theorem 9.1(2). If  $E$  is stable,  $h^0(E) = n+1$  and  $E$  is generically generated, then  $(E, H^0(E)) \in U(n, d, n+1)$ , which is smooth of dimension  $\beta(n, d, n+1)$  by Theorem 3.1(4). The result follows from [5, Theorem 11.4(iv)].  $\square$

We know that this corollary has genuine content since the flip loci at  $\alpha_l = \alpha_L$  have dimension  $\leq \beta(n, d, n+1) - 1$  (Corollary 5.2 and Proposition 5.4).

We now turn to our second application. Suppose that  $\mathcal{L}$  is a generated line bundle of degree  $d > 0$  and let  $V$  be a linear subspace of  $H^0(\mathcal{L})$  of dimension  $n+1$  which generates  $\mathcal{L}$  (in other words,  $(\mathcal{L}, V)$  is a generated coherent system of type  $(1, d, n+1)$ ). We have an evaluation sequence

$$(9.1) \quad 0 \longrightarrow M_{V, \mathcal{L}} \longrightarrow V \otimes \mathcal{O} \longrightarrow \mathcal{L} \longrightarrow 0.$$

This is also known as the *dual span* construction (see [12]) and has been used in the context of coherent systems in [5, 8] and also in the proof of Proposition 3.7. The following is a special case of [12, Conjecture 2].

**Conjecture 9.5.** *Let  $X$  be a Petri curve of genus  $g \geq 3$ . Suppose that  $\beta := \beta(1, d, n+1) \geq 0$  and that  $\mathcal{L}$  is a general element of  $B(1, d, n+1)$  (when  $\beta = 0$ ,  $\mathcal{L}$  can be any element of the finite set  $B(1, d, n+1)$ ) and let  $V$  be a general subspace of  $H^0(\mathcal{L})$  of dimension  $n+1$ . Then  $M_{V, \mathcal{L}}$  is stable.*

This conjecture is related to our results by the following simple proposition (compare [5, Theorem 5.11]).

**Proposition 9.6.** *Suppose that  $X$  is a Petri curve. The following are equivalent:*

- (1) *there exists a generated coherent system  $(\mathcal{L}, V)$  of type  $(1, d, n+1)$  with  $M_{V, \mathcal{L}}$  stable;*
- (2)  $U(n, d, n+1) \neq \emptyset$ .

*Proof.* For (1) $\Rightarrow$ (2), we note that  $(M_{V, \mathcal{L}}^*, V^*)$  is a generated coherent system of type  $(n, d, n+1)$  with  $M_{V, \mathcal{L}}^*$  stable, so  $(M_{V, \mathcal{L}}^*, V^*) \in U(n, d, n+1)$  by Proposition 6.3. Conversely, suppose  $U(n, d, n+1) \neq \emptyset$ . If  $\beta(n, d, n+1) > 0$ , the generic element of  $U(n, d, n+1)$  is a generated coherent system  $(E, W)$  with  $h^0(E^*) = 0$  and  $E$  stable. If  $\beta(n, d, n+1) = 0$ , then all elements of  $U(n, d, n+1)$  have this property. The dual of the evaluation sequence of  $(E, W)$  can be written as

$$0 \longrightarrow E^* \longrightarrow W^* \otimes \mathcal{O} \longrightarrow \mathcal{L} \longrightarrow 0,$$

where  $\mathcal{L}$  is a line bundle of degree  $d$ . It follows that  $M_{W^*, \mathcal{L}} \cong E^*$  and is therefore stable, proving (1).  $\square$

**Remark 9.7.** By Theorem 8.2 and Proposition 9.6, the conjecture fails for  $g = 2$ ,  $d = 2n$ , but is otherwise true for  $g = 2$ . In fact, although Butler [12, §1] discusses the question of whether  $M_{V,\mathcal{L}}$  is stable, [12, Conjecture 2] actually has the weaker conclusion that  $(M_{V,\mathcal{L}}^*, V^*) \in G_0(n, d, n+1)$ . In this form the conjecture is true for  $g = 2$  (see Theorem 8.2).

Using Proposition 9.6, we can now begin to form a list of cases for which Conjecture 9.5 holds. In the list we have noted where each case was proved.

- $g + n - \lfloor \frac{g}{n+1} \rfloor \leq d \leq g + n$  ([12], [8, Proposition 4.1]);
- $g \geq n^2 - 1$  ([12], [8, Proposition 4.6]);
- $d \geq d_1$  (Proposition 6.6, [30]);
- $d \leq 2n$  ([20, 22, 7]);
- $n = 3, 4$  (Theorems 7.2, 7.3)

The first and fourth items in this list can be expanded further by the use of Remark 2.3 and Proposition 6.8. According to the analysis in section 7, the following cases for  $n = 3$  and  $n = 4$  depend on the use of extensions of coherent systems (possibly in conjunction with other methods):

- $n = 3, g = 5, d = 9, 12$ ;
- $n = 4, g = 3, d = 10$ ;
- $n = 4, g = 4, d = 10, 14$ ;
- $n = 4, g = 5, d = 10, 14$ ;
- $n = 4, g = 6, d = 11, 12, 15, 16$ ;
- $n = 4, g = 7, d = 12, 13, 16, 17, 20$ ;
- $n = 4, g = 8, d = 14, 18$ ;
- $n = 4, g = 9, d = 14, 18, 22$ ;
- $n = 4, g = 11, d = 16, 20, 24, 28$ ;
- $n = 4, g = 13, d = 18, 22, 26, 30$ .

All of these cases, and those depending on Propositions 6.6 and 6.8, are (so far as we are aware) new.

Of the methods we have used, the only ones capable of further development appear to be elementary transformations (using direct sums of higher rank vector bundles) and extensions of coherent systems (using more refined calculations). The methods of [30] could also yield improved results for general  $X$ .

## REFERENCES

- [1] E. Arbarello, M. Cornalba, P. A. Griffiths and J. Harris, *Geometry of Algebraic curves* Volume I, Springer-Verlag, New York 1985.
- [2] E. Ballico, Coherent systems with many sections on projective curves, preprint.
- [3] A. Beauville, Some stable vector bundles with reducible theta divisor, *Manuscripta Math.* **110** (2003) 343–349.



- [4] S. B. Bradlow and O. Garcia-Prada, An application of coherent systems to a Brill-Noether problem, *J. Reine Angew Math.* **551** (2002) 123–143.
- [5] S. B. Bradlow, O. Garcia-Prada, V. Muñoz and P. E. Newstead, Coherent systems and Brill-Noether theory, *Internat. J. Math.* **14** (2003) 683–733.
- [6] S. B. Bradlow, O. Garcia-Prada, V. Mercat, V. Muñoz and P. E. Newstead, On the geometry of moduli spaces of coherent systems on algebraic curves, *Internat. J. Math.* **18** (2007) 411–453.
- [7] S. B. Bradlow, O. Garcia-Prada, V. Mercat, V. Muñoz and P. E. Newstead, Moduli spaces of coherent systems of small slope on algebraic curves, preprint, arXiv:0707.0983.
- [8] L. Brambila-Paz, Non-emptiness of moduli spaces of coherent systems, *Internat. J. Math.*, to appear.
- [9] L. Brambila-Paz and A. Ortega, Tensor product of coherent systems, preprint, arXiv:0711.3944
- [10] L. Brambila-Paz, I. Grzegorzcyk and P. E. Newstead, Geography of Brill-Noether loci for small slopes, *J. Alg. Geom.* **6** (1997) 645–669.
- [11] D. C. Butler, Normal generation of vector bundles over a curve, *J. Diff. Geom.* **39** (1994) 1–34.
- [12] D. C. Butler, Birational maps of moduli of Brill-Noether pairs, preprint, arXiv:alg-geom/9705009.
- [13] L. Ein and R. Lazarsfeld, Stability and restrictions of Picard bundles with an application to the normal bundles of elliptic curves, *Complex projective geometry (Trieste 1989/Bergen 1989)*, ed. G. Ellingsrud, C. Peskine, G. Sacchiero and S. A. Stromme, LMS Lecture Notes Series **179** 149–156, CUP, Cambridge 1992.
- [14] M. L. Green, Koszul Cohomology and the Geometry of Projective Varieties, *J. Diff. Geom.* **19** (1984) 125–171.
- [15] M. He, Espaces de modules de systèmes cohérents, *Internat. J. Math.* **9** (1998) 545–598.
- [16] A. King and P. E. Newstead, Moduli of Brill-Noether pairs on algebraic curves *Internat. J. Math.* (1995) 733–748.
- [17] H. Lange and P. E. Newstead, Coherent systems of genus 0, *Internat. J. Math.* **15** (2004) 409–424.
- [18] H. Lange and P. E. Newstead, Coherent systems on elliptic curves, *Internat. J. Math.* **16** (2005) 787–805.
- [19] J. Le Potier, Faisceaux semi-stables et systèmes cohérents, *Vector bundles in Algebraic Geometry*, Durham 1993, ed N. J. Hitchin, P. E. Newstead and W. M. Oxbury, LMS Lecture Notes Series **208**, 179–239.
- [20] V. Mercat, Le problème de Brill-Noether pour les fibrés stables de petite pente, *J. Reine Angew. Math.* **506** (1999), 1–41.
- [21] V. Mercat, Le problème de Brill-Noether et la théorème de Teixidor, *Manuscripta Math.* **98** (1999) 75–85.
- [22] V. Mercat, Fibrés stables de pente 2, *Bull. London Math. Soc.* **33** (2001) 535–542.
- [23] E. Mistretta, Stability of line bundles transforms on curves with respect to low codimensional subspaces, preprint, arXiv:math/0703465.
- [24] M. S. Narasimhan and S. Ramanan, Deformations of the moduli space of vector bundles over an algebraic curve *Ann. of Math.* **101** (1975) 391–417.
- [25] K. Paranjape and S. Ramanan, On the canonical ring of a curve, *Algebraic geometry and Commutative Algebra in Honor of Masayoshi Nagata* (1987) 503–516.
- [26] N. Raghavendra and P. A. Vishwanath, Moduli of pairs and generalized theta divisors, *Tôhoku Math. J.* **46** (1994) 321–340.
- [27] R. Re, Multiplication of sections and Clifford bounds for stable vector bundles on curves, *Comm. Algebra* **26** (1998) 1931–1944.
- [28] N. Sundaram, Special divisors and vector bundles, *Tôhoku Math. J.* **39** (1987) 175–213.
- [29] M. Teixidor i Bigas, Brill-Noether theory for stable vector bundles, *Duke Math. J.* **62** 9(1991) 385–400.
- [30] M. Teixidor i Bigas, Existence of coherent systems, *Internat. J. Math.*, to appear.

TIFR, HOMI BHABHA ROAD, MUMBAI 400005, INDIA

*E-mail address:* usha@math.tifr.res.in

CIMAT, APDO. POSTAL 402, C.P. 36240. GUANAJUATO, GTO, MÉXICO

*E-mail address:* lebp@cimat.mx

DEPARTMENT OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF LIVERPOOL, PEACH STREET,  
LIVERPOOL L69 7ZL, UK

*E-mail address:* newstead@liv.ac.uk