

Morphological and molecular characterization of a new microsporidian (Protozoa: Microsporidia) isolated from *Spodoptera litura* (Fabricius) (Lepidoptera: Noctuidae)

S. JOHNY^{1,2}, S. KANGINAKUDRU¹, M. C. MURALIRANGAN² and J. NAGARAJU^{1*}

¹Laboratory of Molecular Genetics, Centre for DNA Fingerprinting and Diagnostics, ECIL Road, Nacharam, Hyderabad 500 076, India

²G.S. Gill Research Institute, Guru Nanak College, Chennai 600 042, India

(Received 10 October 2005; revised 8 December 2005; accepted 8 December 2005)

SUMMARY

A microsporidium was isolated from larvae of *Spodoptera litura* (Fabricius) collected from Tamil Nadu, India. This microsporidian species is monomorphic, disporous and develops in direct contact with the cytoplasm of the host cell. The nuclear configuration of merogonic and sporogonic stages was diplokaryotic. The merogonic proliferative stage was unusual that normal development with 1, 2 and 4 binucleated forms were common, while large multinucleate meronts containing 8 and 12 small compact horseshoe-like diplokaryotic nuclei were also observed. The fresh spores were typically ovocylindrical in shape, with a mean size of $3.91 \times 1.91 \mu\text{m}$ and the polar filament length was $\sim 90 \mu\text{m}$. Infection was systemic with mature spores produced in the midgut, nervous tissue, muscles, labial glands, gonads, tracheae, epidermis, Malpighian tubules and, most extensively, fat body tissues. The new isolate was highly pathogenic to *S. litura* larvae. Host specificity tests performed on 37 non-target hosts of 5 different insect orders revealed that the new isolate is pathogenic only to lepidopteran insects. We sequenced the 16S small subunit rRNA (SSU rRNA) gene of the isolate and compared it with 72 non-redundant microsporidian sequences from the GenBank. Based on the light microscopic studies and phylogenetic analyses, the new isolate is assigned to the genus *Nosema*. Significant differences in the SSU rRNA sequence were identified when compared with the type species *Nosema bombycis* and other closely related species viz., *Nosema spodopterae*. Structural differences were also observed in the 16S SSU rRNA between the new isolate and the two above-mentioned microsporidian pathogens. We conclude that the microsporidian isolate reported here is distinctly different from the other known species and is likely to be a new species.

Key words: Microsporidia, *Nosema*, *Spodoptera litura*, 16S SSU rDNA, phylogeny.

INTRODUCTION

Microsporidia are obligate, intracellular parasites belonging to the phylum *Microsporidia*, traditionally placed in the Kingdom Protista. However, evidence from phylogenetic analyses using protein coding genes, especially α - and β -tubulins (Keeling and Fast, 2002; Keeling, 2003), and on LSU rRNA sequences (Van de Peer *et al.* 1998) now suggest that microsporidia share a common origin with fungi. The phylum *Microsporidia* consists of approximately 143 genera and over 1200 species (Wittner and Weiss, 1999), which infect a wide range of vertebrate and invertebrate taxa (Didier *et al.* 2000). Currently, almost half of the described genera of microsporidia have an insect as the host (Becnel and Andreadis, 1999). They may infect the gut, fat body,

reproductive, muscle, nervous, and silk gland tissues and Malpighian tubules, or can be systemic (Becnel and Andreadis, 1999; Solter and Becnel, 2000). There is renewed interest in studying microsporidia because some species cause opportunistic infections in immunocompromised humans, mainly human immunodeficiency virus-positive subjects (Desportes, Le Charpentier and Galian, 1985; Franzen and Muller, 1999; Kotler and Orenstein, 1999; Molina *et al.* 2000), but also in travellers, the elderly, and malnourished children (Weber *et al.* 1994; Muller *et al.* 2001). Other microsporidia are important in agriculture as pathogens of beneficial insects such as silkworm and honeybees (Becnel and Andreadis, 1999; Nageswara Rao *et al.* 2004), or as biological control agents for use against medical or agricultural pests (Lockwood, Bomar and Ewen, 1999; Solter and Becnel, 2000; Briano and Williams, 2002). The members of the genus *Nosema* are often considered the most important and widely distributed group of microsporidia, particularly in the Lepidoptera (Tsai, Lo and Wang, 2003) and most '*Nosema*' isolates from hosts of other insect orders are not closely related

* Corresponding author: Laboratory of Molecular Genetics, Centre for DNA Fingerprinting and Diagnostics, ECIL Road, Nacharam, Hyderabad 500 076, India. Tel: +91 40 27171427. Fax: +91 40 27155610. E-mail: jnagaraju@cdfd.org.in

genetically and genus designations are being changed (Slamovits, Williams and Keeling, 2004; Sokolova *et al.* 2005).

The tobacco caterpillar, *Spodoptera litura* is a polyphagous noctuid with high reproductive potential and the ability to migrate long distances as adults. These factors contribute to the role of *S. litura* as a major pest of many agricultural crops throughout its geographical range (Armes *et al.* 1997). In India, it has attained a major pest status on agricultural crops such as cotton, groundnut and cauliflower. Failure to control *S. litura* in the major cotton and groundnut growing regions of Andhra Pradesh and Tamil Nadu states of India has resulted in an unexpected population outbreak and development of resistance to several insecticides (Armes *et al.* 1997; Johny and Muralirangan, 2000). In order to avoid resistance problems, as well as to address environmental concerns regarding chemical pesticide use, it is important to identify and develop suitable alternate control strategies. In this context, pathogens may be key biological control agents against major agricultural pests because they are eco-friendly and may be manipulated to effectively control insect pests. In an earlier study, Johny (2002) reported a microsporidian pathogen from the field population of *S. litura*. In this paper, we provide a developmental and molecular characterization of this microsporidian isolate from *S. litura*. We have studied the developmental stages including merogony and sporogony, host specificity data, pathogenicity, and the small subunit rRNA (SSU rRNA) gene sequence. The phylogenetic status of the new isolate based on SSU rRNA analyses is also presented.

MATERIALS AND METHODS

Isolation of spores

The microsporidian spores were originally collected from *S. litura* larvae on 18 January 2000 from a tobacco field in the Dindigul District (10°22'N 78°00'E) of Tamil Nadu, India. The spores were harvested and propagated in *S. litura* larvae reared on castor leaves. The spores were purified from infected fat body tissues of dead or moribund *S. litura* larvae collected from the field. The tissues were homogenized in sterile deionized water and the homogenate was filtered through cheesecloth and also cleaned in several cycles of water washes followed by centrifugation. The concentrated spore pellet was further purified by 90% Percoll gradient centrifugation (Undeen and Vavra, 1997). The pelleted spores were collected, washed with sterile distilled water to remove the Percoll content and resuspended in distilled water, quantified by haemocytometer, and stored at 4 °C until further use.

Life-cycle study

S. litura larvae, maintained on castor leaves as a microsporidian-free stock, were used for the present study. Castor leaf discs, each 7 mm diameter, were smeared with 10 µl of water/spore suspension containing 2.4×10^3 spores and allowed to dry at room temperature. Third-instar larvae were starved for 6 h and each of them was fed with 1 leaf disc. The larvae that consumed the entire disc only were retained. Thirty larvae were used for treatment and were maintained separately on fresh castor leaves in plastic Petri dishes at 28 ± 2 °C, light regime 14L:10D.

To study the microsporidian life-cycle, 1 or 2 larvae were dissected every 24 h following the initial infection, up to the completion of the life-cycle. The gut and fat body tissues were removed and placed in a drop of physiological saline (Yeager's solution) on a glass slide, macerated with forceps, fixed in absolute methanol for 2 min, air dried and stained with 10% (v/v) Giemsa staining solution (1 ml of Giemsa stock solution mixed with 9 ml of phosphate buffer, pH 7.4) for about 20–25 min at room temperature. The slides were then rinsed in running water for 5 min, air dried and then observed using a Leitz compound microscope under bright-field conditions for the presence of different stages of the microsporidia. Fresh spores were spread on water agar (Undeen and Vavra, 1997) and measured using an ocular micrometer under phase-contrast microscopy. All the measurements are presented in micrometers as mean values followed parenthetically by range, standard deviation and sample size.

Isolation of DNA, PCR amplification and sequencing of 16S SSU rDNA

The spores of the microsporidian isolate were suspended in 100 mM NaCl, pH 9.5, glycine-NaOH buffer (10 mM) at 30 °C for 15 min to elicit germination (Undeen and Cockburn, 1989). After germination, the emptied content of the spores was used for DNA extraction by standard phenol-chloroform method. The 16S small subunit (SSU) rRNA gene was amplified using the primers 18f 5'-CACCAG-GTTGATTCTGCC-3' and 1537r 5'-TTATGATCCTGCTAATGGTTC-3' designed by Baker *et al.* (1995). PCR amplification was carried out in 200 µl microfuge tubes, using 10 ng DNA, 5 pmol of each primer, 0.2 mM of each dNTP, 2 mM MgCl₂, and 1 U of Taq Polymerase (MBI Fermentas, USA). The amplification conditions used were: 94 °C denaturing for 3 min, followed by 30 cycles of 94 °C for 30 s, 54 °C for 45 s, and extension at 72 °C for 90 s, with a final extension of 10 min on a thermal cycler (Eppendorf). PCR products were directly cloned into PCR 2.1 vector (Invitrogen). Three clones were sequenced using Big dye terminator chemistry on a ABI PRISM 377 automated sequencer.

Phylogenetic analyses

Seventy-two non-redundant microsporidian 16S SSU-rRNA gene sequences were obtained from GenBank (Table 1) aligned with Clustal-X programme (Thompson *et al.* 1997) and manually edited using the programme GeneDoc (Nicholas and Nicholas, 1997). To identify the best-fit model of nucleotide substitution, MODELTEST programme (Posada and Crandall, 1998) was run as implemented in HYPHY programme (Sergei *et al.* 2005). Based on the output, GTR+G model was selected as the best-fit model. A maximum likelihood (ML) tree was constructed using the programme DNAML of Phylip package (Felsenstein, 1989) version 3.64, using a gamma shape parameter (alpha) value of 0.892351. The sequence of *Amblyospora connecticus* was used as an outgroup.

Structural analyses of 16S SSU rRNA

The secondary structure of 16S SSU-rRNA was predicted using the online software 'mfold' (Zuker, 2003) for 3 accessions, L39111 (*N. bombycis*), AY211392 (*N. spodopterae*) and the new isolate reported here. Structures were predicted using the default parameters and the thermodynamically stable and most conserved structure amongst all possible outputs were selected and compared.

Host range

Insects belonging to different orders, collected from both the agroecosystem and other ecosystems (Table 2), were exposed to 10 μ l of water/spore suspension containing 2.4×10^8 spores per insect by contaminating their diet. The inoculated larvae and adult insects were dissected from 10 to 15 days post-inoculation (p.i.), and the tissues were examined microscopically for the presence of vegetative stages and spores in wet smears as well as in Giemsa-stained specimens.

Pathogenicity

Freshly moulted third-instar larvae of *S. litura*, maintained on castor leaves as a microsporidian-free stock, were inoculated with 10 μ l of spore suspension containing 2.4×10^2 , 2.4×10^3 and 2.4×10^4 spores, as described under 'Life-cycle study'. Thirty larvae were used for each treatment and were maintained individually in plastic Petri dishes at 28 ± 2 °C, light regime of 14L:10D, and mortality was recorded daily up to 20 days p.i.. Larval mortality in each treatment was corrected using Abbott's (1925) formula. The spore dosage required to cause 50% mortality (LD_{50}) was calculated by Probit Analysis (Finney, 1971).

RESULTS

Life-cycle of *Nosema sp.*

Hereafter referred to as NSD (*Nosema sp.* isolated from *S. litura* of Dindigul population).

First merogony and sporogony cycle

The first merogonial stage was observed in midgut epithelial cells at 24 h post-inoculation (p.i.) and was characterized by spherical binucleate meronts measuring 1.88μ m ($1.6-2.3$, ± 0.2 ; $n=10$) in diameter (Fig. 1A). Growth of the meronts was followed by a nuclear division, resulting in quadrinucleate meronts, measuring 5.94μ m ($5.3-6.6$, ± 0.45 ; $n=10$) in diameter (Fig. 1B). These meronts have diffused nuclei and an additional division resulted in an octanucleate form with a diameter of 7.09μ m ($6.6-7.5$, ± 0.34 ; $n=10$). This stage eventually divided into tetra- and binucleate daughter diplokarya (Fig. 1C-E). Division of meronts was by binary fission of tetranucleate forms. After 48 h p.i., stages with 1, 2 or 4 diplokarya were the most common in the midgut, and nuclei were large and irregular and occupied about half of the area of the meronts (Fig. 1F-H). Giemsa stained the cytoplasm of meronts moderately blue and the nuclei purple. We were unable to identify sporonts, sporoblasts and primary spores with light microscopy in the midgut tissues.

Second merogony and sporogony cycle

After 72 h p.i., large multinucleate meronts with small compact horseshoe-like nuclei of uniform size and shape were observed in the fat body tissues (Fig. 1I, J). These forms developed by repeated karyokinesis without immediate cytokinesis, resulting in the production of 8 and 12 nucleated forms. They subsequently gave rise to binucleated daughter meronts which developed into sporonts. In addition, bi- and tetranucleate meronts similar to that of primary merogony cycle were also observed. This probably represents a second phase of the merogony since we did not observe any infection in the fat body tissues at 48 h p.i.. After 96 h p.i., binucleate sporonts measuring 4.84μ m ($4.05-5.84$, ± 0.54 ; $n=10$) in diameter, and with deeply stained compact nuclei that occupied a central position in the cells, were observed (Fig. 1K). Further nuclear and cytoplasmic divisions resulted in tetranucleated sporonts of size 5.59μ m ($4.64-6.52$, ± 0.70 , $n=10$) (Fig. 1L). Subsequently, the tetranucleate sporonts divided into 2 sporoblasts measuring 5.4μ m ($4.52-6.82$, ± 0.45 , $n=10$) in length and 2.07μ m ($1.84-2.34$, ± 0.25 , $n=10$) in width, characterized by the presence of 2 small intensely stained and centrally placed nuclei (Fig. 1M). After 144 h p.i., binucleate sporonts, sporoblasts and immature spores were

Table 1. 16S SSU rRNA sequences of Microsporidia used for phylogenetic analyses

Microsporidian name	Host name	Class: Order	Acc. no.
<i>Amblyospora connecticus</i>	<i>Aedes cantator</i>	Insecta: Diptera	AF025685
<i>Antonospora locustae</i>	Grasshopper (Melanoplus)	Insecta: Orthoptera	AY376351
<i>Brachiola algerae</i>	<i>Anopheles stephensi</i>	Insecta: Diptera	AF069063
<i>Culicosporella lunata</i>	<i>Culex pilosus</i>	Insecta: Diptera	AF027683
<i>Encephalitozoon cuniculi</i> 1	<i>Homo sapiens</i>	Mammalia: Primates	L17072
<i>Encephalitozoon cuniculi</i> 2	<i>Homo sapiens</i>	Mammalia: Primates	L39107
<i>Encephalitozoon cuniculi</i> 3	<i>Mus musculus</i>	Mammalia: Rodentia	X98467
<i>Encephalitozoon cuniculi</i> 4	<i>Canis familiaris</i>	Mammalia: Carnivora	X98469
<i>Encephalitozoon cuniculi</i> 5	<i>Oryctolagus cuniculus</i>	Mammalia: Lagomorpha	Z19563
<i>Encephalitozoon hellem</i> 1	<i>Homo sapiens</i>	Mammalia: Primates	AF118142
<i>Encephalitozoon hellem</i> 2	<i>Homo sapiens</i>	Mammalia: Primates	AF118143
<i>Encephalitozoon hellem</i> 3	<i>Homo sapiens</i>	Mammalia: Primates	AF338365
<i>Encephalitozoon hellem</i> 4	<i>Homo sapiens</i>	Mammalia: Primates	AF338366
<i>Encephalitozoon hellem</i> 5	<i>Homo sapiens</i>	Mammalia: Primates	L39108
<i>Encephalitozoon intestinalis</i> 1	<i>Homo sapiens</i>	Mammalia: Primates	L19567
<i>Encephalitozoon intestinalis</i> 2	<i>Homo sapiens</i>	Mammalia: Primates	L39113
<i>Endoreticulatus bombycis</i>	<i>Bombyx mori</i>	Insecta: Lepidoptera	AY009115
<i>Endoreticulatus schubergi</i>	<i>Homo sapiens</i>	Mammalia: Primates	L39109
<i>Endoreticulatus</i> sp. 1	<i>Bombyx mori</i>	Insecta: Lepidoptera	AF240355
<i>Endoreticulatus</i> sp. 2	<i>Ocinara lida</i>	Insecta: Lepidoptera	AY502944
<i>Endoreticulatus</i> sp. 3	<i>Lymantria dispar</i>	Insecta: Lepidoptera	AY502945
<i>Enterocytozoon bienewisi</i>	<i>Homo sapiens</i>	Mammalia: Primates	L16868
<i>Enterocytozoon salmonis</i>	<i>Oncorhynchus tshawytscha</i>	Actinopterygii: Salmoniforms	U10883
NIK-3 h	<i>Bombyx mori</i>	Insecta: Lepidoptera	AY017212
NIK-4 m	<i>Bombyx mori</i>	Insecta: Lepidoptera	AY017213
<i>Nosema apis</i> 1	<i>Apis mellifera</i>	Insecta: Hymenoptera	U26534
<i>Nosema apis</i> 2	<i>Apis</i> sp.	Insecta: Hymenoptera	X73894
<i>Nosema bombi</i>	<i>Bombus terrestris</i>	Insecta: Hymenoptera	AY008373
<i>Nosema bombycis</i> 1	<i>Bombyx mori</i>	Insecta: Lepidoptera	AB093008
<i>Nosema bombycis</i> 2	<i>Bombyx mori</i>	Insecta: Lepidoptera	AB093009
<i>Nosema bombycis</i> 3	<i>Bombyx mori</i>	Insecta: Lepidoptera	AB093010
<i>Nosema bombycis</i> 4	<i>Spodoptera exigua</i>	Insecta: Lepidoptera	AB093011
<i>Nosema bombycis</i> 5	<i>Bombyx mori</i>	Insecta: Lepidoptera	AB093012
<i>Nosema bombycis</i> 6	<i>Bombyx mori</i>	Insecta: Lepidoptera	AB097401
<i>Nosema bombycis</i> 7	<i>Bombyx mori</i>	Insecta: Lepidoptera	AB125666
<i>Nosema bombycis</i> 8	<i>Bombyx mori</i>	Insecta: Lepidoptera	AF240347
<i>Nosema bombycis</i> 9	<i>Bombyx mori</i>	Insecta: Lepidoptera	AY017210
<i>Nosema bombycis</i> 10	<i>Bombyx mori</i>	Insecta: Lepidoptera	AY017211
<i>Nosema bombycis</i> 11	<i>Helicoverpa armigera</i>	Insecta: Lepidoptera	AY259631
<i>Nosema bombycis</i> 12	<i>Bombyx mori</i>	Insecta: Lepidoptera	L39111
<i>Nosema carpocapsae</i>	<i>Cydia pomonella</i>	Insecta: Lepidoptera	AF426104
<i>Nosema ceranae</i>	<i>Apis cerana</i>	Insecta: Hymenoptera	U26533
<i>Nosema furnacalis</i>	<i>Ostrinia furnacalis</i>	Insecta: Lepidoptera	U26532
<i>Nosema granulosis</i>	<i>Gammarus duebeni</i>	Eumalacostraca: Amphipoda	AJ011833
<i>Nosema oulemae</i>	<i>Oulema melanopus</i>	Insecta: Coleoptera	U27359
<i>Nosema</i> sp. isolate 1	<i>Antheraea mylitta</i>	Insecta: Lepidoptera	AB009977
<i>Nosema</i> sp. isolate 2	Noctuid moth	Insecta: Lepidoptera	AF141130
<i>Nosema</i> sp. isolate 3	<i>Calospilos suspecta</i>	Insecta: Lepidoptera	AF240348
<i>Nosema</i> sp. isolate 4	<i>Bombyx mori</i>	Insecta: Lepidoptera	AF240349
<i>Nosema</i> sp. isolate 5	<i>Bombyx mori</i>	Insecta: Lepidoptera	AF240350
<i>Nosema</i> sp. isolate 6	<i>Bombyx mori</i>	Insecta: Lepidoptera	AF240351
<i>Nosema</i> sp. isolate 7	<i>Phyllobrotica armata</i>	Insecta: Coleoptera	AF240353
<i>Nosema</i> sp. isolate 8	<i>Pieris rapae</i>	Insecta: Lepidoptera	AF240354
<i>Nosema</i> sp. isolate 9	<i>Pieris rapae</i>	Insecta: Lepidoptera	AY383655
<i>Nosema</i> sp. isolate 10	<i>Bombyx mori</i>	Insecta: Lepidoptera	D85501
<i>Nosema spodopterae</i>	<i>Spodoptera litura</i>	Insecta: Lepidoptera	AY211392
<i>Nosema tyriae</i>	<i>Tyria jacobaeae</i>	Insecta: Lepidoptera	AJ012606
<i>Nosema</i> sp.	<i>Pieris rapae</i>	Insecta: Lepidoptera	AF485270
<i>Pleistophora</i> sp.	<i>Bombyx mori</i>	Insecta: Lepidoptera	D85500
<i>Vairimorpha cheracis</i>	<i>Cherax destructor destructor</i>	Malacostraca: Decapoda	AF327408
<i>Vairimorpha imperfecta</i> 1	<i>Plutella xylostella</i>	Insecta: Lepidoptera	AJ131645
<i>Vairimorpha imperfecta</i> 2	<i>Plutella xylostella</i>	Insecta: Lepidoptera	AJ131646
<i>Vairimorpha lymantriae</i> 1	<i>Lymantria dispar</i>	Insecta: Lepidoptera	AF033315
<i>Vairimorpha lymantriae</i> 2	<i>Lymantria dispar</i>	Insecta: Lepidoptera	AF141129
<i>Vairimorpha necatrix</i> 1	<i>Apis cerana</i>	Insecta: Hymenoptera	U11051
<i>Vairimorpha necatrix</i> 2	<i>Pseudaletia unipuncta</i>	Insecta: Lepidoptera	Y00266

Table 1 (Cont.)

Microsporidian name	Host name	Class: Order	Acc. no.
<i>Vairimorpha</i> sp. 1	<i>Solenopsis richteri</i>	Insecta: Hymenoptera	AF031539
<i>Vairimorpha</i> sp. 2	<i>Plutella xylostella</i>	Insecta: Lepidoptera	AF124331
<i>Vairimorpha</i> sp. 3	<i>Bombyx mori</i>	Insecta: Lepidoptera	D85502
<i>Vairimorpha</i> sp. 4	<i>Bombyx mori</i>	Insecta: Lepidoptera	L39114
<i>Vavraia oncoperae</i>	<i>Wiseana</i> sp.	Insecta: Lepidoptera	X74112
<i>Vittaforma corneae</i>	<i>Homo sapiens</i>	Mammalia: Primates	U11046
NSD*	<i>Spodoptera litura</i>	Insecta: Lepidoptera	DQ323510

* The new isolate reported in the present study.

simultaneously observed in the fat body tissues. The cytoplasm of the sporoblasts stained lightly with Giemsa.

Mature spores

At 168 h p.i., sporoblasts were observed in the fat body. The binucleate condition was apparent in immature spores (Fig. 1N), while in the mature environmentally resistant ('environmental') spores nuclei were not distinguishable (Fig. 1O). However, on hydrolysis with 1 M HCl at 60 °C for 1 min, the binucleate state became clear (Fig. 1P). In fresh smears of infected tissues, different stages of spore development were observed. Immature spores were relatively large, with a conspicuous vacuole at one end. After 10 days p.i., environmental spores were observed in large numbers in the cytoplasm of fat body cells. They were highly refractive, ovocylindrical in shape, measuring 3.91 µm (3.8–4.0, ±0.0123, *n*=10) in length and 1.91 µm (1.8–2.0, ±0.0143, *n*=10) in width. Mature spores completely replaced the fat body tissue at 15 days p.i.. The polar filament of the spore was extruded upon rehydration of air-dried spores (Kellen and Lindegren, 1969; Watanabe, 1976) and the mean length of the extruded polar filaments was found to be approximately 90 µm.

Sequencing and phylogenetic analyses of 16S SSU-rRNA gene

The 1231 bp 16S SSU rRNA gene of NSD was sequenced using the primers 18f and 1537r. The sequences of 72 non-redundant microsporidian isolates were downloaded from the GenBank and aligned along with the sequence of NSD and the best-fit model of evolution was identified using MODELTEST, which selected the 'TIM+G' model as the best-fit model of nucleotide substitution based on Akaike Information Criterion (AIC) (Akaike, 1974). However, because this model was not available in the Phylip package, we used the nearest alternative GTR+G model (AIC=33501.762) for phylogenetic analyses. The 50% consensus ML tree

using gamma correction resulted in a single large cluster of *Nosema/Vairimorpha* clades (Fig. 2). The SSU rRNA gene sequence of NSD strongly suggested that this species is closely related to the typical *Nosema* subgroup within the *Nosema/Vairimorpha* clade as identified by Baker *et al.* (1994). The phylogenetic analyses using the sequence data of 72 microsporidia (Table 1) indicated that NSD is closely related to *N. bombycis* and *N. spodopterae*. The sequence of NSD differed by 18 bases from *N. bombycis* (GenBank Accession no. L39111). Altogether there were 2 insertions, 5 deletions, 8 transitions and 3 transversions. NSD also differed by 11 bases from *N. spodopterae* (AY211392), including 2 insertions, 3 deletions and 6 transitions.

Structure analyses of 16S SSU rRNA

Secondary structure prediction tool, mfold was used to obtain the folding patterns of the 16SSU-rRNA sequences. The default parameters were used for obtaining the most feasible structure at a minimum free energy (ΔG value of -260 ± 4). The result indicated that the secondary structures of *N. bombycis* (L39111) and *N. spodopterae* (AY211392) were similar to each other whereas the structure of the NSD differed from those of both *N. bombycis* and *N. spodopterae* (Fig. 3). Such a difference is obvious because of a 3 nucleotide deletion in the new isolate at the position 918.

Host range

Cross-transmission tests to insects of 5 different insect orders (Table 2) showed that NSD was infective to all the 9 lepidopteran species tested namely, *Bombyx mori* (L.), *Chilo infuscatellus* Sn, *Earias vitella* (F.), *Spodoptera exigua* (H.), *Helicoverpa armigera* (H.), *Leucinodes orbonalis* Gn., *Pericallia ricini* (Fab.), *Plutella xylostella* (L.), and *Tricoplusia ni* (Hb.), although lower percentages of some species were infected. *B. mori*, *S. litura*, and *P. ricini* showed 100% infection. However, other insects of Orthoptera, Coleoptera, Hemiptera, and Diptera did not develop any infection.

Table 2. Host range of the new isolate, *Nosema* sp. from *Spodoptera litura*

Order	Insect species	Stage of inoculation 2.4×10^3 spores/insect	Percentage infected (N^*)
Orthoptera	<i>Acrida exaltata</i> (Walk.)	Adult	0.0 (5)
	<i>Ailopus t. thalassinus</i> (Fab.)	Adult	0.0 (3)
	<i>Atractomorpha crenulata</i> (Fab.)	Adult	0.0 (10)
	<i>Cyrtacanthacris t. tartarica</i> (Linn.)	Adult	0.0 (4)
	<i>Diabolocantantops pinguis</i> (Walk.)	Adult	0.0 (5)
	<i>Epistaurus sinetyi</i> Bol.	Adult	0.0 (4)
	<i>Eyprepocnemis a. alacris</i> (Serv.)	Adult	0.0 (10)
	<i>Heteracris pulcher</i> (Bol.)	Adult	0.0 (3)
	<i>Orthacris maindroni</i> Bol.	Adult	0.0 (5)
	<i>Oxya fuscovittata</i> (Marsh.)	Adult	0.0 (10)
	<i>Oxya nitidula</i> (Walk.)	Adult	0.0 (10)
	<i>Spathosternum p. prasiniferum</i> (Walk.)	Adult	0.0 (5)
	<i>Trilophidia annulata</i> (Thunb.)	Adult	0.0 (5)
	<i>Tristria pulvinata</i> (Uv.)	Adult	0.0 (3)
	<i>Poecilocerus pictus</i> (Fab.)	Adult	0.0 (5)
	Coleoptera	<i>Rhizopertha dominica</i> (F)	Adult
<i>Tribolium castaneum</i> (Herbst)		Adult	0.0 (15)
<i>Sitophilus oryzae</i> (L)		Adult	0.0 (10)
<i>Atactogaster finitimus</i> Fst		Adult	0.0 (5)
<i>Mylabris pustulata</i> (Thunp.)		Adult	0.0 (3)
Hemiptera	<i>Pyrilla perpusilla</i> Wlk.	Adult	0.0 (4)
	<i>Nephotettix nigropictus</i> Stal	Adult	0.0 (7)
	<i>Nilaparvata lugens</i> Stal.	Adult	0.0 (10)
	<i>Spilostethus hospes</i> (Fab.)	Adult	0.0 (5)
	<i>Spilostethus pandurus</i> (Scopoli)	Adult	0.0 (5)
	<i>Dysdercus cingulatus</i> (F.)	Adult	0.0 (5)
Lepidoptera	<i>Bombyx mori</i> (L.)	Larva	100 (10)
	<i>Chilo infuscatellus</i> Sn.	Larva	57.1 (7)
	<i>Earias vitella</i> (F.)	Larva	60.0 (10)
	<i>Helicoverpa armigera</i> (Hubner)	Larva	88.8 (9)
	<i>Leucinodes orbonalis</i> Gn.	Larva	80.0 (9)
	<i>Pericallia ricini</i> (Fab.)	Larva	100 (10)
	<i>Plutella xylostella</i> (L)	Larva	40.0 (10)
	<i>Spodoptera litura</i> (Fab)	Larva	100 (30)
	<i>Spodoptera exigua</i> (H.)	Larva	++
	<i>Trichoplusia ni</i> (Hb.)	Larva	++
Diptera	<i>Anopheles</i> sp.	Larva	0.0 (25)

* Number of insects tested; ++ Known to cause disease (Solter, personnel communication).

Pathogenicity

The inoculation of NSD spores to third-instar larvae at 3 different concentrations of 2.4×10^2 , 2.4×10^3 , and 2.4×10^4 spores/larva resulted in larval mortality rates of 13.33%, 46.66%, and 73.33% respectively. The LD_{50} was 3885 spores/larva and the 95% limit lies between 2558.51–6110.06 spores/larva.

DISCUSSION

The microsporidium isolated from *S. litura* from the Dindigul, Tamil Nadu State, India is an obligatory intracellular parasite, multiplying in host cells in the form of small binucleate meronts and sporonts. Heavy infection resulted in change of larval colour to pink, frequently accompanied by swelling caused by hypertrophy of infected fat body cells. Occasionally, the body of the insect appeared deformed. The

spores were found in the faecal material of the infected larvae, which probably provide the primary means of transmission to the healthy individuals due to contamination of the food source. Though cannibalistic behaviour is uncommon in *S. litura*, the parasitized larvae feed on the dead diseased larvae in the laboratory, suggesting that transmission of disease could also be due to cannibalism. Although experiments were not conducted to establish the transovarial transmission, presence of spores in ovarian tissues suggested that the parasite is probably incorporated into the eggs as most of the lepidopteran *Nosema*, for example *N. pyraustae* (Andreadis, 1987), *N. heliothidis* (Brooks, 1968) *N. bombycis* (Hatakeyama and Hayasaka, 2002) and also *Vairimorpha* pathogenic to lepidopterans (Haque, Canning and Wright, 1999; Nageswara Rao *et al.* 2004). The new microsporidian isolate appears to be apansporoblastic since the spores were in direct




Fig. 1. Giemsa-stained light micrograph of merogonic and sporogonic stages of *Nosema* sp. in *Spodoptera litura*. Scale bar on A applies also to B–P. (A) Early diplokaryotic meront. (B,C) Diplokaryotic meronts with 2 and 4 nuclei. (D) Dividing octanucleate meront undergoing cytoplasmic division. (E) Meront with 6 nuclei. (F) Binucleate meront. (G, H) Dividing meronts. (I) Octanucleate meront with horseshoe-like diplokaryotic nuclei. (J) Meront with 12 horseshoe-like diplokaryotic nuclei. (K) Binucleated sporont. (L) Tetranucleate sporont. (M) Early sporoblast. (N) Sporoblast (arrow). (O) Mature spores. (P) Spores after treatment with 0.1N HCl showing diplokarya.

contact with the host cell cytoplasm possessing no sporophorous vesicle. All developmental stages were observed to be diplokaryotic. NSD shares all the salient features of the genus *Nosema* namely, apansporoblastic, and diplokaryotic developmental stages.

Sprague and Vernick (1971) characterized *Nosema* as having sporonts that divide into 2 sporoblasts and

distinguished *Nosema* from the genera *Glugea* and *Encephalitozoon* by their nuclear arrangement. In the present study, the sporonts were binucleate and developed into 2 sporoblasts. Our observations show the NSD to be very similar in its development to *N. bombycis*, as described by Ishihara (1969), and therefore we placed it in the genus *Nosema*. The




Fig. 2. The 50% majority consensus maximum likelihood tree of 73 non-redundant microsporidian 16S SSU rRNA gene sequences. *Amblyospora connecticus* was included as an outgroup. Numbers on the nodes indicate the number of times a particular branch was recorded per 100 bootstrap replications following 1000 replicates. The new *Nosema* sp. isolate (NSD) is shown in bold.

frequent occurrence of multinucleate meronts with 12 nuclei, however, easily distinguished NSD from *N. bombycis*.

A few reports have described the microsporidia from *S. litura*. Watanabe (1976) reported a *Nosema* sp. (NSW) from *S. litura* with development of




Fig. 3. Comparison of the 16S SSU rRNA structure of (A) *Nosema* sp. from *Spodoptera litura* (NSD) (B) *Nosema bombycis* (L39111) and (C) *Nosema spodopterae*. Only a portion of the structure (875–950) corresponding to the deleted region in NSD is shown in the figure. For all the species, the most conserved structure with similar minimum free energy ($\Delta G = -260 \pm 4$ Kcal/mol) was used.

binucleated meronts very similar to NSD. But the monosporous nature (sporont developed into single sporoblast) and uninucleate meronts of NSW distinctly differentiate it from NSD. Hsu, Hsu and Yen (1991, 1992), isolated a microsporidian species from *S. litura* in Taiwan and named it as *Nosema spodopterae* since they observed dimorphism of sporoblasts in their transmission electron microscopic study. The authors reported that one type of sporoblast possessed a plasmalemma inside of plasma membrane, and the other appeared as an amoeba-like shape without a plasmalemma within the spores. In general, the plasma membrane and plasmalemma are two names of the same structure. The internal layer of the spore wall is always referred to as a plasma membrane. The sporoblast derives its plasma membrane from the membranes of the polaroplast during the process of eversion through the polar filament during germination. No microsporidia have an amoeba-like stage in the spores. The sporoblast is the stage most difficult to fix properly for electron microscopy and electron microscopy fixation often destroys the shape of the sporoblast (Larsson, personal communication) leading to a star-like form in fixed tissues (Sokalova and Lange, 2002). Hence the

two kinds of sporoblasts reported by Hsu *et al.* (1992) in *N. spodopterae* are probably an artifact arising due to fixation. Though the size of NSD sporoblasts ranges from 4.52 to $6.52 \mu\text{m}$, the shape of the sporoblasts did not vary. The fresh spore size of NSD ($3.91 \pm 0.012 \times 1.91 \pm 0.014 \mu\text{m}$) was similar to that of *N. spodopterae* ($4.0 \pm 0.26 \times 1.9 \pm 0.12 \mu\text{m}$). Li and Wenn (1987) described a microsporidium, *Nosema liturae*, obtained from *S. litura* collected from China. The fresh spore size of *N. liturae* ($4.34 \pm 0.27 \times 1.99 \pm 0.14 \mu\text{m}$) exceeds the size of NSD ($3.91 \pm 0.012 \times 1.91 \pm 0.014 \mu\text{m}$). *Bombyx mori* is refractive to *N. liturae*, while the NSD is highly pathogenic to the same host species. The multinucleated meronts with up to 12 compact horseshoe-like nuclei reported in NSD were not reported in *N. spodopterae* and *N. liturae*.

Since many attempts to amplify large subunit rDNA did not succeed across all microsporidian species (Vossbrinck *et al.* 1993; Kent *et al.* 1996), 16S SSU rDNA has become the standard sequence for molecular characterization of microsporidia (Weiss and Vossbrinck, 1999; Vossbrinck and Debrunner-Vossbrinck, 2005). Here we aligned the SSU rRNA gene sequence of NSD with those of other known

microsporidia that infect lepidopteran as well as non-lepidopteran hosts. The results lead us to confirm that NSD is the member of true *Nosema* namely, *N. bombycis* complex and belongs to the genus *Nosema* Nageli, 1857.

Comparison of the 16S rRNA gene sequence of NSD with the corresponding sequence from several *Nosema* species showed that the parasite is closer to *N. bombycis*. We used *N. bombycis* 'L39111' as a reference strain in the present study since it is the most studied isolate for phylogenetic analyses (Baker *et al.* 1995; Canning *et al.* 2002; Choi *et al.* 2002; Vossbrinck and Debrunner-Vossbrinck, 2005). In NSD, the absence of 3 nucleotides at position 918 as compared to L39111 and other *N. bombycis* suggested a possible difference between these two isolates. The published sequence of *N. spodopterae* is, in fact, closer to that of L39111 than to NSD, suggesting that the new isolate is possibly a new species of *Nosema* from *S. litura*. Consistently, the secondary structure of *N. bombycis* and *N. spodopterae* 16S SSU rRNA was identical (Huang *et al.* 2004; Tsai, Huang and Wang, 2005) whereas the structure of NSD was different. The significance of these structural differences needs to be experimentally verified, which is out of purview of the present manuscript. These observations compel us to consider that the microsporidian isolate NSD as a new species, confirmation of which awaits further ultrastructural characterization.

The pathogenicity of NSD appears to be proportional to the inoculation dosage. The LD₅₀ for larvae inoculated in third-instar was calculated to be 3.88×10^3 spores/larva at 20 days p.i. and is comparable with the published reports of other *Nosema* isolates from *S. litura*. Tsai and Wang (2001) obtained an LD₅₀ of 1.13×10^4 spores/larva after 21 days when *S. litura* third-instar larvae were treated with *N. spodopterae* spores. The LD₅₀ of Taiwan isolate in third-instar *S. litura* larvae was lower, 1.78×10^3 spores/larvae at 30 days p.i. (Tsai *et al.* 2003).

Host specificity is an important consideration for pathogens that are used for biological control. Microsporidia usually do not infect species in other insect orders when larvae are fed spores, but inter-ordinal infections have been reported in laboratory infectivity studies (Fantham and Porter, 1958; Undeen and Maddox, 1973). All the lepidopteran hosts tested were susceptible to NSD indicating that this species appear to have a wider host range under laboratory conditions. Previous host specificity studies under lab conditions (Undeen and Maddox, 1973; Solter, Maddox and McManus, 1997; Solter and Maddox, 1998) suggest that terrestrial microsporidia have a broader laboratory host range within the taxonomic order of the natural host, but atypical microsporidia/host interactions suggested that the pathogens often do not reproduce optimally in

non-target hosts, and data from surveys of microsporidian pathogens of various native and introduced insect hosts from field collections suggest that the ecological host ranges of entomopathogenic microsporidia are relatively narrow (Campbell and Podgwaite, 1971; Andreadis *et al.* 1983; Siegal *et al.* 1988; Jeffords *et al.* 1989). In the present study, lower percentages of infection rate observed in some lepidopteran species like *P. xylostella* and *C. infuscatellus* also reflected that these may be non-target hosts to NSD under field conditions. Hence, the use of NSD as a biological control agent against *S. litura* should be investigated.

Financial support to J.N. by the Department of Biotechnology, New Delhi under the project 'Molecular Ecology and Systematics (Grant no. BT/012/01/NBDB/2004)' is greatly acknowledged. We thank Dr Leellen F. Solter, University of Illinois, Urbana, USA, for her valuable help in life-cycle studies.

REFERENCES

- Abbott, W. S.** (1925). A method for computing the effectiveness of an insecticide. *Journal of Economic Entomology* **18**, 265–267.
- Akaike, H.** (1974). A new look at the statistical model identification. *IEEE Transactions on Automatic Control* **19**, 716–723.
- Andreadis, T. G.** (1987). Horizontal transmission of *Nosema pyrausta* (Microsporidia: Nosematidae) in the European corn borer, *Ostrinia nubilalis* (Lepidoptera: Pyralidae). *Environmental Entomology* **16**, 1124–1129.
- Andreadis, T. G., Dubois, N. R., Moore, R. E. B., Anderson, J. F. and Lewis, F. B.** (1983). Single applications of high concentrations of *Bacillus thuringiensis* for control of gypsy moth (Lepidoptera: Lymantriidae) populations and their impact on parasitism and disease. *Journal of Economic Entomology* **76**, 1417–1422.
- Armes, N. J., Wightman, J. A., Jadhav, D. R. and Ranga Rao, G. V.** (1997). Status of insecticide resistance in *Spodoptera litura* in Andhra Pradesh, India. *Pesticide Science* **50**, 240–248.
- Baker, M. D., Vossbrinck, C. R., Didier, E. S., Maddox, J. V. and Shaddock, J. A.** (1995). Small subunit ribosomal DNA phylogeny of various microsporidia with emphasis on AIDS related forms. *Journal of Eukaryotic Microbiology* **42**, 564–570.
- Baker, M. D., Vossbrinck, C. R., Maddox, J. V. and Undeen, A. H.** (1994). Phylogenetic relationships among *Vairimorpha* and *Nosema* species (Microspora) based on ribosomal RNA sequence data. *Journal of Invertebrate Pathology* **64**, 100–106.
- Becnel, J. J. and Andreadis, T. G.** (1999). Microsporidia in insects. In *The Microsporidia and Microsporidiosis* (ed. Wittner, M. and Weiss, L. M.), pp. 447–501. American Society for Microbiology Press, Washington, D.C.
- Briano, J. A. and Williams, D. F.** (2002). Natural occurrence and laboratory studies of the fire ant pathogen *Vairimorpha invictae* (Microsporida: Burenellidae) in Argentina. *Environmental Entomology* **31**, 887–894.

- Brooks, W. M.** (1968). Transovarian transmission of *Nosema heliothidis* in the corn earworm, *Heliothis zea*. *Journal of Invertebrate Pathology* **11**, 510–512.
- Campbell, R. W. and Podgwaite, J. D.** (1971). The disease complex of the gypsy moth. I. Major components. *Journal of Invertebrate Pathology* **18**, 101–107.
- Canning, E. U., Refardt, D., Vossbrinck, C. R., Okamura, B. and Curry, A.** (2002). New diplokaryotic microsporidia (Phylum Microsporidia) from freshwater bryozoans (Bryozoa, Phylactolaemata). *European Journal of Protistology* **38**, 247–265.
- Choi, J. Y., Kim, J. G., Choi, Y. C., Goo, T. W., Chang, J. H., Je, Y. H. and Kim, K. Y.** (2002). *Nosema* sp. isolated from cabbage white butterfly (*Pieris rapae*) Collected in Korea. *Journal of Microbiology* **40**, 199–204.
- Desportes, I., Le Charpentier, Y. and Galian, A.** (1985). Occurrence of a new microsporidian: *Enterocytozoon bieneusi* n. g., n. sp. in the enterocytes of a human patient with AIDS. *Journal of Protozoology* **32**, 250–254.
- Didier, E. S., Didier, P. J., Snowden, K. F. and Shadduck, J. A.** (2000). Microsporidiosis in mammals. *Microbes and Infection* **2**, 709–720.
- Fantham, H. B. and Porter, A.** (1958). Some pathogenic bacteriform microsporidia from Crustacea and Insecta. *Proceedings of the Zoological Society of London, B* **130**, 153–168.
- Felsenstein, J.** (1989). PHYLIP – Phylogeny Inference Package (Version 3.2). *Cladistics* **5**, 164–166.
- Finney, D. J.** (1971). *Probit Analysis*, 3rd Edn. Cambridge University Press, Cambridge.
- Franzen, C. and Muller, A.** (1999). Molecular techniques for detection, species differentiation, and phylogenetic analyses of microsporidia. *Clinical Microbiology Reviews* **12**, 243–285.
- Hatakeyama, Y. and Hayasaka, S.** (2002). Specific amplification of microsporidian DNA fragments using multiprimer PCR. *Japan Agricultural Research Quarterly* **36**, 97–102.
- Haque, M. A., Canning, E. U. and Wright, D. J.** (1999). Entomopathogenicity of *Vairimorpha* sp. (Microsporidia) in the diamondback moth, *Plutella xylostella* (Lepidoptera: Yponomeutidae). *Bulletin of Entomological Research* **89**, 147–152.
- Hsu, T. H., Hsu, E. L. and Yen, D. F.** (1991). Morphogenesis of microsporidian *Nosema* sp. from the tobacco cutworm *Spodoptera litura*. *Plant Protection Bulletin, Taichung* **33**, 395–409.
- Hsu, T. H., Hsu, E. L. and Yen, D. F.** (1992). *Nosema spodopterae* n. sp. a new species of microsporidia from the tobacco cutworm *Spodoptera litura*. *Journal of the Agricultural Association of China* **157**, 81–90.
- Huang, H. W., Tsai, S. J., Lo, C. F., Soichi, Y. and Wang, C. H.** (2004). The novel organization and complete sequence of the ribosomal gene of *Nosema bombycis*. *Fungal Genetics and Biology* **41**, 473–481.
- Ishihara, R.** (1969). The life cycle of *Nosema bombycis* as revealed in tissue culture cells of *Bombyx mori*. *Journal of Invertebrate Pathology* **14**, 316–320.
- Jeffords, M. R., Maddox, J. V., McManus, M. L., Webb, R. E. and Wieber, A.** (1989). Evaluation of the overwintering success of two European microsporidia inoculatively released into gypsy moth populations in Maryland, USA. *Journal of Invertebrate Pathology* **53**, 235–240.
- Johny, S.** (2002). Resistance to insecticides among different populations of *Spodoptera litura* (Fab.) (Lepidoptera: Noctuidae) and their control with a new microsporidian parasite. Ph.D. thesis, University of Madras, Chennai, India.
- Johny, S. and Muralirangan, M. C.** (2000). Monitoring susceptibility to selected insecticides in *Spodoptera litura* (Fabricius) (Lepidoptera: Noctuidae) in Tamil Nadu (India). *Pestology* **24**, 32–36.
- Keeling, P. J.** (2003). Congruent evidence from a α - tubulin and β - tubulin gene phylogenies for a zygomycete origin of microsporidia. *Fungal Genetics and Biology* **38**, 298–309.
- Keeling, P. J. and Fast, N. M.** (2002). Microsporidia: biology and evolution of highly reduced intracellular parasites. *Annual Review of Microbiology* **56**, 93–116.
- Kellen, W. R. and Lindgren, J. E.** (1969). Host-pathogen relationships of two previously undescribed microsporidian from the Indian-meal moth, *Plodia interpunctella* (Hübner), (Lepidoptera: Phycitidae). *Journal of Invertebrate Pathology* **14**, 328–335.
- Kent, M. L., Hervio, D. M., Docker, M. F. and Devlin, R. H.** (1996). Taxonomy studies and diagnostic tests for myxosporean and microsporidian pathogens of salmonid fishes utilising ribosomal DNA sequence. *Journal of Eukaryotic Microbiology* **43**, 98S–99S.
- Kotler, D. P. and Orenstein, J. M.** (1999). Clinical syndromes associated with microsporidiosis. In *The Microsporidia and Microsporidiosis* (ed. Wittner, M. and Weiss, L. M.), pp. 258–292. American Society for Microbiology Press, Washington, D.C.
- Li, S. P. and Wenn, C. S.** (1987). *Nosema liturae* sp. n., a new species of microsporidia from the cotton leafworm, *Spodoptera litura*. *Scientia Agricultura Sinica* **20**, 71–74.
- Lockwood, J. A., Bomar, C. R. and Ewen, A. B.** (1999). The history of biological control with *Nosema locustae*: lessons for locust management. *Insect Science and its Application* **19**, 333–350.
- Molina, J. M., Goguel, J., Sarfati, C., Michiels, J. F., Desportes-Livage, I., Balkan, S., Chastang, C., Cotte, L., Maslo, C., Struxiano, A., Derouin, F. and Decazes, J. M.** (2000). Trial of oral fumagillin for the treatment of intestinal microsporidiosis in patients with HIV infection. ANRS 054 Study Group. Agence Nationale de Recherche sur le SIDA. *AIDS* **14**, 1341–1348.
- Muller, M., Bialek, R., Kamper, A., Fatkenheuer, G., Salzberger, B. and Franzen, C.** (2001). Detection of Microsporidia in Travelers with Diarrhea. *Journal of Clinical Microbiology* **39**, 1630–1632.
- Nageli, C.** (1857) *Nosema bombycis* Nageli. *Botanische Zeitung* **15**, 760–761.
- Nageswara Rao, S., Muthulakshmi, M., Kanginakudru, S. and Nagaraju, J.** (2004). Phylogenetic relationships of three new microsporidian isolates from the silkworm, *Bombyx mori*. *Journal of Invertebrate Pathology* **86**, 87–95.
- Nicholas, K. B. and Nicholas, H. B. Jr.** (1997). GeneDoc: a tool for editing and annotating multiple sequence alignments. Distributed by the author, available at: <http://www.psc.edu/biomed/genedoc>

- Posada, D. and Crandall, K. A.** (1998). Modeltest: testing the model of DNA substitution. *Bioinformatics* **14**, 817–818.
- Sergei, L., Pond, S. L. K., Frost, S. D. W. and Muse, S. V.** (2005). *HyPhy: hypothesis testing using phylogenies*. *Bioinformatics. Advance Access published on March 1, 2005*, *Bioinformatics* **21**, 676–679.
- Siegel, J., Maddox, J. V. and Ruesink, W. G.** (1988). Seasonal progress of *Nosema pyrausta* in the European corn borer, *Ostrinia nubilalis*. *Journal of Invertebrate Pathology* **52**, 130–136.
- Slamovits, C. H., Williams, B. A. and Keeling, P. J.** (2004). Transfer of *Nosema locustae* (Microsporidia) to *Antonospora locustae* n. comb. based on molecular and ultrastructural data. *Journal of Eukaryotic Microbiology* **51**, 207–13.
- Sokolova, Y. Y., Issi, I. V., Morzhina, E. V., Tokarev, Y. S. and Vossbrinck, C. R.** (2005). Ultrastructural analysis supports transferring *Nosema whitei* Weiser 1953 to the genus *Paranosema* and creation a new combination, *Paranosema whitei*. *Journal of Invertebrate Pathology* **90**, 122–126.
- Sokolova, Y. Y. and Lange, C. E.** (2002). An ultrastructural study of *Nosema locustae* Canning (Microsporidia) from three species of acrididae (Orthoptera). *Acta Protozoologica* **41**, 229–337.
- Solter, L. F. and Becnel, J. J.** (2000). Entomopathogenic microsporidia. In *Field Manual of Techniques for the Evaluation of Entomopathogens* (ed. Lacey, L. A. and Kaya, H.), pp. 231–254. Kluwer Academic Publishers, Dordrecht, The Netherlands.
- Solter, L. F. and Maddox, J. V.** (1998). Physiological host specificity of microsporidia as an indicator of ecological host specificity. *Journal of Invertebrate Pathology* **71**, 207–216.
- Solter, L. F., Maddox, J. V. and McManus, M. L.** (1997). Host specificity of microsporidia (Protista: Microspora) from European populations of *Lymantria dispar* (Lepidoptera: Lymantriidae) to indigenous North American Lepidoptera. *Journal of Invertebrate Pathology* **69**, 139–150.
- Sprague, V. and Vernick, S. H.** (1971). The ultrastructure of *Encephalitozoon cuniculi* (Microsporidia: Nosematidae) and its taxonomic significance. *Journal of Protozoology* **18**, 566–569.
- Thompson, J. D., Gibson, T. J., Plewniak, F., Jeanmougin, F. and Higgins, D. G.** (1997). The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Research* **25**, 4876–4882.
- Tsai, S. J., Huang, W. F. and Wang, C. H.** (2005). Complete sequence and gene organization of the *Nosema spodopterae* rRNA gene. *Journal of Eukaryotic Microbiology* **52**, 52–54.
- Tsai, S. J., Lo, C. F. and Wang, C. H.** (2003). The characterization of microsporidian isolates (Nosematidae: *Nosema*) from five important lepidopteran pests in Taiwan. *Journal of Invertebrate Pathology* **83**, 51–59.
- Tsai, S. J. and Wang, C. H.** (2001). Interaction of the microsporidium and nucleopolyhedrovirus in *Spodoptera litura*. *Formosan Entomologist* **21**, 183–195.
- Undeen, A. H. and Cockburn, A. F.** (1989). The extraction of DNA from microsporidia spores. *Journal of Invertebrate Pathology* **54**, 132–133.
- Undeen, A. H. and Maddox, J. V.** (1973). The infection of nonmosquito hosts by injection with spores of the microsporidian *Nosema algerae*. *Journal of Invertebrate Pathology* **22**, 258–265.
- Undeen, A. H. and Vavra, J.** (1997). Research methods for entomopathogenic protozoa. In *Manual of Techniques in Insect Pathology* (ed. Lacey, L. A.), pp. 117–151. Academic Press, San Diego, USA.
- Van de Peer, Y., Caers, A., De Rijk, P. and De Wachter, R.** (1998). Database on the structure of small ribosomal subunit RNA. *Nucleic Acids Research* **26**, 179–182.
- Vossbrinck, C. R., Baker, M. D., Didier, E. S., Debrunner-Vossbrinck, B. A. and Shadduck, J. A.** (1993). Ribosomal DNA sequences of *Encephalitozoon hellem* and *Encephalitozoon cuniculi*: species identification and phylogenetic construction. *Journal of Eukaryotic Microbiology* **40**, 354–362.
- Vossbrinck, C. R. and Debrunner-Vossbrinck, B. A.** (2005). Molecular phylogeny of the Microsporidia: ecological, ultrastructural and taxonomic considerations. *Folia Parasitologica* **52**, 131–142.
- Watanabe, H.** (1976). A *Nosema* species of the Egyptian cotton leaf worm *Spodoptera litura* (Lepidoptera): its morphology, development, host range, and taxonomy. *Journal of Invertebrate Pathology* **28**, 321–328.
- Weber, R., Bryan, R. T., Schwartz, D. A. and Owen, R. L.** (1994). Human microsporidial infections. *Clinical Microbiology Reviews* **7**, 426–461.
- Weiss, L. M. and Vossbrinck, C. R.** (1999). Molecular biology, molecular phylogeny, and molecular diagnostic approaches to the Microsporidia. In *The Microsporidia and Microsporidiosis* (ed. Wittner, M. and Weiss, L. M.), pp. 129–171. American Society for Microbiology Press, Washington, D.C.
- Wittner, M. and Weiss, L. M.** (1999). *The Microsporidia and Microsporidiosis*. American Society for Microbiology Press, Washington, D.C.
- Zuker, M.** (2003). Mfold web server for nucleic acid folding and hybridization prediction. *Nucleic Acids Research* **31**, 3406–3415.