

Glycine Supports *in Vivo* Reduction of Nitrate in Barley Leaves

Received for publication May 17, 1988 and in revised form August 15, 1988

POLUMETLA A. KUMAR, T. V. R. NAIR, AND YASH P. ABROL*

Nuclear Research Laboratory and Division of Plant Physiology, Indian Agricultural Research Institute, New Delhi 110 012, India

ABSTRACT

Glycine, a photorespiratory intermediate, enhanced the *in vivo* reduction of nitrate in barley (*Hordeum vulgare* L.) leaf slices, when included in the assay medium. Isonicotinyl hydrazide, an inhibitor of glycine oxidation, partially reduced NO_2^- production. The enhancement caused by glycine treatment was reversed by isonicotinyl hydrazide when both were present together in the medium. Similar effects were observed when the excised leaves were preincubated with the metabolite and the inhibitor. Glycine also partially relieved the inhibition of nitrate reduction caused by malonate, an inhibitor of the tricarboxylic acid cycle. The results support the hypothesis that glycine decarboxylation activity is a source of NADH for nitrate reductase activity.

done. Wherever preincubation was involved, the excised leaves were incubated with different treatment solutions for 3 h under light ($800 \mu\text{E m}^{-2} \text{S}^{-1}$) at 30°C .

RESULTS AND DISCUSSION

Glycine enhanced nitrate reduction in barley leaves when included in the *in vivo* assay medium (Fig. 1). The maximum enhancement was attained at 10 mM concentration. Three possible explanations could be attributed to explain this enhancement: a. possible induction of NR by the amino acid; b. increased stability of the enzyme in the presence of ammonium (13) released during glycine oxidation; c. increased supply of reducing power to the enzyme. It has been observed by many workers that amino acids either repress the synthesis or reduce the activity of NR (10). We observed no considerable change in NR activity by using a wide range of amino acids other than glycine in the

NR^1 (EC 1.6.6.1) reduces nitrate to nitrite utilizing NADH as reducing power (2). Earlier studies by Klepper *et al.* (8) have shown that NADH for nitrate reduction in leaves is derived from photosynthetically produced glyceraldehyde 3-P, which after migrating from chloroplasts, generates NADH in the cytoplasm on being oxidized to PGA by triosephosphate dehydrogenase. On the other hand, Sawhney *et al.* (11) have also demonstrated that NADH generated in the TCA cycle can be used for the reduction of nitrate. A third source of NADH could be the process of glycine oxidation in mitochondria. Green leaves of C_3 plants are known to synthesize substantial quantities of glycine during photorespiration (15). Two molecules of glycine form one molecule each of serine, CO_2 and NH_3 with concomitant reduction of one molecule of NAD^+ by the action of glycine decarboxylase complex in the mitochondria (7). The turnover of glycine in glycolate pathway is quite rapid, often attaining rates of about $80 \mu\text{mol}$ of serine formed per g tissue per h (16). In this report we show that glycine can act as a source of reducing power for nitrate reduction in leaves.

MATERIALS AND METHODS

Barley plants (*Hordeum vulgare* L. cv DL-157) grown in sandy loam soil in pots were used. Third and fourth leaves from 15 to 20 d old plants were selected. Nitrate reductase *in vivo* was assayed according to Klepper *et al.* (8) with slight modification (6). In experiments where various metabolites and inhibitors were included in the assay medium, vacuum infiltration was




FIG. 1. Changes in NR activity (% of control) caused by glycine and INH when included in the preincubation medium (---) or in the assay medium (—). Assay medium consisted of 0.1 M phosphate buffer (pH 6.0) + 0.2 M KNO_3 .

¹ Abbreviations; NR, Nitrate reductase; PGA, glycerate 3-phosphate; INH, isonicotinylhydrazide; α -HPMS, α -hydroxy-2-pyridine methane sulfonic acid.

Table I. NR Activity in Vivo in Barley Leaf Slices as Affected by Glycine

Preincubation medium in the case of control was distilled water. Assay medium consisted of 0.1 M phosphate buffer (pH 6.0) with 0.2 M KNO₃.

Treatment	NR Activity			
	Preincubation medium		Assay medium	
	$\mu\text{mol NO}_2^- \text{ g}^{-1} \text{ fresh wt h}^{-1}$	% of control	$\mu\text{mol NO}_2^- \text{ g}^{-1} \text{ fresh wt h}^{-1}$	% of control
Control	2.57	100	2.43	100
INH, 35 mM	1.19	46	1.22	60
Glycine, 10 mM	3.21	125	3.02	124
Glycine 10 mM + INH, 35 mM	1.58	61	1.57	64

Table II. NR Activity in Vivo in Barley Leaf Slices as Affected by Glycolate and α -HPMS

Glycolate and α -HPMS were included in the assay medium consisting of 0.1 M phosphate buffer (pH 6.0) and 0.2 M KNO₃.

Treatment	NR Activity	
	$\mu\text{mol NO}_2^- \text{ g}^{-1} \text{ fresh wt h}^{-1}$	% of control
Control	2.67	100
α -HPMS, 10 mM	0.86	33
Glycolate, 10 mM	2.97	116
Glycolate, 10 mM + α -HPMS, 10 mM	1.39	54

Table III. NR Activity in Vivo in Fresh and Dark (24 h) Treated Barley Leaves as Affected by Malonate, Glycine and INH

Malonate (5 mM), glycine (10 mM), and INH (35 mM) were included in the assay medium consisting of 0.1 M phosphate buffer (pH 6.0) and 0.2 M KNO₃.

Treatment	NR Activity			
	Fresh leaves		Dark-treated leaves	
	$\mu\text{mol NO}_2^- \text{ g}^{-1} \text{ fresh wt h}^{-1}$	% of control	$\mu\text{mol NO}_2^- \text{ g}^{-1} \text{ fresh wt h}^{-1}$	% of control
Control	2.47	100	1.16	100
Malonate	1.03	42	1.07	92
Malonate + glycine	1.90	77	1.73	149
Malonate + glycine + INH	0.52	21	0.55	47

assay medium (results not shown). Second, free ammonium content in the leaves does not alter considerably even after incubation of leaves with glycine (9). This is due to the presence of an efficient system of reamination of photorespiratory ammonia by glutamine synthetase (14). The third possibility has been examined by using INH, a potent inhibitor of glycine oxidation. Figure 1 shows the effect of INH on NR activity. There has been a 40 to 50% reduction in the enzyme activity by either mode of treatment. Table I shows further that the enhancement of NR activity by glycine is effectively reversed by INH when it is included along with glycine. This clearly shows that glycine oxidation is responsible for the increase in NR activity. Similar results have been obtained using leaves kept in dark for 24 h to deplete the carbohydrate reserves (data not shown).

In another experiment we have provided glycolate (10 mM), a precursor of glycine, to the excised leaves. Glycolate treatment slightly enhanced NR activity (Table II). α -HPMS, an inhibitor of glycolate oxidation, reduced nitrate reduction when provided alone or together with glycolate.

Table III shows the effect of malonate, a competitive inhibitor of succinate dehydrogenase, on nitrate reduction in fresh and 24 h dark treated leaves. Malonate inhibited nitrate reduction in fresh leaves by 60%. This inhibition was substantially reversed by the addition of glycine. Malonate, glycine, and INH together reduced NR activity by 80%. This clearly demonstrates that glycine partially restores the NR activity inhibited by malonate. This restoration is nullified by INH.

In dark treated leaves NR activity persisted though at a reduced level. This activity was resistant to malonate which indicate sources of NADH other than the TCA cycle. The malonate

resistant component was found in fresh leaves as well. Glycine substantially increased nitrate reduction when added with malonate. INH effectively inhibited this enhancement.

It is now well established that in green leaves in light, NADH for nitrate reduction can be provided by two sources *viz.*, metabolism of glyceraldehyde 3-P and oxidation of TCA cycle intermediates (1, 11). In the present investigation we demonstrate that glycine oxidation also supports the *in vivo* reduction of nitrate by providing reducing power. This is supported by the partial inhibition of nitrate reduction by INH and α -HPMS, inhibitors of glycine and glycolate oxidation, respectively. Exogenous supply of glycine to the leaves enhanced nitrate reduction and this enhancement was curtailed by INH. Further evidence is provided by the experiment wherein glycine reversed, though not completely, the inhibition of nitrate reduction by malonate and INH blocked this reversal.

It has been suggested that glycine oxidation is coupled to the mitochondrial electron transport (5). However, high phosphorylation potential which exists in the cell during photosynthesis may be inhibitory to the mitochondrial electron transport (4) thus making NADH available for nitrate reduction (12). Thus, the NADH generated during glycine oxidation can also be available for nitrate reduction in the cytosol. The export of NADH out of mitochondria could be via the malate-oxaloacetate shuttle proposed by Woo *et al.* (17) or via the transmembrane transhydrogenase system present in plant mitochondria (3). Woo *et al.* (18) have demonstrated that glycine can support nitrate reduction in a reconstituted system, but they held the view that glycine oxidation is not linked to nitrate reduction *in vivo*. Our results conclusively prove that glycine oxidation can also support nitrate

reduction *in vivo*, thus making it yet another source of reducing power for NR. This raises the question again whether photorespiration long considered to be a wasteful process is really so.

Acknowledgments—We are grateful to Prof. Ann Oaks for critically reading the manuscript. We thank Mr. D. Gangadhar for helpful discussions.

LITERATURE CITED

1. ABROL YP, SK SAWHNEY, MS NAIK 1983 Light and dark assimilation of nitrate in plants. *Plant Cell Environ* 6: 595–599
2. BEEVERS L, RH HAGEMAN 1980 Nitrate and nitrite reduction. *In* BJ Mifflin, ed, *The Biochemistry of Plants*, Vol 5. Academic Press, New York, pp 115–118
3. DAY DA, JT WISKICH 1978 Pyridine nucleotide interactions with isolated plant mitochondria. *Biochim Biophys Acta* 501: 396–404
4. DOUCE R 1985 Mitochondria in Higher Plants: Structure, Function and Biogenesis. Academic Press, Orlando, FL
5. GARDESTRÖM P, A BERGMAN, I ERICSON 1980 Oxidation of glycine via the respiratory chain in mitochondria prepared from different parts of spinach. *Plant Physiol* 65: 389–391
6. GROVER HL, TVR NAIR, YP ABROL 1978 Nitrogen metabolism of the upper three leaf blades of wheat at different soil nitrogen levels. *Physiol Plant* 42: 287–292
7. KEYS AJ 1980 Synthesis of glycine and serine. *In* BJ Mifflin, ed, *The Biochemistry of Plants*, Vol. 5. Academic Press, New York, pp 359–374
8. KLEPPER L, P FLESHER, RH HAGEMAN 1971 Generation of reduced nicotinamide adenine dinucleotide for nitrate reduction in green leaves. *Plant Physiol* 48: 580–590
9. KUMAR PA, TVR NAIR, YP ABROL 1984 Effect of photorespiratory metabolites, inhibitors and methionine sulphoximine on the accumulation of ammonia in the leaves of mungbean and *Amaranthus*. *Plant Sci Lett* 33: 303–307
10. OAKS A, B HIREL 1985 Nitrogen metabolism in roots. *Annu Rev Plant Physiol* 36: 345–365
11. SAWHNEY SK, MS NAIK, DJD NICHOLAS 1978 Regulation of NADH supply for nitrate reduction in green plants via photosynthesis and mitochondrial respiration. *Biochem Biophys Res Commun* 81: 1209–1216
12. SAWHNEY SK, MS NAIK, DJD NICHOLAS 1978 Regulation of nitrate reduction by light, ATP and mitochondrial respiration in wheat leaves. *Nature* 272: 647–648
13. SIHAG RK, S GUHA-MUKHERJEE, SK SOPORY 1978 Regulation of nitrate reductase level in pea: *in vivo* stability by ammonium. *Biochem Biophys Res Commun* 85: 1017–1024
14. SINGH P, PA KUMAR, YP ABROL, MS NAIK 1985 Photorespiratory nitrogen cycle—a critical evaluation. *Physiol Plant* 66: 169–176
15. TOLBERT NE 1980 Photorespiration. *In* DD Davies, ed. *The Biochemistry of Plants*, Vol 2. Academic Press, New York, pp 487–523
16. WOO KC, CB OSMOND 1976 Glycine decarboxylation in mitochondria isolated from spinach leaves. *Aust J Plant Physiol* 3: 771–785
17. WOO KC, M JOKINEN, DT CANVIN 1980 Nitrate reduction by a dicarboxylate shuttle in a reconstituted system from spinach leaves. *Aust J Plant Physiol* 7: 123–130
18. WOO KC, M JOKINEN, DT CANVIN 1980 Reduction of nitrate via a dicarboxylate shuttle in a reconstituted system of supernatant and mitochondria from spinach leaves. *Plant Physiol* 65: 433–436