

Evolutionary Questions Raised by Cellular Slime Mould Development

SONIA KAUSHIK and VIDYANAND NANJUNDIAH*

Centre for Ecological Sciences, Indian Institute of Science and Jawaharlal Nehru Centre for Advanced Scientific Research, Bangalore 560012

(Received on 9 September 2003; Accepted after revision on 15 October 2003)

The cellular slime moulds (CSMs) are amoeboid organisms whose life cycle can be viewed in two ways. Firstly, because free-living amoebae come together to build bodies, they are ideal models for studying multicellular development in terms of the properties of single cells. Secondly, coming together and participating in an integrated unit implies social behaviour. Consequently differentiation (especially in the advanced CSMs) can be seen as a form of division of labour in which only some amoebae get to transmit their genes to the next generation. Viewed thus, their life cycle is ideally suited for studying the evolutionary basis of cooperation with some members of the cooperating group exhibiting altruistic behaviour. The present review takes the second approach. We examine alternative explanations for social behaviour (i.e., multicellular development) based on individual-level and group (including kin-group) selection. Non-clonal fruiting bodies are likely to be common in nature; we show a case with at least nine genotypes. The CSMs display both individual and group-level adaptations and both levels of selection operate in their appropriate contexts. The review ends with questions for the future and indicates how studies of CSM development might help to explain the evolution of altruism in this group.

Key Words: Evolution of development, *Dictyostelium*, Microbial social behaviour, Altruism, Cheating

Introduction

Studies on the evolution of development tend to focus on features such as metamerism, organs of specialisation and body patterns. With some exceptions, they gloss over what has been acknowledged, ever since Weismann (1893), as the most striking feature of multicellularity, division of labour between the germ line and soma. Somatic cells contribute to building bodies and define what we think of as the phenotype. In due course they die and their genes disappear with the death of the body that they have built. The germ line does not contribute to the visible phenotype but the genes that it contains are transmitted to the next generation. Developmental biologists do not often ask *why* there is a germ line-soma distinction at all. The reason could be that it is so familiar; or, that it is hardly possible to conceive of how things could be otherwise. The usual approach to the difference between the germ line and soma is to look at it as a *developmental* problem; that is to say, as something that demands an explanation in terms of the causal pathways of gene expression and cellular

physiology. That it also poses an *evolutionary* problem is a relatively recent realisation and can be traced to two influences. The first influence, an indirect one, was the incisive analysis of conceptual issues related to the evolution of social behaviour carried out by Hamilton (1964), Williams (1966) and Trivers (1971). They showed the importance of making a distinction between the individual and the group as units of selection and discussed the roles that kinship might or might not play in the evolution of group-level adaptations, especially adaptations in which an individual seemed to display altruistic behaviour. The other influence has been direct. In a number of books and papers, Bonner has hammered home the point that to understand development, especially morphogenesis and differentiation, in a fundamental sense, it has to be looked at as an evolutionary problem involving cooperative behaviour with division of labour (Bonner 1958, 1965, 2001). His message is that (a) the unit of selection is the life cycle, (b) size plays a central role in evolution, and (c) all multicellular organisms shrink dramatically in size for part of their life cycle (during the germ cell phase).

*Corresponding author. E-mail: vidya@ces.iisc.ernet.in

Therefore, if we want to understand the origin of somatic cells in a background of unicellular reproductives, we must take into account size increase as a preadaptation. More specifically, almost every one of his publications ranging over the past sixty years illuminates some adaptive aspect of development in the cellular slime moulds (CSMs), the system of interest to us here (Bonner 1991 contains a selection that needs to be updated). The present review touches on something akin to the germ line-soma dichotomy that is found in the asexual life cycle of the CSMs. In these social amoebae the individual genotype has a persistence possessed only by genes, or by DNA segments that form units of recombination, in sexually reproducing organisms.

The evolutionary problem posed by the germ line-soma distinction is this: How can it benefit the genes that are in one cell (a somatic cell) to help genes in another cell (a germ cell) to be transmitted to future generations? There are two reasons behind the lack

of attention paid to this problem. Firstly, development usually involves changes in size, shape and organisation within a confluent group of cells all of which are in constant and, as it were, enforced cooperation with one another. The extent of mutual dependence in the process leaves no room for cellular phenotypes to vary outside narrowly defined limits; development in the higher metazoans is 'frozen in'. A mutation that results in a somatic cell entering the germ line or a primordial germ cell becoming part of the soma is likely to be so harmful to the body carrying it that it is subject to strong negative selection. Secondly, the problem of division of labour between the germ line and soma seems to have a facile solution. The fertilised egg develops into the embryo and the adult via a series of mitotic divisions; multicellular organisms are clones. Thus the genetic makeup of somatic cells is identical to that of the germ line and the division of labour can be viewed as a mechanism by which somatic cells make use of germ cells as vehicles for passing on their own genes to the next generation (Dawkins 1982).




Figure 1. a-i Life cycle of *Dictyostelium discoideum* a, Fruiting body; b, Spores; c, Amoebae; d, Loose aggregate; e, Streaming aggregate; f, Mound; g, Slug; h, Early culminants; i, Two macrocysts formed by *D. giganteum* stains. Scales: the mature fruiting body and slug are about 1mm; spores and amoebae are about 10 μm ; and the macrocysts are about 100 μm .

These limitations do not hold in the cellular slime moulds (CSMs). The CSMs are free-living soil amoebae that grow and divide by mitosis so long as food is plentiful and enter a developmental stage without cell division when the food supply is exhausted (Bonner 1967). In one form of development amoebae form unicellular cysts; in another they construct a much larger cyst after aggregation; and in a third, they build a migratory multicellular structure known as the slug, also after aggregation (figure 1). In the species that have been well studied, the end or terminal state of differentiation, called the fruiting body, consists of an elevated column of dead cells (stalk, equivalent to somatic tissue) that supports a globular mass of live cells (spores, comparable to the germ line) on its top and thereby aids in their dispersal. In some species, well before terminal differentiation has taken place, one can identify presumptive ('pre') stalk and spore cells. The presumptive state remains reversible until the fruiting body is built. Pre-stalk and pre-spore cells can transdifferentiate; they can also revert to the feeding stage if food is made available. Because cellular slime mold amoebae form truly multicellular structures, they serve as paradigms for both the evolution of multicellularity and the evolution of altruism in social groups.

One might say that CSM amoebae come together by choice and not because they are born in close proximity. Such being the case, there is no reason why they should be genetically identical; aggregates can contain amoebae of more than one genotype. This raises more starkly the question of why (in an evolutionary sense) only some of the cells that join an aggregate contribute to the next generation whereas the rest die, apparently for the sake of furthering the genetic interests of the survivors. The death of the amoebae that contribute to the stalk is thus an example of behavioural altruism. To the extent that cellular behaviour, including social behaviour, has a genetic basis, the altruism would seem to demand an explanation in terms of the genes that influence cell behaviour. Our aim in this review is to (a) reiterate that in the CSMs, multicellular development and social behaviour are the same thing; and (b) set up a framework within which to discriminate between different explanations for altruistic behaviour in the CSMs such as group selection, kin selection and individual level selection.

Social behaviour and division of labour in the CSMs shows features that are common to other systems in which too a tension exists between individual and group advantage. Only a few instances can be listed here. There are microbial systems (Crespi 2001, Velicer 2003), foremost being the myxobacteria, which display an amazing extent of convergent evolution with the CSMs in that single cells aggregate and build a fruiting body (Kaiser 1986, 1993), a form of social behaviour can be exploited or lost (Velicer et al. 2000, 2002). The volvocine algae display a size-dependent evolutionary trend with a clear somatic segregation being apparent only in the larger-sized species (Kirk 1998, Michod & Nedelcu 2003, Solari et al. 2002), a trend that can also be seen in the CSMs (Bonner 2003a). In the colonial ascidian *Botryllus*, the fusion of two individuals can result in a competition between cell lineages for populating the germ line or soma (Stoner et al. 1999), resembling a possible competition between CSM amoebae for becoming pre-spores or pre-stalk cells (Buss 1999). The most fascinating analogy to the CSMs is to be found in the social insects, especially the primitively eusocial wasps (Gadagkar & Bonner 1994). The wasp worker is comparable to a pre-stalk amoeba and the queen to a pre-spore amoeba. The analogy extends further. Caste differentiation in *Ropalidia marginata* can be described in terms that are similar to those used for describing the developmental strategies adopted by starved amoebae of *Dictyostelium discoideum*. What has been termed 'coin-tossing' in the case of *D. discoideum* (Nanjundiah & Lokeshwar 1984, Nanjundiah & Bhogle 1995) is 'gambling' in *R. marginata* (Gadagkar 1990). The difference is that in the former case the outcome of coin-tossing biases the probability that an amoeba that has joined a group becomes a stalk cell or a spore, whereas in the latter case the gamble has to do with whether being a founder is better or worse than joining a group. The choice of differentiation pathways is influenced by pre-aggregation biases in *D. discoideum* (reviewed in Nanjundiah & Saran 1992), paralleling pre-imaginal caste biasing in the wasp (Gadagkar et al. 1988). Reproductive dominance of workers by the queen wasp can be compared to dominance by high-quality amoebae (Atzmony et al. 1997).

Many workers have looked at division of labour in the CSMs from an evolutionary viewpoint (Filosa 1962, Bonner 1967, 1982, Armstrong 1984, Nanjundiah

1985, DeAngelo et al. 1990, Matsuda & Harada 1990, Gadagkar & Bonner 1994, Hilson et al. 1994, Atzmony et al. 1997, Strassmann et al. 2000, Hudson et al. 2002, Queller et al. 2003). Bonner (2003a,b) has presented an integrated hypothesis for the evolution of development in the CSMs in which aggregate size plays a crucial role. On the whole, though, the literature on social behaviour in CSMs is richer in theoretical considerations than in empirical data, especially observations made under natural conditions. The present work reviews previous discussions about CSM life cycles as evolutionary problems; sometimes the same aspect is discussed under different heads. In particular, we address the issue of altruism during fruiting body formation with reference to the best-studied species, *Dictyostelium discoideum*. Unless stated otherwise, our discussions will be with reference to it. Simply because so much more is known about *D. discoideum* than any other species, we tend to think of it as if its properties are illustrative of CSMs in general; it must be kept in mind that this need not be true. Other CSMs may differ in essential respects from it because they occupy niches that differ in ways unknown to us. Further information can be found in books (Bonner 1967, Raper 1984, Kessin 2001).

Natural History and Phylogeny

The CSMs are a group of eukaryotic amoebae found in soils all over the world; the number of recorded species approaches 100. Oskar Brefeld discovered them in 1869 when he isolated *D. mucoroides* from horse dung and followed its life cycle. The most commonly used laboratory species, *D. discoideum*, was isolated from North Carolina (USA) forest soil by Raper in 1935. The size of a CSM amoeba varies from 10 to 15 μm . The amoebae are motile and move by means of pseudopodia. Bacteria are their natural prey but they can also live off yeasts. Reproduction is asexual and leads to an increase in cell numbers by binary fission. It is characteristic of the CSMs that growth and cell division are separated in time from development and differentiation; the latter phases follow after starvation. The conventional end of one developmental cycle and the beginning of the next, the fruiting body, is the stage of hibernation and potential dispersal. The means of dispersal may include passing soil invertebrates, flowing water and, in the case of *Acrasis*, wind (Bonner 1982). Soil nematodes are predators of the CSMs; but the

nematodes can digest only amoebae, not spores (Kessin et al. 1996), and one can find a food chain that ends with CSMs being dispersed over very long distances by birds (Suthers 1985). To cope with a range of osmotic conditions, CSM amoebae are equipped with contractile vacuoles and lysosomal enzymes help in the rapid digestion of food. They occur in nature as microcysts, spores or vegetative amoebae in the soil and on tree barks; the relative percentages of the different forms exhibit seasonal variations (Kuserk 1980). Visual evidence of CSM fruiting bodies in nature – on deer scat – has been provided recently by Velicer (2003).

Raper (1984) classified the CSMs under the kingdom Mycetae (fungi), in the division Myxomycota, and the class Acrasiomycetes. The correctness of this classification is now in doubt. Though they have been thought of as resembling 'true' slime moulds such as *Physarum* and many fungi, the CSMs are very different from both. Unlike true slime moulds, at no stage in their life cycle do CSM amoebae fuse with each other; rather, they maintain a cellular identity throughout development. Also, the absence of vegetative hyphae and non-saprophytic feeding differentiates them from fungi, as stressed by Olive (1975) who emphasised their animal-like characteristics. A phylogeny based on the small ribosomal RNA subunit indicates that *Dictyostelium* diverged from the main animal lineage before yeasts; but a phylogeny based on protein sequence data says that *Dictyostelium* diverged much later than yeast (Loomis & Smith 1995). Recent phylogenies based on combined proteins sequences (Baldauf et al. 1997, 2000) place the CSMs, along with *Physarum* in a group of their own known as the Mycetozoa, which was also the name used by Olive (1975). The Mycetozoa form part of a clade just before the animal and fungal clades and after the plants. *D. discoideum* contains six haploid chromosomes and the total nuclear DNA content amounts to 34 Mb. It also has a multi-copy 90 Kb extra-chromosomal fragment containing rRNA genes and a 55 Kb mitochondrial genome. Approximately 8000 protein-coding genes are estimated to be present. The coding regions of the *D. discoideum* genome have relatively low A-T content (60-65%) compared to the rest of the genome (73-77%); the fully sequenced second chromosome has an A-T content of about 80% (DNA-related data are from Loomis & Kuspa 1997 and Glöckner et al. 2002). This should be compared to the A-T content

Table 1 Characteristic features of *Acrasids* and *Dictyostelids* (adapted from Bonner 1967, Olive 1975 and Raper 1984).

Group	Type example	Characteristics
Acrasids		No clear division of labour among cells, fruiting bodies about 1mm in height. No chemoattractant reported.
	<i>Guttulina</i>	Stalk cells are viable, globular mass of spores atop small stalk.
	<i>Copromyxa</i>	Stalk cells are viable; spores are borne on branched structures.
	<i>Guttulinopsis</i>	Stalk cells are dead, spores are somewhat differentiated.
	<i>Fonticula</i>	Mound of cells secretes extracellular stalk tube through which cells propel themselves upward.
	<i>Acrasis</i>	All cells are alive; spores are borne on branched structures.
Dictyostelids		Fruiting bodies with division of labour, size range 1-5 mm. Aggregation with chemotaxis, can include signal delay.
	<i>Dictyostelium discoideum</i>	Chemoattractant, cAMP. No stalk during slug migration. Stalk bears single sorus on the top. Presence of basal disc. Lemon-coloured spore mass.
	<i>D. giganteum</i>	Chemoattractant, cAMP. Slugs show stalked migration, mature stalks slender and longer than in <i>D. discoideum</i> .
	<i>Polysiphondylium violaceum</i>	Chemoattractant, Glorin (a dipeptide). Stalked migration of slugs, branched fruiting bodies. Violet-coloured spore mass.
	<i>Acytostelium</i> spp.	All cells secrete an cellular stalk and all differentiate into spores. Height of fruiting bodies varies from 0.1 to 1.2 mm.

of other metazoans- approximately 59% in *Drosophila melanogaster*, 54% in the mouse and 57% in the human genome (Campbell et al. 1999). The CSMs fall into two major groups, *Acrasids* and *Dictyostelids* (table 1). Of the two, the *Acrasids* have been little investigated compared to the *Dictyostelids*.

Acrasids

Acrasids aggregate without breaking up into secondary streams and lack a clear demarcation between spores and stalk cells. To date, no chemoattractant has been identified as being responsible for their aggregation. In the genera *Acrasis* and *Copromyxa*, sorocysts (amoebae) pile on top of each other, come above the surface and branch out. All terminally differentiated cells are viable. In *Guttulinopsis*, the stalk is made of dead cells (Olive 1975). *Fonticula* shows no division of labour in the sense that every cell differentiates into spores, which rise in the air while extruding an extracellular stalk. Coming together and sticking to each other appears to have the function of raising the mass of amoebae above the substrate level. To the extent that every amoeba participates in sporulation, all get a survival and eventual reproduction (the chances may or may not be the same for all). It may be that the *Acrasids* retain

relatively primitive evolutionary features and that the appearance of the *Dictyostelids*, with their morphologically differentiated stalk and spore cells, represents a later evolutionary step. Other than in *Guttulinopsis*, cell death does not seem to be part of the developmental cycle (Olive 1975, Raper 1984). The majority of *Acrasid* fruiting bodies consist of viable spores and viable stalk cells. The absence of reproductive division of labour in most of the *Acrasids* may imply that they lack an advanced communication system such as the *Dictyostelids* possess (Bonner 1967).

Dictyostelids

Dictyostelid amoebae aggregate in response to self-generated chemoattractants and differentiate into two distinct cell types, spore and stalk, which taken together constitute the fruiting body. The broad features of development and differentiation in *Dictyostelid* species such as *D. mucoroides*, *D. purpureum*, *D. giganteum*, *D. lacteum*, and *P. violaceum* are similar to those in *D. discoideum*. However, there are differences too (Raper 1984). The chemoattractant used for aggregation, the geometry of aggregation, pattern formation in the slug and fruiting body morphology can all vary.

In many species of Dictyostelids the spores contain refractile granules. These can be found on both extremities of spores and also in a central position and are known as polar granules (Hagiwara 1989). Traub and Hohl (1976) drew attention to a curious feature, namely that species with polar granules do not use cyclic AMP as the chemoattractant while species lacking polar granules do. One does not know how general the correlation is.

An elaborate morphogenetic process involving division of labour precedes fruiting body formation and distinguishes the Dictyostelids from the Acrasids. *Acytostelium* is an exception to the rule. Members of this genus are characterized by the absence of division of labour; following aggregation, all cells secrete an extracellular stalk and differentiate into viable spores (Bonner 1967, Raper 1984). An immediate advantage of coming together and sticking to each other is that a large size protects against the environment by decreasing the surface area relative to the volume. Later, because of large numbers, a mass of amoebae (differentiated or otherwise) can be raised above the substrate level, thereby helping dispersal. Though they may remain viable, in the Acrasids many amoebae end up as stalk cells and not as part of an elevated mass. Therefore on average the chances of dispersal and post-aggregation survival per amoeba that enters an aggregate may be less in the Acrasids than in the Dictyostelids. On the other hand, the fact that there is no cell death probably redresses the balance.

Survival Strategies: Development and Social Behaviour

As long as food is available, amoebae keep growing and dividing and lead a solitary existence. Starvation triggers the multicellular or social phase. During which large numbers of amoebae (typically 10^3 – 10^5 in *D. discoideum*) signal to one other, aggregate and undergo a complex developmental cycle. The result is either a fruiting body made up of a dead stalk and a viable spore mass on top of the stalk, or a macrocyst, which may involve a sexual phase.

As in other microorganisms, sporulation or sexual reproduction are responses to the stress of starvation. But CSMs can respond to harsh conditions in one other way, which is by forming microcysts, a solitary means

of survival. Microcyst formation obviously does not involve division of labour. Macrocyst formation begins with aggregation. This is followed by similar but complementary behaviour on the part of two cells that fuse. The fused giant cell becomes a diploid and enters into meiosis, but before doing that it cannibalises the others. Therefore there is some division of labour here. The physical environment and biotic factors may influence the choice of a particular life cycle strategy by an amoeba (Raper 1984).

Microcysts

An amoeba can respond to starvation by encysting itself. Unlike spores, which are surrounded by a three-layered wall structure, microcysts have a less firm double-layered structure around them. Raper (1984) has drawn attention to intriguing correlations. Delicate species such as *D. polycarpum*, *D. mexicanum*, *D. lavandulum* and *Acytostelium leptosomum* form microcysts whereas robust ones, for example *D. discoideum*, *D. purpureum*, *D. giganteum* and *Polysphondylium violaceum*, do not; however, *P. pallidum* does. A natural assumption is that fruiting body formation elevates the spore mass above the ground level, favours dispersal of spores and thereby improves the chances of successful germination. There being no food at the site of aggregation, the chances of food being available elsewhere must be better, or at any rate no worse. It may be that the less robust species can resort to both fruiting body formation and microcyst formation under the appropriate set of conditions (exactly what the conditions are, is unknown). Delicate species may not be capable of forming a sturdy stalk – in particular *Acytostelium spp.*, which makes an extracellular stalk. Even in a species such as *D. polycarpum*, which forms a cellular stalk, the fragility of the stalk may make it difficult for the spores to remain in place long enough to have a chance of dispersing very far. Microcysts would seem to have a poor ability to disperse. Nevertheless, depending on the circumstances, the strategy of forming a microcyst may be preferred. Firstly, it could improve the average chance of long-term survival of the cell, or better put, of the genotype. Secondly, if for some reason clonal populations of amoebae are rare, forming a microcyst may be preferable to chancing the survival of the genotype by forming a chimaeric fruiting body in association with unrelated individuals; the cost of losing the

opportunity to disperse may be traded off against the benefit of allowing for a higher probability of individual survival. Overall, too little is known about the microcyst cycle for us to do more than speculate about its evolutionary connotations.

Macrocyts

Besides the asexual modes of propagation, namely via, fruiting body and microcyst formation, Dictyostelids also possess a sexual mode of reproduction. This involves what is known as the macrocyst cycle. Amoebae can undergo sexual development under conditions of high humidity and darkness when there are individuals of opposite mating types present in the same aggregate. Macrocyst formation begins with aggregation via chemotaxis and the fusion of cells of opposite mating types within the aggregate (Blaskovics & Raper 1957, O'Day 1979). The fused cell, normally binucleate, proceeds to engulf and digest the other cells and develops into an encysted structure known as the macrocyst. Bozzone and Bonner (1982) have shown that the mere presence of a small fraction of cells belonging to a different mating type can cause cells of the majority genotype to fuse. If this is common in nature, the macrocyst is an interesting way of survival but is not a sexual stage. Given appropriate conditions, the macrocyst can germinate. A diploid cell results from nuclear fusion in the binucleate cell. The diploid nucleus undergoes meiosis and gives rise to four haploid progeny, of which one survives to give rise to an amoeba (for details see Urushihara 1992, 1997). Macrocysts seem to have two functions: (i) zygote formation and recombination, and (ii) an alternative resistant stage, without any recombination, probably in response to sudden flooding and especially in homothallic forms (see Bonner 1982, Bozzone & Bonner 1982).

Different mating types have been discovered in many species of cellular slime molds (Clark et al. 1973, Erdős et al. 1975). O'Day and Lewis (1975) showed that diffusible mating type factors mediate macrocyst formation. In the laboratory, *D. discoideum* can form macrocyts in various combinations including that of the two naturally occurring strains NC4 and V12. Getting them to germinate is generally difficult, though certain pairs of strains readily yield recombinant progeny (Francis 1998). Erdős et al. (1975) tested 42 natural

isolates of *D. giganteum*, 50 % of which formed macrocyts. In contrast, Clark et al. (1973) say "... very few of the many strains of cellular slime molds that have been isolated from nature actually form macrocyts". Evans et al. (1988) used the technique of restriction fragment-length polymorphism (RFLP) to distinguish between *Dictyostelium* genotypes. They found that wild isolates of *D. discoideum* shared a minimum of 80% of the observed restriction fragments with NC4, the commonly used laboratory strain originally isolated by Raper in 1935. Francis and Eisenberg (1993) examined *D. discoideum* strains from the same forest soil from which Raper had originally isolated it. Based on RFLP data, one of the strains seemed to be genetically identical to Raper's original NC4 isolate - an unlikely outcome if recombination had been frequent in the wild. These results imply that genetic exchange is rare among *D. discoideum* strains in the wild and that macrocyst formation may not be a common mode of reproduction in nature in this species. Transient diploids can be generated artificially under laboratory conditions and rare mitotic crossovers can be utilized for parasexual genetic analysis (Katz & Sussman 1972). It is not known whether anything similar occurs under natural conditions. According to Kessin (2001), because (i) the multicellular mode in CSMs very likely derived from one or more unicellular ancestors, (ii) starved cells aggregate by chemotaxis before a macrocyst is formed and (iii) the fruiting body is a far more complex structure than the macrocyst, the micro- and macrocyst life cycles could have served as pre-adaptations that favoured the evolution of fruiting bodies. In this regard, Mutzel (1991) drew attention to the occurrence of predation and cannibalism in the macrocyst cycle and said that predatory behaviour might have led to sociality.

Aggregation and Fruiting Body Formation

Aggregation by chemotaxis (Bonner 1947) is the prelude to forming a fruiting body. Hagiwara (1989) divides the aggregation pattern into four types: *mucoroides*-type, *violaceum*-type, *minutum*-type, and *microsporum*-type. In the *mucoroides* type, amoebae move towards the centre in continuous streams and one or more fruiting bodies are formed at the centre. Species lacking polar granules, for example

D. discoideum, *D. mucoroides* and *D. purpureum*, come under this category. Sometimes streams break up to give rise to more than one slug. The *violaceum* type is similar except that streams break up frequently and secondary centres are formed. Species like *P. violaceum* and *D. aureo-stipes* display this kind of aggregation. In *minutum*-type aggregates, amoebae usually aggregate without streaming. This results in solitary or clustered fruiting body formation; *D. minutum* undergoes this kind of aggregation. The *microsporum*-type, exemplified by *D. microsporum*, is similar to the *minutum*-type during the early stages, but later secondary centres are formed around the main one; the amoebae go on to form tertiary centres around the secondary ones.

The well-known second messenger cyclic AMP is the chemoattractant that drives aggregation in *D. discoideum* and several other Dictyostelid species; *P. violaceum* (and probably *P. pallidum*) uses a dipeptide, glorin (Shimomura et al. 1982) and *D. lacteum* uses pterin (van Haastert et al. 1982). Bonner (1982) alludes to at least 8 distinct chemoattractants in the CSMs and tries to link the diversity of attractants with isolating mechanisms. In proposing that the aggregation signal was pulsatile, relayed from cell to cell and subject to rapid degradation, Shaffer (1961, 1962) anticipated the essential features of what we know today about the pathway which links cAMP signaling, its reception and directed movement by the receiving cell (Parent & Devreotes 1999). Relaying extends the influence of the original centre of signaling and increases the size of the aggregating mass. A fascinating finding is that the cAMP signal in *D. discoideum* is released in regular periodic bursts every 7 min or so and relayed from cell to cell (Gerisch & Wick 1975, Tomchik & Devreotes 1981); oscillatory movements, presumably driven by oscillatory signal propagation, persist through later stages (Dormann et al. 1998). One can show that given the parameters appropriate to *D. discoideum*, periodic signaling increases the range of the signal well beyond the range of a steady signal under comparable conditions – that is, given that both require the same expenditure of energy in the long run; other parameter values can tilt the balance in favour of steady signaling (Nanjundiah 1973). Interestingly, periodicity per se appears to be adaptive in another manner. When supplied externally for the purpose of stimulating an aggregationless mutant of *D. discoideum* to

differentiate, a periodic train of cAMP pulses is significantly more effective than an aperiodic train of the same strength per pulse and the same mean period (Nanjundiah 1988).

D. discoideum is characterized by the stalkless migration of slugs and a well-developed basal disc in fruiting bodies. Slugs exhibit phototaxis and thermotaxis (Bonner et al. 1950) and it has been argued that in conjunction with localized ammonia production, these abilities serve as adaptations that enable them to reach the soil surface (Bonner et al. 1988, 1989). In this species and its close relatives, pre-spore cells are positioned in the posterior of the slug and pre-stalk cells in the anterior, with the spatial pattern being the result of segregation, or sorting out, of cells with different pre-existing tendencies (Bonner 1959, reviewed in Nanjundiah & Saran 1992). Sorting out must entail a great deal of relative movement within aggregates and slugs. By making use of an elegant technique for inducing flat (two-dimensional) slugs to form, Bonner (1998) showed that the pattern of movement was seemingly uncoordinated and random in pre-stalk cells and coordinated and directed towards the anterior in pre-spore cells. His vivid description is 'gas molecules zipping about inside a balloon' and 'a flock of sheep moving down a narrow street' respectively (Bonner 2000). Internal movement persists through further development and, along with movement of the mass as a whole, is responsible for shaping the fruiting body during culmination (Dormann et al. 1996, 1998). A functional spatial segregation of cell types is achieved very late in development in other species, if achieved at all.

Fruiting bodies of *D. discoideum* have a clear basal disc and a tapered stalk; the sori of are colourless to yellowish. Fruiting bodies of *D. purpureum* are solitary, unbranched and phototrophic. Sori are light to deep purple in colour. *D. giganteum* is characterized by long creeping stalks (approximately 17-25mm) and small white sori (Singh 1947); *Polysphondylium* has branched fruiting bodies. In species other than *D. discoideum*, stalks can be very long and slender. Raper (1984) lists these and many other morphological differences between species. The reason for spending time on them here is that on the one hand they are striking, and on the other hand we have no clue as to what the differences mean. How many of these reflect adaptations, how

many are traits that have been fixed by drift, how many are side effects of selection for something else or reflect developmental constraints (Bonner 1982), is unknown.

The speed of slug movement increases with size and presumably makes dispersal more efficient, which is yet another adaptive reason for getting into an aggregate and inducing others to get in (Bonner et al. 1953). According to Raper (1984) three species besides *D. discoideum* have a stalkless migratory stage and one (*D. sphaerocephalum*) is stalkless when migrating at low temperatures; this means that the majority of known species leave a stalk behind during the migrating slug stage. Often the stalk is aerial. We do not know whether stalked migration is a primitive or an advanced feature. But we can speculate on the possible advantages and disadvantages of leaving behind a stalk during migration. It may be that the loss of the cells that form the stalk is compensated by the ability of the remaining cells to travel a greater distance than they would if all cells conserved their energy and none differentiated into a stalk during migration; Bonner (1982) mentions a fruiting body 22cm long. But this cannot be the whole story, because as said larger aggregates move faster than smaller ones (Bonner et al. 1953). It may be that under natural conditions the long stalk that is left behind means that the period of migration is lengthened and more time is had for seeking out a suitable habitat for fruiting (Bonner 1982). Stalked slugs soar above the substrate before they start culminating, so that besides aiding dispersal the stalk raises cells away from possible toxins in the soil (Gadagkar & Bonner 1994) and reduces the chances of their being eaten by soil predators such as nematodes (Kessin et al. 1996).

Fitness and the Fruiting Body

Forming a microcyst is self-evidently a means of tiding over harsh times and a macrocyst, in addition to fulfilling the same role, can promote genetic exchange, which – for a variety of reasons, not all of them fully clarified (Bell 1982) – is believed to be generally advantageous to the participants. Here we confine our attention to the fruiting body.

For the sake of definiteness let us start with a single amoeba that has just germinated from a spore. Its fitness measured over the entire life cycle has many components. To begin with, there is cell division: the efficiency with which the amoeba makes copies of

itself during the vegetative phase is its fitness component during growth. Starvation makes the daughter cells stop dividing and prepare themselves to withstand the unfavourable condition. The viability of a starved amoeba, the probability that it enters an aggregate, the probability that the slug remains cohesive and migrates to the surface, all depend on single cell traits that contribute to fitness. Fitness can be measured after this stage in terms of the average number of spores that result from an amoeba that gave rise to one clone of cells within the same aggregate – that is, all the spores that can be traced back to the single amoeba that we began with. This takes into account the fact that not all the clonal products of an amoeba may survive starvation. The likelihood that a differentiated spore germinates, which is partly related to the probability that it disperses successfully, must also form a component of fitness. Finally, fitness depends on the environment in which it is measured. Ponte et al. (1998) demonstrated that the same strain behaves differently on different substrates. They discovered that a mutant having a deletion in the *csA* gene (the *csA* molecule mediates EDTA-resistant adhesion between cells) resembled its wild type parent when observed on a standard agar surface. However, when soil was used as the substratum the wild type sporulated more efficiently than the *csA* null mutant.

When there is no division of labour, one can think of fitness in at least two different ways. If all the amoebae in an aggregate secrete a stalk and form viable spores (e.g. *Fonticula*, *Acytostelium*), the simplest assumption is that they are all on a par (this ignores any jockeying that might occur to get into a favoured position within the spore mass). Then, everything else being equal, the fitness of the founder would be proportional to the number of amoebae that aggregate. This would place a premium on aggregate size. Counter-selection would become significant once considerations of energy expenditure and mechanical stability begin to disfavour large spore masses. It can be argued (see later) that this form of fruiting body construction would benefit each founder whether the co-aggregating cells are derived from a single founder or many genetically distinct founders. On the other hand, if the fruiting body consists of a stalk and spores but both are viable (e.g., *Guttulina*), one would imagine (on grounds of improved chances of dispersal) that spore cells have an advantage over

stalk cells. Therefore some of the amoebae in an aggregate are more likely to transmit the genes of the founders than the others, and one already has the beginnings of a primitive form of division of labour. An operational definition of fitness will have to incorporate the relative degrees of reproductive success achieved via the spore and stalk pathways.

CSMs without Division of Labour

Protostelium is an asocial species with the same terminal morphology as the others except that here a single cell transforms itself into a spore and extrudes a stalk (Raper 1984). This shows that the evolution of a dispersive stage that was raised above the soil may have preceded the evolution of multicellularity and division of labour.

As has been said, in most Acrasids there is no evidence for division of labour. In some, for instance *Guttulina*, all terminally differentiated cells are potentially capable of reproduction but it is not known whether their fitnesses are the same. Cells occupy different positions in the multicellular structure and it may be that those situated at the bottom of the fruiting body do not get to disperse as far as the ones at the top. Unfortunately there is no report of the dispersal or survival of the cells in Acrasid fruiting bodies. *Acytostelium* and *Protostelium* provide other examples of fruiting body formation without division of labour. In *Acytostelium* all cells sporulate and secrete an extracellular stalk, which keeps them raised above the substrate.

Purely on mechanical grounds, there are advantages to a cell that can sporulate and secrete an extracellular stalk, to do so as one of many cells (as in *Fonticula*) rather than by itself (as in *Protostelium*). The principle depends on a scaling law and is illustrated by the fable 'Unity in Strength' (Aesop 1954). Imagine that a starved amoeba has a fixed amount of resource, measurable in terms of energy, to be partitioned between making a spore and secreting a stalk. It helps if the spore rests on a tall stalk. But there is an upper limit to how much of the resource can be invested in making the stalk, because spore viability drops as the fraction of the energy allocated to building the stalk goes up. A stalk of volume v is the most that a cell can manage. v is given by the formula $\pi \cdot r^2 \cdot l$ where l is the height and r the radius of the cross-section. Now, given a cylindrical stalk of length l and radius r , the

maximum load (which is to say, the maximum spore weight) that it can carry before it becomes unstable with respect to an infinitesimal amount of random bending is given by $T_{\max} = \pi^2 E I_2 / 4 l^2$. Here I_2 , which equals $\pi r^4 / 4$, stands for the moment of inertia of the cross-section and E is the modulus of elasticity of the stalk (which we take to be a characteristic of the material of which it is made) (Landau & Lifshitz 1970, p.98). Also, r^2 equals $v / \pi l$. Substituting first for I_2 and next for r^2 , we get $T_{\max} = \pi E v^2 / 16 l^4$. This is the relation between the minimum viable weight of a single spore and the maximum length of stalk that it can secrete. Assume now that instead of one spore we have n spores forming one spore mass and n stalks, each of them of volume v and length l , forming one large and approximately cylindrical bundle. The volume of this cylinder will be $V = n \cdot v$ and its radius will be R where $\pi \cdot R^2 \cdot l = n \cdot v$. What is the maximum load that can be carried by this collective cylinder before it becomes unstable? The answer, as shown by simple substitution, is n^2 times, and not n times, the maximum load that can be carried by one cylinder. In other words, a bundle of ten stalks can carry 100 times the load that a single stalk can before instability sets in. Equivalently, to be assured of the same degree of mechanical stability, each one of ten amoebae that takes part in laying down one stalk tube needs to invest just $\sqrt{1/10}$, or about 30%, in making a stalk relative to what it needs to invest when sporulating by itself. The extra 70% can be re-directed towards improving spore viability. The viability of the spore would be still higher, of course, if it did not have to use its resources in making a stalk at all but could rest on a stalk made by other amoebae.

The costs and benefits of fruiting body formation with reproductive division of labour, relative to forming a fruiting body without it, are unknown. A rigid, layered and cellular stalk would resist loading better and offers more stability than a stalk made up of an extracellular extrusion. Taken by itself, the benefit conferred by a cellular stalk would be expected to become significant as the aggregate size increases. This is just what one sees in the CSMs; *D. lacteum* sometimes forms tiny fruiting bodies, and when it does so, the stalk is partly acellular (Bonner 2003a). An increased stability of the stalk could mean a longer period of stability of the spore mass, giving more time for dispersal

to occur. This may be advantageous if the average duration between successive episodes of significant bacterial growth in the same patch of soil is a long one. On the other hand, the natural ecology might be such that there is a relatively short interval between bouts of bacterial growth in the same location. In that case, it might be unimportant to insure for long-term stability of the structure and a less sturdy extracellular stalk might do. Also, the secretion of stalk material must require energy. Other things being equal, then, under natural conditions *Acrasid* and *Acytostelium* spores should also be less capable of withstanding long-term starvation than Dictyostelid spores. Perhaps in nature they do not need to remain viable as long as the latter. But all these arguments remain purely speculative in the absence of a detailed knowledge of the relevant ecological conditions.

CSMs with Division of Labour

Division of labour is most clearly exhibited when a sub-set of cells within multicellular groups of Dictyostelid aggregations forms a dead stalk in the course of terminal differentiation. The amoebae that die appear to improve the chances of reproductive success of those that sporulate. When they can be recognised well in advance, as in the case of *D. discoideum*, they are known as presumptive stalk or pre-stalk cells. Not all CSMs contain identifiable pre-stalk cells (Bonner 1952), but in those that do, pre-stalk cells may also exhibit traits that seem to benefit pre-spore cells, and so benefit spores. For example, in *D. discoideum* pre-stalk cells are predominantly the providers of the motive force for slug movement and dispersal (Inouye & Takeuchi 1980).

In those cases in which some amoebae die and form a cellular stalk, one must explicitly take into account a possible indirect role of stalk cells in enhancing fitness (at a level to be discussed below). One way to do so is to make the plausible and convenient assumption that pre-stalk and stalk cells aid in dispersal, and that the extent of the aid is a monotonic function of their number. If the whole fruiting body is one clone, a reasonable expression for the fitness of the original amoeba that gave rise to it would be the number of spore cells in the spore mass multiplied by an appropriate function of the number of stalk cells (Nanjundiah 1985, Matsuda & Harada 1990). Symbolically, let N be the total number of amoebae in an aggregate and n the number of

spores that it gives rise to. Ignoring any cells that are lost during migration, the number of stalk cells will be $N-n$. Given that the stalk aids in dispersal and the spores give rise to amoebae in the next generation, the fitness W will be positively correlated with both n and $N-n$. The simplest such correlation is expressed by the equation $W = n \times f(N-n)$. The choice of a functional form for f is not obvious, nor can we be sure that it makes sense to choose one particular form. What is important is that by representing fitness in this way we see why both stalk and spore cells are required. Too many spores reduce the efficiency of dispersal; a small number of spores make dispersal efficient (because the stalk becomes taller) but at a cost, because the number of propagules is reduced. There is a trade-off between allocating progeny to the stalk and spore pathways and the optimum solution is to direct some to both.

If f is proportional to the number of stalk cells raised to some power, it is easy to show that at the optimum (maximum) value of W , n and N , and therefore stalk and spore cells, are in a fixed ratio with respect to each other. Nanjundiah (1985) attempted to derive a specific form for W by assuming that the dispersal of a spore mass was ballistic, driven by a gust of wind, say. This may not be the way spore dispersal works for fruiting bodies that form on the soil surface except, as stated by Bonner (1982), in the case of *Acrasis*. But the form that W takes reflects a generic way of modelling dispersal. The model may mimic what actually happens in the case of arboreal fruiting bodies and for *Acrasis*. In other cases, it is an example of how a wrong assumption can lead to the right result – in this case, to the result that the relative proportions of the two cell types in the fruiting body are constant over a huge range of total cell numbers (Bonner 1967). In one sense (i.e., in terms of what happens if a slug is depleted of one or the other cell type) proportioning is equivalent to regulative embryogenesis and its astonishing constancy has been long thought of the central problem in CSM development (Bonner 1967). The correct explanation for proportioning may have to do with many factors besides the requirement of optimising the chances of spore dispersal, for example the stability of tensile columnar or cylindrical stalks loaded on the top. It is tempting to think that the right mechanical and

ecological model may in fact yield a power-law for dispersal. The limitation of our analysis is that we have assumed that an aggregate is a single clone. Matsuda and Harada (1990) showed that proportioning persists under somewhat relaxed assumptions about the dispersal function f and that the essence of the argument remains the same as the one sketched above when aggregates are multi-clonal. But before coming to that we need to examine the issue of altruism.

Altruistic Behaviour and Levels of Selection

We discuss how the altruism displayed by amoebae that die and contribute to the stalk might be viewed most obviously as an adaptation displayed at the level of the group, and less obviously as an adaptation at the level of the individual amoeba. Group adaptations can be usefully sub-divided into those in which kinship plays an essential role (kin selection) and those in which kinship may exist but has a minor role (group selection).

Group Selection

One can try to account for altruistic behaviour as a group-level adaptation that overrides a tendency to selfish behaviour at the individual level *even under circumstances in which the group does not consist of related individuals*. We have already given two reasons why joining a group may be better for a starved amoeba than staying alone and both remain valid. Firstly, a large aggregate can migrate faster than a small one. Secondly, it is more cost-effective (in the sense of the benefit to be gained by mechanical stability relative to the resources to be invested) for many spores to rest on a combined stalk than to do so on separate stalks. The second inference was drawn with reference to the case of an extracellular stalk tube but applies just as well for a cellular stalk, though the details of the stability argument can not be the same. Bonner (1982) showed that in *D. discoideum* the mid-point diameter and height of the stalk are proportional to one another, unlike the conclusion we reached earlier for a secreted cylindrical stalk tube. Both extracellular and cellular stalks provide an additional advantage, that of improved dispersal, which we consider now.

Let us consider a situation in which starved amoebae are present; assume that they can form neither microcysts nor macrocysts. The amoebae aggregate, and we wish to compare two sorts of

aggregates. In one, every amoeba sporulates but remains at the ground level or at the bottom of a crevice in the soil. In another, a few amoebae die and form a stalk, in the process permitting the others, which sporulate, to rise above the ground. Post-aggregative migration might occur in both situations. Suppose that aerial spores are better situated for dispersal than those on the ground. The difference will be meaningful if dispersal is essential in order to ensure that spore germination takes place before the spore viability falls too low. In that case, putting aside the question of whether a group consists of related individuals or not, a group of amoebae that participates in division of labour and building a fruiting body can do better than a group that does not.

One can make the point with the help of symbols. We will simplify matters by assuming that other things being equal, dispersal is a measure of reproductive success (thereby implying that the distance of dispersal is immaterial). This has no bearing on the essence of the argument. Suppose an aggregate consists of N genetically distinct amoebae. Let us say that when all of them sporulate, and therefore stay at ground level, the probability of dispersal of the spore mass is d_1 . Therefore the chance that any given amoeba in the aggregate will disperse successfully as a spore is $d_1 \cdot N \times 1/N$, namely once again d_1 , which we can now think of as the probability that the average amoeba will disperse in the form of a spore. In the second situation, out of the N amoebae in the aggregate a fraction f (less than 1) sporulate. In other words, any given amoeba sporulates with a probability f and forms a stalk cell with a probability $1-f$. Thanks to stalk formation, the spore mass is now elevated above ground level. Suppose that in consequence its probability of dispersal is now d_2 . Then the average probability that an amoeba will disperse as a spore is $d_2 \times f \cdot N \times 1/N$, which works out to $d_2 \times f$. Therefore the second option is better whenever $f \cdot d_2 > d_1$. (The extension of the argument to fruiting bodies with stalks of different lengths is trivial and, as before, leads to the inference that there must be a trade-off between stalk and spore production.) Note that the smaller the value of d_1 (i.e., the poorer the chances of a spore mass on the ground to disperse), the stronger is the force of selection for the group-level adaptation, namely for division of labour and altruistic behaviour.

Kinship has not entered the picture so far; what we are looking at is the expected benefit that an amoeba (i.e., the genotype whose representative is an amoeba) obtains by taking part in building a stalked fruiting body relative to the expected benefit to it of forming spores alone. In both cases the expectations are to be averaged over many life cycles, in some of which the genotype of interest goes into a spore and in others dies and differentiates into a stalk cell. Kinship can be taken into account in one of two ways. The role of kinship could be passive: that is to say, an amoeba automatically gains additional benefit when many of its daughters participate in building a common fruiting body rather than just one of them doing so. The role of kinship could also be active. Genetic relatedness could influence the probability of sporulation vis-à-vis becoming a stalk cell; in other words, f could vary between clones and depend on how many clones there are. Once again altruistic behaviour will be favoured in selection between groups. However, in the absence of additional reinforcement (for example, an enhancement of overall group reproduction by the presence of an altruist genotype within the group), a group selection model is inherently unstable. If a mutant that differentiates constitutively into a spore, a 'cheater', arises, it will spread unchecked. As we will see, kin selection, which is the name given to group selection in which the group consists of closely related individuals, has the same difficulty. The reason for this is that the success of both group and kin selection depends on a group-level adaptation, in this case the construction of a fruiting body with division of labour.

Kin Selection

Models for kin selection are usually framed in terms of the probability that two individuals in a group share a certain fraction of their genes by common descent. This framework is not so useful if what we are looking at is an asexual life cycle. Of course mutation will generate variation at a slow rate. But while building models, in the absence of genetic exchange, it is difficult to think of relatedness as a continuous variable (however easy it may be to come up with measurements of similarity between two genomes). It is more convenient to think of just two alternatives, individuals sharing all their genes - because they belong to the same clone - or not sharing all their genes.

We will now compare two situations: one in which aggregates are made up of closely related amoebae - ideally, members of a clone, and another

in which many different clones participate in building the same aggregate. In either case fitness will mean inclusive fitness, that is, it will refer to the clone or genotype as unit. In the case of reasonably-sized fruiting bodies of *D. discoideum*, about 20% of the cells give up their lives and in the process the remaining 80% survive (Nanjundiah & Bhogle 1995). It seems that when the entire fruiting body is built by a single clone, the cost incurred by the founder of the clone as a result of the death of some of its progeny is small relative to the benefit obtained - especially if, as we have just seen, *not* incurring the cost would imply a very low chance of survival overall. But what if there is more than one clone in a fruiting body?

Division of Labour as an Evolutionary Stable Strategy (ESS)

Matsuda and Harada (1990) modelled the decision made by amoebae to follow a stalk or spore pathway to see whether division of labor could be an evolutionary stable strategy (ESS) (Maynard Smith 1982). The ESS was a conditional or mixed strategy, and was dependent on the ability on the part of the amoebae to sense the total number of cells in an aggregate as well as each other's genetic relatedness. It turns out that in order to accurately model the spore:stalk ratios measured in small fruiting bodies, one needs to assume that the cells that make up an aggregate can sense how many of them there are (Nanjundiah & Bhogle 1995). We do not know how size-sensing might work, but it is not unlikely that it exists, because it is known that amoebae of *D. discoideum* possess quorum-sensing mechanisms (Brock & Gomer 1999, Okuwa et al. 2001). As shown by the outcome of mixing experiments, CSM amoebae have the ability to distinguish self from non-self (reviewed in Kaushik 2002); but it is not known whether they can also distinguish between members of their clone and others, or sense how many clones there are.

There were three main predictions of the ESS analysis. Firstly, for a given aggregate size, genotypes in a chimaeric aggregate that were represented by very few cells would contribute only spores until they attained a critical number; beyond that they would form a fixed number of spores which would be the same as that of every genotype whose representation was super-critical, the rest becoming stalk cells. Secondly, the relative number of amoebae that differentiated into spores would be an increasing function of the number of clones within an aggregate;

the more genetically diverse that an aggregate was, the more 'spory' would be the fruiting body that it gave rise to. Thirdly (in effect explaining the second prediction), the relative number of cells belonging to any single genotype that was allotted to forming spores would increase with the degree of genetic heterogeneity within an aggregate: the level of altruism would fall off as a function of the number of clones in a chimaera.

Hudson et al. (2002) have carried out a comparable but more extensive analysis of the outcomes to be expected in chimaeric aggregates. They too assume that the fitness of a founder cell is proportional to the product of two factors as before. The first factor (fecundity) is the number of spores that differentiate in aggregates consisting of its descendants and the second factor (dispersibility) is an increasing function of the number of stalk cells formed by the same aggregates. On the grounds that spore masses have some expectation of non-zero dispersal by themselves, they explicitly allow for the possibility of a small but positive contribution to fitness by the second factor even in the absence of a stalk. A puzzling feature of their model is that the fecundity and dispersibility both depend on the *fraction* of cells that form stalk, not – as one would expect – their absolute number. However, since Hudson et al. do not explore the issue of proportioning, their qualitative predictions are similar to those of Matsuda and Harada (1990). But the two models differ with regard to what they have to say about the coexistence of altruist and cheater genotypes (see below).

To sum up, reasonable-looking fitness functions can account for the allocation of some cells to a non-reproductive pathway (those which form the stalk). With the help of subsidiary assumptions whose plausibility is unknown they can also ensure an invariant spore:stalk ratio at the ESS. The ESS is derived by maximizing the fitness of each clone subject to the constraint that every other clone's fitness is also maximized. It exists, the ESS is automatically stable with respect to invasion by any other genotype that also allocates amoebae to both the spore and stalk pathways. Whether it is also stable against a 'cheater' that differentiates constitutively into spores remains to be examined.

Individual Level Selection

The underlying hypothesis here is that entering an aggregate and participating in the formation of a

fruiting body is an adaptation at the level of the individual cell (Atzmony et al. 1997). The test of the hypothesis of individual selection is to see whether it holds good when an aggregate contains many different genotypes, perhaps as many as the number of cells in it (to take the most extreme case). Then the assertion would be that both an amoeba that sporulates and one that ends up as a stalk cell benefits *as an individual amoeba* by coming together with the others and building a fruiting body. For this to be true in the case of an amoeba that contributes to the stalk, the assertion can be valid only if one condition is met. Namely, it must be the case that on average, for any amoeba, not joining an aggregate is a poorer option than joining one. This condition can be justified. Firstly, a starved isolated amoeba has extremely limited powers of dispersal, and is guaranteed to die in a relatively short time (Gregg 1971). Secondly, after joining an aggregate, many factors influence the choice of what an amoeba differentiates into. The factors can be summed up as an intrinsic 'quality' relative to the qualities of the other amoebae (discussed in Kawli & Kaushik 2001, Kaushik 2002). It is the relative aspect that makes participation in an aggregate an act of 'coin-tossing', a gamble on the part of an amoeba. Gambling in the hope of differentiating into a spore is preferable to remaining solitary, because it is impossible for an amoeba to predict in advance whether the other cells in its group will tend to be of higher or lower quality than itself. For this model to work, (i) amoebae in the wild must be heterogeneous with respect to their phenotypic qualities, (ii) the amoebae in an aggregate should be capable of assessing each other's quality by means of cell-cell signaling and (iii) after aggregation, amoebae should adopt a prestalk or a prespore fate depending on their relative quality.

Quality

'Quality' is defined as a cellular property that exists prior to aggregation and is correlated with fitness. It is expressed as a dominance hierarchy, a pecking order; the hierarchy can be discerned at the onset of aggregation and is physiological, not genetic (Bonner 1996). In our case, this means that in the normal course of things, the quality of a cell reflects the *relative* probability of spore formation by that cell. For the concept to be useful it should be measurable in the cell itself or in a genetically and (if at all possible) phenotypically homogeneous cell population and should be a reliable predictor of cell behaviour in social

groups. For example, the quality of an amoeba could be reflected in its ability to survive prolonged starvation when kept in isolation. Other correlates of quality exist; in the case of some, but not all, one can see right away that the relevant trait might have something to do with fitness.

Amoebae of *D. discoideum* that are grown in an axenic medium containing high levels of glucose, and are therefore nutritionally well-endowed, tend to form spores when combined with amoebae grown without glucose, which are in comparison nutritionally poorly endowed; the latter tend to form stalk cells (Leach et al. 1973). The important thing is for the two sets of amoebae to have different reserves of metabolisable sugar when they are starved (Takeuchi et al. 1986). Similarly, amoebae that are in the mid- to late G2 phase of the cell cycle when food runs out tend to form spores in mixtures with amoebae that are in late S or early G2, which show a tendency to stalk differentiation (McDonald & Durston 1984, Weijer et al. 1984). Amoebae with relatively low calcium levels at starvation are predisposed to a pre-spore pathway relative to those with higher calcium level, which are predisposed to a pre-stalk pathway (Azhar et al. 2001). It seems intuitively likely that an amoeba that has more energy reserves should be more likely to become a spore than one with relatively lower reserves of energy. However, cell size seems to act opposite to the way in which one would guess: in the beginning, pre-stalk cells can be larger than pre-spore cells (Bonner 1959, Bonner et al. 1985, Saran et al. 1994).

D. discoideum amoebae produce toxic molecules that are membrane-permeable chlorinated phenolic derivatives generically called DIF for 'differentiation-inducing factor'; the most potent of the DIF isoforms identified, DIF-1 (MW~ 300), induces stalk cell differentiation when supplied at an appropriate concentration (Kay 1997). In the original individual selection model it was hypothesized that the capacity to make DIF-1 and the tendency to resist its effects might be indications of high quality (Atzmony et al. 1997). Both predictions have been borne out in the sense that DIF-1 seems to be made by pre-spore cells and is broken down by pre-stalk cells, and nutritionally well-endowed cells are less sensitive to DIF-1 than nutritionally poorly endowed cells (Kay et al. 1993, Thompson & Kay 2000a, 2000b). However, the DIF story is more complicated because a mutant that is unable to make DIF-1 can form slugs

which contain cells that are of pre-stalk as well as pre-spore type; however, it lacks a particular subset of pre-stalk cells (pstO cells; Thompson & Kay 2000b). It may be that in the mutant background, some other molecule takes over the role normally performed by DIF-1 (the role of a coercive agent made by high-quality amoebae that induces low-quality amoebae to differentiate into pre-stalk cells). On the other hand, DIF-1 and related compounds may be merely one among a variety of factors used by cells to signal their qualities.

One can try to artificially flatten the quality profile by raising the quality of every member of a population. This has been done by growing amoebae in a sugar-rich medium (Leach et al. 1973), by synchronizing their growth and then starving them at G2-M boundary of the cell cycle (Azhar et al. 2001) and by lowering their calcium levels with the help of an ionophore (Baskar et al. 2000). Amoebae that have been treated in any of these fashions aggregate and form abnormally 'spory' fruiting bodies. The complementary experiment of lowering all qualities results in 'stalky' fruiting bodies. This lends strength to the supposition that quality is a reflection of a cell's intrinsic fitness. At the same time, because more or less normal development ensues in spite of the perturbations, intercellular interactions must play an important role in determining the fate of a cell. Since pre-aggregation biases can be shown to exist even when genetically identical amoebae are raised in a uniform environment, the biases, which are indicators of quality differences, must depend on micro-heterogeneities between amoebae (Azhar et al. 2001).

As we will see now, an attractive feature of the individual-selection model is that its outcome is stable with respect to invasion by cheaters.

Cheating

Irrespective of our explanation for the evolution of social behaviour, its persistence implies that sociality is stable with respect to exploitation by an individual that manages to take advantage of its benefits without incurring all of the costs. If such an individual, termed a cheater, arises as the result of a genetic alteration, cheating poses a threat to the long-term stability of the social unit. (One says 'individual' because the implicit assumption is that the cheater is a rare mutant.

The possibility of simultaneous exploitation by a clonal group of invaders is not implausible and its consequences need to be examined.) How well do the arguments that we have made for altruistic behaviour in the CSMs fare in the presence of cheaters?

A 'cheater' can be of many types. It could be an amoeba that (i) invariably differentiates into a spore, never to a stalk cell, and does not influence the behaviour of a co-aggregating partner, (ii) can sporulate and simultaneously manipulate, or exploit, co-aggregating partners so that they contribute more cells than usual to the stalk, or (iii) forms a spore when in a chimaeric aggregate with the wild type 'altruist' but, when developing on its own, can form a spore or a stalk cell - (i) and (ii) are illustrative of what might be called unsophisticated cheating and (iii) represents sophisticated cheating. As we have seen earlier, a model of fitness enables us to infer that group fitness is maximised at a certain ratio of spore cell number to stalk height (Nanjundiah 1985). In a clonal group the outcome can also be interpreted as maximising individual fitness. A cheater that differentiates constitutively into spores and invades a wild type aggregate will always have a higher fitness than the fitness of an average wild type cell. It will therefore increase its representation from one generation to another. But as the cheater genotype spreads it reduces the fitness of the group as a whole because proportioning becomes non-optimal. If it is a constitutive, spores-only cheater, it may drive the group to extinction.

One way for the wild type to hold its own is by trying to ensure that in every generation at least some aggregates are made up of wild type clones. Armstrong (1984) discussed conditions that could either favour or inhibit the evolution of cheaters (in his definition, mutant cells which joined wild type aggregates and invariably sporulated). He performed computer simulations of growth and aggregation on a rectangular grid, using very small aggregate sizes (of the order of 100 cells). The wild type could persist in the long run if, following spore dispersal, germination and feeding, amoebae that were derived from different spores were not in close proximity when starvation set in. The inference was that cheating was unlikely to be successful when aggregate sizes were small (aggregates would tend to be founded by single spores and so would consist of clones, a factor which favours division of labour)

or if amoebae migrated large distances in a random fashion while feeding (in which case, once again, clonal groups were likely to form). This line of reasoning was extended by Matapurkar and Watve (1997) who demonstrated that by juggling with the relevant parameters it was theoretically possible to have a weakly oscillatory coexistence of wild type and cheater genotypes.

How do the ESS models discussed above fare in the presence of a cheater? A sophisticated cheater may try to always allocate relatively fewer cells to the stalk than the wild type does. But, as we have seen, this strategy stops working once the proportion of wild type cells falls below some critical value, because then the ESS logic implies that all wild type cells differentiate into spores. In short, the wild type recovers and eventually coexists stably with the cheater (which is thus a cheater only in name). An unsophisticated, constitutive cheater will invariably increase in frequency until there are so few wild type cells that all cells in the aggregate differentiate into spores. In the long term, there can be one of three outcomes. One, the system crashes because of extinction (if spore masses without a stalk have zero fitness on account of a very low dispersibility). Two, some wild type clones form and develop into aggregates in the next generation, allowing both genotypes to co-exist (a version of the Matapurkar & Watve picture). Three, the two genotypes persist at a stable relative frequency that is affected only by drift (if a spore mass on its own has some chance of dispersal and an average fitness per spore of greater than zero). In every case, if the cheater is also a manipulator, that is, if the cheater induces the wild type to form more stalk cells than otherwise, the long-term fate of the population, assuming that it persists at all, will be biased even more in favour of the cheater.

Contrary to these assertions, Hudson et al. (2002) claim to show that stalk-making (i.e., altruistic) and stalkless (i.e., cheater) strains can coexist stably in fixed proportions and that their coexistence can be explained as an ESS for each genotype. This result is said to follow on the basis of assuming that stalkless clones have low dispersibility and that the advantage of a stalk for dispersal increases very rapidly (i.e., as a concave function) as the number of stalk cells increases from zero. However, their proof depends on other assumptions of uncertain validity. Firstly, as already mentioned, they assume

that dispersibility is a function of the relative proportion of cells allocated to the stalk pathway, not of the actual number of stalk cells. Secondly, the chimaeras that they consider do not consist of (for example) a single wild type cell and a number of cheaters. Rather, the chimaeras are made up of the progeny of a single wild type founder cell and the progeny of one or more cheaters. Thirdly, and most difficult to understand, it is assumed that when they form a common fruiting body, wild type and cheater spores have different dispersibilities, the latter being characteristic of a stalkless fruiting body and the former of a stalked fruiting body.

The individual selection model for altruism rests on the assumption that pre-aggregation amoebae are heterogeneous in a functional sense and that they compete for the chance of sporulating (Atzmony et al. 1997). An amoeba's success in accomplishing this is correlated with where it falls on a scale of quality, in other words on certain aspects (perhaps a large number of aspects) of its phenotype in relation to the rest. Those who fail in the competition are coerced to die and form a stalk. If this is a reasonable picture, the very concept of a cheater loses meaning irrespective of whether there are genetic differences between amoebae or not. To be sure, if a cell arises that has, because of its genotype, a higher quality than the others, its frequency in the population will certainly increase. But, given that small differences between cells are sufficient to discriminate between them, environmental factors will ensure that the qualities of the amoebae belonging to the successful genotype will not be identical. Phenotypic selection will continue to act between the cheater individuals themselves. They too will compete with one another for sporulation and will form a fruiting body in the process. What if the cheater's genotype imposes constitutive sporulation? Within the framework of an individual selection model such a cheater is better termed a loser; its frequency increases in the short term but declines in the long term for the reasons discussed earlier. A plethora of evidence supports the hypothesis that phenotypic heterogeneity is the fundamental cause of differentiation in *D. discoideum* (see later). This lends strength to the belief that there must be something to the individual selection-based analysis of why altruistic behaviour is stable. Hudson et al. (2002) use the phrase 'anticheater adaptations' to describe why cheating

may not pay in *D. discoideum*. Thereby they seem to imply something that arose after the evolution of altruistic behaviour. If the individual-selection model is correct, 'anticheater adaptations' reflect cellular properties correlated with quality that existed prior to the evolution of sociality (and may have become reinforced later).

Experimental Evidence

Behaviour in genetic chimaeras

Genetically diverse amoebae can form chimaeric fruiting bodies in the laboratory (Bonner & Adams 1958, Filosa 1962). They can also contribute disproportionately to the spore population (Buss 1982, Strassmann et al. 2000, Kaushik 2002), raising the possibility of behavioural parasitism. Filosa (1962) studied *D. mucoroides* cultures obtained from a single sorus collected in the wild and maintained in the laboratory for eight years. When sub-clones were generated from them, morphologically different clones were obtained. On mixing the morphological variants, all deriving from a single strain originally, Filosa saw that cells belonging to different clones mixed to form a common slug and fruiting body. When mixed with the wild type too, the variants gave rise to a common fruiting body, this time with the wild type phenotype. In one case a variant that could form slugs but not fruit by itself did so when mixed with the wild type and preferentially formed spores. Buss (1982) obtained eleven *D. mucoroides* strains from natural soil within a very small distance (1mm) of each other. On sub-culturing, one strain gave rise to a normally proportioned fruiting body; another one formed just a mass of spores on the substrate. In other words, the latter was a stalkless form. When the stalkless form began in a small minority (0.1%), competition between the two strains resulted in an increase in its frequency; also, in the presence of the stalkless form, the spore contribution of the stalked form decreased by 20%. Thus 'cheater' strains do occur in nature and can take advantage of the 'honest' cell types. Interestingly, except for the one strain just discussed, other *D. mucoroides* strains that appeared to develop in a manner similar to the standard wild type did not mix at all with this stalkless form. This points to CSMs having defensive mechanisms to guard against invasion by cheaters.

De Angelo et al. (1990) measured the sorus (spore mass) diameter and stalk length in pure and chimaeric fruiting bodies constructed by two

laboratory strains of *D. discoideum*; the spore:stalk ratio so defined was taken to reflect the degree of selfishness exhibited by the cells taken as a whole. They concluded that a mixture of the two strains displayed more selfishness (and so a lesser extent of altruism) than either pure strain by itself. The results appeared to indicate an increased tendency to selfish behaviour in groups of unrelated individuals. This was taken as support for the simple ESS-based prediction (Matsuda & Harada 1990), at least qualitatively. But it turned out later that the results were reversed if a rich nutrient medium was used (Hilson et al. 1994). Using microsatellite DNA probes, Strassmann et al. (2000) found that when different wild genotypes of *D. discoideum* were forced to co-aggregate in the laboratory, sometimes their contributions to the spore population was disproportionate to their representation in the initial mix. In other words, amoebae belonging to one clone could behave more selfishly when mixed with other strains than when by themselves. Recent studies have found no link between genetic heterogeneity and selfish behaviour as reflected by an enhanced spore-forming tendency (Foster et al. 2002, Kaushik 2002).

The potential for 'cheater'-like behaviour was probed in some detail in a particular mutant-wild type chimaera involving a mutant gene (*chtA*) which appeared to convert mutant cells that found themselves in the company of their wild type counterparts into cheaters (Ennis et al. 2000). By itself, the mutant strain did not develop beyond the slug stage. However, when mixed with wild type cells, mutant amoebae differentiated normally and when they did, they contributed disproportionately to the spore population. At the same time, in the presence of the mutant, wild type amoebae displayed an increased tendency towards the stalk pathway. The *chtA* gene encodes a protein harbouring an F box and WD 40 repeats. Such proteins are known to regulate the removal of other proteins by targeting them for the ubiquitination-mediated proteolytic degradation pathway. Considering that the *chtA* strain is unable to differentiate by itself, it may be more appropriate to call the mutant an obligate parasite or an unsophisticated cheater.

The relevance of the findings listed above hinges on the degree of genetic similarity between slime mould populations existing in close proximity in

nature. We discuss this aspect now, and in particular look at how many distinct genotypes might contribute to building one fruiting body.

Genetic Heterogeneity in Nature

Very few studies have been carried out on the degree of genetic relatedness among cellular slime mould populations found in close proximity in the wild (Eisenberg 1976, Kaushik 2002, Fortunato et al. 2003) and, as far as we know, just one on the relatedness between members of an aggregate that forms under reasonably natural conditions (Kaushik 2002, reported below). Eisenberg (1976) found a patchy distribution of CSMs in the soil with up to four species occurring within a 6mm circle. Kaushik (2002), while confirming this finding, extended it by demonstrating that different strains of the same species could be isolated from single particles (ca 1mm) of undisturbed forest soil and moreover, could form genetically heterogeneous aggregates when mixed in the laboratory. Fortunato et al. (2003) report that different clones of *D. discoideum* co-exist over very small spatial scales in nature (within a soil sample weighing 0.2 gm) and can form chimaeric aggregates when brought together in the laboratory. Indirect but compelling evidence indicates that genetic heterogeneity cannot be uncommon in Dictyostelid aggregates under natural conditions (Filosa 1962, Buss 1982, Francis & Eisenberg 1993, Strassmann et al. 2000), and our own observations support this (see below). The experiments described so far show that there is a potential for conflict between coaggregating amoebae in terms of contributing to the next generation. However, direct evidence for the likelihood of conflict under natural conditions was lacking. An experiment to be described now shows that up to 9 genotypes of *Dictyostelium* spp. (tentatively identified as *D. giganteum*) can come together to form a single fruiting body under simulated natural conditions (Kaushik 2002).

The experiment was carried out with isolates derived from the campus of the Indian Institute of Science in the following manner. A sample of soil was picked up from a wooded area on the IISc campus with the broad end of a sterile micropipette tip of 6 mm diameter and tapped lightly on to the surface of a non-nutrient agar plate. Care was taken to see that as little mixing as possible occurred during the process of transferring the soil. In some experiments the soil was left undisturbed on the plate. In other

experiments 50µl of *Klebsiella aerogenes* was gently pipetted on to the soil; this was just sufficient to moisten it. In the undisturbed situation *Dictyostelium* fruiting bodies appeared on plates after 2 to 3 weeks. Presumably bacteria existing in the soil provided whatever food was necessary and the moisture in the agar supplied the required humidity. In those cases in which bacteria were added, fruiting bodies appeared within a week. Spores were isolated from single fruiting bodies, counted and diluted for cloning. Fruiting bodies were generated from individual spores and DNA was extracted from their spores. The DNA extracted was analyzed with the help of the polymerase chain reaction using 4 randomly generated primers (opa01, opa03, opa14 and opa18 from Operon Technology, USA). The results were highly reproducible: a clone yielded the same pattern of bands with a given primer. This made it possible to establish that the genotypes present in the original fruiting body were not all the same.

A typical RAPD profile is shown in figure 2. The spores in this particular fruiting body contained at least 9 genotypes. In general, 6 to 9 genetic variants were present in samples of 10 randomly cloned spores from a single fruiting body. In none of the fruiting bodies were all the examined spores from the same clone. Also the degree of genetic relatedness between genotypes was estimated. For the case shown in figure 2, the average degree of relatedness is 0.73 ± 0.15 (using 4 RAPD primers). Interestingly, the value is comparable to the mean within-species relatedness in *Dictyostelids* isolated from the wild (Kaushik 2002 and unpublished observations), which makes it likely that the co-aggregating amoebae belong to the same species. However, this is only an average value. Spores with as low a level of relatedness as 0.41 (comparable to that between two *Dictyostelid* species) can be found in the same fruiting body. This shows that fruiting bodies obtained under quasi-natural conditions can originate from a large number of genotypes – and indirectly, that there must be a fair amount of mixing during the phase that covers spore germination, cell division and pre-aggregation movement.

Conclusions: Implications for the Evolution of Altruism

We have shown how a number of features of CSM development can be understood as adaptations. One can write down more than one plausible

sequence of steps through which altruistic behaviour could have evolved (Bonner 1982, 2003b, Nanjundiah 1985). For example: beginning with the sporulation of a single amoeba that secreted an extracellular stalk (Step 1; as in *Protostelium*), one can think of aggregation followed by sporulation and the production of an extracellular stalk (Step 2; as in *Fonticula*), a cellular or extracellular stalk (Step 3; as in *D. lacteum*) and a constitutively cellular stalk as in the other *Dictyostelids* that we know of (Step 4). In this picture aggregation (Step 2) could be selectively advantageous even if the aggregating cells belonged to different genotypes. This can also be true of Steps 3 and 4 if the likelihood that an amoeba sporulates after aggregation is finite and the chance that a solitary amoebae survives the period of starvation is much smaller or zero. In the long run, meaning if we average the genotypic fitness over a great many generations, joining an aggregate would benefit the genotype irrespective of whether the aggregate consists of relatives or not. The fitness advantage conferred by Steps 3 or 4 would be higher in the case of aggregates formed by genetically related individuals, because the inclusive fitness of even those individuals that did not succeed in sporulating – in becoming the reproductives – would be positive (the argument resembles the one put forward to explain why lekking takes place in spite of there being gross disparities in individual reproductive successes). In general, any advantage conferred by a group adaptation becomes a bigger advantage in a kin group. Bonner (1967) has put forward two hypotheses to explain the origin of aggregation. On the one hand, it could be a means for bringing together amoebae belonging to different genotypes that had complementary phenotypes. Thus it could have a quasi-sexual function ('recombination on a cellular level'). On the other hand, its function could be to build a multicellular structure.

The monophyly of the CSMs, along with the astonishing degree of convergence that they exhibit, makes it attractive to think in terms of an evolutionary sequence such as we have sketched. Convergence can be thought of as similar strategies that have evolved independently in different lineages. The way the




Figure 2. A sample of soil was picked up from a wooded area on the Indian Institute of Science campus with the broad end of a sterile micropipette tip of 6mm diameter and tapped lightly on to the surface of a non-nutrient agar plate. 50 μ l of a thin suspension of *Klebsiella* aerogenes was added so as to moisten the soil. Spores were isolated after a week from the individual fruiting bodies and cloned. DNA was extracted from the secondary fruiting bodies formed by these clones. RAPD-PCR was carried out using 4 random primers (opa01, opa03, opa14 and opa18 from Operon Technologies Inc. U.S.A). (a) A typical RAPD profile of wild genotypes Wi1.1 to Wi1.10. The extremes show molecular size markers, the control lane has no DNA and C.DNA had DNA from a known *D. giganteum* strain. Note that Wi1.5 and Wi1.9 are identical. (b) Distribution of genetic relatedness between pairs of spores, defined as the number of shared bands relative to the total number of bands, averaged over four primers.

strategies are implemented can be thoughts of as tactics that reflect the unique evolutionary history of each species. The variety of chemical attractants in Dictyostelids is an example of different tactics for implementing the common strategy of aggregation. The use of a particular tactic may be attributable to an intrinsic fitness advantage or to constraints imposed by development (Bonner 1982), besides those imposed by physics and chemistry (Newman 1992). A priori, the distinction is not an easy one to make.

Consider the use of cyclic AMP as a chemoattractant. To account for it, one can come up with at least three different evolutionary explanations. (i) It could have been chosen by accident, as a message from one cell that says 'come here' to another cell. The reason for sending the message is that starvation has set in and joining a group has become advantageous; the reason for responding to the message is the same. The message is entirely symbolic and could have been conveyed just as well by any other molecule so long as the appropriate machinery for signaling and responding developed in parallel. (ii) A rise in cAMP could be something that happens in a starved cell (for example, as a correlate of sugar deprivation). Therefore a cell that sensed external cAMP would sense an external correlate of an internal state that implied starvation; the signal would be meaningful, so to speak, and not

merely symbolic. (iii) The cAMP molecule is energetically expensive to make, a fact based on chemistry and thermodynamics. The reason why a cell uses *it* as a signal, and not any other molecule, may be because – by virtue of its being costly – it is an honest signal and therefore a reliable indicator of the quality of the sender (Zahavi 1977). This would imply that the other known chemoattractants must also be molecules that convey honest signals of cellular quality. Note that the first two explanations are the more attractive if the sender and receiver are clonal relatives and operate as a functional whole. Then the evolution of cAMP as an agent of intercellular communication would involve kin selection. But as far as the third explanation goes, the sender and receiver could be independent units of selection.

One can extend this way of categorizing alternative explanations to other intercellular signals. Further, one can ask how much the feature of CSM development of interest to us, division of labour, depends on intercellular interactions and how much can be explained by intrinsic properties of cells. This is an old and well-studied question (reviewed in Nanjundiah & Saran 1992) and the short answer is that single-cell traits and cell-cell interactions are equally important. The fact that a pre-aggregation pecking order exists at all shows that it makes sense to think of division of labour in

terms of cellular heterogeneity, while the fact that isolates of pre-stalk or pre-spore cells can transdifferentiate and make up the missing cell type (Raper 1940) shows that it also makes sense to think of division of labour as a form of collective behaviour. As far as evolutionary hypotheses go, these findings mean that both group selection and individual selection could have been at work. The evidence for the occurrence of genetically heterogeneous aggregations in nature is strong and it is a good bet that amoebae of different genotypes encounter each other fairly regularly. However, sorting out between different clones is common (Kaushik 2002), implying that amoebae prefer to stay in physical proximity to their kin. Foster et al. (2002) discovered that a slug consisting of two clones migrates more efficiently than a slug consisting of a single clone and interpreted this as evidence for an advantage of chimaera formation. But because their chimaeric slugs were twice as large as the clonal slugs, what they really found was the well-known size effect (Bonner et al. 1953); indeed, Foster et al. saw that in a given time chimaeric slugs traveled a shorter distance than same-sized slugs of a single genotype. In spite of abundant evidence that deficient clones can exhibit functional complementation when mixed (Sussman 1954, Sussman & Lee 1955), i.e., recover the wild type phenotype, there is no evidence that chimaera formation is generally favoured under natural conditions.

If one can show that amoebae belonging to the same species can distinguish between each other on the basis of kinship alone and shape their behaviour accordingly, it would favour the kin selection hypothesis. Queller et al. (2003) claim that they have found a protein in *D. discoideum* that enables kin to be recognized as such. This is the contact site A (csA) molecule, which mediates homophilic adhesion. A cell that lacks csA should be unable to adhere as efficiently to another cell (either *csA* or wild type) as one wild type cell to another. The observation is that in wild type- *csA* mixtures that are made to develop on soil, the wild type cells sporulate preferentially. The authors interpret this as the consequence of selective altruism by wild type cells towards one another (mediated in this case by stronger cohesion). There are three problems with this interpretation. Firstly, the outcome is reversed when the

experiments are carried out on agar – paradoxically, in mixtures that develop on agar, *csA* amoebae are *more* efficient than the wild type. Secondly, the most likely explanation for the presence of the *csA* protein is that it stabilizes aggregates by helping cells to adhere to each other. In the absence of evidence bearing on the degree of variation in the *csA* protein between genotypes and the consequences of such variation, it is difficult to conclude that it is made use of as a mark of identity, a ‘green beard’, that aids amoebae in directing altruistic acts towards clonal relatives.

The third problem is more general and extends to designations such as ‘cheater’ and ‘green beard’. A large number of *D. discoideum* mutations, perhaps the majority, show aberrant differentiation in the sense that when they form fruiting bodies at all, they make disproportionately more spores or more stalk cells than the wild type (independent of the phenotype that has been used to identify the presence of the mutation). This is exactly what is to be expected if genetic networks are complex and genes have pleiotropic effects. In a mixture with the wild type, assuming that it mixes well, such a mutant would be expected to influence the behaviour of wild type cells. Depending on the nature of the influence, it may form more spores or fewer spores than expected. For example, on mixing a slight majority of stalkless (variant) *D. mucoroides* cells with the wild type, Filosa (1962) found that the spore proportions were overwhelmingly biased in favour of the mutant. Should one conclude from this observation alone that (other than in a metaphorical sense) the mutant is a cheater, or exhibits altruistic behaviour towards its own kind? If the mutant had come off worse in sporulation efficiency, would one be justified in saying that it was being manipulated by the wild type? The use of these terms carries with it the suggestion that the evolutionary function of the gene in question is reflected in just the role implied by either term, an inference which may be unwarranted.

Are there any general conclusions that can be drawn on the basis of what we have discussed in this review? First and foremost, it is clear that the correct way to look at division of labour and altruism in the CSMs is to acknowledge that both selection at the level of the single cell and selection at the level of the group (including kin groups) must

Table 2. Illustrative outcomes to be expected under various models when amoebae belonging to different genotypes co-aggregate. 'Kin selection' also involves intercellular interactions; the predictions refer to the ESS model. 'Complex interactions' allow for many more outcomes than indicated here. (See text for details.)

Hypothesis	Predictions	Expected Phenotype
<i>Kin selection (ESS)</i>		
(a) Strains will tend to be more altruistic towards each other when they are closely related.	(a) Cells will try to maximize the over all fitness of a clonal fruiting body.	(a) Under certain assumptions, constant proportions between cell types
(b) Unrelated strains will behave selfishly when they co-aggregate	(b) in chimaeras, cells from different strains will optimize their fitnesses separately.	(b) Larger spore to stalk ratio in chimaeric fruiting bodies as compared to pure ones.
<i>No interaction</i>		
(a) In chimaeras, co-aggregating strains behave as they would in the absence of the other strain.	(a) The strain that has a higher spore to stalk ratio when on its own contributes relatively more to the spores in the chimaera.	(a) The spore to stalk ratio of a chimaeric fruiting body will be higher than that of the strain with the smaller ratio and smaller than that of the one with the higher ratio.
<i>Complex interactions</i>		
(a) One strain negatively regulates the spore contribution by the other strain (by inducing the latter to form stalk)	(a) Relative to its initial frequency, the majority of the spores are contributed by one strain.	(a) Spore to stalk ratio may vary depending on the contribution of spores by each strain.
(b) Both strains try to contribute more to the spore population.	(b) Both strains contribute more than their normal share of spores.	(b) Larger spore to stalk ratio in chimaeric fruiting bodies than in pure ones.
(c) One strain interferes in the development of the other strain.	(c) Only one of the strains develops normally.	(c) Fruiting bodies of only one strain are formed.

have played a role. Given that many clones can participate in building a single fruiting body, division of labour between the amoebae that constitute a single social unit is unlikely to be based solely on kin selection. Rather, it may depend on a combination of cooperation between identical genotypes and conflict between non-identical ones (Kawli & Kaushik 2001). In the latter situation, the altruism exhibited by stalk cells may be the result of individual-level selection that makes it advantageous for a starving amoeba to participate in forming a fruiting body at the risk of dying in the attempt. Experimentally, there are ways in which one can try to distinguish between different models for altruism (table 2). Many, or even all, of the outcomes listed may in fact occur.

One needs to know more about the natural ecology of the CSMs. Equally necessary is an appreciation of just what the evolutionary implications are of the detailed information building up from laboratory studies. There are many important questions for which answers are as yet not available. (1) What accounts for the presence of so many kinds of SMs in what seems superficially to be the same environment? All CSMs

have life history stages of approximately the same size, utilize the same food and have very similar life cycles. One assumes that they occupy different niches; but what are the niches? Horn (1971) made a beginning by showing that co-occurring species exhibit feeding preferences between different bacteria. (2) What biotic interactions with other organisms (besides that of predator and prey) are significant for CSMs? Ellison and Buss (1983) discovered a naturally occurring strain of *D. mucoroides* that was induced to go through normal morphogenesis by a diffusible substance released by the fungus *Mucor hiemalis* which was isolated together with the CSM. Bhavani Prasanna et al. (1998) found that the development of *D. discoideum* is inhibited by the isoflavonoid phytoalexins. They hypothesized that this, taken together with the observation CSMs are common in the rhizosphere (Agnihotrudu 1956), might mean that leguminous plants manipulate CSM amoebae so that they remain in a feeding state and get rid of potentially harmful bacteria. (3) What are the relative frequencies of clonal and non-clonal aggregates in nature? And when aggregates are non-clonal, what is the degree of genetic overlap between amoebae? (4) Can amoebae discriminate between kin and

non-kin? (5) What is the relative importance of genetic and epigenetic influences bearing on the likelihood that amoebae belonging to different strains associate with each other? For example, given a choice between fruiting in a combination with genetically different cells (with which they have been raised together) and genetically identical cells (which were grown separately), would amoebae display a preference? As a continuation of this, can one make a distinction between genotypic and phenotypic components of quality among wild isolates? (6) From laboratory experiments we know that division of labour can be found in clonal groups and take it for granted that genetic differences are irrelevant. Is this true of the wild as well? Julian et al. (2002) and Volny and Gordon (2002) have shown that there can be a genetic basis for division of labour in ants; in one case in a hybrid zone within which two otherwise distinct genotypes coexist sympatrically and interbreed, and in the other case based on a presumed difference between heterozygotes and homozygotes at a single locus (so far identified as a microsatellite locus). If chimaeric aggregates are common in nature, it is important to see whether there are systematic genetic differences between the cells that sporulate and those that form stalk cells in natural aggregates. It could be, that in some species the necessary genetic polymorphism is maintained by sexual reproduction as in the ant case. (7) In many species a starved CSM amoeba seems to have the option of either leading a solitary existence as a microcyst or entering a social phase by aggregating with others. On what basis is the option exercised – what factors tilt the balance in favour of solitary behavior vis-à-vis sociality? If this question can be explored under controlled conditions we would be able to speculate usefully about the evolutionary origins of group living and division of labour.

The development of *Dictyostelium discoideum* is being intensively studied at the level of genes and gene products. There is a seemingly unending stream of information, most of it fascinating, and no doubt essential if one wants to decipher the details of cell behaviour (Kessin 2001). But it is not easy to make out how much of the information is of broader significance for understanding the

evolutionary forces behind the *D. discoideum* life cycle, besides the life cycles of other CSMs. Sadly, the extreme concentration on one species has meant that the comparative biology of CSM development is by and large a neglected subject. Still, one can ask, what evolutionary insights can be expected from developmental studies? From the individual selection versus kin selection standpoints it would be important to know how far one can push the link between pre-aggregation heterogeneity and cell fate. Can one identify a simple basis for the assignment of relative quality? Does it form a transitive ('linear') hierarchy? Can it be looked at as if it was a quasi-continuous input (e.g., a variation in cellular calcium) that gets resolved into two outputs, that is, two cell types (Nanjundiah & Saran 1992)? There are pointers. Durston's (1976) experiments showed that slug tip formation could be explained in terms of an inhibitory gradient. Lokeshwar and Nanjundiah (1983) found that as judged by the time taken for a tip to regenerate after it was amputated, the pre-spore region exhibited a steady variation – the time was taken quite little when regeneration occurred towards the previous anterior end and it kept increasing as one moved posteriorly. Haberstroh and Firtel (1990), working with different promoter constructs of the *cotC* gene, found results consistent with a spatially varying signal that activated the promoter differentially in the front and back of the pre-spore zone. In a recent study Kibler et al. (2003) find that cells that are mutated in the *comD* gene can form spores when they are derived from the posterior portion of the pre-spore zone in slugs but not from the anterior portion, as if there were at least two types of pre-spore cells.

Both kin- and group selection presuppose the existence of group-level adaptations that are not trivial extensions of individual adaptations. One can search for the genes that underlie group adaptations by looking within the class of 'non-cell autonomous' mutations. In such mutations the genotype of one cell influences the phenotype of another, a necessary but not sufficient condition if the gene product contributes to a trait that benefits the group. In complex organisms it may be difficult to find genes that affect only the cells in which they are active. But the prospects are better

in a simple organism, especially one in which the communal phase is separated in time from the solitary phase. Among the genes that one is looking for would be those whose direct or indirect products are involved in intercellular signalling pathways, and their presence can be inferred by phenotypic complementation experiments (Sussman 1954, Filosa 1962). One can narrow the search further and ask, is there genetic evidence from which one can infer that the activity of a gene in a pre-stalk cell promotes the spore pathway in a pre-spore cell? Consider these examples. SDF-2 is a small peptide produced by pre-stalk cells of *D. discoideum* that helps in the encapsulation of pre-spore cells, an essential step in spore maturation (Anjard et al. 1998). A mutation in the gene that encodes the regulatory subunit of the cAMP-dependent protein kinase and is expressed only in pre-stalk cells is compatible with spore production if mutant and wild type cells are

mixed, but there is no sporulation if the mutant is allowed to develop on its own (Harwood et al. 1992). The *comD* gene appears to be active in prestalk cells but influences the differentiation of pre-spore cells (Kibler et al. 2003). Here are signs that pre-stalk cells are needed for pre-spore cells, and it is hard to see how that could be the case unless the evolution of the two classes was favoured as a combined unit. This brings us back to the point that both individual and group (kin) selection must have been important in the evolutionary past of the CSMs.

Acknowledgements

Anonymous reviewers made extremely useful comments on an earlier version of the paper their contribution is gratefully acknowledged. Thanks are due to CSIR for financial support. Part of this work is based on a Ph.D. thesis submitted by SK to the Indian Institute of Science.

References

- Aesop 1954 *Fables of Aesop* (Penguin Books)
- Agnihotrudu V 1956 Occurrence of Dictyosteliaceae in the rhizosphere of plants in Southern India; *Experientia* **12** 149-150
- Anjard C, Zeng C, Loomis W F and Nellen W 1998 Signal transduction pathways leading to spore differentiation in *Dictyostelium discoideum*; *Dev. Biol.* **193** 146-155
- Armstrong D P 1984 Why don't cellular slime molds cheat? *J. Theor. Biol.* **45** 119-129
- Atzmony D, Zahavi A and Nanjundiah V 1997 Altruistic behaviour in *Dictyostelium discoideum* explained on the basis of individual selection; *Curr. Sci.* **72** 142-145
- Azhar M, Kennady P K, Pande G, Espiritu M, Holloman W, Brazil D, Gomer R H and Nanjundiah V 2001 Cell cycle phase, cellular Ca²⁺ and development in *Dictyostelium discoideum*; *Int. J. Dev. Biol.* **45** 405-413
- Baldauf S L and Doolittle W F 1997 Origin and evolution of the slime molds (Mycetozoa); *Proc. Natl. Acad. Sci. USA.* **94** 12007-12012
- _____, Roger A J, Wenk-Siefert I and Doolittle W F 2000 A Kingdom-Level Phylogeny of Eukaryotes Based on Combined Protein Data; *Science* **290** 970-977
- Baskar R, Chhabra P, Mascarenhas P and Nanjundiah V 2000 A cell type-specific effect of calcium on pattern formation and differentiation in *Dictyostelium discoideum*; *Int. J. Dev. Biol.* **44** 491-498
- Bell G 1982 *The Masterpiece of Nature: The Evolution and Genetics of Sexuality* (California: University of California Press)
- Bhavani Prasanna T, Vairamani M and Kasbekar D P 1998 Effects of pisatin on *Dictyostelium discoideum*: its relationship to inducible resistance to nystatin and extension to other isoflavonoid phytoalexins; *Arch. Microbiol.* **170** 309-312
- Blaskovics J C and Raper K B 1957 Encystment Stages of *Dictyostelium*; *Biol. Bull.* **113** 58-88
- Bonner J T 1947 Evidence for the formation of cell aggregates by chemotaxis in the development of the slime mold *Dictyostelium discoideum*; *J. Exptl. Zool.* **106** 1-26
- _____, 1950 Observations on polarity in the slime mold *Dictyostelium discoideum*; *Biol. Bull.* **99** 143-151
- _____, 1952 The pattern of differentiation in amoeboid slime molds; *Am. Nat.* **86** 79-89
- _____, 1958 *The Evolution of Development*; (UK: Cambridge Univ. Press)
- _____, 1959 Evidence for the sorting out of cells in the development of the cellular slime mold; *Proc. Natl. Acad. Sci. USA.* **45** 379-384
- _____, 1965 *Size and Cycle*; (Princeton: Princeton Univ. Press)
- Bonner J T 1967 *The Cellular Slime Molds.* 2nd ed. (Princeton: Princeton: Univ. Press)
- _____, 1982 Evolutionary strategies and developmental constraints in the cellular slime molds; *Am. Nat.* **119** 530-552
- _____, 1991 *Researches on Cellular Slime Moulds*; (Bangalore: Indian Academy of Sciences)

- _____. 1992 The fate of a cell is a function of its position and *vice-versa*; *J. Biosci.* **17** 95-114
- _____. 1996 *Sixty years of Biology*; (Princeton: Princeton Univ. Press)
- _____. 1998 A way of following individual cells in the migrating slugs of *Dictyostelium discoideum*; *Proc. Natl. Acad. Sci. USA* **95** 9355-9359
- _____. 2000 *First Signals. The Evolution of Multicellular Development*; (Princeton: Princeton Univ. Press)
- _____. 2003a On the origin of differentiation; *J. Biosci.* **28** 523-528
- _____. 2003b Evolution of development in the cellular slime molds; *Evolution & Development* **5** 305-313
- _____, Clarke W W jr., Neely C L jr. and Slifkin M K 1950 The orientation to light and the extremely sensitive orientation to temperature gradients in the slime mold *Dictyostelium discoideum*; *J. Cellular Comp. Physiol.* **36** 149-158
- _____, Koontz P G jr. and Paton D 1953 Size in relation to the rate of migration in the cellular slime mold *Dictyostelium discoideum*; *Mycologia* **45** 235-240
- _____ and Adams M S 1958 Cell mixtures of different species and strains of cellular slime moulds; *J. Embryol. Exp. Morphol.* **6** 346-356
- _____, Joyner B D, Moore A A, Suthers H B and Swanson J A 1985 Successive asexual life cycles of starved amoebae in the cellular slime mould, *Dictyostelium mucoroides* var. *stoloniferum*; *J. Cell Sci.* **76** 23-30
- _____, Chiang A, Lee J and Suthers H B 1988 The possible role of ammonia in phototaxis of migrating slugs of *Dictyostelium discoideum*; *Proc. Natl. Acad. Sci. USA* **85** 3885-3887
- _____, Har D and Suthers H B 1989 Ammonia and thermotaxis: Further evidence for a central role of ammonia in the directed cell mass movements of *Dictyostelium discoideum*; *Proc. Natl. Acad. Sci. USA* **86** 2733-2736
- Bozzone D M and Bonner J T 1982 Macrocyt formation in *Dictyostelium discoideum*: mating or selfing? *J. Exptl. Zool.* **220** 391-394
- Brock D A and Gomer R H 1999 A cell-counting factor regulating structure size in *Dictyostelium*; *Genes Dev.* **13** 1960-1969
- Buss L W 1982 Somatic cell parasitism and the evolution of somatic tissue compatibility; *Proc. Natl. Acad. Sci. USA*, **79** 5337-5341
- Buss L W 1999 Slime molds, ascidians and the utility of evolutionary theory; *Proc. Natl. Acad. Sci. USA*, **96** 8801-8803
- Campbell A, Mrázek J and Karlin S 1999 Genome signature comparisons among prokaryote, plasmid and mitochondrial DNA; *Proc. Natl. Acad. Sci. USA* **96** 9184-9189
- Clark M, Francis D and Eisenberg R 1973 Mating types in cellular slime molds; *Bioch. Biophys. Res. Com.* **52** 672-678
- Crespi B J 2001 The evolution of social behavior in microorganisms; *Trends Ecol. Evol.* **16** 178-183
- Dawkins R 1982 *The Extended Phenotype*; (Oxford: Oxford University Press)
- DeAngelo M J, Kish V M and Kolmes S A 1990 Altruism, selfishness and heterocytosis in cellular slime molds; *Ethol. Ecol. Evol.* **2** 439-443
- Dormann D, Siegert F and Weijer C J 1996 Analysis of cell movement during the culmination phase of *Dictyostelium* development; *Development* **122** 761-769
- _____, Vasiev B and Weijer C J 1998 Propagating waves control *Dictyostelium discoideum* morphogenesis; *Biophys. Comm.* **71** 22-35
- Durston A J 1976 Tip formation is regulated by an inhibitory gradient in the *Dictyostelium discoideum* slug *Nature* **263** 126-129
- Eisenberg R M 1976 Two-dimensional microdistribution of cellular slime molds in forest soil; *Ecology* **57** 380-384
- Ellison A M and Buss L W 1983 A naturally occurring developmental synergism between the cellular slime mold, *Dictyostelium mucoroides* and the fungus, *Mucor hiemalis*; *Amer. J. Bot.* **70** 298-302
- Ennis H L, Dao D N, Pukatzi S U and Kessin R H 2000 *Dictyostelium* amoebae lacking an F-box protein form spores rather than stalk in chimeras with wild type; *Proc. Natl. Acad. Sci. USA* **97** 3292-3297
- Erdős G W, Raper K B and Vogen L K 1975 Sexuality in the Cellular Slime Mold *Dictyostelium giganteum*; *Proc. Nat. Acad. Sci. USA* **72** 970-973
- Evans W B, Hughes J E and Welker D L 1988 The use of DNA probes for taxonomic study of *Dictyostelium* wild isolates; *Genetics* **119** 561-569
- Filosa M F 1962 Heterocytosis in cellular slime molds; *Am. Nat.* **96** 79-91
- Fortunato A, Strassmann J E, Santorelli L and Queller D C 2003 Co-occurrence in nature of different clones of the social amoeba, *Dictyostelium discoideum*; *Mol. Ecol.* **12** 1031-1038
- Foster K R, Fortunato A, Strassmann J E and Queller D C 2002 The costs and benefits of being a chimera; *Proc. R. Soc. Lond. B* **269** 2357-2362
- Francis D 1998 High frequency recombination during the sexual cycle of *Dictyostelium discoideum*; *Genetics* **148** 1829-1832
- Francis D and Eisenberg R 1993 Genetic structure of a natural population of *Dictyostelium discoideum*, a cellular slime mold; *Mol. Ecol.* **2** 385-391
- Gadagkar R, Vinutha C, Shanubhogue A and Gore A P 1988 Pre-imaginal biasing of caste in a primitively eusocial insect; *Proc. Soci. Lond. B* **233** 175-189
- _____. 1990 Origin and evolution of eusociality: A perspective from studying primitively eusocial wasps; *J. Genetics* **69** 113-125

- _____ and Bonner J T 1994 Social insects and social amoebae; *J. Biosci.* **19** 219-245
- Gerisch G and Wick U 1975 Intracellular oscillations and release of cyclic AMP from *Dictyostelium* cells; *Biochem. Biophys. Res. Comm.* **65** 365-370
- Glöckner et al. 2002 Sequence and analysis of chromosome 2 of *Dictyostelium discoideum*; *Nature* **418** 79-85
- Gregg J H 1971 Developmental potential of isolated *Dictyostelium myxamoebae*; *Dev. Biol.* **26** 476-485
- Gross J D 1994 Developmental decisions in *Dictyostelium discoideum*; *Microbiol. Rev.* **58** 330-351
- Haberstroh L and Firtel R A 1990 A spatial gradient of expression of a cAMP-regulated prespore cell-type specific gene in *Dictyostelium discoideum*; *Genes Dev.* **4** 5996-6125
- Hagiwara H 1989 *The Taxonomic Study of Japanese Dictyostelid Cellular Slime Molds*; (Tokoyo: National Science Museum)
- Hamilton W D 1964 The genetical evolution of social behaviour (I & II); *J. Theor. Biol.* **7** 1-16, 17-52
- Harwood A J, Hopper N A, Simon N M, Driscoll D M, Veron M and Williams J G 1992 Culmination in *Dictyostelium* is regulated by the cAMP-dependent protein kinase; *Cell* **69** 615-624
- Hilson J A, Kolmes S A and Nellis L F 1994 Fruiting body architecture, spore capsule contents, selfishness, and heterocytosis in the cellular slime mold, *Dictyostelium discoideum*; *Ethol. Ecol. Evol.* **6** 529-535
- Horn E G 1971 Food competition among the cellular slime molds (Acrasidae); *Ecology* **52** 475-484
- Hudson R E, Aukema J E, Rispe C and Roze D 2002 Altruism, cheating and anticheater adaptations in cellular slime molds; *Amer. Natur.* **160** 31-43
- Inouye K and Takeuchi I 1980 Motive force of the migrating pseudoplasmodium of the cellular slime mould *Dictyostelium discoideum*; *J. Cell Sci.* **41** 53-64
- Julian G E, Fewell J H, Gadau J, Johnson R A and Larrabee D 2002 Genetic determination of the queen caste in an ant hybrid zone; *Proc. Natl. Acad. Sci. USA* **99** 8157-8160
- Kaiser D 1986 Control of multicellular development: *Dictyostelium* and *Myxococcus*; *Ann. Rev. Genet.* **20** 539-566
- _____ 1993 Roland Thaxter's legacy and the origins of multicellular development; *Genetics* **135** 249-254
- Katz E R and Sussman M 1972 Parasexual recombination in *Dictyostelium discoideum*: selection of stable heterozygotes and stable haploid segregants; *Proc. Natl. Acad. Sci. USA.* **69** 495-498
- Kaushik S 2002 Genetic heterogeneity and social behaviour in cellular slime moulds;. Ph. D. thesis, Indian Institute of Science, Bangalore
- Kawli T S and Kaushik S 2001 Cell fate choice and social evolution in *Dictyostelium discoideum*: Interplay of morphogens and heterogeneities; *J. Biosciences* **26** 130-133
- Kay R R, Large S, Traynor D and Nayler O 1993 A localized differentiation-inducing-factor sink in the front of the *Dictyostelium* slug; *Proc. Natl. Acad. Sci. USA.* **90** 487-491
- _____ 1997 DIF signaling; in *Dictyostelium - A Model System for Cell and Developmental Biology* pp279-292 eds. Y Maeda, K Inouye and I Takeuchi (Tokoyo: Universal Academy Press)
- Kessin R H 2001 *Dictyostelium - Evolution, Cell Biology, and the Development of Multicellularity* (Cambridge, UK: Cambridge Univ. Press)
- _____, Gundersen G G, Zaydfudim V, Grimson M and Blanton R L 1996 How cellular slime molds evade nematodes. *Proc. Natl. Acad. Sci. USA* **93** 4857-4861
- Kibler K, Nguyen T-L, Svetz J, Driessche N V, Ibarra M, Thompson C, Shaw C and Shaulsky G 2003 A novel developmental mechanism in *Dictyostelium* revealed in a screen for communication mutants; *Dev. Biol.* **259** 193-208
- Kirk D L 1998 *Volvox: Molecular-Genetic Origins of Multicellularity and Cell Differentiation*; (UK: Cambridge Univ. Press)
- Kuserk F T 1980 The relationship between cellular slime molds and bacteria in forest soil; *Ecology* **61** 1474-1485
- Landau L D and Lifshitz E M 1970 *Theory of Elasticity* 2nd ed. (Oxford : Pergamon Press)
- Leach C K, Ashworth J M and Garrod D R 1973 Cell sorting out during the differentiation of mixtures of metabolically distinct populations of *Dictyostelium discoideum*; *J. Embryol. Exp. Morph.* **29** 647-661
- Lokeshwar B L and Nanjundiah V 1983 Tip Regeneration and Positional Information in the Slug of *Dictyostelium discoideum*; *J. Embryol. exp. Morphol.* **73** 151-162
- Loomis W F and Kuspa A 1997 The Genome of *Dictyostelium discoideum*; in *Dictyostelium – A Model System for Cell and Developmental Biology* pp 15-32 eds Y Maeda, K Inouye and I Takeuchi (Tokyo: Yamada Science Foundation and Universal Academic Press)
- _____ and Smith D W 1995 Consensus phylogeny of *Dictyostelium*; *Experientia.* **51** 1110-1115
- Matapurkar A K and Watve M G 1997 Altruist-cheater dynamics in *Dictyostelium*: aggregated distribution gives stable oscillations; *Am. Natur.* **150** 790-797
- Matsuda H and Harada Y 1990 Evolutionary stable stalk to spore ratio in cellular slime molds and the law of equalization in net incomes; *J. Theor. Biol.* **147** 329-344
- Maynard Smith J 1982 *Evolution and the Theory of Games* (Princeton: Princeton Univ. Press)
- McDonald S A and Durston A J 1984 The cell-cycle and sorting in *Dictyostelium discoideum*; *J. Cell. Sci.* **66** 195-204

- Michod R E and Nedelcu A M 2003 Cooperation and conflict during the unicellular multicellular and prokaryotic-eukaryotic transitions; in *Evolution: from Molecules to Ecosystems* pp. 195-208 eds A Moya and E Font (Oxford: Oxford University Press)
- Mizutani A, Hagiwara H and Yanagisawa K 1990 A killer factor produced by the cellular slime mold *Polysphondylium pallidum*; *Arch. Microbiol.* **153** 413-416
- _____ and Yanagisawa K 1990 Cell division inhibitor produced by killer strain of cellular slime mold *Polysphondylium pallidum*; *Dev. Growth Differ.* **32** 397-402
- Mutzel R 1991 Cellular slime molds: why and how to become pluricellular; *Bull. Inst. Pasteur.* **89** 51-58
- Nanjundiah V 1973 Chemotaxis, signal relaying, and aggregation morphology; *J. Theor. Biol.* **42** 63-105
- _____ 1985 The evolution of communication and social behaviour in *Dictyostelium discoideum*; *Proc. Indian Acad. Sci. (Animal Sci.)* **94** 639-653
- _____ 1988 Periodic stimuli are more successful than randomly spaced ones for inducing development in *Dictyostelium discoideum*; *Biosci. Reports* **8** 571-577
- _____ and Bhogle A S 1995 The precision of regulation in *Dictyostelium discoideum*: implications for cell-type proportioning in the absence of spatial pattern; *Indian J. Biochem. Biophys.* **32** 404-416
- _____ and Lokeshwar B L 1984 Biological Patterns and the Problem of Regulation; in *The Living State* pp 308-322 ed. R K Mishra (Delhi: Wiley-Eastern)
- _____ and Saran S 1992 The determination of spatial pattern in *Dictyostelium discoideum*; *J. Biosci.* **17** 353-394
- Newman S A 1992 Generic physical mechanisms of morphogenesis and pattern formation as determinants in the evolution of multicellular organization; *J. Biosci.* **17** 193-215
- O'Day D H 1979 Aggregation during sexual development in *Dictyostelium discoideum*; *Can. J. Microbiol.* **25** 1416-1426
- _____ and Lewis K E 1975 Diffusible mating type factors induce macrocyst development in *Dictyostelium discoideum*; *Nature Lond.* **754** 431-432
- Okuwa T, Katayama T, Takano A, Kodaira K and Yasukawa H 2001 Two cell-counting factors regulate the aggregate size of the cellular slime mould *Dictyostelium discoideum*; *Develop. Growth Differ.* **43** 735-744
- Olive L S 1975 *The Mycetozoans* (New York: Academic Press)
- Parent C and Devreotes P N 1999 A cell's sense of direction; *Science* **284** 765-770
- Ponte E, Bracco E, Faix J and Bozzaro S 1998 Detection of subtle phenotypes: the case of the cell adhesion molecule csA in *Dictyostelium*; *Proc. Natl. Acad. Sci. USA* **95** 9360-9365
- Queller D C, Ponte E, Bozzaro S and Strassmann J E 2003 Single-gene greenbeard effects in the social amoeba *Dictyostelium discoideum*; *Science* **299** 105-106
- Raper K B 1940 Pseudoplasmodium formation and organization in *Dictyostelium discoideum*; *J. Elisha Mitchell Sci. Soc.* **56** 241-282
- _____ 1984 *The Dictyostelids* (Princeton: Princeton University Press)
- Roisin-Bouffay C, Jang W, Caprette D R and Gomer R H 2000 A precise group size in *Dictyostelium* is generated by a cell-counting factor modulating cell-cell adhesion; *Mol. Cell* **6** 953-959
- Shaffer B M 1961 Species differences in the aggregation of the Acrasieae. *Recent Advances in Botany; Proc. 9th Int. Bot Congress* pp 294-298 (Toronto: Univ. Toronto Press)
- _____ 1962 The Acrasina; *Adv. Morphogenesis* **2** 109-182
- Shimomura O, Suthers H L B and Bonner J T 1982 Chemical identity of the acrasin of the cellular slime mould *Polysphondylium violaceum*; *Proc. Natl. Acad. Sci. USA* **79** 7376-7379
- Singh B N 1947 Studies on soil Acrasieae. 2. The active life of species of *Dictyostelium* in soil and the influence thereon of soil moisture and bacterial food; *J. Gen. Microbiol.* **1** 361-367
- Solari C A, Nedelcu A M and Michod R E 2003 Fitness and complexity in volvocalean green algae; in *Computational Synthesis: from Basic Building Blocks to High-level Functionality* eds H Lipson, E K Antonsson and J R Koza (USA, Stanford: AAAI Press)
- Stoner D S, Rinkevich B and Weissman I L 1999 Heritable germ and somatic cell lineage competitions in chimeric colonial protochordates; *Proc. Natl. Acad. Sci. USA.* **96** 9148-9153
- Strassmann J E, Zhu Y and Queller D C 2000 Altruism and social cheating in the social amoeba *Dictyostelium discoideum*; *Nature* **408** 965-967
- _____, Zhuy and Queller D C 2000 Altruism and social cheating in the social amoeba *Dictyostelium discoideum*; *Nature* **408** 965-967
- Sussman M 1954 Synergetic and antagonistic interactions between morphogenetically deficient variants of the slime mould *Dictyostelium discoideum*; *J. Gen. Microbiol.* **10** 110-120
- _____ and Lee F 1955 Interactions among variant and wild-type strains of cellular slime mold across thin agar membranes; *Proc. Natl. Acad. Sci. USA* **41** 70-78
- Suthers H B 1985 Ground-feeding migratory songbirds as cellular slime mold distribution vectors; *Oecologia (Berlin)* **65** 526-530

- Takeuchi I, Noce T and Tasaka M 1986 Prestalk and prespore differentiation during development of *Dictyostelium discoideum*; *Curr.Top. Dev. Biol.* **20** 243-256
- Thompson C R and Kay R R 2000a Cell-fate choice in *Dictyostelium*: intrinsic biases modulate sensitivity to DIF signalling; *Dev. Biol.* **227** 56-64
- _____ and _____ 2000b The role of DIF-1 signalling in *Dictyostelium discoideum*; *Mol. Cell* **6** 1509-1514
- Tomchik K J and Devreotes P N 1981 Adenosine 3',5'-monophosphate waves in *Dictyostelium discoideum*: a demonstration by isotope dilution-fluorography technique; *Science* **212** 443-446
- Traub F and Hohl R 1976 A new concept for the taxonomy of the family *Dictyosteliaceae* (cellular slime molds); *Am. J. Bot.* **63** 664-672
- Trivers R L 1971 The evolution of reciprocal altruism; *Q. Rev. Biol.* **46** 35-57
- Urushihara H 1992 Sexual development of cellular slime mold; *Devel. Growth Differ.* **34** 1-8
- _____ 1997 Cell recognition in the sexual development of *Dictyostelium discoideum*; in *Dictyostelium - A Model System for Cell and Developmental Biology*, pp. 137-142 Y Maeda, K Inouye and I Takeuchi. (Tokyo: Universal Academy Press)
- van Haastert P J M, de Wit R J W, Grijpma Y and Konijn T M 1982 Identification of a pterin as the acrasin of the cellular slime mold *Dictyostelium lacteum*; *Proc. Natl. Acad. Sci. USA* **79** 6270-6274
- Velicer G J 2003 Social strife in the microbial world; *Trends in Microbiology* **11** 330-337
- _____, Kroos L and Lenski R E 2000 Developmental cheating in the social bacterium *Myxococcus xanthus*; *Nature* **404** 598-601
- _____, Lenski R E and Kroos L 2002 Rescue of social motility lost during evolution of *Myxococcus xanthus* in an asocial environment; *J. Bacteriol.* **184** 2719-2727
- Volny V P and Gordon D M 2002 Genetic basis for queen-worker dimorphism in a social insect; *Proc. Natl. Acad. Sci. USA* **99** 6108-6111
- Waddell D R and Duffy K T I 1986 Breakdown of self/nonsel self recognition in cannibalistic strains of the predatory slime mold, *Dictyostelium caveatum*; *J. Cell Biol.* **102** 298-305
- Weijer C J, Duschl G and David C N 1984 Dependence of cell-type proportioning and sorting on cell cycle phase in *Dictyostelium discoideum*; *J. Cell Sci.* **70** 111-131
- Weismann A 1893 *The Germ-Plasm: A Theory of Heredity*; translated by W Newton Parker and H Ronnfeld. (London : Walter Scott Ltd.)
- Williams G C 1966 *Adaptation and Natural Selection*; (Princeton: Princeton University Press)
- Yamamoto A and Takeuchi I 1983 Vital staining of autophagic vacuoles in differentiating cells of *Dictyostelium discoideum*; *Differentiation* **24** 83-87
- Zahavi A 1977 Reliability in communication systems and the evolution of altruism; in *Evolutionary Ecology* pp 253-259 eds. B Stonehouse and C M Perrins