

BIOLOGY LETTERS

rsbl.royalsocietypublishing.org



Research

Cite this article: Pike TW. 2015 Interference coloration as an anti-predator defence. *Biol. Lett.* 20150159.

<http://dx.doi.org/10.1098/rsbl.2015.0159>

Received: 1 March 2015

Accepted: 20 March 2015

Subject Areas:

behaviour

Keywords:

iridescence, Japanese quail, greenbottles, *Lucilia* spp., motion dazzle

Author for correspondence:

Thomas W. Pike

e-mail: tpike@lincoln.ac.uk

Electronic supplementary material is available at <http://dx.doi.org/10.1098/rsbl.2015.0159> or via <http://rsbl.royalsocietypublishing.org>.

THE ROYAL SOCIETY
PUBLISHING

Animal behaviour

Interference coloration as an anti-predator defence

Thomas W. Pike^{1,2}

¹Centre for Ecology and Conservation, University of Exeter, Penryn, Cornwall TR10 9EZ, UK

²School of Life Sciences, University of Lincoln, Riseholme Park, Lincoln LN2 2LG, UK

Interference coloration, in which the perceived colour varies predictably with the angle of illumination or observation, is extremely widespread across animal groups. However, despite considerable advances in our understanding of the mechanistic basis of interference coloration in animals, we still have a poor understanding of its function. Here, I show, using avian predators hunting dynamic virtual prey, that the presence of interference coloration can significantly reduce a predator's attack success. Predators required more pecks to successfully catch interference-coloured prey compared with otherwise identical prey items that lacked interference coloration, and attacks against prey with interference colours were less accurate, suggesting that changes in colour or brightness caused by prey movement hindered a predator's ability to pinpoint their exact location. The pronounced anti-predator benefits of interference coloration may explain why it has evolved independently so many times.

1. Introduction

Interference coloration, in which the perceived colour and brightness varies predictably with the angle of illumination or observation [1], is extremely widespread across animal groups, having evolved independently several times in insects such as beetles and butterflies, as well as in some birds, fish, reptiles and cephalopods, and at least one mammal [2–4]. However, despite widespread interest in the mechanisms underlying the production of interference colours in animals [5], its function is still unclear [3]. It has been suggested, for example, that because interference colours are often visually striking they may function in sex or species recognition, or have evolved as sexually selected signals [2,5–8]. Alternatively, the structures that produce interference coloration may also have non-communicative functions such as enhancing water repellence, friction reduction or thermoregulation, and may even alter light reaching the retina and so play a role in vision [2,9,10]. A further role of interference coloration may be as an anti-predator mechanism [2].

Although interference colours often exhibit conspicuous changes in colour and brightness when observed outside their natural context, some forms of interference colour have been suggested to have an anti-predator function by allowing animals to appear cryptic against their background [11–14]; for example, the most common interference colour in beetles is green [11,13]. However, animals with interference colours that contrast sharply with the background would seem poorly adapted for crypsis, leading to the suggestion that such colours may function as an aposematic signal [15,16] or as an active anti-predator mechanism [17,18]. In particular, interference colours can produce bright flashes of colour or sudden changes in brightness that might briefly startle a potential predator and thereby increase the prey's probability of escape [17], or because movement-induced changes in colour or brightness may hinder a predator's ability to pinpoint the prey's exact location in an attempted strike [18]. Such mechanisms have been proposed as a function of interference colour in beetles [15,19–21], butterflies [22], birds [2] and fish [23]. However, whether having interference

coloration decreases the chance of predation for mobile prey has never been empirically tested. Here, I experimentally tested this using Japanese quail (*Coturnix japonica*) predating virtual insects.

2. Material and methods

(a) Experimental design

The subjects were female Japanese quail ($n = 7$) from laboratory stock. They were aged between six and 18 months, and had all participated in a number of previous operant experiments involving visual search and detection of static objects (e.g. locating cryptic items against a heterogeneous greyscale background). They had not, however, participated in a task involving moving or coloured stimuli. All birds had the same prior experience.

Experiments took place in a wooden cage (160×65 cm and 25 cm high), at one end of which was a calibrated flat-screen CRT monitor (Iiyama VisonMaster 513) equipped with a 30×23 cm infrared touch screen bezel (CarrollTouch Smart Frame, Elo TouchSystems, Menlo Park, CA) to record pecks, and an automatic feeder. Stimuli, which were presented on the monitor, consisted of a circular moving 'prey', 20 mm in diameter (chosen because stimuli of this size were invariably attacked and caught, regardless of their appearance). Each prey item moved in a straight line at a constant speed of approximately 150 mm s^{-1} until it reached the edge of the screen, when it turned round and moved off at the angle of incidence. No attempt was made to replicate the locomotive behaviour of any given species, but rather to provide a target that was challenging for quail to catch and that did not introduce any 'behaviours' that may interact with colour in affecting its chances of predation, such as random movement patterns. When subjects made a 'successful' choice, defined as a peck within 10 mm of the centre of a prey item (i.e. within its body), they were rewarded with 10 s access to a feeder containing ad libitum whole dried mealworms (*Tenebrio molitor* larvae), a favourite food. There was no attempt to food-deprive the birds, as the mealworm reward provided sufficient motivation to complete the task.

Subjects were initially trained to peck at black moving target stimuli on a uniform grey background. Training continued until a subject caught six consecutive stimuli within three pecks, after which it progressed to the test phase of the experiment. The mean \pm s.e. number of trials required to reach the training criterion was 23.9 ± 4.5 .

After successfully completing training, each subject was presented sequentially with 20 stimuli, 10 with interference coloration (treatment) and 10 without (control), in a random order, against a uniform grey background. Stimuli were constructed as polygon models consisting of one half sphere, with the convex face pointing towards the observer. For treatment stimuli, all vertices on the polygon model were assigned a colour based on the angle between a virtual light source, the vertex and a fixed point in front of the centre of the screen representing the observer, and the faces of the polygon coloured by interpolation of these vertex colours. The colour assigned to a vertex was assumed to be produced by a simple planar multilayer composed of stacked chitin plates of equal thicknesses, separated by very thin air spaces. This alternating chitin/air multilayer structure is a common configuration in insects, producing colour through constructive interference [24]. Treatment stimuli were modelled on greenbotflies, *Lucilia sericata* [24], which appear metallic green at normal incidence, turning to blue at large angles. As they moved around the screen, they, therefore, underwent angle-dependent changes in colour and brightness. The colour of each control stimulus was chosen at random from within the gamut of colours that could be produced by the treatment stimuli. These stimuli were

therefore uniformly coloured, and colour was independent of spatial position. Note that the colours displayed on the screen only loosely matched those produced by real flies. However, the aim was not to exactly replicate colours produced by real insects or replicate how real colours would be perceived by quail predators. Rather, the aim was to produce target stimuli that exhibited the salient characteristics of interference coloration, namely chromatic and achromatic shifts in response to changes in viewing angle as the target moved around the screen.

The effectiveness of the treatment as an anti-predator strategy was assessed using (i) the number of pecks needed to successfully catch a prey item, (ii) the mean distance of all the unsuccessful pecks from the centre of the prey item prior to capture (a measure of peck accuracy) and (iii) the latency to make the first attack. Full details of the experimental set-up, stimulus design, training and testing procedures are given in the electronic supplementary material.

(b) Statistical analyses

To test whether the number of pecks needed to successfully catch a stimulus differed between treatment and control stimuli, I used a generalized linear mixed-effects model with a negative binomial error distribution to account for overdispersion [25]. Stimulus type (treatment or control) was included as a fixed factor, and there was a random effects term of subject identity. I also included trial number, and the interaction between trial number and stimulus type, in order to test for improvements in predator performance over successive trials (i.e. learning). To test whether peck accuracy and latency to peck differed between stimuli, I used linear mixed-effects models, parametrized as described above. Both peck latency and accuracy were normalized using a log-transformation prior to analysis. For all models, p -values were calculated by comparing the full model to a reduced model lacking the fixed effect term using likelihood ratio tests [25], and models simplified using backwards stepwise elimination of non-significant terms. Minimum adequate models are presented. All analyses were conducted in R (R Core Development Team, v. 2.15.2) using either the 'lmer' function in the lme4 package [26] or the 'glmmadmb' function in the glmmADMB package [27], and models validated following [28].

3. Results

The number of pecks needed to catch a treatment (interference-coloured) stimulus was significantly greater than the number needed to catch a control (non-interference-coloured) stimulus ($\chi^2_1 = 7.89$, $p = 0.005$; figure 1a) and peck accuracy was significantly poorer for unsuccessful pecks at treatment than control stimuli ($\chi^2_1 = 10.99$, $p < 0.001$; figure 1b). There was no difference between stimulus types in the latency to attack (mean \pm s.d., treatment: 24.15 ± 10.62 s; control: 23.06 ± 9.78 s; $\chi^2_1 = 0.29$, $p = 0.592$). In all models, trial number and the interaction between trial number and stimulus type were non-significant (all $p > 0.3$), suggesting that performance did not improve with learning.

4. Discussion

The results of this experiment show that predators attacking treatment prey items with interference-like colours require significantly more pecks, and unsuccessful pecks are significantly less accurate, than when attacking otherwise identical control prey. These findings suggest that significant anti-predator benefits may be gained by having interference




Figure 1. (a) Mean \pm s.e. number of pecks needed to successfully catch interference-coloured (treatment) and non-interference-coloured (control) stimuli. (b) Mean \pm s.e. distance of unsuccessful pecks at treatment and control stimuli, measured in mm from the centre of the prey item at the time of pecking. ** $p < 0.01$; *** $p < 0.001$.

coloration, and may explain why interference colours are so widespread [15,16,19–23]. Because the accuracy of prey-directed pecks, but not the latency to peck, was significantly

reduced when attacking treatment prey items, it is likely that perceived changes in colour or brightness caused by the movement of animals with interference coloration hinders a predator's ability to pinpoint the prey's exact location in an attempted strike [18], rather than startling them through sudden chromatic or achromatic changes [17,29]. This is consistent with recent work on motion dazzle, in which high-contrast colour patterns can impair a predator's ability to judge the speed and direction of moving prey, making them harder to catch [29–32]. The sudden changes in brightness perceived by predators viewing rapidly moving prey with interference colours may therefore act as a form of dazzle coloration. More generally, this finding raises the intriguing possibility that changing appearance per se may be important in predator avoidance (see [33,34]), and that interference colours are a special case of this phenomenon.

While this study presents evidence that, at least in some cases, interference coloration has evolved as an anti-predator defence, whether this mechanism acts to protect prey in natural situations and how this relates to its other putative roles [2,3,5–10] remains to be tested.

Ethics statement. The experiment was carried out with the approval of the University of Exeter's local ethics committee.

Data accessibility. Raw data can be found in the electronic supplementary material.

Acknowledgements. Thanks to Nicole Westbury-Harris for animal care, and to three anonymous referees whose constructive comments greatly improved the manuscript.

Funding statement. This work was supported by a Natural Environment Research Council fellowship (NE/F016514/1).

Author contributions. All the work described here is my own.

Conflict of interests. I have no competing interests.

References

- Osorio D, Ham AD. 2002 Spectral reflectance and directional properties of structural coloration in bird plumage. *J. Exp. Biol.* **205**, 2017–2027.
- Doucet SM, Meadows MG. 2009 Iridescence: a functional perspective. *J. R. Soc. Interface* **6**, S115–S132. (doi:10.1098/rsif.2008.0395.focus)
- Meadows MG, Butler MW, Morehouse NI, Taylor LA, Toomey MB, McGraw KJ, Rutowski RL. 2009 Iridescence: views from many angles. *J. R. Soc. Interface* **6**, S107–S113. (doi:10.1098/rsif.2009.0013.focus)
- Vukusic P. 2004 Natural photonics. *Phys. World* **17**, 35–39.
- Vukusic P, Kelly R, Hooper I. 2009 A biological sub-micron thickness optical broadband reflector characterized using both light and microwaves. *J. R. Soc. Interface* **6**, S193–S201. (doi:10.1098/rsif.2008.0345.focus)
- Loyau A, Gomez D, Moureau BT, They M, Hart NS, Saint Jalme M, Bennett ATD, Sorci G. 2007 Iridescent structurally based coloration of eyespots correlates with mating success in the peacock. *Behav. Ecol.* **18**, 1123–1131. (doi:10.1093/beheco/arm088)
- Rutowski RL, Macedonia JM, Merry JW, Morehouse NI, Yturralde K, Taylor-Taft L, Gaalema D, Kemp DJ, Papke RS. 2007 Iridescent ultraviolet signal in the orange sulphur butterfly (*Colias eurytheme*): spatial, temporal and spectral properties. *Biol. J. Linn. Soc.* **90**, 349–364. (doi:10.1111/j.1095-8312.2007.00749.x)
- Rutowski RL, Macedonia JM, Morehouse N, Taylor-Taft L. 2005 Pterin pigments amplify iridescent ultraviolet signal in males of the orange sulphur butterfly, *Colias eurytheme*. *Proc. R. Soc. B* **272**, 2329–2335. (doi:10.1098/rspb.2005.3216)
- Mathger LM, Denton EJ, Marshall NJ, Hanlon RT. 2009 Mechanisms and behavioural functions of structural coloration in cephalopods. *J. R. Soc. Interface* **6**, S149–S163. (doi:10.1098/rsif.2008.0366.focus)
- Seago AE, Brady P, Vigneron JP, Schultz TD. 2009 Gold bugs and beyond: a review of iridescence and structural colour mechanisms in beetles (Coleoptera). *J. R. Soc. Interface* **6**, S165–S184. (doi:10.1098/rsif.2008.0354.focus)
- Crowson RA. 1981 *The biology of the Coleoptera*. New York, NY: Academic Press.
- Knisley CB, Schultz TD. 1997 *The biology of tiger beetles and a guide to the species of the South Atlantic States*. Martinsville, VA: Virginia Museum of Natural History.
- Parker AR, Mckenzie DR, Ah Yong ST. 1998 A unique form of light reflector and the evolution of signalling in *Ovalipes* (Crustacea: Decapoda: Portunidae). *Proc. R. Soc. Lond. B* **265**, 861–867. (doi:10.1098/rspb.1998.0371)
- Pearson D, Vogler A. 2001 *Tiger beetles: the evolution, ecology, and diversity of the cicindelids*. Ithaca, NY: Cornell University Press.
- Vulinec K. 1997 Iridescent dung beetles: a different angle. *Flora Entomol.* **80**, 132–141. (doi:10.2307/3495550)
- Fabricant SA, Exnerova A, Jezova D, Stys P. 2014 Scared by shiny? The value of iridescence in aposematic signalling of the hibiscus harlequin bug. *Anim. Behav.* **90**, 315–325. (doi:10.1016/j.anbehav.2014.01.021)
- Hinton HE. 1973 Some recent work on the colours of insects and their likely significance. *Proc. Br. Ent. Nat. Hist. Soc.* **6**, 43–54.
- Robinson MH. 1969 Defenses against visually hunting predators. *Evol. Biol.* **3**, 225–259.

- 190 19. Acorn JH. 1988 Mimetic tiger beetles and the
191 puzzle of cicindelid coloration (Coleoptera:
192 Cicindelidae). *Coleopt. Bull.* **42**, 28–33.
- 193 20. Schultz TD. 1986 Role of structural colors in
194 predator avoidance by tiger beetles of the genus
195 Cicindela (Coleoptera: Cicindelidae). *Bull. Entomol.*
196 *Soc. Am.* **32**, 142–146. (doi:10.1093/besa/32.3.142)
- 197 21. Schultz TD. 2001 Tiger beetle defenses revisited:
198 alternative defense strategies and colorations of two
199 neotropical tiger beetles, *Odontocheila nicaraguensis*
200 Bates and *Pseudoxycheila tasis* Bates (Carabidae:
201 Cicindelinae). *Coleopt. Bull.* **55**, 153–163. (doi:10.
202 1649/0010-065X(2001)055[0153:TBDRAD]2.0.CO;2)
- 203 22. Clench HK. 1966 Behavioral thermoregulation in
204 butterflies. *Ecology* **47**, 1021–1034. (doi:10.2307/
205 1935649)
- 206 23. Denton EJ. 1970 On the organization of reflecting
207 surfaces in some marine animals. *Phil. Trans. R. Soc.*
208 *Lond. B* **258**, 285–313. (doi:10.1098/rstb.
209 1970.0037)
- 210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
24. Martinec I, Pudis D, Satka A, Janigova I, Csomorova
K, Cernobila F. 2008 Temperature effect on optical
properties of the cuticle of *Lucilia sericata*. *Optik*
119, 523–527. (doi:10.1016/j.ijleo.2007.02.009)
25. Crawley MJ. 2005 *Statistics: an introduction using R*.
Chichester, UK: John Wiley.
26. Bates D, Maechler M, Bolker B, Walker S. 2013
lme4: Linear mixed-effects models using Eigen and
S4. R package v. 1.0–4.
27. Fournier DA, Skaug HJ, Ancheta J, Ianelli J,
Magnusson A, Maunder M, Nielsen A, Sibert J. 2012
AD model builder: using automatic differentiation
for statistical inference of highly parameterized
complex nonlinear models. *Optim. Methods*
Softw. **27**, 233–249. (doi:10.1080/10556788.
2011.597854)
28. Zuur AF, Ieno EN, Elphick CS. 2010 A protocol for
data exploration to avoid common statistical
problems. *Methods Ecol. Evol.* **1**, 3–14. (doi:10.
1111/j.2041-210X.2009.00001.x)
29. Stevens M, Yule DH, Ruxton GD. 2008 Dazzle
coloration and prey movement. *Proc. R. Soc. B* **275**,
2639–2643. (doi:10.1098/rspb.2008.0877)
30. Scott-Samuel NE, Baddeley R, Palmer CE, Cuthill IC.
2011 Dazzle camouflage affects speed perception.
PLoS ONE **6**, e20233. (doi:10.1371/journal.pone.
0020233)
31. Stevens M, Searle WTL, Seymour JE, Marshall KLA,
Ruxton GD. 2011 Motion dazzle and camouflage as
distinct anti-predator defenses. *BMC Biol.* **9**, 81.
(doi:10.1186/1741-7007-9-81)
32. von Helversen B, Schooler LJ, Czienskowski U. 2013
Are stripes beneficial? Dazzle camouflage influences
perceived speed and hit rates. *PLoS ONE* **8**, e61173.
(doi:10.1371/journal.pone.0061173)
33. Papageorgis C. 1975 Mimicry in neotropical
butterflies. *Am. Sci.* **63**, 522–532.
34. Endler JA. 1978 A predator's view of animal color
patterns. *Evol. Biol.* **11**, 319–364. (doi:10.1007/
978-1-4615-6956-5_5)