



GESDAV

Social patterning of alcohol consumption among mothers with infants in the UK

Sarah Baker¹, Hilary Graham²

¹Department of Nutrition and Dietetics, Faculty of Health and Social Sciences, Leeds Metropolitan University, Leeds, United Kingdom, ²Department of Health Sciences, University of York, York, United Kingdom

Address for correspondence:

Dr. Sarah Baker, Department of Nutrition and Dietetics, Faculty of Health and Social Sciences, Leeds Metropolitan University, Leeds, United Kingdom.
E-mail: sarah.baker@leedsmet.ac.uk

Received: June 16, 2014

Accepted: July 08, 2014

Published: July 16, 2014

ABSTRACT

Background: We examined patterns of alcohol use among mothers with infants in the UK focusing on (i) common (“majority”) patterns of alcohol use (frequency and quantity) and (ii) associated social factors. **Methods:** We analyzed data from 15,510 mothers who took part in waves 1 and 2 of the millennium cohort study in 2000/1. Nine months after birth, mothers reported their frequency and quantity of alcohol use, along with information on social circumstances. Logistic regression, carried out in 2011, helped identify the social factors associated with majority patterns of alcohol use. **Results:** Majority patterns of alcohol use were: Infrequent drinking (never/<1/week), infrequent light drinking (1 unit/day, <1/week), and frequent light drinking (<14 units/week). In mutually adjusted models, infrequent drinking was associated with childhood, educational, and income disadvantage, and younger age at first birth. Infrequent light drinking was associated with educational and income disadvantage, economic inactivity, and marriage. Frequent light drinking was associated with marriage and fewer children in the household. **Conclusions:** Among mothers in the UK, the majority drank infrequent small quantities. Positive social gradients were evident for frequency of alcohol use among socially advantaged mothers.

KEY WORDS: Alcohol, motherhood, social circumstances, social patterning

INTRODUCTION

Alcohol is measured in units and “One unit is 10 ml or 8 g of pure alcohol” [1]. The maximum recommended intake of alcohol for women in the UK is 2-3 units/day; these have remained unchanged since 1995 when recommendations based on weekly quantities (units) were dropped in favour of daily quantities (units). The recommendations are comparable to international standards [2].

The majority of adults in the UK report that they drink alcohol and for most adults, alcohol consumption is not considered to be health-damaging [3]. However, research has focused on the minority of the population whose patterns of consumption are associated with a range of health and social problems [4,5]. As a result, more is known about “risky” alcohol use than about the drinking patterns of the majority of the population. In contrast, this paper takes a population perspective that examines the underlying social determinants of alcohol use rather than focusing on a small proportion of “risky” drinking individuals in order to identify potential areas for widespread public health intervention [6].

UK surveys provide important information on majority drinking patterns among women. They indicate that alcohol use is more common among white British women; women from minority ethnic groups are more likely to report never drinking or drinking less than once per week. They point, too, to positive gradients in the frequency, and negative gradients in the quantity, of alcohol use among socially advantaged groups [7-10]. Similarly, studies in other high-income countries report more frequent alcohol use among employed women with higher educational levels and living in higher income households [11,12]. The opposite has been found for drinking quantity. Women who are advantaged with respect to education, employment status, occupational class and household income consume smaller quantities of alcohol on drinking occasions than women from more disadvantaged backgrounds [11-13].

Domestic circumstances are also associated with women’s patterns of alcohol use. Compared with non-married women, married women report lower alcohol consumption in terms of both frequency and quantity [12]. Having children in the household is also associated with alcohol use, with decreased alcohol consumption among parents [4,12]. In addition, there

is evidence that the timing and circumstances of entry into motherhood are related to patterns of alcohol use [14-18].

However, few studies have looked specifically at women living with children [16]; in consequence, little is known about the frequency and quantity of alcohol use among mothers. This is despite the fact that most women become mothers, and the home is the primary location in which children are introduced to alcohol [3,19]. As with other health-related behaviours, children's subsequent drinking patterns are associated with the drinking behaviours of their parents [15,20].

The millennium cohort study (MCS) is a unique source of UK data on the social circumstances and lifestyles of mothers and has a sample size that permits analysis of the social patterning of the frequency and quantity of alcohol use among mothers. We focus on alcohol use 9 months after birth (wave 1). Although questions on frequency of alcohol use were also asked at wave 2 (3 years after birth), questions on quantity were only asked of a sub-sample of mothers.

METHODS

Participants

The MCS is a survey based on a sample of children born in the UK in 2000/01, disproportionately sampled in favour of disadvantaged groups. The cohort was recruited when the child was 9 months old (UK response rate 68%; $n = 18,552$), with a pre-school follow-up at 3 years (UK response rate 78% of original sample; $n = 15,590$) [21,22]. At first contact, mothers reported their ethnicity and natural mothers from a white British ethnic background [23] who were resident in the house in which the cohort baby lived were considered eligible for inclusion. Preliminary analysis pointed to marked ethnic differences in alcohol use. Compared to white British mothers, a much larger proportion of mothers from ethnic minorities never drank or drank less than once per week, but the small numbers prevented ethnic-group specific analyses.

We included eligible mothers who participated in interviews wave 1 (9 months) and wave 2 (3 years), when a question on childhood social circumstances was asked ($n = 15,510$).

Ethics

This study was approved by the University of York's Department of Health Sciences Research Governance Committee, York, UK.

Outcome Measures

Frequency of alcohol consumption was measured by a single item ("which of these best describes how often you drink?") with fixed categorical responses. We combined the lowest two frequency categories (never and <1 week) because preliminary analyses indicated that they had the same socio-economic distribution (data not shown). We also combined the highest two categories (5/6 days a week and everyday) because of small

Table 1: Alcohol use amongst mothers

Alcohol use	N	Weighted %
A) Frequency		
Never/<1/week	9086	59
1-2/week	4285	28
3-4/week	1379	9
5-6/everyday/week	757	5
Missing	3	
	15510	100
B) Quantity (if frequency <1/week)		
1 unit/day on each occasion	3595	40
2-3 units/day on each occasion	2542	28
>3 units/day on each occasion	2949	33
	9086	100
C) Quantity (if frequency >1/week)		
<14 units/week	5868	91
14-21 units/week	454	7
>21 units/week	99	2
	6421	100

numbers in the latter category. The distribution of mothers across these groups is given in Table 1 (Block A).

For quantity of consumption, mothers drinking <1/week were asked about the average number of units they consumed on each drinking occasion; those drinking ≥ 1 /week were asked about the average number of units they drank a week.

Mothers who drank infrequently (<1/week) were grouped into those who drank under (1 unit/day), within (2-3 units/day) and over (>3 units/day) the daily recommendations (Table 1 Block B). We examined the social patterning of units within these categories (e.g. 2 units and 3 units in the 2-3 units category) to check that social variations in the quantity of alcohol use were not being obscured by the 3-fold categorization (data not shown).

Mothers who frequently drank (>1/week) were categorized into those who drank under (<14 units/week), within (14-21 units/week) and over (>22 units/week) the weekly recommendations (Table 1 Block C).

The analysis focused on these three most commonly reported patterns: Drinking infrequently (never/<1/week), infrequent light quantities (1 unit/day, <1/week) and frequent light quantities (<14 units/week).

Socio-demographic Factors

The mother's childhood circumstances were measured at wave 2 by her father's occupational class when she was aged 14, using the collapsed 3-category NS-SEC classification (managerial/professional, intermediate and routine/manual) [24]. Fathers who were not working were included in the routine/manual category because of small numbers. In addition, "unknown" category was constructed for women who did not know what job their father did. All other measures were derived from wave 1 data. Educational level was measured by age of leaving full-time education and employment status was split into two categories (economically active/inactive). Equivalised household income was based on MCS income bands, with "unknown" category for women who

did not know their household income. Along with these standard socioeconomic measures, we included maternal age at first birth, current cohabitation status (married, cohabiting, lone) and the number of children in the household, including the cohort child.

Statistical Analysis

The analysis, carried out in 2011, was conducted using the statistical software package STATA version 10.1 [25], with survey commands to account for the sample design, and survey weights to correct for disproportionate sampling in favour of disadvantaged groups.

Using logistic regression, bivariate analyses of mother's alcohol use were undertaken for each social factor adjusted only for age. Those predictors found to be significant were included in a mutually adjusted model. Odds ratios were calculated for the three outcomes of interest: Infrequent drinking (never/<1/week), infrequent light drinking (1 unit/day, <1/week), and frequent light drinking (<14 units/week).

RESULTS

Table 2 describes the social profile of mothers. Table 3 describes the associations between social circumstances and the majority

Table 2: Social profile of the sample

Social profile of the sample (n=15510)	N	Weighted %
Childhood circumstances		
Managerial/prof	3138	20.2
Intermediate	2659	17.1
Economically inactive/lowest	5480	35.3
Unknown	4233	27.3
Age left education		
>22	1443	9.3
19-21	1739	11.2
17-18	4498	29.0
<16	7830	50.5
Employment status		
Economically active	7957	51.3
Economically inactive	7553	48.7
Household income		
£31,200+	3021	19.9
£20,800-31,200	3038	20.0
£10,400-20,800	4741	31.3
£0-10,400	3640	24.0
Unknown	728	4.8
Missing	342	
Age at first live birth		
>30	3735	24.3
25-29	4298	27.9
20-24	4068	26.4
14-19	3286	21.4
Missing	123	
Cohabitation status		
Married	8671	56.2
Cohabiting	4120	26.7
Lone parent	2638	17.1
Missing	81	
Number of children		
>3	3403	21.9
2	5535	35.7
1	6572	42.4

patterns of alcohol use: Infrequent drinking (never/<1/week), infrequent light quantities (1 unit/day, <1/week), and frequent light quantities (<14 units/week).

Infrequent drinking (never/<1/week) was associated in the bivariate analyses with all measures of disadvantage, as well as with having more children in the household. In a multivariate analysis, disadvantaged childhood circumstances, educational disadvantage, income disadvantage, and younger age at first birth remained significant predictors.

Infrequent light drinking (1 unit/day, <1/week) was also associated with disadvantage. In the bivariate analyses, educational disadvantage, economic inactivity and income disadvantage increased the odds of being an infrequent light drinker, as did being married and having more children in the household. In a multivariate analysis, number of children in the household was no longer a significant predictor.

Frequent light drinking (<14 units/week) was associated with younger age at first birth in the bivariate analyses, as was being married and having fewer children living in the household in both the bivariate and multivariate analyses.

Strengths and Limitations

The MCS relies on self-reported levels of alcohol use and includes only a limited set of measures. Self-reports underestimate consumption when compared with the sale of taxed alcohol [26], but provide a reliable means of comparing drinking within population groups [27]. Studies have found that women are more likely than men to under-report their alcohol use and mothers represent a group who may be more inclined to report conservative estimates of alcohol consumption [28]. Moreover, it is likely that mothers who are problematic drinkers will be under-represented in the MCS, as has been found to be the case in the general population surveys [29]. Nevertheless, self-reported measures of alcohol use are the only option for large scale population surveys [30].

Quantity was measured in units of alcohol in line with UK recommendations. Studies indicate that people do not understand how many units of alcohol relate to everyday measures like glass size and pints or what the recommendations are. For example, a British study indicated that, of the 75% of individuals who had heard of daily limits, only 44% and 52% knew the correct daily limits for men and women respectively. Furthermore, of these individuals, only 12% of men and 14% of women kept track of the number of units they consumed [31]. Unit-based measures are, therefore, likely to lack precision; our use of broad-based categories of consumption accommodates this.

In addition, data from the MCS were collected over a decade ago and may not reflect contemporary patterns of alcohol use. However, more recent studies suggest that the patterns uncovered in our analyses remain valid [11,13]. The demography of motherhood has also changed over time. Nevertheless, evidence suggests that levels of employment and household income have increased at a modest rate [32,33]. Similarly,

Table 3: The OR of drinking patterns according to: Social circumstances and level of disadvantage

	Infrequent drinking (never/<1/week)		Infrequent light drinking (1 unit/day, <1/week)		Frequent light drinking (<14 units/week)	
	Bivariate age adjusted	Mutually adjusted (n=14975)	Bivariate age adjusted	Mutually adjusted (n=8828)	Bivariate age adjusted	Mutually adjusted (n=6397)
Childhood circumstances	(n=15503)		(n=9083)		(n=6420)	
Managerial/prof	1.00	1.00	1.00		1.00	
Intermediate	OR: 1.40	OR: 1.21	OR: 1.09		OR: 1.24	
	(CI: 1.25-1.58)	(CI: 1.08-1.36)	(CI: 0.91-1.29)		(CI: 1.03-1.50)	
Economically inactive/lowest	OR: 1.87	OR: 1.42	OR: 0.97		OR: 1.13	
	(CI: 1.65-2.12)	(CI: 1.27-1.59)	(CI: 0.84-1.12)		(CI: 0.96-1.32)	
Unknown	OR: 1.85	OR: 1.39	OR: 1.16		OR: 0.90	
	(CI: 1.63-2.10)	(CI: 1.23-1.57)	(CI: 0.98-1.36)		(CI: 0.74-1.09)	
	P=0.000	P=0.000	P=0.206		P=0.412	
Age of leaving education	(n=15503)		(n=9083)		(n=6420)	
22 and over	1.00	1.00	1.00	1.00	1.00	
19-21	OR: 1.30	OR: 1.15	OR: 0.82	OR: 0.78	OR: 0.94	
	(CI: 1.08-1.56)	(CI: 0.95-1.38)	(CI: 0.62-1.09)	(CI: 0.58-1.04)	(CI: 0.76-1.17)	
17-18	OR: 2.05	OR: 1.52	OR: 0.74	OR: 0.70	OR: 1.09	
	(CI: 1.71-2.46)	(CI: 1.28-1.80)	(CI: 0.58-0.95)	(CI: 0.54-0.89)	(CI: 0.86-1.33)	
16 and under	OR: 2.84	OR: 1.70	OR: 0.67	OR: 0.58	OR: 0.95	
	(CI: 2.36-3.43)	(CI: 1.41-2.03)	(CI: 0.53-0.85)	(CI: 0.45-0.75)	(CI: 0.79-1.14)	
	P=0.000	P=0.000	P=0.000	P=0.000	P=0.739	
Employment status	(n=15503)		(n=9083)		(n=6420)	
Economically active	1.00	1.00	1.00	1.00	1.00	
Economically inactive	OR: 1.40	OR: 1.08	OR: 1.55	OR: 1.55	OR: 1.06	
	(CI: 1.30-1.51)	(CI: 0.98-1.18)	(CI: 1.39-1.74)	(CI: 1.37-1.74)	(CI: 0.93-1.20)	
	P=0.000	P=0.104	P=0.000	P=0.000	P=0.421	
Household income	(n=15164)		(n=8885)		(n=6279)	
£31,200+	1.00	1.00	1.00	1.00	1.00	
£20,800-31,200	OR: 1.91	OR: 1.64	OR: 0.84	OR: 0.87	OR: 1.31	
	(CI: 1.65-2.22)	(CI: 1.42-1.88)	(CI: 0.70-1.01)	(CI: 0.72-1.06)	(CI: 1.11-1.56)	
£10,400-20,800	OR: 2.98	OR: 2.24	OR: 1.06	OR: 1.11	OR: 1.22	
	(CI: 2.56-3.46)	(CI: 1.93-2.59)	(CI: 0.90-1.24)	(CI: 0.93-1.32)	(CI: 1.03-1.44)	
£0-10,400	OR: 3.42	OR: 2.38	OR: 1.18	OR: 1.32	OR: 0.80	
	(CI: 2.88-4.07)	(CI: 1.98-2.87)	(CI: 0.97-1.43)	(CI: 1.06-1.64)	(CI: 0.63-1.01)	
Unknown	OR: 2.23	OR: 1.81	OR: 1.21	OR: 1.25	OR: 1.28	
	(CI: 1.80-2.78)	(CI: 1.45-2.25)	(CI: 0.95-1.55)	(CI: 0.97-1.60)	(CI: 0.94-1.73)	
	P=0.000	P=0.000	P=0.002	P=0.001	P=0.670	
Age at first live birth	(n=15382)		(n=9007)		(n=6375)	
30+	1.00	1.00	1.00		1.00	1.00
25-29	OR: 1.70	OR: 1.44	OR: 0.83		OR: 1.09	OR: 1.18
	(CI: 1.50-1.92)	(CI: 1.28-1.63)	(CI: 0.70-0.98)		(CI: 0.93-1.28)	(CI: 1.00-1.39)
20-24	OR: 2.40	OR: 1.52	OR: 0.93		OR: 0.90	OR: 1.16
	(CI: 2.08-2.76)	(CI: 1.31-1.77)	(CI: 0.77-1.13)		(CI: 0.74-1.10)	(CI: 0.91-1.47)
14-19	OR: 2.96	OR: 1.61	OR: 0.89		OR: 0.65	OR: 0.95
	(CI: 2.51-3.50)	(CI: 1.32-1.97)	(CI: 0.72-1.09)		(CI: 0.50-0.85)	(CI: 0.70-1.29)
	P=0.000	P=0.000	P=0.429		P=0.018	P=0.525
Cohabitation status	(n=15422)		(n=9025)		(n=6397)	
Married	1.00	1.00	1.00	1.00	1.00	1.00
Cohabiting	OR: 1.24	OR: 0.99	OR: 0.81	OR: 0.77	OR: 0.73	OR: 0.72
	(CI: 1.13-1.36)	(CI: 0.90-1.08)	(CI: 0.71-0.92)	(CI: 0.67-0.88)	(CI: 0.62-0.85)	(CI: 0.61-0.85)
Lone parent	OR: 1.61	OR: 0.98	OR: 0.82	OR: 0.64	OR: 0.50	OR: 0.52
	(CI: 1.40-1.85)	(CI: 0.84-1.14)	(CI: 0.69-0.97)	(CI: 0.52-0.79)	(CI: 0.38-0.64)	(CI: 0.39-0.68)
	P=0.000	P=0.149	P=0.006	P=0.000	P=0.000	P=0.000
Number of children in household	(n=15503)		(n=9083)		(n=6420)	
3+	1.00	1.00	1.00	1.00	1.00	1.00
2	OR: 0.69	OR: 0.92	OR: 0.82	OR: 0.84	OR: 1.13	OR: 1.12
	(CI: 0.62-0.77)	(CI: 0.82-1.03)	(CI: 0.71-0.96)	(CI: 0.72-0.99)	(CI: 0.95-1.34)	(CI: 0.93-1.35)
1	OR: 0.60	OR: 1.00	OR: 0.79	OR: 0.86	OR: 1.32	OR: 1.34
	(CI: 0.54-0.68)	(CI: 0.88-1.15)	(CI: 0.67-0.92)	(CI: 0.73-1.02)	(CI: 1.11-1.56)	(CI: 1.09-1.65)
	P=0.000	P=0.541	P=0.009	P=0.231	P=0.001	P=0.002

OR: Odds ratio, CI: Confidence interval, P=Statistical significance

relatively small increases in the age of leaving education and first birth have been witnessed, as has a steady decline in birth within marriage [34,35]. Therefore, the results of our study remain comparable to current demographic groups in the UK.

Notwithstanding these limitations, our analyses provide a “snapshot” of mothers’ alcohol use at key life stage and describe how mothers’ social circumstances may influence their patterns of alcohol use.

CONCLUSIONS

In the UK, alcohol consumption is considered a social norm [3]. As such, alcohol is deeply ingrained in society, with evidence that alcohol use tracks over time and across generations [36,37]. However, there have been few studies focused on mothers [16]. We, therefore, undertook an analysis of the UK's largest study of mothers to investigate patterns of alcohol use around 9 months after birth. We focused on the most common patterns of use: Infrequent drinking (never/<1/week), infrequent light quantities (1 unit/day, <1/week), and frequent light quantities (<14 units/week). We used the rich data on social circumstances collected at 9 months and, for the mother's childhood circumstances, at 3 years to describe the social profile of these three patterns of use.

Our findings suggest that the patterns of alcohol use reported by the wider population are also evident among mothers with young children. Studies of women and of young adults suggest that increasingly advantaged social circumstances are associated with increased frequency of alcohol consumption [11,13]. There is also evidence for adult women, young adults, and mothers that social disadvantage are associated with higher quantities of consumption on each drinking occasion [11,13,17].

The study adds to this evidence by employing a more comprehensive range of indicators of mothers' social background. Focusing on the most common patterns of alcohol use among mothers with infants, it confirms that the frequency and quantity of alcohol use vary in line with their socioeconomic and domestic circumstances. These findings have implications for future research and public health interventions from a population perspective with regards to the underlying social determinants of alcohol use among mothers and its potential effect on children.

ACKNOWLEDGMENTS

We are grateful to The Centre for Longitudinal Studies, Institute of Education for the use of these data and to the UK Data Archive and Economic and Social Data Service for making them available. However, they bear no responsibility for the analysis or interpretation of these data. This study was undertaken as part of an ESRC-funded doctoral studentship, award code: A0052802.

REFERENCES

1. Drink aware. Frequently asked questions about alcohol. What is a unit? 2013. Available from: <http://www.drinkaware.co.uk/check-the-facts/what-is-alcohol/frequently-asked-questions>. [Accessed on 2014 Apr 22].
2. International Center for Alcohol Policies, ICAP Reports 14. International Drinking Guidelines. Washington, DC: ICAP; 2003.
3. Smith L, Foxcroft D. Drinking in the UK. An Exploration of Trends. York: Joseph Rowntree Foundation; 2009.
4. Kuntsche S, Gmel G, Knibbe RA, Kuendig H, Bloomfield K, Kramer S, *et al*. Gender and cultural differences in the association between family roles, social stratification, and alcohol use: A European cross-cultural analysis. *Alcohol Alcohol Suppl* 2006;41:i37-46.

5. Bloomfield K, Grittner U, Kramer S, Gmel G. Social inequalities in alcohol consumption and alcohol-related problems in the study countries of the EU concerted action 'Gender, Culture and Alcohol Problems: A Multi-national Study'. *Alcohol Alcohol Suppl* 2006;41:i26-36.
6. Rose G. Sick individuals and sick populations. *Int J Epidemiol* 1985;14:32-8.
7. Fuller E. Health and lifestyles: Adult alcohol consumption. In: Craig R, Hirani V, editors. *The Health Survey for England 2009*. Vol. 1., Ch. 10. England: NHS Information Centre; 2010.
8. Sharp C. Adult health: Alcohol consumption. In: Bromley C, Given L, Ormston R, editors. *The Scottish Health Survey 2009*. Vol. 1., Ch. 3. Edinburgh: Scottish Executive; 2010.
9. The Welsh Assembly Government. Health-related lifestyle. In: Walters L, Kingdon A, Roberts R, editors. *The Welsh Health Survey 2009*. Ch. 4. Wales: The Welsh Assembly Government; 2010.
10. Department of Health Social Services and Public Safety, Health Survey Northern Ireland: Results from the 2010/2011 Survey. Ireland: Information and Analysis Directorate: Health Survey Northern Ireland; 2011.
11. Giske K, Turrell G, Bentley R, Kavanagh A. Individual and household-level socioeconomic position is associated with harmful alcohol consumption behaviours among adults. *Aust N Z J Public Health* 2011;35:270-7.
12. Christie-Mizell CA, Peralta RL. The gender gap in alcohol consumption during late adolescence and young adulthood: Gendered attitudes and adult roles. *J Health Soc Behav* 2009;50:410-26.
13. Casswell S, Pledger M, Hooper R. Socioeconomic status and drinking patterns in young adults. *Addiction* 2003;98:601-10.
14. Kokko K, Pulkkinen L, Mesiaenen P. Timing of parenthood in relation to other life transitions and adult social functioning. *Int J Behav Dev* 2009;33:356-65.
15. Little M, Handley E, Leuthe E, Chassin L. The impact of parenthood on alcohol consumption trajectories: Variations as a function of timing of parenthood, familial alcoholism, and gender. *Dev Psychopathol* 2009;21:661-82.
16. Maloney E, Hutchinson D, Burns L, Mattick R. Prevalence and patterns of problematic alcohol use among Australian parents. *Aust N Z J Public Health* 2010;34:495-501.
17. Stroup-Benham CA, Treviño FM, Treviño DB. Alcohol consumption patterns among Mexican American mothers and among children from single- and dual-headed households: Findings from HHANES 1982-84. *Am J Public Health* 1990;80:36-41.
18. Waterson EJ. Women and alcohol: The social context of changing patterns of use during pregnancy and early motherhood. In: London School of Economics and Political Science. London: University of London; 1992.
19. Valentine G, Holloway SL, Jayne M, Knell C. *Drinking places: Where people drink and why*. York: Joseph Rowntree Foundation; 2007.
20. Vermeulen-Smit E, Koning IM, Verdurmen JE, Van der Vorst H, Engels RC, Vollebbergh WA. The influence of paternal and maternal drinking patterns within two-partner families on the initiation and development of adolescent drinking. *Addict Behav* 2012;37:1248-56.
21. Dex S, Joshi H. *Children of the 21st Century: From Birth to Nine Months*. Bristol: The Policy Press; 2005.
22. Plewis I, Ketende S. editors. *Millennium Cohort Study Technical Report on Response*. 2nd ed. London: Centre for Longitudinal Studies, University of London; 2007.
23. Office for National Statistics. *Ethnic Group Statistics: A Guide for the Collection and Classification of Ethnicity Data*, 2003. Available from: <http://www.statistics.gov.uk>. [Accessed on 2014 Apr 22].
24. Rose D, Pevalin D, O'Reilly K. *The National Statistics Socio-Economic Classification: Origins, Development and Use*. London: Palgrave; 2005.
25. StataCorp, *Stata Statistical Software: Release 10*. College Station, TX: StataCorp LP; 2007.
26. Goddard E. *Obtaining Information about Drinking through Surveys of the General Population*. National Statistics Methodology Series No. 24, London: ONS; 2001.
27. Del Boca FK, Daries J. The validity of self-reports of alcohol consumption: State of the science and challenges for research. *Addiction* 2003;98 Suppl 2:1-12.
28. Sommers MS, Howe SR, Dyehouse JM, Fleming M, Fargo JD, Schafer JC. Patterns of drinking four weeks prior to an alcohol-related

- vehicular crash. *Traffic Inj Prev* 2005;6:110-6.
29. Zhao J, Stockwell T, Macdonald S. Non-response bias in alcohol and drug population surveys. *Drug Alcohol Rev* 2009;28:648-57.
 30. NHS Information Centre. Lifestyles statistics. *Statistics on Alcohol: England, 2011*. London: The Health and Social Care Information Centre; 2011.
 31. Office for National Statistics. *Drinking: Adult's behaviour and knowledge in 2009*. London: NHS Information Centre for Health and Social Care, OPSI; 2010. p. 1-137.
 32. Office for National Statistics. *Labour Market Statistics, 2013*. Available from: http://www.ons.gov.uk/ons/dcp171778_297429.pdf. [Accessed on 2014 Jul 02].
 33. Cribb J, Hood A, Joyce R, Phillips D. *Living standards, poverty and inequality in the UK*. IFS Reports, No. R81. London: Institute for Fiscal Studies; 2013.
 34. Department for Education and Skills. *Participation in Education, Training and Employment by 16-18 Year Olds in England: 2005 and 2006 and Participation in Education and Training by 16 and 17 Year Olds in each Local Area in England: 2004 and 2005*. London: DFES; 2007.
 35. Office of National Statistics, Statistical Bulletin. *Live Births in England and Wales by Characteristics of Mother 1, 2011, 2013*. Available from: http://www.ons.gov.uk/ons/dcp171778_296157.pdf. [Accessed on 2014 Jul 02].
 36. Robinson S, Harris H. *Smoking and drinking among adults, 2009*. In: Dunstan S, editor. *A Report on the 2009 General Lifestyle Survey*. Office for National Statistics. 2011. p. 1-84.
 37. Fuller E. *Adult alcohol consumption*. In: Craig R, Hirani V, editors. *Health Survey for England 2009: Health and Lifestyles*. Vol. 1., Ch. 10. The NHS Information Centre for Health and Social Care; 2011. p. 167-76.

© GESDAV; licensee GESDAV. This is an open access article licensed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted, non-commercial use, distribution and reproduction in any medium, provided the work is properly cited.

Source of Support: Nil, Conflict of Interest: None declared.