

Enterococcus gallinarum carrying the *vanA* gene cluster: first report in Brazil

I.L.B.C. Camargo¹,
A.L. Barth², K. Pilger²,
B.G.S. Seligman²,
A.R.L. Machado² and
A.L.C. Darini¹

¹Departamento de Análises Clínicas, Toxicológicas e Bromatológicas,
Faculdade de Ciências Farmacêuticas de Ribeirão Preto,
Universidade de São Paulo, Ribeirão Preto, SP, Brasil
²Unidade de Pesquisa Biomédica, Serviço de Patologia Clínica,
Hospital de Clínicas de Porto Alegre, Porto Alegre, RS, Brasil

Abstract

Correspondence

A.L.C. Darini
Departamento de Análises Clínicas
Toxicológicas e Bromatológicas
Faculdade de Ciências Farmacêuticas
de Ribeirão Preto, USP
Av. do Café, s/n
14040-903 Ribeirão Preto, SP
Brasil
E-mail: aldarini@fcrp.usp.br

A preliminary version of this study was presented at the I South Brazilian Congress of Clinical Microbiology, Curitiba, PR, Brazil, August 9-11, 2002.

Research supported by FAPESP (No. 99/12233-0) and FIPE/HCPA (No. 01/203). I.L.B.C. Camargo was the recipient of a CNPq fellowship (No. 130486/2001-9).

Received March 10, 2004
Accepted July 15, 2004

In 2000, *Enterococcus faecalis* resistant to vancomycin was first reported at a tertiary hospital in Porto Alegre, southern Brazil. The resistance spread to other hospitals and surveillance programs were established by hospital infection committees to prevent the spread of vancomycin-resistant enterococci. In February 2002, an isolate initially identified at the genus level as *Enterococcus* was obtained by surveillance culture (rectal swab) from a patient admitted to a hospital for treatment of septic arthritis in the shoulder. The isolate proved to be resistant to vancomycin by the disc diffusion method and confirmed by an E-test resulting in a minimal inhibitory concentration of ≥ 256 $\mu\text{g/ml}$. This isolate was sent to a reference laboratory (Laboratório Especial de Bacteriologia e Epidemiologia Molecular, Faculdade de Ciências Farmacêuticas de Ribeirão Preto, USP) for further study and proved to be an *E. gallinarum* by the polymerase chain reaction (PCR) using specific primers for the species. Due to the phenotype of unusually high vancomycin resistance, the isolate presumably had the resistance genes (*vanA* and *vanB*) and this was confirmed by PCR, which indicated the presence of the *vanA* gene. A 10.8-kb Tn1546-related transposon was also identified by long-PCR. Interspecies transfer of the vancomycin-resistance gene from the donor *E. gallinarum* was performed in a successful conjugation experiment *in vitro*, using *E. faecium* GE-1 and *E. faecalis* JH22 as receptors. This is the first report of the detection of a *vanA* determinant naturally acquired by *E. gallinarum* in Brazil, indicating the importance of characterizing VRE by both phenotype and genotype methods.

Key words

- *Enterococcus gallinarum*
- VanA phenotype
- Vancomycin
- Resistance

Enterococci have emerged as increasingly important nosocomial and community-acquired pathogens. First described in 1988 in France and England (1,2), glycopeptide-resistant enterococci have been isolated in Brazil since 1996, when the first case of vancomycin-resistant enterococci (VRE) due to a *Enterococcus faecium* VanD₄ was reported in Curitiba (Paraná State) (3,4). Further VRE

isolates (*E. faecium* with the *vanA* gene) were reported in São Paulo State (5,6).

Glycopeptide resistance in enterococci is associated with a variety of phenotypes and genotypes. VanA and VanB are the two major genetically distinct forms of acquired resistance. The VanA phenotype displays a high level of resistance to vancomycin and teicoplanin but the VanB phenotype displays

a high level of resistance only to vancomycin. However, both have been found primarily in *E. faecalis* and *E. faecium* (7). In contrast, *E. gallinarum*, *E. casseliflavus* and *E. flavescens* have an intrinsic low-level resistance to vancomycin due to the VanC phenotype (7). The *vanC-1* gene is specific for *E. gallinarum*, and *vanC-2/3* is specific for *E. casseliflavus* and *E. flavescens*. Nevertheless, a few cases of *vanA* and *vanB* genes acquired by *E. gallinarum* have been reported: in Switzerland (8), Australia (9), Italy (10), Belgium (11), and Taiwan (12).

In 2000, vancomycin-resistant *E. faecalis* was first described at a tertiary hospital in Porto Alegre, southern Brazil. The resistance spread to other hospitals and surveillance programs were established by the hospital infection committees in order to prevent the spreading of VRE. In February 2002, an isolate initially identified at the genus level as *Enterococcus* by a few basic tests (Gram, bile esculin, catalase, growth on 6.5% NaCl and L-pyrrolidonyl- β -naphthylamide, PYR) was obtained at Hospital de Clínicas de Porto Alegre by surveillance culture (rectal swab) from a patient admitted to the hospital due to septic arthritis in the shoulder. The isolate proved to be resistant to vancomycin by the disc diffusion method




Figure 1. Conjugation experiments. A, Plasmid extraction product in 1% agarose gel; B, Southern blotting hybridization of *vanA* probes in a nylon membrane. Lane 1, *E. gallinarum* ET193; lane 2, transconjugant of *E. faecium* GE-1; lane 3, transconjugant of *E. faecalis* JH22. crDNA = chromosomal DNA.

carried out according to NCCLS (13) and this was confirmed by the E-test (AB Biodisk, Sweden) resulting in a minimal inhibitory concentration of ≥ 256 $\mu\text{g/ml}$.

This isolate was sent to a reference laboratory (Laboratório Especial de Bacteriologia e Epidemiologia Molecular, Faculdade de Ciências Farmacêuticas de Ribeirão Preto) for further study and proved to be an *E. gallinarum* (termed ET193) by the polymerase chain reaction (PCR) using specific primers for the species (11). Due to the phenotype of unusually high level of vancomycin resistance, the isolate was tested for the presence of specific resistance genes (*vanA* and *vanB*) by PCR, which indicated the presence of the *vanA* gene (14,15).

The high level of vancomycin resistance among enterococcus isolates is often mediated by a self-transferable plasmid that has acquired the Tn1546-related transposons, which carry the *vanA* gene cluster (16). We demonstrated a Tn1546-related 10.8-kb transposon which was found to be identical to the Tn1546 prototype in the *E. gallinarum* ET193 isolate by long-PCR (17).

Interspecies transfer of the vancomycin resistance determinant from the *E. gallinarum* ET193 donor was successfully carried out by a conjugation experiment *in vitro* using *E. faecium* GE-1 and *E. faecalis* JH22 as receptors (18). After successful conjugation, several transconjugants were obtained by selection on plates containing 4 $\mu\text{g/ml}$ vancomycin, 200 $\mu\text{g/ml}$ rifampicin, and 100 $\mu\text{g/ml}$ fusidic acid. Plasmid extraction was performed and a plasmid of about 70 kb was found in the transconjugants and in the *E. gallinarum* ET193 donor isolate. Eventually, the electrophoretic profile was analyzed by Southern blot and hybridization using a digoxigenin-labeled *vanA* gene probe in order to locate the presence of the gene. The gene-specific probe was bound to both plasmids and chromosomal DNA of the several transconjugants and donor *E. gallinarum* ET193 (Figure 1).

This is the first case of *E. gallinarum* carrying the *vanA* gene cluster isolated in Brazil. As the patient was also colonized by a *vanA* genotype vancomycin-resistant *E. faecalis*, it is reasonable to speculate that this gene cluster was transferred from *E. faecalis* to *E. gallinarum* *in vivo*. This was, most probably, a natural acquisition of a peculiar high level resistance gene by the *E. galli-*

narum species and indicates the importance of characterizing VRE by both phenotype and genotype methods.

Acknowledgments

The authors thank Izabel Cristina Vanzato Palazzo for technical assistance during the transfer experiments.

References

- Uttley AHC, Collins CH, Naidoo J & George RC (1988). Vancomycin-resistant enterococci. *Lancet*, 1: 57-58.
- Leclercq R, Derlot E, Duval J & Courvalin P (1988). Plasmid-mediated resistance to vancomycin and teicoplanin in *Enterococcus faecium*. *New England Journal of Medicine*, 319: 157-161.
- Dalla Costa LM, Souza DC, Martins LTF, Zanella RC, Brandileone MC, Bokermann S, Sader HS & Souza APHM (1998). Vancomycin-resistant *Enterococcus faecium*: First case in Brazil. *Brazilian Journal of Infectious Diseases*, 3: 160-163.
- Dalla Costa LM, Reynolds PE, Souza HA, Souza DC, Palepou MF & Woodford N (2000). Characterization of a divergent VanD-type resistance element from the first glycopeptide-resistant strain of *Enterococcus faecium* isolated in Brazil. *Antimicrobial Agents and Chemotherapy*, 44: 3444-3446.
- Cereda RF, Gales AC, Silbert S, Jones RN & Sader HS (2002). Molecular typing and antimicrobial susceptibility of vancomycin-resistant *Enterococcus faecium* in Brazil. *Infection Control and Hospital Epidemiology*, 23: 19-22.
- Zanella RC, Valdetaro F, Lovgren M, Tyrrel GJ, Bokermann S, Almeida SCG, Vieira VSD & Brandileone MCC (1999). First confirmed case of a vancomycin-resistant *Enterococcus faecium* with Van A phenotype from Brasil: isolation from a meningitis case in São Paulo. *Microbial Drug Resistance*, 5: 159-162.
- Woodford N (1998). Glycopeptide-resistant enterococci: a decade of experience. *Journal of Medical Microbiology*, 47: 849-862.
- Liassine N, Frei R, Jan I & Auckenthaler R (1998). Characterization of glycopeptide-resistant Enterococci from a Swiss Hospital. *Journal of Clinical Microbiology*, 36: 1853-1858.
- Schooneveldt JM, Marriott RK & Nimmo GR (2000). Detection of *vanB* determinant in *Enterococcus gallinarum* in Australia. *Journal of Clinical Microbiology*, 38: 3902.
- Biavasco F, Paladini C, Vignaroli C, Foglia G, Manso E & Varaldo PE (2001). Recovery from a single blood culture of two *Enterococcus gallinarum* isolates carrying both *vanC-1* and *vanA* cluster genes and differing in glycopeptide susceptibility. *European Journal of Clinical Microbiology and Infectious Diseases*, 20: 309-314.
- Dutka-Malen S, Blaimont B, Wauters G & Courvalin P (1994). Emergence of high-level resistance to glycopeptides in *Enterococcus gallinarum* and *Enterococcus casseliflavus*. *Antimicrobial Agents and Chemotherapy*, 7: 1675-1677.
- Lu JJ, Wu JC, Chiueh TS, Perng CL, Chi WM & Lee WH (2000). Characterization of a highly glycopeptide-resistant *Enterococcus gallinarum* isolate. *Journal of the Formosan Medical Association*, 4: 305-310.
- NCCLS (2002). *Performance Standards for Antimicrobial Susceptibility Testing. Twelfth Informational Supplement M100-S12*. National Committee for Clinical Laboratory Standards, Villanova, PA, USA.
- Woodford N, Morrison D, Johnson AP, Briant V, George RC & Cookson B (1993). Application of DNA probes for rRNA and *vanA* genes to investigation of a nosocomial cluster of vancomycin-resistant Enterococci. *Journal of Clinical Microbiology*, 31: 653-658.
- Woodford N, Morrison D, Johnson AP, Bateman AC, Hasting JGM & Cookson B (1995). Plasmid-mediated *vanB* glycopeptide resistance in enterococci. *Microbial Drug Resistance*, 1: 235-240.
- Molinas AM, Depardieu F & Courvalin P (1993). Characterization of Tn1546, a Tn3-related transposon conferring glycopeptide resistance by synthesis of depsipeptide peptidoglycan precursors in *Enterococcus faecium* BM4147. *Journal of Bacteriology*, 175: 117-127.
- Woodford N, Watson AP & Chadwick PR (1997). Investigation by long PCR of genetic elements mediating VanA glycopeptide resistance in enterococci from uncooked meat in South Manchester. In: Horaud T, Bouvet A, Leclercq R, de Montelos H & Sicard M (Editors), *Streptococci and the Host* 1997. Plenum Press, New York, 406-412.
- Tremlett CH, Brown DFJ & Woodford N (1999). Variation in structure and location of VanA glycopeptide resistance elements among enterococci from a single patient. *Journal of Clinical Microbiology*, 17: 818-820.