



University for the Common Good

## Yoga for stroke rehabilitation

Lawrence, Maggie; Celestino Junior, Franciso T.; Matozinho, Hemilianna H.S.; Govan, Lindsay; Booth, Jo; Beecher, Jane

*Published in:*  
Cochrane Database of Systematic Reviews

*DOI:*  
[10.1002/14651858.CD011483.pub2](https://doi.org/10.1002/14651858.CD011483.pub2)

*Publication date:*  
2017

*Document Version*  
Publisher's PDF, also known as Version of record

[Link to publication in ResearchOnline](#)

### *Citation for published version (Harvard):*

Lawrence, M, Celestino Junior, FT, Matozinho, HHS, Govan, L, Booth, J & Beecher, J 2017, 'Yoga for stroke rehabilitation', *Cochrane Database of Systematic Reviews*, vol. 2017, no. 12, CD011483. <https://doi.org/10.1002/14651858.CD011483.pub2>

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

### **Take down policy**

If you believe that this document breaches copyright please view our takedown policy at <https://edshare.gcu.ac.uk/id/eprint/5179> for details of how to contact us.



**Cochrane**  
**Library**

Cochrane Database of Systematic Reviews

## Yoga for stroke rehabilitation (Review)

Lawrence M, Celestino Junior FT, Matozinho HHS, Govan L, Booth J, Beecher J

Lawrence M, Celestino Junior FT, Matozinho HHS, Govan L, Booth J, Beecher J.

Yoga for stroke rehabilitation.

*Cochrane Database of Systematic Reviews* 2017, Issue 12. Art. No.: CD011483.

DOI: 10.1002/14651858.CD011483.pub2.

[www.cochranelibrary.com](http://www.cochranelibrary.com)

---

Yoga for stroke rehabilitation (Review)

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

**WILEY**

[Intervention Review]

# Yoga for stroke rehabilitation

Maggie Lawrence<sup>1</sup>, Francisco T Celestino Junior<sup>2</sup>, Hemilianna HS Matozinho<sup>2</sup>, Lindsay Govan<sup>3</sup>, Jo Booth<sup>4</sup>, Jane Beecher<sup>5</sup>

<sup>1</sup>Department of Nursing and Community Health, School of Health and Life Sciences, Glasgow Caledonian University, Glasgow, UK. <sup>2</sup>School of Health and Life Sciences, Glasgow Caledonian University, Glasgow, UK. <sup>3</sup>Health Economics and Health Technology Assessment, Institute of Health and Wellbeing, University of Glasgow, Glasgow, UK. <sup>4</sup>Department of Health & Community Sciences, Glasgow Caledonian University, Glasgow, UK. <sup>5</sup>Rotational Physiotherapy, Queen Elizabeth University Hospital, Glasgow, UK

Contact address: Maggie Lawrence, Department of Nursing and Community Health, School of Health and Life Sciences, Glasgow Caledonian University, A101f, Govan Mbeki Building, Glasgow, G4 0BA, UK. [maggie.lawrence@gcu.ac.uk](mailto:maggie.lawrence@gcu.ac.uk).

**Editorial group:** Cochrane Stroke Group.

**Publication status and date:** New, published in Issue 12, 2017.

**Citation:** Lawrence M, Celestino Junior FT, Matozinho HHS, Govan L, Booth J, Beecher J. Yoga for stroke rehabilitation. *Cochrane Database of Systematic Reviews* 2017, Issue 12. Art. No.: CD011483. DOI: 10.1002/14651858.CD011483.pub2.

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

## ABSTRACT

### Background

Stroke is a major health issue and cause of long-term disability and has a major emotional and socioeconomic impact. There is a need to explore options for long-term sustainable interventions that support stroke survivors to engage in meaningful activities to address life challenges after stroke. Rehabilitation focuses on recovery of function and cognition to the maximum level achievable, and may include a wide range of complementary strategies including yoga.

Yoga is a mind-body practice that originated in India, and which has become increasingly widespread in the Western world. Recent evidence highlights the positive effects of yoga for people with a range of physical and psychological health conditions. A recent non-Cochrane systematic review concluded that yoga can be used as self-administered practice in stroke rehabilitation.

### Objectives

To assess the effectiveness of yoga, as a stroke rehabilitation intervention, on recovery of function and quality of life (QoL).

### Search methods

We searched the Cochrane Stroke Group Trials Register (last searched July 2017), Cochrane Central Register of Controlled Trials (CENTRAL) (last searched July 2017), MEDLINE (to July 2017), Embase (to July 2017), CINAHL (to July 2017), AMED (to July 2017), PsycINFO (to July 2017), LILACS (to July 2017), SciELO (to July 2017), IndMED (to July 2017), OTseeker (to July 2017) and PEDro (to July 2017). We also searched four trials registers, and one conference abstracts database. We screened reference lists of relevant publications and contacted authors for additional information.

### Selection criteria

We included randomised controlled trials (RCTs) that compared yoga with a waiting-list control or no intervention control in stroke survivors.

### Data collection and analysis

Two review authors independently extracted data from the included studies. We performed all analyses using Review Manager (RevMan). One review author entered the data into RevMan; another checked the entries. We discussed disagreements with a third review author until consensus was reached. We used the Cochrane 'Risk of bias' tool. Where we considered studies to be sufficiently similar, we conducted a meta-analysis by pooling the appropriate data. For outcomes for which it was inappropriate or impossible to pool quantitatively, we conducted a descriptive analysis and provided a narrative summary.

---

Yoga for stroke rehabilitation (Review)

Copyright © 2017 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

WILEY

## Main results

We included two RCTs involving 72 participants. Sixty-nine participants were included in one meta-analysis (balance). Both trials assessed QoL, along with secondary outcomes measures relating to movement and psychological outcomes; one also measured disability.

In one study the Stroke Impact Scale was used to measure QoL across six domains, at baseline and post-intervention. The effect of yoga on five domains (physical, emotion, communication, social participation, stroke recovery) was not significant; however, the effect of yoga on the memory domain was significant (mean difference (MD) 15.30, 95% confidence interval (CI) 1.29 to 29.31,  $P = 0.03$ ), the evidence for this finding was very low grade. In the second study, QoL was assessed using the Stroke-Specific QoL Scale; no significant effect was found.

Secondary outcomes included movement, strength and endurance, and psychological variables, pain, and disability.

Balance was measured in both studies using the Berg Balance Scale; the effect of intervention was not significant (MD 2.38, 95% CI -1.41 to 6.17,  $P = 0.22$ ). Sensitivity analysis did not alter the direction of effect. One study measured balance self-efficacy, using the Activities-specific Balance Confidence Scale (MD 10.60, 95% CI -7.08, to 28.28,  $P = 0.24$ ); the effect of intervention was not significant; the evidence for this finding was very low grade.

One study measured gait using the Comfortable Speed Gait Test (MD 1.32, 95% CI -1.35 to 3.99,  $P = 0.33$ ), and motor function using the Motor Assessment Scale (MD -4.00, 95% CI -12.42 to 4.42,  $P = 0.35$ ); no significant effect was found based on very low-grade evidence.

One study measured disability using the modified Rankin Scale (mRS) but reported only whether participants were independent or dependent. No significant effect was found: (odds ratio (OR) 2.08, 95% CI 0.50 to 8.60,  $P = 0.31$ ); the evidence for this finding was very low grade.

Anxiety and depression were measured in one study. Three measures were used: the Geriatric Depression Scale-Short Form (GDS15), and two forms of State Trait Anxiety Inventory (STAI, Form Y) to measure state anxiety (i.e. anxiety experienced in response to stressful situations) and trait anxiety (i.e. anxiety associated with chronic psychological disorders). No significant effect was found for depression (GDS15, MD -2.10, 95% CI -4.70 to 0.50,  $P = 0.11$ ) or for trait anxiety (STAI-Y2, MD -6.70, 95% CI -15.35 to 1.95,  $P = 0.13$ ), based on very low-grade evidence. However, a significant effect was found for state anxiety: STAI-Y1 (MD -8.40, 95% CI -16.74 to -0.06,  $P = 0.05$ ); the evidence for this finding was very low grade.

No adverse events were reported.

## Quality of the evidence

We assessed the quality of the evidence using GRADE. Overall, the quality of the evidence was very low, due to the small number of trials included in the review both of which were judged to be at high risk of bias, particularly in relation to incompleteness of data and selective reporting, and especially regarding the representative nature of the sample in one study.

## Authors' conclusions

Yoga has the potential for being included as part of patient-centred stroke rehabilitation. However, this review has identified insufficient information to confirm or refute the effectiveness or safety of yoga as a stroke rehabilitation treatment. Further large-scale methodologically robust trials are required to establish the effectiveness of yoga as a stroke rehabilitation treatment.

## PLAIN LANGUAGE SUMMARY

### Yoga for stroke rehabilitation

#### Review question

We wanted to know if yoga helps to improve quality of life for stroke survivors.

#### Background

Stroke is a major health issue worldwide, which affects people in many different ways. For example, stroke survivors may have problems moving around, and communicating and socialising with other people. Stroke may also affect how people feel. It may cause problems with memory and concentration. After discharge from hospital or other stroke services, stroke survivors have to cope with the long-

term effects of stroke. Research has shown that yoga can help people with other long-term conditions to cope better. Yoga can improve quality of life (QoL).

#### **Search date**

We searched for studies published to July 2017.

#### **Study characteristics**

We found two research studies that had assessed yoga for stroke survivors. Seventy-two people took part in the two studies. One study was in the USA and one was in Australia. On average, the stroke survivors were between 60 and 63 years old and it had been between four years three months and nine years since they had had a stroke. In the American study, yoga classes were held twice a week for eight weeks. In the Australian study, yoga classes were held once a week for 10 weeks. Both studies encouraged people to practice yoga at home, in their own time. Both studies used waiting-list control groups. This means that people in the control group could go to yoga classes at the end of the study.

#### **Funding sources**

The American study was funded by the US Government. The Australian study was funded by the National Stroke Foundation (Australia).

#### **Key results**

We were able to analyse study data from 69 participants. No significant benefit was found on measures of QoL, balance, strength, endurance, pain, disability scores. No significant benefit was found on measures of movement, although one study reported a significant benefit in improving aspects of range of movement. One study reported a significant benefit in reducing anxiety. Neither study reported on measures of patient harm.

#### **Quality of the evidence**

We assessed the quality of the evidence using GRADE. Overall, the quality of the evidence was very low, due to the small number of trials included in the review, both of which we judged to be at high risk of bias, particularly in relation to incompleteness of data and selective reporting, and especially regarding the representative nature of the sample in one study.

#### **Conclusion**

The review could not identify enough high-quality evidence on the benefits and safety of yoga in stroke rehabilitation. More good-quality research studies are needed to be sure that yoga has benefits for stroke survivors.