

## Tests of top compositeness at hadron colliders

C. DEGRANDE<sup>(\*)</sup>*Centre for Cosmology, Particle Physics and Phenomenology (CP3)  
Université catholique de Louvain - Chemin du Cyclotron, 2  
B-1348, Louvain-la-Neuve, Belgium*

(ricevuto il 16 Luglio 2010; approvato il 16 Luglio 2010; pubblicato online il 15 Ottobre 2010)

**Summary.** — Top pair production can be used to probe composite top models. Associated with 4-top and  $t\bar{t}b\bar{b}$  productions, it can be used to distinguish different hypotheses.

PACS 14.65.Ha – Top quarks.

PACS 12.60.Rc – Composite models.

### 1. – Introduction

The effects of top compositeness can be parametrized by adding order by order higher-dimensional operators to the SM Lagrangian,

$$(1) \quad \mathcal{L} = \mathcal{L}_{\text{SM}} + \frac{1}{\Lambda^2} \sum_i c_i \mathcal{O}_6^i + \mathcal{O} \left( \frac{1}{\Lambda^4} \right).$$

There are two main reasons to do so. On the one hand, the usual perturbative expansion cannot be used because the new interaction is strong. On the other hand, this effective approach is able to describe a large class of models even beyond composite top models. Naive dimensional analysis (NDA) [1] has been used to classify the operators. Each coefficient  $c_i$  is then a function of  $\xi$ , with  $\xi \sim 4\pi$  for strongly coupled theories.

### 2. – Top pair production

**2.1. Lagrangian.** – Only a few operators contribute to the top pair production [2]:

$$(2) \quad \mathcal{L}_{t\bar{t}} = \mathcal{L}_{t\bar{t}}^{\text{SM}} + \frac{1}{\Lambda^2} [g_h \mathcal{O}_{hg} + c_R \mathcal{O}_{Rg} + a_R \mathcal{O}_{Ra}^{\text{S}} + \text{h.c.} + (R \leftrightarrow L)],$$

<sup>(\*)</sup> E-mail: [celine.degrande@uclouvain.be](mailto:celine.degrande@uclouvain.be)




Fig. 1. – In light (dark) gray, Tevatron allowed region by the total cross-section measurement (the  $M_{tt}$  distribution). LHC allowed regions if the measured total cross-section is the one predicted by the SM are delimited by the thick black lines.

where

$$(3) \quad \mathcal{O}_{hg} = [(H\bar{Q}) \sigma^{\mu\nu} T^A P_R t] G_{\mu\nu}^A, \quad \mathcal{O}_{Ra}^8 = [\bar{t} \gamma^\mu T^A P_R t] \sum_q [\bar{q} \gamma_\mu T^A \gamma_5 q],$$

$$(4) \quad \mathcal{O}_{Rg} = [\bar{t} \gamma^\mu T^A D^\nu P_R t] G_{\mu\nu}^A.$$

If only  $t_R$  is composite,  $c_R \sim a_R \sim 1$ ,  $c_L = a_L = 0$  and  $g_h \sim 1/\xi$ . If, on the contrary, only  $Q_L$  is composite,  $c_L \sim a_L \sim 1$ ,  $c_R = a_R = 0$  and  $g_h \sim 1/\xi$ . If both chiralities are composite,  $c_R \sim a_R \sim c_L \sim a_L \sim g_h \sim 1$ . So, the magnitude of  $g_h$  compared to the other coefficients counts the number of composite fields.

**2.2. Total cross-section and  $M_{tt}$  distribution.** – These observables only depend on  $g_h$  and the combination  $c_V \equiv c_R + c_L$ . The  $t\bar{t}$  production via gluon fusion only depending on  $g_h$ , the Tevatron and the LHC are rather complementary as shown in fig. 1. The distortion of the invariant mass distribution is mainly due to the operators  $\mathcal{O}_{R,Lg}$  and is thus less visible at the LHC. It should be noted that, for low values of  $\Lambda$ , Tevatron measurements already imply that  $g_h \sim c_V$ , and suggest 2 composite fields.

**2.3. Forward-backward asymmetry.** – The forward-backward asymmetry measured at the Tevatron,  $A_{\text{FB}}^t = 0.19 \pm 0.065(\text{stat}) \pm 0.024(\text{syst})$  [3], is about  $2\sigma$  away from the SM value,  $A_{\text{FB}}^t = 0.05 \pm 0.015$ . This large deviation could be explained by the compositeness of the top if  $\Lambda \sim 1$  TeV and  $a_A \equiv a_R - a_L$  is  $\mathcal{O}(1)$ :

$$(5) \quad \delta A_{\text{FB}}^t = 0.0342_{-0.009}^{+0.016} a_A \left( \frac{1 \text{ TeV}}{\Lambda} \right)^2.$$

**2.4. Spin correlation.** – Left- and right-handed composite top cannot be distinguished using only kinematic observables. Fortunately there is a strong correlation between the top spin and the direction of the lepton coming from its decay [4]. We have shown that the distribution as a function of the charged leptons directions is not only sensitive to  $g_h$  and  $c_V$  but also to the combination  $c_A \equiv c_R - c_L$  and is thus able to disentangle between the left- and right-handed composite top.

### 3. – 4-top and $t\bar{t}b\bar{b}$ productions

Contrary to top pair production, the 4-top one can probe the dominant  $\mathcal{O}(\xi^2)$  operators for a right-handed composite top,

$$(6) \quad \mathcal{O}_R = [\bar{t}\gamma^\mu P_R t] [\bar{t}\gamma_\mu P_R t]$$

and for a left-handed composite top,

$$(7) \quad \mathcal{O}_L = [\bar{Q}\gamma^\mu P_L Q] [\bar{Q}\gamma_\mu P_L Q], \quad \mathcal{O}_L^8 = [\bar{Q}\gamma^\mu T^A P_L Q] [\bar{Q}\gamma_\mu T^A P_L Q].$$

When both chiralities are composite, there are 2 additional operators:

$$(8) \quad \mathcal{O}_S = [\bar{Q}^\alpha P_R t] [\bar{t} P_L Q^\alpha], \quad \mathcal{O}_S^8 = [\bar{Q}^\alpha T^A P_R t] [\bar{t} T^A P_L Q^\alpha].$$

In the above expressions, we have assumed that  $SU(2)_L$  is unbroken by the new interaction, *i.e.* the full doublet, including  $b_L$ , is composite. Yet, all these operators lead to similar cross-sections between 2 and 20 pb for  $g = 4\pi$  and  $\Lambda = 1$  TeV. They can thus all fit any measurement by a slight change of their coefficient  $g^2/\Lambda^2$  since  $\sigma \sim \left(\frac{g}{\Lambda}\right)^4$ . If  $Q_L$  is composite, there will also be a modification of  $t\bar{t}b\bar{b}$  cross-section. The ratio of the 2 processes can be used to identify the operator because it is independent of  $g/\Lambda$  but strongly dependent on the operator. Its coefficient can then be extracted.

### 4. – Conclusion

Top pair production is a good probe for new physics. First, the total cross-section and the invariant mass distribution can pin down the number of composite fields. Secondly, the forward-backward asymmetry could be the first hint of top compositeness from which we can estimate the scale. Thirdly, the spin correlation differentiates left- from right-handed composite top. The  $t\bar{t}t\bar{t}$  and  $t\bar{t}b\bar{b}$  productions can probe the hierarchy among the dominant and subdominant operators.

\* \* \*

These are preliminary results of a work done in collaboration with J.-M. GÉRARD, C. GROJEAN, F. MALTONI and G. SERVANT. This work was supported by the Fonds National de la Recherche Scientifique, by the Belgian Federal Office for Scientific, Technical and Cultural Affairs through the Interuniversity Attraction Pole No. P6/11 and by the European Research Council Starting Grant Cosmo@LHC.

### REFERENCES

- [1] MANOHAR A. and GEORGI H., *Nucl. Phys. B*, **234** (1984) 189; GEORGI H., *Phys. Lett. B*, **298** (1993) 187.
- [2] DEGRANDE C., GÉRARD J.-M., GROJEAN C., MALTONI F. and SERVANT G., in preparation.
- [3] STRYCKER G. *et al.*, CDF note 9724 (2009).
- [4] MAHLON G. and PARKE S. J., *Phys. Rev. D*, **53** (1996) 4886.