

IL NUOVO CIMENTO  
DOI 10.1393/ncc/i2011-10802-5

VOL. 33 C, N. 6

Novembre-Dicembre 2010

COLLOQUIA: IFAE 2010

## Inclusive W and Z production at LHC startup with the CMS experiment

M. DE GRUTTOLA

*Università "Federico II" and INFN, Sezione di Napoli - Napoli, Italy*

(ricevuto l'8 Ottobre 2010; pubblicato online il 15 Febbraio 2011)

**Summary.** — W and Z candidates decaying into leptons can be observed in the very early phase of the LHC run. The results for the W/Z production presented are obtained for 10 TeV simulated data and are fully valid for 7 TeV center-of-mass energy (via a scale factor of the W and Z signals around 0.7).

PACS 13.38.Be – Decays of W bosons.

PACS 13.38.Dg – Decays of Z bosons.

### 1. – Introduction

W and Z bosons decay into isolated high transverse momentum leptons with a clean experimental signature. The W/Z cross section into muons and electron is given by

$$(1) \quad \sigma(W/Z) = \frac{N_{\text{sig}} - N_{\text{bkgd}}}{A_{W/Z} \times \epsilon_{W/Z} \times \int L dt},$$

where  $N_{\text{sig,bkg}}$  are the number of signal and background in each channel;  $\epsilon_{W/Z}$  are the efficiency to detect and trigger the leptons and the neutrino;  $A_{W/Z}$  is the geometric acceptance of the signal from Monte Carlo.

### 2. – W and Z

High momentum electrons in CMS are reconstructed by matching a cluster in the electromagnetic calorimeter (ECAL) and track in the silicon Tracker. The selection of the  $W \rightarrow e\nu$  [1] decay starts from a cluster with high transverse energy. The neutrino instead results in events with unbalanced energy in the transverse plane ( $E_T^{\text{miss}}$ ). To identify the  $Z \rightarrow e^+e^-$  [1] decay one looks for two isolated and opposite sign clusters with high  $E_T$  and very high Electron ID footprint.

High- $p_T$  muons are reconstructed in CMS matching a track in the muon chambers outside the magnet plus a track in the tracker. The  $W \rightarrow \mu\nu$  [2] selection starts requiring




Fig. 1. –  $E_T^{\text{miss}}$  distribution for  $W \rightarrow e\nu$  and background after selection.




Fig. 2. –  $M_{ee}$  distribution for  $Z \rightarrow e^+e^-$  and background.




Fig. 3. –  $M_T^{\text{miss}}$  distribution for  $W \rightarrow \mu\nu$  and background after selection.




Fig. 4. –  $M_{\mu\mu}$  distribution for  $Z \rightarrow \mu^+\mu^-$  and background.

a global track with high transverse momentum  $p_T$ . The transverse mass of the W,  $M_T$ , is computed by combining the measured muon and the missing transverse energy in the event. The  $Z \rightarrow \mu^+\mu^-$  [2] selection requires two opposite signs tracks with high  $p_T$ .

Estimation of the background from W signal using data is done with the “template method”: the signal missing energy shape is built from the observed missing energy in

Z events, while the QCD shape is obtained applying the full set of selection cuts but reverting the isolation one. The Z signal is instead almost background free.

A common way to measure the efficiencies on data is to use the “Tag & Probe” method on Z events.

Figures 1-4 show the preliminary plots obtained by the CMS experiment using 10 TeV signal and background simulated data for events passing the W decaying into electron and neutrino, Z decaying into electrons, W into muon and neutrino and Z decaying into muons, respectively.

### 3. – Conclusion

LHC aims to deliver  $100 \text{ pb}^{-1}$  of integrated luminosity of pp collision data by the end of 2010. We expect to see about  $3000 \text{ W/pb}^{-1}$  in each leptonic channel for the LHC operation at 7 TeV and about  $300 \text{ Z/pb}^{-1}$  in each leptonic channel. The high-purity and well-known properties of the W/Z signals make them one of the most promising channels for the commissioning of CMS physics and lepton identification and selection.

#### REFERENCES

- [1] THE CMS COLLABORATION, CMS PAS-EWK-09/004.
- [2] THE CMS COLLABORATION, CMS PAS-EWK-09/001.