

IL NUOVO CIMENTO
DOI 10.1393/ncc/i2011-10798-8

VOL. 33 C, N. 6

Novembre-Dicembre 2010

COLLOQUIA: IFAE 2010

Perspectives for the measurement of $pp \rightarrow Z \rightarrow \mu\mu + X$ cross-section with the ATLAS detector at LHC

R. DI NARDO

Università di Roma "Tor Vergata" and INFN, Sezione di Roma "Tor Vergata" - Rome, Italy

(ricevuto l'8 Ottobre 2010; pubblicato online il 15 Febbraio 2011)

Summary. — This work describes the measurement of the inclusive cross-section of the channel $Z \rightarrow \mu\mu$ that will represent one of the first physics measurements in ATLAS. A description of the analysis model used together with a discussion on the overall selection efficiency and acceptance calculation will be given. A discussion on the main source of uncertainty on the cross-section measurement will be given as well.

PACS 13.85.Qk – Inclusive production with identified leptons, photons, or other nonhadronic particles.

PACS 13.38.Dg – Decays of Z bosons.

PACS 14.70.Hp – Z bosons.

PACS 12.10.-g – Unified field theories and models.

1. – Introduction

The LHC operations started successfully in November 2009 at the center-of-mass energy of $\sqrt{s} = 900$ GeV and restarted after the winter shutdown in March 2010 at $\sqrt{s} = 7$ TeV. A new long data-taking period is foreseen in 2010-2011 with the goal to reach 1 fb^{-1} of integrated luminosity. The measurement of $Z \rightarrow \mu\mu$ production cross-section with the ATLAS experiment [1,2] will represent a stringent test of QCD and also the measurement of the Z -boson properties, like reconstructed mass and width, can be used to assess the performance of the detector (*i.e.* trigger and reconstruction efficiencies, muon momentum resolution). The measured total cross-section can be expressed as $\sigma = \frac{N-B}{LA\varepsilon}$, where N is the number of events passing a given set of cuts, L is the integrated luminosity, A is the geometrical acceptance of the signal, ε is the reconstruction efficiency of the signal in within the fiducial acceptance and B is the number of background events. The $Z \rightarrow \mu\mu$ cross-section, calculated with FEWZ program at NNLO, is 989 pb at $\sqrt{s} = 7$ TeV. The overall measurement uncertainty gets contributions from the different terms as $\frac{\delta\sigma}{\sigma} = \frac{\delta N \oplus \delta B}{N-B} \oplus \frac{\delta L}{L} \oplus \frac{\delta A}{A} \oplus \frac{\delta\varepsilon}{\varepsilon}$. In this expression, $\delta N \sim 1/\sqrt{N}$ is of purely statistical origin, and the relative uncertainty decreases with increasing luminosity. The terms δB , δA and $\delta\varepsilon$ are of both theoretical and experimental origin. They are considered as systematic uncertainties in the cross-section measurements, but can be constrained via




Fig. 1. – Acceptance as a function of c.m.e. for three different p_T cuts on the muons coming from the Z -boson decay and $|\eta| < 2.5$ (only statistical errors are shown).

auxiliary measurements that will improve over time. The uncertainty on the luminosity measurement, δL , is expected to be at the beginning at the order of 20% and to decrease with the improved understanding of the LHC beam parameters and of the luminosity detector.

2. – Event selection and background rejection

A single-lepton trigger requiring a muon ($p_T > 10$ GeV) is used in the analysis. Events with two muons with $p_T > 20$ GeV and $|\eta| < 2.5$ and opposite charge are selected. The main backgrounds in the analysis are: $t\bar{t}$ events where muons come from the $t\bar{t}$ quarks decay in $t \rightarrow Wb$ and $W \rightarrow \mu\nu$, $W \rightarrow \mu\nu$ where one muon comes from W decay and the other one is a fake muon, $Z \rightarrow \tau\tau$ due to possibility to have muons from τ decay, jet background due in particular to muons coming from b -hadron decay. In order to minimize the impact of this backgrounds, Inner Detector track isolation cuts are also applied ($\Sigma p_T < 5$ GeV and $N_{\text{tracks}}^{\text{ID}} < 6$ within a cone around the candidate muon with a size $\Delta R = 0.5$). The invariant mass of the muon pair should fulfill $|91.2 \text{ GeV} - M_{\mu\mu}| < 20$ GeV. With this cuts $\sim 70\%$ of the $Z \rightarrow \mu\mu$ events with muons in the acceptance are selected. The residual background fraction is 0.004 ± 0.001 and the dominant component comes from $t\bar{t}$ background.

3. – Acceptance calculation

Already in the first stage of data taking ($\sim 50 \text{ pb}^{-1}$), the uncertainties on the acceptance are relevant for the Z cross-section measurement. The acceptance is calculated through Monte Carlo simulation (MC@NLO) imposing kinematical cuts (p_T and η) on outgoing muons from Z -boson. The contributions to the systematic errors on the acceptance are: the errors on the PDF used, the Initial State Radiation (ISR), the intrinsic p_T of incoming partons, the QED corrections and the spin correlation between incoming partons and final leptons. The dominant contribution comes from the PDF and it has been estimated to be of the order of $\sim 4\%$. Figure 1 shows the acceptance as a function of \sqrt{s} for three different p_T cuts on the muons coming from the Z -boson decay.

REFERENCES

- [1] AAD G. *et al.* (THE ATLAS COLLABORATION), *Expected Performance of the ATLAS Experiment — Detector, Trigger and Physics*, arXiv:0901.0512, CERN-OPEN-2008-020.
- [2] AAD G. *et al.* (THE ATLAS COLLABORATION), *JINST*, **3** (2008) S08003.