

## Research Paper

**Cite this article:** Sándor D, Gyöngy M, Nyeste K, Czeglédi I, Székely C, Buchmann K, Cech G (2020). Digenean *Holostephanus* (Trematoda: Digenea: Cyathocotylidae) metacercariae in common carp (*Cyprinus carpio* Linnaeus, 1758) muscle: zoonotic potential and sensitivity to physico-chemical treatments. *Journal of Helminthology* 1–8. <https://doi.org/10.1017/S0022149X1900110X>

Received: 29 July 2019

Revised: 11 October 2019

Accepted: 5 December 2019

**Key words:**


Common carp; digenean trematodes; zoonosis; viability; muscle metacercariae

**Author for correspondence:**

D. Sandor,

E-mail: [sandor.diana@agrar.mta.hu](mailto:sandor.diana@agrar.mta.hu)

# Digenean *Holostephanus* (Trematoda: Digenea: Cyathocotylidae) metacercariae in common carp (*Cyprinus carpio* Linnaeus, 1758) muscle: zoonotic potential and sensitivity to physico-chemical treatments

D. Sándor<sup>1,2</sup> , M. Gyöngy<sup>3,4</sup>, K. Nyeste<sup>3,4</sup>, I. Czeglédi<sup>5</sup>, C. Székely<sup>1</sup>, K. Buchmann<sup>6</sup> and G. Cech<sup>1</sup>

<sup>1</sup>Institute for Veterinary Medical Research, Centre for Agricultural Research, Hungarian Academy of Sciences, Budapest, Hungary; <sup>2</sup>Eötvös Loránd University, Doctoral School of Biology Program of Zootaxonomy, Animal Ecology and Hydrobiology, Budapest, Hungary; <sup>3</sup>Department of Hydrobiology, University of Debrecen, Debrecen, Hungary; <sup>4</sup>Pál Juhász-Nagy Doctoral School of Biology and Environmental Sciences, University of Debrecen, Debrecen, Hungary; <sup>5</sup>Centre for Ecological Research, Balaton Limnological Institute, Hungarian Academy of Sciences, Tihany, Hungary and <sup>6</sup>Department of Veterinary and Animal Science, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark

**Abstract**

Metacercariae of various species within the genus *Holostephanus* Szidat, 1936 (Trematoda: Digenea: Cyathocotylidae) occur in muscles of both farmed and wild fish, including common carp (*Cyprinus carpio* Linnaeus, 1758). The life cycle includes a snail as first intermediate host, fish as second intermediate host and birds or mammals as final hosts. We studied the zoonotic potential and the viability of *Holostephanus* metacercariae from common carp following exposure to various physical and chemical treatments. Muscle tissue samples of common carp specimens from a fish farm in the north-eastern part of Hungary were examined and metacercariae recovered. The zoonotic potential was evaluated experimentally by using small mammals as models (albino mice,  $n = 2$ ; and Syrian hamsters,  $n = 4$ ) infected *per os* with *Holostephanus* cysts. Parallely, *Metagonimus* metacercariae were used as positive controls. We could not confirm the zoonotic potential of *Holostephanus* metacercariae as they did not survive in the mammalian intestine whereas *Metagonimus* metacercariae developed to the adult stage. We assessed the viability of metacercariae isolated from common carp specimens during exposure to different physical treatments (temperatures of  $-18^{\circ}\text{C}$ ,  $+20^{\circ}\text{C}$ ,  $+40^{\circ}\text{C}$  and  $+60^{\circ}\text{C}$ ) and chemical agents (5% and 10% acetic acid and 10% sodium chloride (NaCl)). Metacercariae lost viability by freezing at  $-18^{\circ}\text{C}$  (2 h), heating at  $60^{\circ}\text{C}$  (20 min), incubation in 5% and 10% acetic acid (5 min) and 10% NaCl (2 h). These methods served as models to investigate the effectiveness of food preparation techniques (such as cold and hot smoking, freezing, salting and pickling) on the survival of metacercariae.

**Introduction**

Worldwide, about 40 million people suffer from trematode infections, most commonly caused by members of the trematode families Echinostomidae, Fasciolidae, Heterophyidae and Opisthorchiidae (Abdussalam *et al.*, 1995; Keiser & Utzinger, 2009; WHO, 2011) associated with the consumption of raw or undercooked fish products (Chai *et al.*, 2005; Grundy-Warr *et al.*, 2012; Pinlaor *et al.*, 2013; Kim *et al.*, 2017). Infections caused by species within the genera *Metagonimus*, *Clonorchis*, *Haplorchis* and *Opisthorchis* usually manifest themselves in mild symptoms including abdominal pain, fever and weight loss (Traverso *et al.*, 2012; Pozio *et al.*, 2013; Pornruseetairatn *et al.*, 2016). However, chronic opisthorchid infections may cause injuries to the common bile duct, eventually leading to cholangitis, choledocholithiasis, pancreatitis and cholangiocarcinomatosis (Watanapa & Watanapa, 2002; Sithithaworn & Haswell-Elkins, 2003; Chai *et al.*, 2005; Rim, 2005; Toledo *et al.*, 2012). The highest prevalence of these trematode infections is in south-east and east Asia (EFSA, 2010), but sporadic cases occur in Europe, as well. In Italy, Armignacco *et al.* (2008) reported two outbreaks of *Opisthorchis felineus* (Rivolta, 1884) in humans from 2007 and recites two former cases from 2003 and 2005 published by Crotti *et al.* (2007). In 2010, a further 45 *O. felineus* cases were confirmed in Italy (Traverso *et al.*, 2012). Human infections have also been reported from Greece (Tselepatiotis *et al.*, 2003) and Germany (Bernhard, 1985; Sängner *et al.*, 1991). Opisthorchiasis was documented earlier in Lithuania (before 1901), Poland (before 1937), Romania (before 1957) and Spain (before 1932), but no cases have been reported since then (Erhardt *et al.*, 1962). Apart from

opisthorchiid and heterophyid metacercariae, also genera within the family Cyathocotylidae (*Holostephanus* spp., *Mesostephanus* spp., etc.) parasitize the musculature of fish, but zoonotic cases caused by them have not been reported yet. The genus *Holostephanus* Szidat, 1936 within the family Cyathocotylidae Poche, 1926 currently includes 12 species distributed in Europe and Asia: *H. luehei* (type species), *H. anHINGA*, *H. calvusi*, *H. corvi*, *H. curonensis*, *H. dubius*, *H. ibisi*, *H. ictaluri*, *H. lutzi*, *H. metorchis*, *H. nipponicus* and *H. phalacrocoraxus* (Seo et al., 2008). According to Gibson et al. (2002), the *Holostephanus* genus is present in Europe and Asia. Recent findings of *Holostephanus* flukes in Europe are documented from the Czechia (Moravec & Scholz, 2016), from Finland (Näreaho et al., 2017), from France (Gettová et al., 2016), from Poland (Kanarek et al., 2003; Sulgostowska, 2007; Mierzejewska et al., 2014), from Slovakia (Ondračková et al., 2009) and from Russia (Kvach et al., 2015). The primary intermediate hosts of *Holostephanus* trematodes are freshwater snails of species belonging to the families Bithyniidae (*Parafossarulus* spp., *Bithynia tentaculata*) and Viviparidae (*Lioplax* spp.) (Erasmus, 1962; Stang & Cable, 1966; Dubois, 1983; Gibson et al., 2002; Seo et al., 2008). Various freshwater fish species act as second intermediate hosts harbouring the *Holostephanus* metacercariae in the muscles. Erasmus (1962) isolated the cysts of *H. luehei* from three-spined stickleback (*Gasterosteus aculeatus* Linnaeus, 1758), while Seo et al. (2008) recovered *H. metorchis* metacercariae from topmouth gudgeon (*Pseudorasbora parva* Temminck & Schlegel, 1846). The natural definitive hosts are usually fish-eating aquatic birds and mammals (Yamaguti, 1939; Gibson et al., 2002). Besprozvannykh (2003) successfully reared the species *H. nipponicus* in chicks, and Seo et al. (2008) conducted similar experiments with *H. metorchis*, suggesting birds as suitable final hosts. Their fate in mammals is poorly described (Erasmus, 1962); however, Chandler (1950) reported a natural infection of a dog with *Mesostephanus longisaccus* (also a member of the family Cyathocotylidae). There are no records of human infection by *Holostephanus* species up to now, but due to the common occurrence of metacercariae in muscle tissue of European fish, it is important to evaluate infectivity to mammals and, thereby, their zoonotic potential.

We examined common carp (*Cyprinus carpio* Linnaeus, 1758) reared in a farm located in the north-eastern part of Hungary and isolated numerous metacercariae from muscle tissue of the fish. We performed a morphological characterization of the *Holostephanus* metacercariae isolated from the muscle tissue of common carp and investigated their infectivity to mammals using rodents (mice and hamsters) as models. We tested the survivability of metacercariae exposed to various physico-chemical conditions to model the effect of different food preparation techniques (freezing, frying, cold and hot smoking, marinating, pickling, salting) and to determine which processes may kill the parasites (Beldsoe & Oria, 2006; Borges et al., 2014; Onsurathum et al., 2016). The European Food Safety Authority (EFSA, 2004, 2010) and the Food and Agriculture Organization (1998) prepared a description about the risk assessment of fish-borne parasites in fishery products and, furthermore, made recommendations on how to prepare them safely.

## Materials and methods

### Sample collection

Metacercariae isolated from the muscle tissue of ten specimens of two-summer-old common carp obtained from a freshwater fish

farm in the north-eastern part of Hungary. The fish were collected by the fish farm personnel using cast-nets or seine nets and then transported to the Institute for Veterinary Medical Research laboratory alive in tanks with oxygenated water. The average weight of the examined fish was  $448.7 \pm 130.5$  (standard deviation (SD)) g and their average length was  $25.7 \pm 2.1$  (SD) cm.

### Artificial digestion

Fish were anaesthetized by adding clove oil to the fish tank water, whereafter fish were decapitated. The muscle tissue (fillets) of the fish (average weight 119.4 g, range 80,189 g) was removed, cut into smaller fragments and used for isolation of the metacercariae by a mechanical (compression between two glass plates) or an enzymatic incubation for 40 min in 0.5% pepsin solution (2 l tap water, 10 g of 1:10,000 NF powder-based pepsin and 16 ml of 25% hydrochloric acid) at a temperature of 40°C while stirring. The intact cysts were pipetted into Petri dishes containing 0.9% physiological saline, counted and examined for viability (presence of dynamic motions) and the dark excretory system under a stereomicroscope (Zeiss, Oberkochen, Germany).

### Morphological examination

Metacercariae were placed on a slide with a pipette in a drop of 0.9% physiological saline, cover-slipped and photographed with an Olympus BH2 photomicroscope (Shinjuku, Tokyo, Japan) at 4×, 10× and 20× magnification. Morphometric characters (body length and width, size of pharynx, oral and ventral suckers and length of caecal branches) were recorded based on 15 specimens and the morphological description of the metacercariae obtained (Erasmus, 1962).

### Experimental infection of mammals

Infection experiments of albino mice and Syrian hamsters were designed to determine the zoonotic potential of the *Holostephanus* trematode species. All metacercariae used for this experiment were isolated from the muscles of ten two-summer-old common carp specimens originating from the fish farm with the most severely infected fish stock. In addition, as positive control we used *Metagonimus* metacercariae collected from River Danube fishes. Trematodes belonging to this genus are well documented zoonotic parasites with an ability to develop in the mammalian intestine (Kang et al., 1983; Chai et al., 1998; Shimazu, 2002; Kudo et al., 2014).

In the first experiment (3 October–12 October 2017), two laboratory albino mice (mice ‘A’ and ‘B’) were orally intubated and infected with 50 metacercariae each pipetted from the saline solution, and then the top of their head was marked with a blue felt-tip pen. The negative control animals (two additional mice) were not marked. Mouse ‘A’ was euthanized on 10 October, while mouse ‘B’ on 11 October, in compliance with the European regulations on animal protection.

In the second experiment (8 July–20 July 2018), four Syrian hamsters (animals ‘C’, ‘D’, ‘E’ and ‘F’) were inoculated, the first two with 50 and the latter two with 100 trematode larvae. Hamster ‘C’, a grey-coated male, and hamster ‘D’, a black female, were not marked as their coat colour clearly differentiated them from the other hamsters. Hamsters ‘E’ and ‘F’ were marked by clipping the hair in a small area on the crown of the head. Hamsters of the control group were also left unmarked. In order to demonstrate the suitability of the hamster infection

model we included two hamsters as positive controls. They were inoculated with 50 *Metagonimus* sp. metacercariae each. The hair of these hamsters was clipped on the crown of the head, in the mediadorsal region and at the tailhead. Hamster 'C' was euthanized with CO<sub>2</sub> gas on 13 July, hamsters 'D', 'E' and the negative control animals on 16 July, while hamster 'F' and the two *Metagonimus*-infected hamsters on 20 July.

Following euthanasia, the intestines of the animals were isolated and each divided into three segments (anterior and middle segment of the small intestine and the large intestine) and placed into Petri dishes containing physiological saline. Then all intestinal segments were cut open longitudinally and mucosal lining with intestinal content scraped into the dish, whereafter the material was sedimented in 500 ml decanter glasses. After 30 min, the intestinal contents and the scrapings collected from the intestinal wall were examined under stereomicroscope. The laboratory mice and the golden hamsters were purchased from Ökomester Ltd. (Budapest, Hungary) after the permissions necessary for the experiment had been obtained (permission number PEI/001/1792-4/2014).

### Sensitivity of metacercariae to physico-chemical treatments

Metacercariae isolated by artificial digestion were placed between two slices of fish muscle (1 × 1 cm wide and 3 mm thick). These double-layer fillets each contained ten metacercariae and were placed into six-well cell-culture plates (one sample per well) and then subjected to different heat and chemical treatments. The treatments were started at 10:00 a.m. on the experimental days and different examination time points were designated (minutes: 1, 5, 10, 15, 20, 30; hours: 1, 2, 3, 4, 6, 12; days: 1, 2, 4, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26) (Borges *et al.*, 2014). At each examination time point, the double fillets were opened with a pair of forceps, whereafter the wells were examined with a Zeiss stereomicroscope. The numbers of surviving and dead metacercariae were recorded. The intact structure and movement of the metacercariae within the capsule were used as viability criteria. Dead metacercariae were characterized by an amorphous body, granular, vesicular morphology or complete lack of motion. All treatment assays were repeated four times.

Thus, four double fillets, each with ten metacercariae, were used for each treatment type, and thus altogether 40 metacercariae per treatment were included in the statistical analysis. The control group also included 40 metacercariae evenly distributed in four double fillets, which were kept in 0.9% physiological saline at 5°C.

We investigated how physical (freezing, heat) and chemical (acetic acid and saline solutions) treatments (procedures commonly applied in the industry) affected the survival of metacercariae. The freezing experiment was performed at -18°C because most household freezers work at this temperature. The double fillets with metacercariae were kept in plastic plates without the addition of physiological saline, and then they were thawed at 27°C for 20 min to record viability of metacercariae. To imitate cold- and hot-smoking procedures, the fillet sandwiches were placed in wells (without saline) and then they were subjected to heat treatment at 20, 40 and 60°C temperature, in an INCO CO<sub>2</sub> Incubator (Memmert, Büchenbach, Germany).

To model the effect of acetic acid (e.g. pickled fish), the double fillets were treated with 5% and 10% vinegar at room temperature. Four sandwiches were also treated with 10% sodium chloride (NaCl) solution to determine the outcome of conservation (e.g. canned fish).

### Procedures used for the statistical analysis

Surviving and dead metacercariae were recorded at the predetermined examination time points, and after reaching 100% mortality these data were summed up. The effectiveness of the different procedures within a treatment type (physical experiments on -18°C, +20°C, +40°C and +60°C and chemical experiments with 5% and 10% acetic acid, 10% NaCl) was compared with the help of survival curves. We compared each curve within a treatment type following pairwise multiple comparisons with Bonferroni correction in the case of significant difference among the curves. The 'survdiff' function ('survival' package) from R version 3.6.1. (R Core Team, 2015) was used to calculate chi-square distance and the corresponding *P*-value (Harrington & Fleming, 1982). Only treatment types with a *P*-value threshold of <0.05 were considered as having a significant effect on the survival of metacercariae. Each metacercaria was a replicate. Since four fillets were used for each treatment type, a preliminary survival analysis was carried out in order to detect the effect of fillets on the survival or death of metacercariae (fillet effect) in each heat or chemical treatment group.


## Results

### Identification

After manual isolation and artificial digestion, the presence of numerous trematode metacercariae was recorded in the musculature of ten two-summer-old common carp specimens from the north-eastern fish farm (fig. 1). Cysts were detected in all common carp specimens; thus, the prevalence of infection was 100% and the mean intensity of infection was 144.6 ± 55.1 (SD) metacercariae per individual.


The species identification of encysted metacercariae was done by morphometric examinations. Light microscopic examination revealed that the cysts of the trematode species detected in the muscles of infected common carp specimens belonged to the prohemistomulum type of metacercariae. This type is characterized by a round or oval body protected by a thick outer wall and a thin inner membrane (fig. 2). They have real ventral suckers but the pseudosucker is missing. The trematode species belonging to this type are further characterized by the presence of a dark excretory system densely enmeshing the body (Gibson *et al.*, 2002; fig. 3). The dimensions of easily detectable morphological traits (body length and width, pharynx, oral and ventral sucker, caecal branches) were determined with the help of a scale built in the ocular lens of the microscope. Based on the study of Erasmus (1962), we recorded and compared the parameters of a total of 15 cysts. The body surface of encysted metacercariae was smooth and spikes were not found on it. Body length was 324.7 ± 35.6 (SD) µm and body width 245.3 ± 52.2 (SD) µm. The oral sucker was 32.3 ± 7.3 (SD) µm long and 32.3 ± 7.3 (SD) µm wide. After a short prepharynx, the oral sucker was closely followed by the small-sized pharynx, which was 39.3 ± 4.6 (SD) µm long and 33.4 ± 4.8 (SD) µm wide. The ventral sucker was 75 ± 9.3 (SD) µm long and 75 ± 9.3 (SD) µm wide. The length of the caecal branches was 274.7 ± 33.1 (SD) µm. The recorded characteristics were highly similar to those described by Erasmus (1962) as the main feature of *Holostephanus* sp. metacercariae. Additional sequence analysis of the internal transcribed spacer (ITS) region was executed, which confirmed the identity of metacercariae – namely, all of them were members of the Cyathocotylidae family, most probably species of *Holostephanus*

Fig. 1 - Colour online



**Fig. 1.** Intact cyathocotyloid metacercariae in the muscle. Scale bar: 400 µm.

Fig. 2 - Colour online



**Fig. 2.** Cyathocotyloid metacercaria from carp muscle (*Cyprinus carpio*) following manual isolation. The figure shows the outer, thick wall typical of prohemistomulum metacercariae. Scale bar: 50 µm.

genus; however, identification to species level was problematic as very few sequence data were available in the database of Genbank (National Center for Biotechnology Information) during the analysis ('Extensive monitoring of Digenean trematodes in European fresh water aquacultures', unpublished manuscript).

### Infection experiments with mammals

The two albino mice and the four Syrian hamsters were euthanized and their intestines examined 7–12 days after oral inoculation. Trematodes were neither detected in any of the inoculated animals nor the non-infected control animals, and the mucous membranes of the intestines were intact and healthy. At the same time, a zoonotic potential was demonstrated for the *Metagonimus* species used as positive control: 22 sexually mature trematodes were isolated from the midgut of one inoculated hamster and five from the large intestine of the other.

### Survival (viability) experiments

The survival analysis did not detect significant differences among the survival of metacercariae in fillets within each heat and



**Fig. 3.** Cyathocotyloid metacercaria after artificial digestion. Scale bar: 50 µm.

chemical treatment group (DF = 3,  $P > 0.05$ ); thus, 'fillet effect' was omitted from the further analysis of data. The survival analysis has showed that there were significant differences among the different heat ( $\chi^2 = 389$ , DF = 4,  $P < 0.01$ ) and chemical treatments ( $\chi^2 = 238$ , DF = 3,  $P < 0.01$ ). According to the pairwise multiple comparisons with Bonferroni correction, the differences between any of the treated groups and the untreated groups were significant (tables 1 and 2). The number of surviving metacercariae in the case of each treatment are presented in table 2 and figs 4 and 5.

Heat treatments at +60°C can be considered the most effective for killing the metacercariae, as they resulted in 100% mortality after just 30 min. The second most effective heat treatment type was that of -18°C, where all metacercariae died after 2 h of exposure (tables 1 and 2, fig. 4).

However, during the chemical treatments, exposure to 5% and 10% acetic acid killed the metacercariae within a few minutes; the 10% acetic acid was significantly more effective than other chemical treatments. Furthermore, the effect of the treatment with 10% NaCl was significantly lower than that of the acetic acid treatments, as during exposure to 10% NaCl solution the mortality of all metacercariae occurred 2 h later (tables 1 and 2, fig. 5).

### Discussion

Common carp reared in fish farms in the north-eastern part of Hungary were recently reported infected with metacercariae in the muscle tissue, and due to the well-known zoonotic potential of muscle-dwelling metacercariae (Healy, 1970; Abdussalam *et al.*, 1995; Fried *et al.*, 2004; Phan *et al.*, 2010), we found it relevant to elucidate the identity and zoonotic potential of the reported parasites from Hungarian carp. Metacercariae were isolated and characterized morphologically and, based on the taxonomical characteristics provided by Erasmus (1962), it was established that they belong to the genus *Holostephanus* within the digenean trematode family Cyathocotyliidae. Sequence data of the ITS region also confirmed the identification (under publication). We then investigated if the metacercariae were infective to rodents (mice and hamsters), but none of the *Holostephanus* metacercariae established or matured in these mammals. Another well-known fish-borne zoonotic trematode species

190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251

**Table 1.** Pairwise comparison of different heat and chemical treatments:  $\chi^2$  and *P*-values (pairwise multiple comparisons with Bonferroni correction, degrees of freedom = 1).

Compared treatments	$\chi^2$	<i>P</i> -value
Heat treatments		
C-(−18°C)	78.7	<0.001
C-20°C	75.9	<0.001
C-40°C	83.1	<0.001
C-60°C	77.3	<0.001
(−18°C)-20°C	78.7	<0.001
(−18°C)-40°C	33.3	<0.001
(−18°C)-60°C	77.3	<0.001
20-40°C	60.1	<0.001
20-60°C	77.3	<0.001
40-60°C	77.3	<0.001
Chemical treatments		
C-5% AcOH	75.6	<0.001
C-10% AcOH	79.0	<0.001
C-10% NaCl	81.0	<0.001
5% AcOH-10% AcOH	50.0	<0.001
5% AcOH-10% NaCl	75.6	<0.001
10% AcOH-10% NaCl	79.0	<0.001

C, control (0.9%, 5°C); AcOH, acetic acid; NaCl, sodium chloride.

(*Metagonimus* sp.) established in the control experiments, suggesting that the model was suitable to elucidate zoonotic potentials. The present study could, therefore, not confirm that the metacercariae from common carp are zoonotic. We also investigated methods to inactivate the *Holostephanus* metacercariae. Numerous studies on parasite elimination methods for parasites in fish and shellfish have been published (Fan, 1998; Abdallah *et al.*, 2009; Borges *et al.*, 2014; Kim *et al.*, 2017). At the same time, it is important to mention that the resistance of different trematode species may vary. Rác & Zemankovics (2002) subjected *Metagonimus yokogawai* metacercariae collected from River Danube fishes to freezing at −26°C, desiccation and acetic acid treatment. In their experiments, the first cysts died after 24 h and the last ones after 27 h after freezing. In contrast, during storage at room temperature without water the survival time of *M. yokogawai* cysts was a maximum of 2–3 h. With regard to resistance to acetic acid, it was found that 100% mortality of the trematode larvae occurred within a few minutes in 10% acetic acid and after 30 min in 5% acetic acid. Fattakhov (1989) demonstrated that at −28°C, −35°C and −40°C *Opisthorchis* metacercariae survived for 20, 8 and 2 h, respectively. At variance with this finding, Borges *et al.* (2014) found that the metacercariae of *Cryptocotyle lingua*, a trematode of northern distribution, died after just 2 h at −20°C, and after 1 h at −40°C. In 2007, infection with the trematode *Opisthorchis* sp. of two humans was reported in Italy, and in a further 18 cases the coprological examination was positive for trematode eggs. The consumed fish (*Tinca tinca* and *Coregonus* sp.) had been stored at −10°C for three days, then pickled in the mixture of vinegar and wine for 24 h. Based on the findings that the zoonotic *Opisthorchis* metacercariae survived this

**Table 2.** Number of live metacercariae found at the different examination time points during the different heat and chemical treatments.

Heat and chemical treatments	Minutes												Hours												Days											
	1	5	10	15	20	23	27	30	31	1	2	3	4	6	12	24	2	4	6	8	10	12	14	16	18	20	22	24								
60°C	40	40	40	37	40	40	40	40	40	22	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0								
40°C	40	40	40	40	40	40	40	31	28	12	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0								
20°C	40	40	40	40	40	40	40	40	40	37	32	25	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0								
−18°C	40	40	40	40	40	40	40	22	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0								
10% AcOH	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0								
5% AcOH	31	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0								
10% NaCl	40	40	39	38	33	27	16	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0								
Control (0.9%, 5°C)	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	40	38	35	33	26	20	15	6	0									

Numbers with grey shading indicate the total mortality of metacercariae. AcOH, acetic acid; NaCl, sodium chloride.

252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313




Fig. 4. Number of survived metacercariae in different examination time points exposed to different heat treatments and those of the control group.




Fig. 5. Number of survived metacercariae in different examination time points exposed to different chemical treatments and those of the control group.

treatment, the Italian Ministry of Health published official guidelines in 2008 recommending that cyprinids intended for human consumption should undergo thorough frying or freezing at  $-20^{\circ}\text{C}$  for seven days before consumption (Armignacco *et al.*, 2008; Traverso *et al.*, 2012). While *Opisthorchis viverrini* cysts were reported to die within 24 h in 13.6% NaCl solution (Kruatrachue *et al.*, 1982), the metacercariae of *Haplorchis taichui* were able to survive for 14 days at room temperature and for 21 days in the refrigerator in *pla som* fish food (sour fish) stored in 15% salt solution (Kaenjampa *et al.*, 2017).

The European households most often use the heat treatment procedure. In the present experiment studying the survival of

*Holostephanus* metacercariae, heat treatment at  $60^{\circ}\text{C}$  was effective but only at a 30 min exposure time. In contrast, the freezing of fish at  $-18^{\circ}\text{C}$  required 120 min to be effective, which corresponded to the effect of 10% NaCl solution. Of all the treatment methods used, exposure to acetic acid was the most effective, as it caused total mortality of the parasite larvae after 5 min when using the 5% solution and after just 1 min when the 10% acetic acid solution was used. It seems that all of the treatments were effective in six hours; however, it should be noted that the pre-treatment (digestion by pepsin solution) might cause metacercariae to be more sensitive to the physical and chemical effects. However, the digestion was necessary as 'the cysts of

*Holostephalus* are much harder, making it almost impossible to isolate the metacercariae without chemical action' (Kvach *et al.*, 2016, p. 259). Moreover, several recent studies applied this methodology in their viability experiments (Rácz & Zemankovics, 2002; Borges *et al.*, 2014).

We conducted infection experiments with small mammals to study the zoonotic potential of metacercariae isolated from the muscles of infected common carp specimens, as we could not find data on the zoonotic potential of *Holostephanus* species in the current international literature. The inoculated mice and hamsters were euthanized and their intestines examined after 7–12 days; however, mature parasite stages were not found, which suggests that this trematode parasitizing the muscles of fish might not have zoonotic potential. In the positive control hamsters, the *Metagonimus* trematode developed to the mature stage, which supports that small mammals can be suitable model organisms for the study of the zoonotic potential of trematodes. Despite the negative experimental results, the definitive host of *Holostephanus* trematodes is suggested to be a higher vertebrate species, as it follows from the general developmental cycle of this group of digeneans (Erasmus, 1962; Dubois, 1983). Mammalian species are presumably not susceptible to *Holostephanus* species. To date, no zoonosis attributable to *Holostephanus* species has been reported in the literature (Seo *et al.*, 2008). The negative result of the small mammal infection experiments in the present study is consistent with this; however, the evolutionary distance between humans and rodent species should naturally be taken into consideration when making such a statement. Moreover, the complete lack of history of human infection in the scientific literature also supports the non-zoonotic nature of these trematodes. There are only a few studies that acknowledge *Holostephanus* species to parasitize mammals. Gibson *et al.* (2002) mention avian and mammalian definitive hosts in their comprehensive work, which is in conformity with the findings of Chandler (1950), who reported the natural infection of a dog with *M. longisaccus*, a species also belonging to the Cyathocotylidae family, thus providing an example of when a mammal is successfully infected with a trematode of this family. However, the culinary practice and preservation methods commonly used in Hungary and in most European countries can prevent the survival and possible transfer of metacercariae present in fish fillets.

**Financial support.** This work was supported by the European Union's Horizon 2020 Research and Innovation Programme (ParaFishControl; grant number 634429); the European Regional and Development Fund and the Government of Hungary within the project GINOP-2.3.2-15-2016-00025; the ÚNKP-18-3 New National Excellence Program of the Ministry of Human Capacities (grant number ÚNKP-18-3-I-ELTE-410); and by the Higher Education Institutional Excellence Programme (NKFIH-1150-6/2019) of the Ministry of Innovation and Technology in Hungary, within the framework of the fourth thematic programme of the University of Debrecen.

**Conflicts of interest.** None.

**Ethical standards.** The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional guides on the care and use of laboratory animals.

## References

Abdallah KF, *et al.* (2009) Effect of different temperatures on viability of seven encysted metacercariae recovered from freshwater fishes in Qalyobia, Egypt. *Journal of the Egyptian Society of Parasitology* 39(2), 413–420.

Abdussalam M, *et al.* (1995) Food safety measures for the control of food-borne trematode infections. *Food Control* 6, 71–79.

Armignacco O, Caterini L, Marucci G, *et al.* (2008) Human illnesses caused by *Opisthorchis felineus* flukes, Italy. *Emerging Infectious Diseases* 14, 1902–1905.

Beldsoe EG and Oria M (2006) Potential hazards in cold-smoked fish: parasites. *Journal of Food Science* 66(7), 1100–1103.

Bernhard K (1985) Nachweis besonderer Helminthenarten bei DDR-Bürgern. *Angewandte Parasitologie* 26, 223–224.

Besprozvannykh VV (2003) Life cycle of a trematode *Holostephanus nipponicus* (Cyathocotylidae) in conditions of Primor'e. *Parazitologiya* 37, 346–351.

Borges JN, *et al.* (2014) Viability of *Cryptocotyle lingua* metacercariae from Atlantic cod (*Gadus morhua*) after exposure to freezing and heating in the temperature range from -80 °C to 100 °C. *Food Control* 50, 371–377.

Chai JY, Murrell KD and Lymbery AJ (2005) Fish-borne parasitic zoonoses: status and issues. *International Journal for Parasitology* 35, 1233–1254.

Chai JY, *et al.* (1998) Surface ultrastructure of *Metagonimus miyagai* metacercariae and adults. *The Korean Journal of Parasitology* 36, 217–225.

Chandler AC (1950) *Mesostephanus longisaccus*, a new cyathocotylid trematode from a dog. *The International Journal for Parasitology* 36, 90.

Crotti D, D'Annibale ML and Crotti S (2007) Opisthorchiasi autoctona del Lago Trasimeno (Perugia): descrizione di due episodi epidemici da *Opisthorchis felineus* e problematiche diagnostiche differenziali. *Microbiologia Medica* 42, 36–41.

Dubois G (1983) Le genre *Holostephanus* Szidat, 1936 (Trematoda: Strigeata: Cyathocotylidae). Commentaire sur les espèces et érection de *Holostephanoides* gen. nov. Bull. Soc. neuchâtel. *The Science of Nature* 106, 87–99.

EFSA (European Food Safety Authority) (2004) Regulation (EC) N° 853/2004 of the European parliament and council of 29 of April 2004 laying down specific hygiene rules for food of animal origin. Available at [https://www.fsai.ie/uploadedFiles/Food\\_Business/Reg853\\_2004.pdf](https://www.fsai.ie/uploadedFiles/Food_Business/Reg853_2004.pdf) (accessed 19 March 2014).

EFSA (European Food Safety Authority) (2010) Panel on Biological Hazards (BIOHAZ) scientific opinion on risk assessment of parasites in fishery products. *EFSA Journal* 8, 1543.

Erasmus DA (1962) Studies on the adult and metacercaria of *Holostephanus lühei* Szidat, 1936. *Parasitology* 52, 353–374.

Erhardt A, Germer WD and Hörning B (1962) *Die Opisthorchiasis, hervorgerufen durch den Katzenleberegel Opisthorchis felineus* (Riv.). *Parasitologische Schriftenreihe*, vol. 15. Jena, Veb Gustav Fischer Verlag.

Fan PC (1998) Viability of metacercariae of *Clonorchis sinensis* in frozen or salted freshwater fish. *International Journal for Parasitology* 28(4), 603–605.

FAO (Food and Agriculture Organization) (1998) Seafood safety e Economics of hazard analysis and critical control point (HACCP) programs. FAO Fisheries Technical Papers, T381. Available at <http://www.fao.org/DOCREP/003/X0465E/X0465E00.HTM> (accessed 18 March 2014).

Fattakhov RG (1989) Low-temperature regimes for decontamination of fish of the larvae *Opisthorchis*. *Medicine Parasitology (Mosk)* 5, 63–65 (in Russian).

Fried B, Graczyk TK and Tamang L (2004) Food-borne intestinal trematodiasis in humans. *Parasitology Research* 93, 159–170.

Gettová L, Gilles A and Šimková A (2016) Metazoan parasite communities: support for the biological invasion of *Barbus barbus* and its hybridization with the endemic *Barbus meridionalis*. *Parasites and Vectors* 9, 588.

Gibson DI, Jones A and Bray RA (2002) *Key to the Trematoda. Volume 1*. Wallingford, CABI Publishing, 206.

Grundy-Warr C, *et al.* (2012) Raw attitudes, wetland cultures, life-cycles: socio-cultural dynamics relating to *Opisthorchis viverrini* in the Mekong Basin. *Parasitology International* 61, 65–70.

Harrington DP and Fleming TR (1982) A class of rank test procedures for censored survival data. *Biometrika* 69, 553–566.

Healy GR (1970) Trematodes transmitted to man by fish, frogs, and crustacea. *Journal of Wildlife Diseases* 6, 255–261.

Kaenjampa P, *et al.* (2017) Elimination of *Haplochis taichui* metacercaria in cyprinoid fish with freezing temperature and soured fish (plasom) with salinity. *The Southeast Asian Journal of Tropical Medicine and Public Health* 48(4), 777–785.

Kanarek G, *et al.* (2003) Digenean fauna of the great cormorant *Phalacrocorax carbo sinensis* (Blumenbach, 1798) in the brackish waters of the Vistula

- Lagoon and the Gulf of Gdańsk (Poland). *Wiadomości Parazytologiczne* **49**, 293–299.
- Kang SY, et al. (1983)** A study on intestinal lesions of experimentally reinfected dogs with *Metagonimus yokogawai*. *The Korean Journal of Parasitology* **21**(1), 58–73.
- Keiser J and Utzinger J (2009)** Food-borne trematodiasis. *Clinical Microbiology Reviews* **22**(3), 466–483.
- Kim TI, et al. (2017)** Inactivation of *Paragonimus westermani* metacercariae in soy sauce-marinated and frozen freshwater crabs. *Parasitology Research* **116**(3), 1003–1006.
- Kruatrachue M, et al. (1982)** Effects of physico-chemical factors on the infection of hamsters by metacercariae of *Opisthorchis viverrini*. *The Southeast Asian Journal of Tropical Medicine and Public Health* **13**, 614–617.
- Kudo N, et al. (2014)** Experimental final hosts of *Metagonimus hakubaensis* (Trematoda: Heterophyidae) and their suitability to the fluke. *Journal of Veterinary Medical Science* **76**(12), 1651–1654.
- Kvach Y, Ondračková M and Jurajda P (2016)** First report of metacercariae of *Cyathocotyle prussica* parasitising a fish host in the Czech Republic, Central Europe. *Helminthologia* **53**(3), 257–261.
- Kvach Y, et al. (2015)** The parasite community of gobiid fishes (Actinopterygii: Gobiidae) from the Lower Volga River region. *Biologia* **70**, 948–957.
- Mierzejewska K, et al. (2014)** Parasites of non-native gobies in the Włocławek Reservoir on the lower Vistula River, first comprehensive study in Poland. *Knowledge and Management of Aquatic Ecosystems* **414**, 1.
- Moravec F and Scholz T (2016)** Helminth parasites of the lesser great cormorant *Phalacrocorax carbo sinensis* from two nesting regions in the Czech Republic. *Folia Parasitologica* **63**, 22.
- Näreaho A, Eriksson-Kallio AM, Heikkinen P, Snellman A, Sukura A and Koski P (2017)** High prevalence of zoonotic trematodes in roach (*Rutilus rutilus*) in the Gulf of Finland. *Acta Veterinaria Scandinavica* **59**, 75.
- Ondračková M, et al. (2009)** The interaction between an introduced fish host and local parasite fauna: *Neogobius kessleri* in the middle Danube River. *Parasitology Research* **105**, 201–208.
- Onsurathum S, Pinlaor P, Charoensuk L, et al. (2016)** Contamination of *Opisthorchis viverrini* and *Haplorchis taichui* metacercariae in fermented fish products in northeastern Thailand markets. *Food Control* **59**, 493–498.
- Phan VT, et al. (2010)** Fish-borne zoonotic trematodes in cultured and wild-caught freshwater fish from the Red River Delta, Vietnam. *Vector-Borne and Zoonotic Diseases* **10**(9), 861–866.
- Pinlaor S, Onsurathum S, Boonmars T, et al. (2013)** Distribution and abundance of *Opisthorchis viverrini* metacercariae in cyprinid fish in north-eastern Thailand. *Korean Journal of Parasitology* **51**(6), 703–710.
- Pornruseaitratn S, Kino H, Shimazu T, et al. (2016)** A molecular phylogeny of Asian species of the genus *Metagonimus* (Digenea)—small intestinal flukes—based on representative Japanese populations. *Parasitology Research* **115**(3), 1123–1130.
- Pozio E, et al. (2013)** *Opisthorchis felineus*, an emerging infection in Italy and its implication for the European Union. *Acta Tropica* **126**(1), 54–62.
- Rác Z and Zemankovics E (2002)** Survival of metacercariae of *Metagonimus yokogawai* (Digenea: Heterophyidae) on fish from River Danube. *Hungarian Veterinary Journal* **124**(7), 437–444.
- R Core Team (2015)** *R: a language and environment for statistical computing*. Vienna, Austria, R Foundation for Statistical Computing. Available at <http://www.R-project.org> (accessed 25 November 2019).
- Rim HJ (2005)** Clonorchiasis: an update. *Journal of Helminthology* **79**, 269–281.
- Sänger R, et al. (1991)** An endemic focus of *Opisthorchis felineus* in Mecklenburg-Pomerania/Germany. *Tropical Medicine and Parasitology* **42**, 463.
- Seo M, et al. (2008)** *Holostephanus metorchis* (Digenea: Cyathocotylidae) from chicks experimentally infected with metacercariae from a fish, *Pseudorasbora parva*, in the Republic of Korea. *Korean Journal of Parasitology* **46**(2), 83–86.
- Shimazu T (2002)** Life cycle and morphology of *Metagonimus miyatai* (Digenea: Heterophyidae) from Nagano, Japan. *Parasitology International* **51**(3), 271–280.
- Sithithaworn P and Haswell-Elkins M (2003)** Epidemiology of *Opisthorchis viverrini*. *Acta Tropica* **88**, 187–194.
- Stang JC and Cable RM (1966)** The life history of *Holostephanus ictaluri* Vernberg, 1952 (Trematoda: Digenea), and immature stages of other North American fresh-water cyathocotylids. *The American Midland Naturalist* **75**, 404–415.
- Sulgostowska T (2007)** Intestinal digeneans of birds (superfamily Diplostomoidea) of the Masurian lakes. *Wiadomości Parazytologiczne* **53**, 117–28 (in Polish).
- Toledo R, Esteban JG and Fried B (2012)** Current status of food-borne trematode infections. *European Journal of Clinical Microbiology and Infectious Diseases* **31**(8), 1705–1718.
- Traverso A, Repetto E, Magnani S, et al. (2012)** A large outbreak of *Opisthorchis felineus* in Italy suggests that opisthorchiasis develops as a febrile eosinophilic syndrome with cholestasis rather than a hepatitis-like syndrome. *European Journal of Clinical Microbiology and Infectious Diseases* **31**, 1089–1093.
- Tselepatiotis E, et al. (2003)** A case of *Opisthorchis felineus* infestation in a pilot from Greece. *Infection* **31**, 430–432.
- Watanapa P and Watanapa WB (2002)** Liver fluke-associated cholangiocarcinoma. *British Journal of Surgery* **89**, 962–970.
- WHO (2011)** *Report of the WHO expert consultation on foodborne trematode infections and taeniasis/cysticercosis*. World Health Organisation. Available at [https://www.who.int/neglected\\_diseases/preventive\\_chemotherapy/WHO\\_HTM\\_NTD\\_PCT\\_2011.3.pdf](https://www.who.int/neglected_diseases/preventive_chemotherapy/WHO_HTM_NTD_PCT_2011.3.pdf).
- Yamaguti S (1939)** Studies on the helminth fauna of Japan. Part 25. Trematodes of birds, IV. *Japanese Journal of Zoology* **8**, 131–210.