

Receiver function mapping of mantle transition zone discontinuities beneath Alaska using scaled 3D velocity corrections

A.M. van Stiphout¹, S. Cottaar², A. Deuss¹

¹*Department of Earth Sciences, Utrecht University, The Netherlands*

²*Bullard Labs, University of Cambridge, United Kingdom*

Accepted 2019 July 30. Received 2019 July 22; in original form 2019 March 20

SUMMARY

The mantle transition zone is the region between the globally observed major seismic velocity discontinuities around depths of 410 and 660 km and is important for determining the style of convection and mixing between the upper and the lower mantle. In this study, P-to-S converted waves, or receiver functions, are used to study these discontinuities beneath the Alaskan subduction zone, where the Pacific plate subducts underneath the North American plate. Previous tomographic models do not agree on the depth extent of the subducting slab, therefore improved imaging of the Earth structure underneath Alaska is required. We use 27,800 high quality radial receiver functions to make common conversion point stacks. Upper mantle velocity anomalies are accounted for by two recently published regional tomographic S-wave velocity models. Using these two tomographic models, we show that the discontinuity depths within our CCP stacks are highly dependent on the choice of velocity model, between which velocity anomaly magnitudes vary greatly. We design a quantitative test to show whether the anomalies in the velocity models are too strong or too weak, leading to over- or under-corrected discontinuity depths. We also show how this test can be used to rescale the 3D velocity corrections in order to improve the discontinuity topography maps.

After applying the appropriate corrections, we find a localised thicker mantle transition zone

and an uplifted 410 discontinuity, which show that the slab has clearly penetrated into the mantle transition zone. Little topography is seen on the 660 discontinuity, indicating that the slab has probably not reached the lower mantle. In the southwest, P410s arrivals have very small amplitudes or no significant arrival at all. This could be caused by water or basalt in the subducting slab, reducing the strength at the 410, or by topography on the 410 discontinuity, preventing coherent stacking. In the southeast of Alaska, a thinner mantle transition zone is observed. This area corresponds to the location of a slab window, and thinning of the mantle transition zone may be caused by hot mantle upwellings.

Key words: Composition and structure of the mantle – Subduction zone processes – Body waves – North America – Seismology

1 INTRODUCTION

The mantle transition zone delineates the upper from the lower mantle, and its characteristic phase transitions both affect and reflect the style of mantle convection. Subducting plates leave an imprint on the boundaries of the mantle transition zone, the seismic discontinuities at approximately 410 and 660 km depth. By studying these we can constrain the depth extent of subducting slabs. In Alaska, tomographic models do not agree on the depth extent of the subducting Alaskan slab. Here, we use receiver functions, which are waves that convert at discontinuities, to image the mantle transition zone to better constrain the Earth structure underneath Alaska.

1.1 The Alaskan subduction zone

Alaska, a U.S. state, is located on the northwestern edge of the North American plate and along the northern part of the Pacific subduction zone. This subduction zone accommodates convergence between the northernmost part of the Pacific plate and the Alaskan part of the North American plate at the Alaskan-Aleutian megathrust (see Fig.1). The Pacific plate moves 5.2 cm/year to the NNW with respect to the North American plate (DeMets and Dixon, 1999). In the east, the Pacific plate terminates against the Fairweather-Queen Charlotte transform system. At the eastern part of the subduction zone lies the Yakutat terrane, a region of unusually thick oceanic crust (Gulick et al.,




Figure 1. Topography map showing the study area with the plate boundaries in black (Bird, 2003). Red dots denote active volcanoes. The orange lines show the subducted Pacific plate at depth with contour intervals of 20 km as imaged in the Slab2 model by Hayes et al. (2018) and the subducted Yakutat crust as imaged by Eberhart-Phillips et al. (2006) is shown by the thin dashed line.

2007), moving with the Pacific oceanic lithosphere and being accreted to the Alaskan margin (Fuis et al., 2008). Collision and underthrusting of the Yakutat terrane is believed to have caused some unusual features, such as a region with absence of seismicity known as the Denali volcanic gap and the Wrangell volcanism (Eberhart-Phillips et al., 2006). This Wrangell volcanic field lies in the north of the Yakutat block, and is a region with a decrease in seismicity below 50 km depth. The existence of a Wrangell slab has been debated in several studies (e.g Martin-Short et al., 2018; Jadamec and Billen, 2012; Jiang et al., 2018; Gou et al., 2019; Bauer et al., 2014), but is thought to only be present in the upper mantle and is not expected to have a signature on the 410 km discontinuity.

Seismic tomographic results are also used to debate the extent of the Pacific slab into and

across the mantle transition zone (MTZ, 400-700 km depth) further west. Qi et al. (2007) found fast velocities associated with the slab reaching depths of around 400-450 km, but also disconnected fast velocities lower in the MTZ, which they propose to be remnants of an older plate, the Kula plate. The models of Martin-Short et al. (2018) indicate that the Pacific slab continues to depths below 400 km, but do not have good resolution below that depth. In the upper mantle Vs model constructed by Jiang et al. (2018) the subducting plate is most prominent at depths until 390 km depth, but weaker fast anomalies appear in the MTZ. In the P-wave velocity model for North America by Burdick et al. (2017) the end of the slab seems to penetrate into the lower mantle, but resolution is lost at greater depth. The P-wave velocity model by Gou et al. (2019) shows the slab down to 450-500 km. Comparing both S and P velocities, "Atlas of the Underworld" (van der Meer et al., 2017) identify the end of the slab within the MTZ.

Plate tectonic reconstructions have long shown that convergence has occurred between the North American plate and the Pacific and previously Kula and Farallon plates for > 100 Myr (Engebretson et al., 1985; Müller et al., 2016; Seton et al., 2012; Madsen et al., 2006). However, the depth extent of the slabs observed in tomographic models is much less than the several thousand km of lithosphere that is expected to be subducted beneath the Aleutian trench based on these plate reconstruction models. This may suggest that the subduction configuration in the past may have been slightly more complex than generally perceived.

The disagreement across different tomographic models on the depth extent of the subducting slab could be due to reducing resolution of the models across the MTZ and the potential smearing along body wave ray paths. Therefore we need additional methods to image the MTZ and the possible extent and existence of slabs in the Alaskan region. In this study, receiver function analysis is used to study the topography of the global mantle discontinuities, to gain more insight in the Earth structure underneath Alaska and the depth extent of the subducting slab.

1.2 Mantle discontinuities

Since tomographic models do not provide a consistent answer to the slab structure underneath Alaska, we study the MTZ in this region. The MTZ divides the Earth's upper and lower mantle

and is characterised by abrupt changes in seismic velocities. The globally-observed major seismic velocity discontinuities occur around depths of 410 and 660 km (hereafter referred to as the '410' and '660') and mark the top and the bottom of the MTZ. These discontinuities have been interpreted as polymorphic phase changes in the Mg_2SiO_4 -system, the olivine system. The 410 is associated with a phase transition from olivine (α - $(\text{Mg,Fe})_2\text{SiO}_4$) to wadsleyite (β - $(\text{Mg,Fe})_2\text{SiO}_4$) in an exothermic reaction (Katsura and Ito, 1989). The 660 is associated with a phase transition from ringwoodite (γ - $(\text{Mg,Fe})_2\text{SiO}_4$) as it disassociates into perovskite ($(\text{Mg,Fe,Al})(\text{Si,Al})\text{O}_3$) and magnesiowüstite ($(\text{Mg,Fe})\text{O}$) in an endothermic reaction (Ringwood et al., 1975; Ito and Takahashi, 1989). The non-olivine or garnet-pyroxene system also predicts phase transitions around the depth of the 660, and it depends on temperature which transition dominates (Yu et al., 2011; Hirose, 2002).

The phase transitions do not occur at the exact same depth everywhere, but they vary depending on temperature, composition (Hernández et al., 2015; Xu et al., 2008; Cobden et al., 2008) and water content (Buchen et al., 2018; Thio et al., 2016; Smyth and Frost, 2002). The behaviour of a phase transition is described by its clapeyron slope, defining the temperature-pressure range in which the transition occurs. For the olivine system, the transition at the 410 has a positive clapeyron slope, while the transition at the 660 has a negative clapeyron slope. Due to the opposite signs of the clapeyron slopes describing the olivine phase transitions around 410 and 660 km depth, the reactions at the top and the base of the MTZ behave opposite to temperature anomalies. This means that in colder regions an uplifted 410 and depressed 660 are expected, which will thicken the MTZ. Conversely, in hotter regions a depressed 410 and uplifted 660 are expected, thus creating a thinner MTZ.

Care should be taken in interpreting the transition zone thickness as temperature anomalies alone, because other factors may also control the reactions. The 660 discontinuity is thought to be especially variable, since not only the transition from ringwoodite to perovskite and magnesiowüstite occurs at this depth, but phase transitions in the non-olivine components play a role as well.

In relatively low temperature regions, the transition in the olivine system is thought to be dom-

inant, but ringwoodite dissociates to ilmenite first before transitioning to perovskite around the depth of the 660 (Stixrude and Lithgow-Bertelloni, 2011). Garnet instead of olivine dominates the phase transition at relatively high temperatures (Stixrude and Lithgow-Bertelloni, 2011). While temperature estimates of where ilmenite or garnet dominates the 660 vary widely, the phase diagram in Stixrude and Lithgow-Bertelloni (2011) suggests the ilmenite transition dominates the 660 at less than ~ 1500 K while the garnet transition dominates the 660 at temperatures larger than ~ 2100 K. Geodynamical estimated temperatures for plumes or slabs in the MTZ also vary. For example, Zhang et al. (2013) find temperatures of 1100 K at transition zone depth for a geodynamical slab model, and Maguire et al. (2018a) model a plume excess temperature of $\delta T=400$ K which relates to temperatures of ~ 2200 K at 660 km.

These phase transitions can cause a complex and broad 660 discontinuity with sometimes multiple signals (Deuss et al., 2006; Andrews and Deuss, 2008). The presence of water can further complicate the phase transitions in the MTZ. In the olivine system, wadsleyite and ringwoodite can store significant amounts of water, which alter the kinetics of the phase transformations (Ohtani et al., 2004).

Receiver functions are P-to-S converted phases at a discontinuity. Many studies have used teleseismic receiver functions to image an anomalous MTZ. Some examples are Maguire et al. (2018b); Tauzin et al. (2017); Jenkins et al. (2016); Owens et al. (2000); Li et al. (2000); Andrews and Deuss (2008) and many more. In this study receiver functions are used to map the topography on the major seismic mantle discontinuities. Receiver functions are sensitive to the local Earth structure underneath a seismic station by representing the response to the incoming compressional (P) wave, and can therefore be used to study discontinuities at different depths.

To use receiver functions to map the topography on the discontinuities, one needs to correct for anomalous crust and upper mantle structure. The exact velocity structure is usually not known, and accurately correcting for all velocity heterogeneities is challenging. Differential times between the arrivals from the 410 and 660 are less sensitive to variations in upper mantle structure, and many studies therefore interpreted the MTZ thickness instead of the discontinuity depths. Tauzin and Ricard (2014) propose a method to separate the competing effects of velocity structure and

temperature on the discontinuities, and use this to estimate the clapeyron slopes of phase changes in the transition zone. Also, some receiver function studies test the robustness of the obtained discontinuity depths using synthetic velocity models (e.g. Jenkins et al., 2016).

Several receiver function studies have been conducted in Alaska. For example, Ferris et al. (2003), Veenstra et al. (2006) and Miller et al. (2018) looked at the subducting plate at shallow depth and crustal thickness and structure, and Ai et al. (2005) studied the crust and upper mantle discontinuities in a small area in central Alaska. Using receiver functions to study the transition zone discontinuities underneath a large part of Alaska has only previously been done by Dahm et al. (2017). They find a broad zone with a thicker than normal MTZ and shallower than normal 410 and conclude that the subducting Alaskan slab has reached the MTZ where it is probably broken into fragments. They used the velocity model by Martin-Short et al. (2016) for the time-to-depth conversion. Much more data has become available compared to their data set which runs to late 2016. This increase is mainly due to the ongoing deployment of the USArray Transportable Array (TA), which has also greatly increased coverage in the more remote corners of Alaska. Besides the improved data coverage, our study also compares using two recently published S-wave velocity models for upper mantle corrections, which have also benefited from the improved data coverage. By comparing these two regional velocity models, we do show the sensitivity of the MTZ discontinuity topography to the velocity models, and particularly the degree of damping in the inversion, used for the mantle corrections.

2 METHOD

2.1 P-to-S receiver functions

When an incoming seismic compressional (P) wave meets a boundary or discontinuity the energy of the wave is partly converted to shear (SV) wave energy. This produces waves referred to as Pds phases, where the d denotes the depth of the conversion. The P410s and P660s, the waves converted at the 410 and 660 discontinuity, arrive 40 to 70 seconds after the direct P-arrival on the radial component of a seismogram. Their delay times can be used to determine the depths of the discontinuities assuming a velocity model between the discontinuity and the seismic station

(Langston, 1979). To highlight the converted phases, we stack the signals. To do this, we first deconvolve the source component. Assuming that the P wave arrival on the vertical component seismogram represents the source, we deconvolve the vertical component from the radial component. The time series resulting from this is called a receiver function and is sensitive to sharp structures underneath a station.

Creating the receiver functions can be done with various deconvolution techniques. We choose to use the iterative time domain deconvolution described by Ligorría and Ammon (1999). In this method the receiver function is constructed by adding Gaussian pulses defined as $G(\omega) = e^{-\frac{\pi^2 \omega^2}{a^2}}$ with $a = 0.4$ rad/s. This filter is also used to pre-filter the two components. The resulting Gaussian pulses in the receiver function in the time domain have a full-width at half maximum of ~ 4 s. Using higher frequencies will result in lower amplitude conversions, for example shown by Jenkins et al. (2017), while using lower frequencies will broaden the pulses further, potentially capturing less detail in the receiver functions.

The timing and amplitude of iteratively added Gaussian pulses to the receiver function is determined with the following steps: The current receiver function is convolved with the vertical component to produce the predicted radial component. This predicted radial component is then subtracted from the observed radial component to produce the unmodelled signal in the data. The new Gaussian pulse is added where the unmodelled signal is largest, determined by cross-correlating the unmodelled signal with the vertical. The Gaussian pulse is scaled by the amplitude of the maximum of the cross-correlated signal. These steps are iterated until adding more pulses shows insignificant improvement or 200 pulses have been added.

2.2 Data acquisition and processing

Data are collected from the Incorporated Research Institutions for Seismology (IRIS) database. ObsPy (Krischer et al., 2015) is used to download and process the data. Stations are selected with latitudes of 56° to 72° N and longitudes between 130° to 170° W. The events used occurred between January 2000 and October 2018 with magnitudes (M_w) between 5.5 and 8.3. Only events within an epicentral distance range of 30° to 90° from the stations are selected. Earthquakes with an epi-




central distance smaller than 30° do not have a ray path with a turning point deep enough to image the MTZ discontinuities, while for events with an epicentral distance larger than 90° the P wave is affected by the core-mantle boundary. The final data set before quality control consists of 405,348 events to station pairs from 477 stations across Alaska. Figs 2a and 2b show the distribution of events, stations and pierce points (the location at depth where the P-to-S conversion occurs).

The trend and mean in the data are removed and the north-south and east-west traces are rotated into radial and transverse component seismograms with 10 samples per second. Data is prefiltered between 0.01 - 0.2 Hz. Seismograms are then cut from 25 s before to 150 s after the predicted direct P arrival and these traces are used to create the receiver functions.

An automated quality check on the created receiver functions is done using the following criteria:

- The seismograms should have a signal-to-noise ratio of 2.5, i.e. the energy of the P wave arrival should be larger than 2.5 times the average energy in the time window used for the receiver functions.
- The receiver function convolved with the vertical component seismogram should at least reproduce 60% of the radial component.
- The direct P arrival in the receiver function needs to be within 1 s from zero, i.e. the main energy on the radial and vertical should be correlated.
- There should be no peaks before the main P wave which are more than 30% of the amplitude of the P wave and no peaks after the main P wave which are more than 70% of the amplitude of the P wave.
- There should be signals after the main P arrival in the receiver function which are at least 4% of the amplitude of the P wave.

Data for all stations are checked visually to ensure the automated quality check did not leave receiver functions with anomalous characteristics in the data set. After the quality check the data set contains 27,800 receiver functions.



2.3 Time-to-depth conversion

Next, a time-to-depth conversion is performed on every trace in order to interpret the receiver functions in terms of discontinuity depths. We have used three velocity models to perform this conversion. First, the 1D radial velocity model PREM (Dziewonski and Anderson, 1981) is used. The seismic travel time calculator Taup (Crotwell et al., 1999) as implemented in ObsPy (Krischer et al., 2015) calculates the predicted arrival times for the direct P-wave and P-to-S conversions. The velocity model calculates the predicted arrival time for a range of Pds phases using PREM and interpolates for all conversion depths, taking into account epicentral distance and event depth.

Because the Alaska region contains 3D upper mantle velocity variations, we need to account for these in the time-to-depth correction. Therefore, two recently published regional 3D velocity models are used:

(i) The shear velocity model by Martin-Short et al. (2018) is an updated version of Martin-Short et al. (2016), which is used in the receiver function study by Dahm et al. (2017). While Martin-Short et al. (2016) is a purely S-wave teleseismic model, Martin-Short et al. (2018) also includes ambient noise, surface waves and P-to-S receiver functions, constraining absolute velocities in the top 200 km as well as the crust. Hereafter we name this model MABPM.

(ii) The model by Jiang et al. (2018) is an upper mantle Rayleigh and S-wave tomography model, constraining absolute velocities in the top 240 km and relative velocities beneath, and is hereafter called model JSWLW.

Where the models give relative velocities, these are converted to absolute velocities using IASP91 (Kennett et al., 1995). The P wave model for the relative velocity models is in both cases obtained by scaling the shear wave velocity perturbation by assuming that $d \ln V_S / d \ln V_P$ increases from 2 at the surface to 3 at the core mantle boundary, following Ritsema et al. (2011). While the Martin-Short et al. (2016) has a corresponding P wave model as well, here we use the scaled model for the updated shear velocity model from Martin-Short et al. (2018) for consistency.

Where absolute V_S values are given in the uppermost 200 km in MABPM and the uppermost 240 km in JSWLW we apply the same $d \ln V_S / d \ln V_p$ scaling as for greater depths, using the




Figure 3. Shear wave velocity variations at 390 km depth for the 3D velocity models MABPM (Martin-Short et al., 2018) and JSWLW (Jiang et al., 2018) used for the time-to-depth correction. The large fast velocity anomaly is interpreted as the subducting Alaskan slab, indicated by the 0.5% $d\ln V_S$ isocontour in the grey dashed line. Active volcanoes at the surface are plotted in red.

average 1D V_S in each model as the reference velocity. The V_P reference velocity is obtained by rescaling the average 1D V_S by the V_P/V_S ratio in IASP91.

The MABPM model includes its own crustal model. For model JSWLW crustal model CRUST1.0 is added (Laske et al., 2013). These two models and PREM are each used to trace back the P and converted S phases within the 2D plane between the station and earthquake. We use the pierce points and delays for the converted phases in our stacking procedure.

Fig. 3 shows a slice through both 3D velocity models at 390 km depth. Note that JSWLW covers a wider area than MABPM. The velocity anomaly patterns of these models are similar, with at the same location the most prominent feature being the fast velocity anomaly in central Alaska, which corresponds to the subducting Pacific slab. However, the anomalies in model JSWLW are approximately twice as strong as the ones in model MABPM. This has large implications for the time-to-depth correction, as we will show in Section 3. In Section 4 we explore rescaling of the time-to-depth corrections for both models.

2.4 Common conversion point stacking

To improve the signal to noise ratio the time-to-depth converted traces are used to make common conversion point (CCP) stacks (Kosarev et al., 1999; Dueker and Sheehan, 1997). CCP stacks are made with receiver functions that sample the same region at depth while recorded at different stations and originating from different sources. We discretize the area underneath Alaska with a grid spacing of 0.5° in latitude and longitude and 2 km in depth, for depths between 60 and 1300 km. Every receiver functions is then traced back from the recording station to the source, using one of the velocity models described above. The amplitudes in the receiver functions at each depth are stacked into nearby grid points using a weighting factor based on the distance to the piercepoints. The weighting factor follows Lekic et al. (2011) and Cottaar and Deuss (2016) and depends on the Fresnel zone half width (FZHW), defined as

$$\text{FZHW}(z) = \sqrt{(\lambda(z)/3 + z)^2 - z^2}$$

where $\lambda(z)$ is the wavelength at depth z . The FZHW is used to normalize the distance between the conversion point at a given depth and the grid point in the stack:

$$\bar{\Delta} = \frac{\Delta}{\text{FZHW}}$$

The weight $\gamma(\bar{\Delta})$ is then given by:

$$\gamma(\bar{\Delta}) = \begin{cases} \frac{3}{4}\bar{\Delta}^3 - \frac{3}{2}\bar{\Delta}^2 & \text{if } \bar{\Delta} \leq 1 \\ \frac{1}{4}(2 - \bar{\Delta})^3 & \text{if } 1 < \bar{\Delta} \leq 2 \\ 0 & \text{if } \bar{\Delta} \geq 2 \end{cases}$$

In this way some smoothing is applied beyond the width of the Fresnel zone, i.e. at the edge of the Fresnel zone the weighting factor is 0.25. The smoothing increases as a function of depth, as the Fresnel zone widens with depth. Additionally, we smooth generously by using the FZHW for a 10-second shear wave, which is a dominant period in shear waves and at the half-width of our Gaussian filter.

The sum of the weights at 410 and 660 km depth can be seen in Fig. 4 for the CCP stack using PREM (Dziewonski and Anderson, 1981). The coverage is wider for 660 km depth, because the




Figure 4. Density of the data is illustrated by the sum of weights in the CCP stack using PREM for conversions at (a) 410 km and (b) 660 km depth. The weights decrease across 2x the Fresnel zone half width of a 10 s wave (with a value of 0.25 at 1x the Fresnel zone half width).

ray piercing points at 660 km depth have a wider distribution around the stations and the Fresnel zones are wider. The coverage is densest beneath the southern part of Alaska, due to the large number of seismic stations in that region. The summed weight maps for the 3D models look very similar.

Errors in the receiver function stacks are calculated using a running standard deviation at each grid point, which is then used to calculate the standard error. We consider arrivals as significant when they have a stacked amplitude larger than twice the corresponding standard error and therefore non-zero with 95% confidence.

3 RESULTS

The receiver function data set is used to make CCP stacks to study the topography on the transition zone discontinuities underneath Alaska using PREM, MABPM and JSWLW. Here we will present our results on the MTZ thickness and the topography of the 410 and the 660, for the different velocity models used.

3.1 Topography of the 410 and 660

Fig. 5 shows the transition zone thickness for the CCP stacks corrected with PREM, MABPM and JSWLW respectively. The MTZ thickness is calculated by subtracting the depth of the 410 from the depth of the 660, represented by the peak amplitudes around those depths in the stacks. The thickness of the MTZ varies considerably (Fig. 5). The most prominent and robust feature is the thicker transition zone beneath central Alaska. Around that region the MTZ thickness seems to be close to the globally considered average thickness of 250 km, with thinner parts in the southeast and southwest.

Fig. 6a shows the depths of the peak amplitudes between 382 and 442 representing the 410 discontinuity. Fig. 6b shows the same for peak amplitudes between 639 and 699, representing the 660 discontinuity. This scaling is chosen in such a way that the mean depth is located in the middle of the colour scheme. The mean depths are 413 and 671 km for MABPM and 411 and 667 km for JSWLW. Since the strongest velocity variations are present in the upper mantle, the transition zone thickness is less dependent on the velocity model used for the mantle corrections than the absolute discontinuity depths of the 410 and 660. Indeed, the transition zone thickness maps corrected with the different models (Fig. 5) are more in agreement with each other than the discontinuity depths maps (Figs 6a and b). The thickness variations seen here are thus robust features.

Next, we compare the discontinuity topography maps, starting with the 410 topography using MABPM with that using PREM (Fig. 6a). The most prominent feature in both images is the significantly uplifted 410 in central Alaska. The location of these uplifted P410s arrivals corresponds well with the location of the slab at 410 km depth in the 3D tomographic models MABPM and JSWLW (see Fig. 3). Since MABPM accounts for the fast velocities inside the slab, the time-to-depth correction shifts the arrivals deeper than with PREM where there is no correction for the fast velocity anomaly. Correcting with this velocity model takes out some of the extreme values present in the PREM results. Due to the positive clapeyron slope of the transition from olivine to wadsleyite an elevated 410 is expected at the location of a cold subducting slab. As this anomaly largely correlates with fast shear velocities in the tomographic models, the uplifted 410 in the stack using the MABPM model is most likely a signature of the slab at that depth. The 410 further has a

deep part in the south east (around 60°N and 135°W) and a shallow part in the north east (around 69°N and 135°W). The 660 discontinuity corrected by MABPM (Fig. 6b), shows less topography than the 410. The P660s arrivals appear slightly shallower beneath central Alaska, and slightly deeper surrounding this area, with an uplifted area in the north east and a deeper part in the south east. Again, if we compare it with the 660 map corrected with PREM, some extreme topography has been subdued resulting in less variation on the corrected 660 topography.

As opposed to MABPM, the JSWLW covers our entire study area. Additionally, the velocity anomalies are stronger in this model, therefore we also observe larger corrections with respect to the topography in the PREM stack, resulting in less topography on the 410 and 660. The 410 does not show a clear uplifted part around the slab location (around 64°N and 152°W), although this area is surrounded by a ring of uplifted 410 correlated with slow seismic velocities in the JSWLW model (see Fig. 3). The topography shows weaker variations in depth overall and does not have large consistent structures. Contrarily the 660 shows a prominent deep part in the south around 152°W , which is not seen when correcting with PREM or MABPM. Around the edges of our study area the JSWLW maps generally show more average depths for the 410 and 660 compared to the other two maps. Shifts are particularly strong in the southeast. The slow velocities in the JSWLW model here, cause what appears to be a deep 410 and deep 660 in the other two maps, to appear as a slightly deep 410 and shallow 660.

Apparently, the resulting depth of the 410 and 660 km discontinuities in the CCP stacked receiver functions is very dependent on the velocity model used for the mantle corrections. Implications of using these different models and interpretation of the imaged features will be discussed in the Section 4.

3.2 Cross sections

We also present two cross sections to show the Pds stacked amplitudes and waveforms (Fig. 7). Relative velocities in MABPM and JSWLW are plotted in the background to locate the large fast velocity anomaly corresponding to the subducting Alaskan slab. Both cross sections intersect the location of the subducting slab in the velocity model. In general, P410s is usually characterised by




Figure 5. Transition zone thickness calculated from the difference between the depth of the 410 and the 660 for the PREM, MABPM and JSWLW velocity model. Values are shown for a minimum summed weight of 40 (see Fig. 4) and only when both the 410 and 660 arrivals are significant, i.e. their amplitudes are more than twice the standard error. Note that MABPM results are only shown where the 3D model is constrained (see Fig. 3).

a simple single peak in the seismic observations, while P660s often has a broader and more complicated waveform (Deuss et al., 2006; Andrews and Deuss, 2008). Indeed, in the cross sections it can be seen that P660s has a broader and more asymmetric peak than P410s, although in the frequency band used here, there is no observable splitting of the P660s.

We start with the cross sections for the stacks corrected with MABPM. On cross section AB, it can be seen that the 410 discontinuity is uplifted around the slab location (between angular distance 4° and 6°), and deeper in the rest of the cross section. Looking at the 660, there is less topography on this discontinuity than on the 410. In the south (near B), both the 410 and the 660 appear deeper than in the rest of the profile. In cross section CD, located to the south of AB, there is again not much topography observed on the 660 discontinuity. In addition, in a part of this cross section amplitudes of the CCP stacked traces decrease significantly around 410 km depth.

The cross sections corrected with JSWLW show differences with the cross sections described above. Around the slab location in cross section AB, we do not see an uplifted 410, but instead the 660 appears deeper there. In the south, the 410 is somewhat deeper, while the 660 is shallower than in the rest of the profile. On cross section CD the 660 also seems to be deeper around the slab location, as opposed to little variation seen on the 660 when using the MABPM model. The 410 arrivals decrease in amplitude in a part of the cross section, just as we observe for the MABPM




Figure 6. Depth of the peak amplitudes in the receiver functions corresponding to (a) the depth of the 410 discontinuity and (b) the depth of the 660 discontinuity for the PREM, MABPM and JSWLW velocity models. Results are plotted if the sum of weights for the CCP stack is more than 40 stacked receiver functions and if the arrivals are significant. Note that MABPM results are only shown where the 3D model is constrained (see Fig. 3). The colour scale is centred around values of 413 and 669 km depth.

results. So, although the depth of the discontinuities shifts when using a different velocity model, the waveforms stay the same.

The CCP stacks show hints of additional arrivals in the upper mantle, within the MTZ, and at the top of the lower mantle, some of which can be seen in the cross sections. We have analysed the incidence angles or slownesses of these arrivals, but the results show they are more likely multiples resulting from shallow complexities in the upper mantle than robust arrivals from depth.




Figure 7. CCP stacked traces for two cross sections AB and CD. The 3D velocity models MABPM by Martin-Short et al. (2018) and JSWLW by Jiang et al. (2018) are plotted in the background, the large fast velocity anomaly is interpreted as the subducting Alaskan slab, indicated by the 0.5% $d\ln V_S$ isocontour in the grey dashed line. Both cross sections sample areas with high data coverage, where data coverage drops below 100 (at shallower depths) the stacked receiver functions are not plotted. The traces are coloured red (positive amplitude) or blue (negative amplitude) when they lie within the 95% confidence interval. The locations of the cross sections are indicated on both velocity models at 390 km depth.

4 DISCUSSION

4.1 Dependency on velocity models

As shown in Section 3, the depth of the discontinuities is very dependent on the velocity model used for the mantle corrections. The strong differences between the two recent models applied here are due to the large amplitude differences in the tomographic models. In this Section we will show through quantitative analysis that one model is under-corrected while the other is over-corrected. We also show how tomographic velocity corrections can be rescaled to improve upper mantle wave speed corrections within CCP stacks using this test.

Looking at the two 3D regional models in Fig. 3, the pattern of fast and slow anomalies is similar, but the anomalies at a given depth are much stronger for JSWLW than for MABPM. For example, at 390 km depth, the largest anomaly in the slab in MABPM is 2.3% fast while the largest anomaly in the slab in JSWLW is 5.1% fast, more than twice as large. At 250 km depth, the largest anomaly in JSWLW has a value of 11.7%, while this is only 3.6% in MABPM. Both models have relative travel-time anomalies at deeper depths (below 200 km for MABPM and 240 km for JSWLW). For such models the mean 1D absolute reference velocities is unknown, and our assumed background model might cause artefacts (Bastow, 2012). For example, what we use in our corrections as slow anomalies, i.e. the anomalies around the fast slab in Fig. 3, might only be slow with respect to the regional mean and not the global 1D reference model we use here. Further, the ill-posed nature of tomographic inversion problems leads to non-unique solutions. For example, different methods (e.g. using a full waveform inversion scheme or not), choices of model parametrisation, the amount of damping and smoothing and the amount of iterations can all lead to different resulting tomographic models, with anomalies of widely varying amplitudes. De Wit et al. (2012) studied the non-uniqueness of tomographic models by examining the null-space of the forward operator and found a broad range of acceptable solutions compatible with the data. Quantitative assessment of the posterior model covariance and synthetic sensitivity analyses could help assess the difference and accuracy in observed amplitudes in the models (Rawlinson and Spakman, 2016).




Figure 8. Correlation tests for the results with time-to-depth conversion using PREM (a), MABPM (b-d), JSWLW (e-f): the depth of the 410 versus the depth of the 660 (a, b, e), the depth of the 410 versus the correction on the 410 (c, f), and the depth of the 660 versus the correction on the 660 (d, g). Pearson's r gives the measure of correlation between the discontinuity depths in the upper left of each subplot. The cartoon in (h) shows the implications of correlated temperatures in the upper mantle and MTZ for the discontinuity depths. The topography on the discontinuities is exaggerated in the cartoon.

Here we present a different way to assess the amplitudes of velocity anomalies in tomographic models through a number of correlation tests. Some receiver function studies use the correlation between the 410 and 660 depths to test whether there are unaccounted for velocity perturbations after mantle corrections (e.g. Gao and Liu, 2014; Chevrot et al., 1999; Dahm et al., 2017; Dueker and Sheehan, 1998). We add to this calculations of the correlation between the discontinuity depths and the correction applied to the discontinuity depths for the different velocity models.

Fig. 8 shows the various correlation plots for discontinuity topography in the CCP stacks for the velocity models PREM (Fig. 8a), MABPM (Fig. 8b-d), and JSWLW (Fig. 8e-f). For this analysis we first retrieve topography values at CCP grid point locations where all three model stacks have significant arrivals for the P410s and the P660s, which occurs at 84% of the grid points with a data coverage over 40. For the tests showing MABPM results, only locations in the area of

the velocity model are used (Fig. 3). Our CCP grid spacing every 0.5° in latitude and longitude direction results in unequal sampling. We therefore interpolate the observed topographies to a grid of 1336 equally spaced points (914 for MABPM) for use in these statistical tests. We used the Pearson correlation coefficient, or Pearson's r , as a measure of correlation. Here we describe the three different correlation tests shown, what we expect for each of these correlations with the help of the cartoon in Fig. 8h, and what we observe in the case of each model.

- *Correlation between the 410 and 660 (Fig. 8a, b and e)*: If the topography on the 410 and 660 is purely due to temperature variations in a olivine-dominated system, we expect anti-correlation between the topography on the 410 and 660 (solid lines in Fig. 8h). In the case that the 410 and 660 are affected by different compositions and temperatures, i.e. anomalies do not extend across the MTZ, there could be an overall lack of correlation. For these reasons we do not know if after 3D corrections the 410 and 660 should be anti-correlated, or decorrelated. Additionally a garnet dominated transition zone at higher temperatures results in local deepening of the 660 and therefore could introduce a degree of positive correlation with the 410 topography.

Lack of corrections for upper mantle structure has the same topographic effect on the 410 and 660 and causes correlation between the two (dashed lines in Fig. 8h). Any incorrect shifts by the velocity model across the upper mantle could increase the correlation between the 410 and 660, while incorrect shifts from velocity anomalies within the MTZ would decorrelate the 410 and 660. As expected, PREM results show the highest correlation between the depths of the 410 and 660 discontinuity, with a Pearson's r of 0.79 (Fig. 8a). Corrections using MABPM lower the Pearson's r to 0.53 (Fig. 8b), but as correlation remains significant, the topography is likely under-corrected. The results corrected with JSWLW have a very low Pearson's R of only 0.08 (Fig. 8e), potentially implying suitable corrections, but this remains inconclusive since the expected correlation given perfect corrections is unknown.

- *Correlation between the 410 and corrections on the 410 (Fig. 8c and f)*: Assuming perfect 3D velocity corrections, we could expect three possible cases here. If there is no correlation between the temperatures in the MTZ and shallow mantle, we expect no correlation between the resulting

topography and the corrections. If there is some correlation between the temperatures in the MTZ and shallower structures, i.e. in the case of continuous hot upwellings or cold downwellings, the positive Clapeyron slope of the 410 phase transition leads to negative correlation with the corrections. In other words, a deep 410 in a hot area was shifted to shallower depths, while a shallow 410 in a cold area was shifted to deeper depths (Fig. 8h). If there is some anti-correlation between the MTZ and upper mantle, i.e. where a slow upper mantle sits above a cold slab in the MTZ, the corrections would be opposite in sign, and a positive correlation is expected.

- *Correlation between the 660 and corrections on the 660 (Fig. 8d and g):* Due to the opposite sign of the Clapeyron slope of the 660 phase transition compared to that for the 410, we expect opposite behaviour in the correlation coefficients. Where there is correlation between temperatures in the MTZ and shallow mantle, we expect positive correlation between the topography of the 660 and the corrections thereof. Where the 660 is deep, corrections have pushed it deeper, and where the 660 is shallow, the corrections have pulled it shallower. When there is anti-correlation between the MTZ and upper mantle, a negative correlation is expected.

In the case that there is a degree of anti-correlation between the temperatures in the MTZ and shallow mantle, we would see positive correlation between the depth of the 410 and corrections thereof, while at the same time negative correlation between the depth of the 660 and corrections on the 660. Since we do not see this in either one of the models when we apply the statistical test, we argue that in this area we do not have such an anti-correlation. We therefore argue that we either have zero, or a positive correlation between temperatures in the MTZ and the shallow mantle.

Looking at the correlation between the 410 and the correction on the 410 for the two models (Figs 8c and f), the MABPM results show a degree of negative correlation, while those for JSWLW show positive correlation. If we take the thought experiment that there is a degree of positive correlation between the MTZ and upper mantle, we would expect to see a negative correlation here. The positive correlation for the JSWLW then indicates that the corrections on the 410 are either

too strong and thus placing the discontinuity too deep (dotted line in Fig. 8h) or are introducing incorrect topography through incorrect anomalies.

For the correlation between the 660 and the correction on the 660 (Figs 8d and g), the Pearson's r for the two models again has an opposite sign. In the thought experiment of positive correlation between the MTZ and upper mantle, the unexpected negative Pearson's r for MABPM means this case is under-corrected (dotted line in Fig. 8h). The Pearson's r value of 0.82 for JSWLW is also suspiciously high, and combined with the positive correlation for the 410 in Fig. 8f could be explained by topography introduced by over-correcting of the model. The strong positive correlation could also, combined with the decorrelation between the 410 and 660 in Fig. 8e, suggest strong corrections across the MTZ, introducing topography on the 660 only.

In summary, the negative correlation coefficient between the 660 and corrections on the 660 for the MABPM results suggests under-correction, while the positive correlation between the 410 and the corrections on the 410 for the JSWLW results suggests over-correction. This is consistent with the stronger amplitudes of the velocity variations seen in the JSWLW over MABPM (see Fig. 3).

We investigate the implications on the correlation coefficients by rescaling the topographic corrections by a range of factors. In order to do so, we take the difference between the topography of the 3D model and PREM, multiply that with a specific factor and add it to the PREM topography again. A major assumption in this approach is that the corrections everywhere are over- or underestimated by the same degree, while in reality resolution and amplitude recovery will vary laterally and with depth in a tomographic model. Fig. 9 shows behaviour of the three correlation coefficients exploring a rescaling factor for the MABPM model between 1.0 and 2.4, and for the JSWLW model between 0.5 and 1.2. The correlations vary with scaling factor in ways we expect. Under-corrected models lead to negative correlations between both the depths of the discontinuities and the corrections on the depths, while over-corrected models lead to positive correlations for both. It should be noted that the correlation coefficient for the 660 increases faster than that for the 410, which is due to there being less topographic variations on the 660 compared to the 410 causing any topography introduced by over-corrections to lead to strong correlations.

As expected, the increase in correction rescaling factor generally leads to a decrease in the correlation between the 410 and 660, but if corrections become too strong these can increase again as seen for the JSWLW case, where the minimum correlation between the 410 and 660 lies around a factor of 0.85-0.9.

We show the range of Goldilocks values for the rescaling factor (grey shaded area in Fig. 9), where the correlation between the depth of the 410 and the correction on the 410 depth is zero or negative *and* the correlation between the depth of the 660 and the correction on the 660 is zero or positive. We suggest the optimal values are where these correlations are roughly of the same magnitude at opposite signs, these correlations result from a degree of correlation between anomalies in the MTZ and shallower structure affecting both the 410 and 660. This gives factors of 1.3 and 0.7 for MABPM and JSWLW, respectively. For these factors the correlation coefficient between the 410 and 660 is 0.39 for MABPM and 0.17 for JSWLW. As we do not expect any positive correlation between the 410 and 660 due to anomalously hot, garnet-dominated, MTZ beneath Alaska, we attribute this residual correlation due to imprecision of the tomographic models in general, and our simplified rescaling in particular. While the rescaling values found are largely driven by the difference in amplitude of the velocity anomalies in the two models, the exact values found also depends on our assumptions when scaling to Vp velocities.

This exploration additionally shows that there are no choices of rescaling that suggest decorrelation or a degree of anti-correlation between the corrections and the topography seen on the MTZ, i.e. there are no values where both the 410 and 660 are decorrelated with the corrections, or values where the 410 has positive correlations with the corrections while at the same time the 660 has negative ones. We could however have chosen a rescaling factor, close to our current chosen factor, where corrections are only correlated with the 410 topography and decorrelated with the 660 topography.

The resulting topographies using these scaling factors are shown in Fig. 10. Compared with Fig. 6, the topographies corrected with the different models are, as expected, more similar. The correlation between the two models changes from 0.66 in the old scaling to 0.86 in the new scaling for the depth of the 410, and from -0.16 to 0.42 for the 660. Interpretation of the final topographies




Figure 9. Correlation coefficients for the three correlation tests as a function of rescaling factors for (a) model MABPM and (b) model JSWLW. Goldilocks range for the rescaling factor are shown by grey area, and optimal values by the dashed line.

is described in the next Section. With the tests above we provide a quantitative assessment of the amplitudes of velocity anomalies in tomographic models based on an independent data set. It should be noted that this assessment is not directly sensitive to the amplitude of the anomalies as such, but to the vertically integrated anomalies, and can therefore have less resolution on the degree of smearing that occurs in a tomographic model. The rescaling approach provides a new avenue to use the tomographic models for a better time-to-depth correction, but needs to be tested at different regional and global scales. In our current application we use a single rescaling factor, ignoring the fact that amplitude recovery can vary laterally and with depth for a tomographic model.

4.2 Interpretation of the imaged features

The two features that have little dependence on the velocity model used for the corrections are the MTZ thickness and the waveforms of the converted phases. Indeed, the MTZ thickness only shows minor changes after rescaling, and is therefore not shown again in Fig. 10. After rescaling we are able to interpret the absolute discontinuity depths with more confidence than without the rescaling before.

As shown in Section 3 the MTZ thickness for all three models increases in the area corre-




Figure 10. Depth of the peak amplitudes in the receiver functions corresponding to (a) the depth of the 410 discontinuity and (b) the depth of the 660 discontinuity for the rescaled MABPM and JSWLW velocity models. Results are plotted if the sum of weights for the CCP stack is more than 40 stacked receiver functions and if the arrivals are significant. Note that MABPM results are only shown where the 3D model is constrained (see Fig. 3). The colour scale is consistent with Fig. 6.

sponding to the location of the subducting slab. A thicker transition zone implies either only an uplift of the 410, or an uplifted 410 and depressed 660. In Fig. 10a it can be seen that both rescaled models show an uplifted 410 discontinuity to depths of 390-395 km around the location of the slab. Since the transition from olivine to wadsleyite has a positive clapeyron slope, this uplifted 410 is interpreted as the interaction of the cold slab with the phase transition with an estimated temperature anomaly of 180 to 290 K, assuming clapeyron slopes between +4 MPa/K (Katsura et al., 2004) and +2.5 MPa/K (Katsura and Ito, 1989).

We do not observe a deeper 660 in the same region as the uplifted 410. Some studies have shown that the clapeyron slope for the 660 might be more gentle than that of the 410 (e.g. Katsura et al., 2003; Fei et al., 2004), leading to less topography on the former discontinuity. However,

potential signatures of slabs on the 660 have been found in various places (e.g. Gao and Liu, 2014; Lee et al., 2014; Suetsugu et al., 2010; Cao and Levander, 2010), with depression of the 660 as much as 35 km. Particularly strong signatures on the 660 have also been found in places of ponding slabs (e.g. Cottaar and Deuss, 2016). Overall in the area studied here, the 660 discontinuity shows little topography (Fig. 10b) with variations up to ~ 10 km, compared the ~ 25 km on the 410. The P660s in the stacks shows consistent amplitudes and waveforms, with no observations of splitting. These observations imply that the slab has not reached the 660.

The results on both discontinuities suggest that the Pacific slab, together with the Yakutat block in the eastern part of the subduction zone, is indeed subducting there and that the slab is penetrating through the 410 discontinuity and ending somewhere in the MTZ. This is in agreement with the depth extent of the slab concluded by Dahm et al. (2017), who used the velocity model by Martin-Short et al. (2016), of which the model MABPM is an update. It is also in agreement with the depth extent of the slab in the velocity models that we use, and with the "Atlas of the Underworld" (van der Meer et al., 2017), and the P-wave velocity model by Gou et al. (2019). These observations raise the question where the thousands of km of subducted lithosphere in common plate reconstruction models (Müller et al., 2016; Seton et al., 2012; Madsen et al., 2006) currently resides in the mantle. Part of the convergence may be taken up by intra-oceanic subduction, as suggested by Domeier et al. (2017) and Vaes et al. (2019). The slab associated with such a subduction zone may correspond to anomalies identified in previous studies (e.g. Müller et al., 2016; Domeier et al., 2017; Sigloch, 2011) such as the North Pacific Slab in the "Atlas of the Underworld" (van der Meer et al., 2017). Our results are consistent with the previously published tomographic images that suggest that the Alaskan slab does not reach the lower mantle and may thus inspire future tectonic reconstructions that reconcile this with large-scale plate convergence.

However, the signal of uplifted 410 extends further to the east (around 65°N and 142°W) than we might expect from tomography or present day surface subduction. Also, looking at the location of the subducting slab in the velocity models (Fig. 3) one would expect the uplifted signal on the 410 to continue west of the Aleutian Island arc. This signal can be observed in JSWLW, but is relatively weak in MABPM.

Further south, beneath the Kenai Peninsula, the area around Katmai National Park and Kodiak Island, shows a different and more confusing topography on the 410. This area lies beneath the subducting slab. A large part of this shows a deep 410, juxtaposed against a shallow 410 beneath the Kodiak Island. Overall the P410s here has weak amplitudes and both maps of MABPM and JSWLW having large regions where no significant arrivals can be picked (shown by the grey areas). The entire area has reasonable data coverage (Fig. 4) and data quality, as exemplified by the quality of the maps for the 660 in this area. In general though, the observed P410s are weaker in amplitude and have narrower arrivals compared to the 660, making this phase more sensitive to incoherent stacking either due to strong topography in the area, or due to incorrect 3D velocity corrections of the complex tectonic region above this area. The entire velocity structure may contribute to amplitude variations, for example velocity gradients near a discontinuity or multiples that contribute energy near the delay times of the Pds phases (Andrews and Deuss, 2008). It is therefore difficult to judge if the topography and the amplitudes in this area are a true representation of a complex discontinuity or an artefact.

In cross section CD in Fig. 7, there is a region where the uplifted P410s has very weak amplitudes for both velocity models. Again the weak amplitudes could be an artefact of stacking across an area of strong topography. If it is a true signature of the discontinuity, a possible explanation for a weak or absent 410 is the presence of water. The existence of water expands the stability field of wadsleyite to a lower pressure. This causes a broader seismic discontinuity at 410 km depth (Litasov et al., 2006; van der Meijde et al., 2003), and thus smaller amplitudes in the arrivals on the receiver functions. Presence of basalt can also reduce the signal at the 410, because it weakens the phase changes in the olivine system. The model by Xu et al. (2008) showed that an increasing basalt fraction leads to a smaller velocity jump around 410 km depth. Further, the presence of ferric Fe in the mantle may play a role in broadening the 410 (Frost and Dolejš, 2007).

A thinner than normal MTZ is observed in the southeast of the area studied, where thicknesses of ~ 230 km are observed. The location corresponds to a low velocity anomaly in the 3D models of Martin-Short et al. (2018), Jiang et al. (2018), Qi et al. (2007) and Burdick et al. (2017). In the geometric plate-tectonic model of Madsen et al. (2006) this location corresponds to the location

of a slab window. Qi et al. (2007) suggest this low velocity anomaly is associated with upwelling mantle in the slab window. Both the velocities in the model, and the thinner MTZ are consistent with the interpretation of a hot mantle upwelling with olivine dominated phase transitions. If we take 20 km thinning of the MTZ and look at an example where the 410 is 10 km deeper and the 660 is 10 km shallower, this corresponds to a temperature increase between 94-150 K on the 410 (assuming clapeyron slopes between +4 MPa/K (Katsura et al., 2004) and +2.5 MPa/K (Katsura and Ito, 1989)) and between 118-160 K on the 660 (assuming clapeyron slopes between -3.4 MPa/K (Hernández et al., 2015) and -2.5 MPa/K (Ye et al., 2014)).

In the northeast, the maps using PREM show an extremely shallow 410 and 660. This area lies at the edge of the North American craton. Cratons consist of highly stable continental crust, which are chemically depleted areas with a lower than average temperature (Yuan and Romanowicz, 2010) and have a higher than average velocity. This area lies outside the region where the velocity model MABPM has good resolution. The JSWLW model shows fast velocities in this region, which, as expected, shifts the 410 and 660 deeper, resulting in average depths in the original stack (Fig. 6) and close to average depth after rescaling (Fig. 10). In all models the MTZ seems slightly thicker than average, around 260 km thick. In the global model UU-P07 (Amaru, 2007; Hall and Spakman, 2015) the fast anomaly below northwestern North America is interpreted as the Yukon slab, visible within the upper part of the lower mantle. Thickening of the MTZ here might be due to the presence of the Yukon anomaly.

5 CONCLUSIONS

Using 27,800 high quality receiver functions to map the MTZ beneath Alaska, we demonstrate that receiver function common conversion point stacks are highly dependent on the velocity model used for the mantle corrections by using two different recent models with differing velocity anomaly amplitudes. We show that correlations between the depth of the discontinuities and the topographic corrections from the 3D velocity model can be used to test the degree of under- or over-correction. We find one model under-corrected and the other model over-corrected, and rescale the topographic corrections to obtain comparable topographic maps. The final maps show the effects of

the slab on the 410, confirming that it has penetrated into the MTZ. As the 660 appears unaffected, the slab must end within the MTZ. In some parts, the P410s arrivals have a very small amplitude, which can be caused by compositional variation in the mantle, or by too much topography on the discontinuity. A thinner than normal transition zone is observed in the southeast, which may be caused by hot mantle upwellings associated with a slab window.

6 ACKNOWLEDGEMENTS

The facilities of IRIS Data Services, and specifically the IRIS Data Management Center, were used for access to waveforms, related metadata, and derived products used in this study. The largest seismic networks contributing data to this research were the USArray Transportable Array from the EarthScope USArray project and the Alaska Regional Network operated by the Alaska Earthquake Center. We thank R. Martin-Short and C. Jiang for providing the 3D velocity models for Alaska, and we thank Alistair Boyce and two anonymous reviewers for helpful comments. AvS and AD were supported by the European Research Council (ERC) under the European Unions Horizon 2020 research and innovation programme (grant agreement No. 681535 - ATUNE) and a Vici award number 016.160.310/526 from the Netherlands organization for scientific research (NWO). SC was supported by the European Research Council (ERC) under the European Unions Horizon 2020 research and innovation programme (grant agreement No. 804071 - ZoomDeep) and the Natural Environment Research Council [NERC grant reference number NE/R010862/1].

References

- Ai, Y., Zhao, D., Gao, X., and Xu, W. (2005). The crust and upper mantle discontinuity structure beneath Alaska inferred from receiver functions. *Physics of the Earth and Planetary Interiors*, 150(4):339–350.
- Amaru, M. (2007). *Global travel time tomography with 3-D reference models*. PhD thesis, Utrecht University.
- Andrews, J. and Deuss, A. (2008). Detailed nature of the 660 km region of the mantle from global receiver function data. *Journal of Geophysical Research: Solid Earth*, 113(B6).

- Bastow, I. D. (2012). Relative arrival-time upper-mantle tomography and the elusive background mean. *Geophysical Journal International*, 190(2):1271–1278.
- Bauer, M. A., Pavlis, G. L., and Landes, M. (2014). Subduction geometry of the Yakutat terrane, southeastern Alaska. *Geosphere*, 10(6):1161–1176.
- Bird, P. (2003). An updated digital model of plate boundaries. *Geochemistry, Geophysics, Geosystems*, 4(3).
- Buchen, J., Marquardt, H., Speziale, S., Kawazoe, T., Ballaran, T. B., and Kurnosov, A. (2018). High-pressure single-crystal elasticity of wadsleyite and the seismic signature of water in the shallow transition zone. *Earth and Planetary Science Letters*, 498:77–87.
- Burdick, S., Vernon, F. L., Martynov, V., Eakins, J., Cox, T., Tytell, J., Mulder, T., White, M. C., Astiz, L., Pavlis, G. L., et al. (2017). Model Update May 2016: Upper-mantle heterogeneity beneath North America from travel-time tomography with global and USArray data. *Seismological Research Letters*, 88(2A):319–325.
- Cao, A. and Levander, A. (2010). High-resolution transition zone structures of the Gorda Slab beneath the western United States: Implication for deep water subduction. *Journal of Geophysical Research: Solid Earth*, 115(B7).
- Chevrot, S., Vinnik, L., and Montagner, J.-P. (1999). Global-scale analysis of the mantle Pds phases. *Journal of Geophysical Research: Solid Earth*, 104(B9):20203–20219.
- Cobden, L., Goes, S., Cammarano, F., and Connolly, J. A. (2008). Thermochemical interpretation of one-dimensional seismic reference models for the upper mantle: evidence for bias due to heterogeneity. *Geophysical Journal International*, 175(2):627–648.
- Cottaar, S. and Deuss, A. (2016). Large-scale mantle discontinuity topography beneath Europe: Signature of akimotoite in subducting slabs. *Journal of Geophysical Research: Solid Earth*, 121(1):279–292.
- Crotwell, H. P., Owens, T. J., and Ritsema, J. (1999). The TauP Toolkit: Flexible seismic travel-time and ray-path utilities. *Seismological Research Letters*, 70(2):154–160.
- Dahm, H. H., Gao, S. S., Kong, F., and Liu, K. H. (2017). Topography of the mantle transition zone discontinuities beneath Alaska and its geodynamic implications: Constraints from receiver

- function stacking. *Journal of Geophysical Research: Solid Earth*, 122(12).
- De Wit, R., Trampert, J., and Hilst, R. (2012). Toward quantifying uncertainty in travel time tomography using the null-space shuttle. *Journal of Geophysical Research: Solid Earth*, 117(B3).
- DeMets, C. and Dixon, T. H. (1999). New kinematic models for Pacific-North America motion from 3 Ma to present, I: Evidence for steady motion and biases in the NUVEL-1A Model. *Geophysical Research Letters*, 26(13):1921–1924.
- Deuss, A., Redfern, S. A., Chambers, K., and Woodhouse, J. H. (2006). The nature of the 660-kilometer discontinuity in Earth’s mantle from global seismic observations of PP precursors. *Science*, 311(5758):198–201.
- Domeier, M., Shephard, G. E., Jakob, J., Gaina, C., Doubrovine, P. V., and Torsvik, T. H. (2017). Intraoceanic subduction spanned the Pacific in the Late Cretaceous–Paleocene. *Science advances*, 3(11):eaao2303.
- Dueker, K. G. and Sheehan, A. F. (1997). Mantle discontinuity structure from midpoint stacks of converted P to S waves across the Yellowstone hotspot track. *Journal of Geophysical Research: Solid Earth*, 102(B4):8313–8327.
- Dueker, K. G. and Sheehan, A. F. (1998). Mantle discontinuity structure beneath the Colorado Rocky Mountains and high plains. *Journal of Geophysical Research: Solid Earth*, 103(B4):7153–7169.
- Dziewonski, A. M. and Anderson, D. L. (1981). Preliminary reference Earth model. *Physics of the earth and planetary interiors*, 25(4):297–356.
- Eberhart-Phillips, D., Christensen, D. H., Brocher, T. M., Hansen, R., Ruppert, N. A., Haeussler, P. J., and Abers, G. A. (2006). Imaging the transition from Aleutian subduction to Yakutat collision in central Alaska, with local earthquakes and active source data. *Journal of Geophysical Research: Solid Earth*, 111(B11).
- Efron, B. and Tibshirani, R. (1991). Statistical data analysis in the computer age. *Science*, 253(5018):390–395.
- Engebretson, D. C., Cox, A., and Gordon, R. G. (1985). Relative motions between oceanic and continental plates in the Pacific basin. *Geological Society of America Special Papers*, 206:1–60.

- Fei, Y., Van Orman, J., Li, J., Van Westrenen, W., Sanloup, C., Minarik, W., Hirose, K., Komabayashi, T., Walter, M., and Funakoshi, K.-i. (2004). Experimentally determined post-spinel transformation boundary in Mg_2SiO_4 using MgO as an internal pressure standard and its geophysical implications. *Journal of Geophysical Research: Solid Earth*, 109(B2).
- Ferris, A., Abers, G. A., Christensen, D. H., and Veenstra, E. (2003). High resolution image of the subducted Pacific (?) plate beneath central Alaska, 50–150 km depth. *Earth and Planetary Science Letters*, 214(3):575–588.
- Frost, D. J. and Dolejš, D. (2007). Experimental determination of the effect of H_2O on the 410-km seismic discontinuity. *Earth and Planetary Science Letters*, 256(1-2):182–195.
- Fuis, G. S., Moore, T. E., Plafker, G., Brocher, T. M., Fisher, M. A., Mooney, W. D., Nokleberg, W. J., Page, R. A., Beaudoin, B. C., Christensen, N. I., et al. (2008). Trans-Alaska crustal transect and continental evolution involving subduction underplating and synchronous foreland thrusting. *Geology*, 36(3):267–270.
- Gao, S. S. and Liu, K. H. (2014). Mantle transition zone discontinuities beneath the contiguous United States. *Journal of Geophysical Research: Solid Earth*, 119(8):6452–6468.
- Gou, T., Zhao, D., Huang, Z., and Wang, L. (2019). Aseismic deep slab and mantle flow beneath Alaska: Insight from anisotropic tomography. *Journal of Geophysical Research: Solid Earth*.
- Gulick, S. P., Lowe, L. A., Pavlis, T. L., Gardner, J. V., and Mayer, L. A. (2007). Geophysical insights into the transition fault debate: Propagating strike slip in response to stalling Yakutat block subduction in the Gulf of Alaska. *Geology*, 35(8):763–766.
- Hall, R. and Spakman, W. (2015). Mantle structure and tectonic history of SE Asia. *Tectonophysics*, 658:14–45.
- Hayes, G. P., Moore, G. L., Portner, D. E., Hearne, M., Flamme, H., Furtney, M., and Smoczyk, G. M. (2018). Slab2, a comprehensive subduction zone geometry model. *Science*, 362(6410):58–61.
- Hernández, E., Brodholt, J., and Alfè, D. (2015). Structural, vibrational and thermodynamic properties of Mg_2SiO_4 and MgSiO_3 minerals from first-principles simulations. *Physics of the Earth and Planetary Interiors*, 240:1–24.

- Hirose, K. (2002). Phase transitions in pyrolitic mantle around 670-km depth: Implications for upwelling of plumes from the lower mantle. *Journal of Geophysical Research: Solid Earth*, 107(B4).
- Ito, E. and Takahashi, E. (1989). Postspinel transformations in the system $\text{Mg}_2\text{SiO}_4 - \text{Fe}_2\text{SiO}_4$ and some geophysical implications. *Journal of Geophysical Research: Solid Earth*, 94(B8):10637–10646.
- Jadamec, M. and Billen, M. (2012). The role of rheology and slab shape on rapid mantle flow: Three-dimensional numerical models of the Alaska slab edge. *Journal of Geophysical Research: Solid Earth*, 117(B2).
- Jenkins, J., Cottaar, S., White, R., and Deuss, A. (2016). Depressed mantle discontinuities beneath Iceland: Evidence of a garnet controlled 660 km discontinuity? *Earth and Planetary Science Letters*, 433:159–168.
- Jenkins, J., Deuss, A., and Cottaar, S. (2017). Converted phases from sharp 1000 km depth mid-mantle heterogeneity beneath Western Europe. *Earth and Planetary Science Letters*, 459:196–207.
- Jiang, C., Schmandt, B., Ward, K. M., Lin, F.-C., and Worthington, L. L. (2018). Upper mantle seismic structure of Alaska from Rayleigh and S wave tomography. *Geophysical Research Letters*, 45(19):10–350.
- Katsura, T. and Ito, E. (1989). The system $\text{Mg}_2\text{SiO}_4 - \text{Fe}_2\text{SiO}_4$ at high pressures and temperatures: Precise determination of stabilities of olivine, modified spinel, and spinel. *Journal of Geophysical Research: Solid Earth*, 94(B11):15663–15670.
- Katsura, T., Yamada, H., Nishikawa, O., Song, M., Kubo, A., Shinmei, T., Yokoshi, S., Aizawa, Y., Yoshino, T., Walter, M. J., et al. (2004). Olivine-wadsleyite transition in the system $(\text{Mg}, \text{Fe})_2\text{SiO}_4$. *Journal of Geophysical Research: Solid Earth*, 109(B2).
- Katsura, T., Yamada, H., Shinmei, T., Kubo, A., Ono, S., Kanzaki, M., Yoneda, A., Walter, M. J., Ito, E., Urakawa, S., et al. (2003). Post-spinel transition in Mg_2SiO_4 determined by high P–T in situ X-ray diffractometry. *Physics of the Earth and Planetary Interiors*, 136(1-2):11–24.
- Kennett, B., Engdahl, E., and Buland, R. (1995). Constraints on seismic velocities in the Earth

- from traveltimes. *Geophysical Journal International*, 122(1):108–124.
- Kosarev, G., Kind, R., Sobolev, S., Yuan, X., Hanka, W., and Oreshin, S. (1999). Seismic evidence for a detached Indian lithospheric mantle beneath Tibet. *Science*, 283(5406):1306–1309.
- Krischer, L., Megies, T., Barsch, R., Beyreuther, M., Lecocq, T., Caudron, C., and Wassermann, J. (2015). Obspy: A bridge for seismology into the scientific Python ecosystem. *Computational Science & Discovery*, 8(1):014003.
- Langston, C. A. (1979). Structure under Mount Rainier, Washington, inferred from teleseismic body waves. *Journal of Geophysical Research: Solid Earth*, 84(B9):4749–4762.
- Laske, G., Masters, G., Ma, Z., and Pasyanos, M. (2013). Update on CRUST1.0 A 1-degree global model of Earth's crust. In *Geophys. Res. Abstracts*, volume 15, page 20132658abstrEGU.
- Lee, S.-H., Rhie, J., Park, Y., and Kim, K.-H. (2014). Topography of the 410 and 660 km discontinuities beneath the Korean Peninsula and southwestern Japan using teleseismic receiver functions. *Journal of Geophysical Research: Solid Earth*, 119(9):7245–7257.
- Lekic, V., French, S. W., and Fischer, K. M. (2011). Lithospheric thinning beneath rifted regions of Southern California. *Science*, 334(6057):783–787.
- Li, X., Kind, R., Priestley, K., Sobolev, S., Tilmann, F., Yuan, X., and Weber, M. (2000). Mapping the Hawaiian plume conduit with converted seismic waves. *Nature*, 405(6789):938.
- Ligorria, J. P. and Ammon, C. J. (1999). Iterative deconvolution and receiver-function estimation. *Bulletin of the seismological Society of America*, 89(5):1395–1400.
- Litasov, K. D., Ohtani, E., and Sano, A. (2006). Influence of water on major phase transitions in the Earth's mantle. *Earth's Deep Water Cycle*, pages 95–111.
- Madsen, J., Thorkelson, D., Friedman, R. M., and Marshall, D. (2006). Cenozoic to recent plate configurations in the Pacific basin: Ridge subduction and slab window magmatism in western North America. *Geosphere*, 2(1):11–34.
- Maguire, R., Ritsema, J., Bonnin, M., van Keken, P. E., and Goes, S. (2018a). Evaluating the resolution of deep mantle plumes in teleseismic travelttime tomography. *Journal of Geophysical Research: Solid Earth*, 123(1):384–400.
- Maguire, R., Ritsema, J., and Goes, S. (2018b). Evidence of subduction-related thermal and

- compositional heterogeneity below the United States from transition zone receiver functions. *Geophysical Research Letters*, 45(17):8913–8922.
- Martin-Short, R., Allen, R., Bastow, I. D., Porritt, R. W., and Miller, M. S. (2018). Seismic imaging of the Alaska subduction zone: Implications for slab geometry and volcanism. *Geochemistry, Geophysics, Geosystems*.
- Martin-Short, R., Allen, R. M., and Bastow, I. D. (2016). Subduction geometry beneath south central Alaska and its relationship to volcanism. *Geophysical Research Letters*, 43(18):9509–9517.
- Miller, M. S., ODriscoll, L. J., Porritt, R. W., and Roeske, S. M. (2018). Multiscale crustal architecture of Alaska inferred from P receiver functions. *Lithosphere*, 10(2):267–278.
- Müller, R. D., Seton, M., Zahirovic, S., Williams, S. E., Matthews, K. J., Wright, N. M., Shephard, G. E., Maloney, K. T., Barnett-Moore, N., Hosseinpour, M., et al. (2016). Ocean basin evolution and global-scale plate reorganization events since Pangea breakup. *Annual Review of Earth and Planetary Sciences*, 44:107–138.
- Ohtani, E., Litasov, K., Hosoya, T., Kubo, T., and Kondo, T. (2004). Water transport into the deep mantle and formation of a hydrous transition zone. *Physics of the Earth and Planetary Interiors*, 143:255–269.
- Owens, T. J., Nyblade, A. A., Gurrola, H., and Langston, C. A. (2000). Mantle transition zone structure beneath Tanzania, East Africa. *Geophysical Research Letters*, 27(6):827–830.
- Qi, C., Zhao, D., and Chen, Y. (2007). Search for deep slab segments under Alaska. *Physics of the Earth and Planetary Interiors*, 165(1):68–82.
- Rawlinson, N. and Spakman, W. (2016). On the use of sensitivity tests in seismic tomography. *Geophysical Journal International*, 205(2):1221–1243.
- Ringwood, A. E. et al. (1975). *Composition and petrology of the earth's mantle*.
- Ritsema, J., Deuss, a. A., Van Heijst, H., and Woodhouse, J. (2011). S40RTS: a degree-40 shear-velocity model for the mantle from new Rayleigh wave dispersion, teleseismic travel-time and normal-mode splitting function measurements. *Geophysical Journal International*, 184(3):1223–1236.

- Seton, M., Müller, R., Zahirovic, S., Gaina, C., Torsvik, T., Shephard, G., Talsma, A., Gurnis, M., Turner, M., Maus, S., et al. (2012). Global continental and ocean basin reconstructions since 200 Ma. *Earth-Science Reviews*, 113(3-4):212–270.
- Sigloch, K. (2011). Mantle provinces under North America from multifrequency P wave tomography. *Geochemistry, Geophysics, Geosystems*, 12(2).
- Smyth, J. R. and Frost, D. J. (2002). The effect of water on the 410-km discontinuity: An experimental study. *Geophysical Research Letters*, 29(10):123–1.
- Stixrude, L. and Lithgow-Bertelloni, C. (2011). Thermodynamics of mantle minerals-II. phase equilibria. *Geophysical Journal International*, 184(3):1180–1213.
- Suetsugu, D., Inoue, T., Obayashi, M., Yamada, A., Shiobara, H., Sugioka, H., Ito, A., Kanazawa, T., Kawakatsu, H., Shito, A., et al. (2010). Depths of the 410-km and 660-km discontinuities in and around the stagnant slab beneath the Philippine Sea: Is water stored in the stagnant slab? *Physics of the Earth and Planetary Interiors*, 183(1-2):270–279.
- Tauzin, B., Kim, S., and Kennett, B. (2017). Pervasive seismic low-velocity zones within stagnant plates in the mantle transition zone: Thermal or compositional origin? *Earth and Planetary Science Letters*, 477:1–13.
- Tauzin, B. and Ricard, Y. (2014). Seismically deduced thermodynamics phase diagrams for the mantle transition zone. *Earth and Planetary Science Letters*, 401:337–346.
- Thio, V., Cobden, L., and Trampert, J. (2016). Seismic signature of a hydrous mantle transition zone. *Physics of the Earth and Planetary Interiors*, 250:46–63.
- Vaes, B., van Hinsbergen, D. J., and Boschman, L. M. (2019). Reconstruction of subduction and back-arc spreading in the NW Pacific and Aleutian Basin: clues to causes of Cretaceous and Eocene plate reorganizations. *Tectonics*.
- van der Meer, D. G., van Hinsbergen, D. J., and Spakman, W. (2017). Atlas of the Underworld: Slab remnants in the mantle, their sinking history, and a new outlook on lower mantle viscosity. *Tectonophysics*.
- van der Meijde, M., Marone, F., Giardini, D., and van der Lee, S. (2003). Seismic evidence for water deep in earth's upper mantle. *Science*, 300(5625):1556–1558.

- Veenstra, E., Christensen, D. H., Abers, G. A., and Ferris, A. (2006). Crustal thickness variation in south-central Alaska. *Geology*, 34(9):781–784.
- Xu, W., Lithgow-Bertelloni, C., Stixrude, L., and Ritsema, J. (2008). The effect of bulk composition and temperature on mantle seismic structure. *Earth and Planetary Science Letters*, 275(1-2):70–79.
- Ye, Y., Gu, C., Shim, S.-H., Meng, Y., and Prakapenka, V. (2014). The postspinel boundary in pyrolytic compositions determined in the laser-heated diamond anvil cell. *Geophysical Research Letters*, 41(11):3833–3841.
- Yu, Y. G., Wentzcovitch, R. M., Vinograd, V. L., and Angel, R. J. (2011). Thermodynamic properties of MgSiO_3 majorite and phase transitions near 660 km depth in MgSiO_3 and Mg_2SiO_4 : A first principles study. *Journal of Geophysical Research: Solid Earth*, 116(B2).
- Yuan, H. and Romanowicz, B. (2010). Lithospheric layering in the North American craton. *Nature*, 466(7310):1063–1068.
- Zhang, Y., Wang, Y., Wu, Y., Bina, C. R., Jin, Z., and Dong, S. (2013). Phase transitions of harzburgite and buckled slab under eastern China. *Geochemistry, Geophysics, Geosystems*, 14(4):1182–1199.