

DES/DEB


Extended Abstract

Treatment of a true CxA-OM bifurcation lesion using a one stent drug-coated balloons provisional technique

 Krešimir Gabaldo*,

 Božo Vujeva,

 Katica Cvitkušić
Lukenda,

 Marijana Knežević
Praveček,

Domagoj Vučić

General Hospital „Dr. Josip
Benčević“ Slavonski Brod,
Slavonski Brod, Croatia

KEYWORDS: bifurcation lesions, acute coronary syndrome, drug coated balloons.**CITATION:** *Cardiol Croat.* 2020;15(3-4):51. | <https://doi.org/10.15836/ccar2020.51>***ADDRESS FOR CORRESPONDENCE:** Krešimir Gabaldo, Opća bolnica Dr Josip Benčević, A. Štampara 42, HR-35000 Slavonski Brod, Croatia. / Phone: +385-98-1398-810 / E-mail: kresimir.gabaldo@gmail.com**ORCID:** Krešimir Gabaldo, <https://orcid.org/0000-0002-0116-5929> • Božo Vujeva, <https://orcid.org/0000-0003-0490-3832>
Katica Cvitkušić Lukenda, <https://orcid.org/0000-0001-6188-0708>Marijana Knežević Praveček, <https://orcid.org/0000-0002-8727-7357>

|||||

Background: Provisional stenting is a favorable option for most bifurcation lesions, while two stent techniques show benefits in true bifurcation performed by experts. Using one stent and DCB in true bifurcation lesions still remains questionable.^{1,2}

Case report: 76-year-old male presented with persisting chest pain. ECG showed no specific ischemic changes, while hsTnI was highly elevated. He had gone PCI LAD 14 years ago. Diagnosis: right coronary angiography revealed occlusion of RCA, while left coronary angiography revealed LLS of 20% in proximal LAD stent, 70% stenosis of mid LAD, and acute occlusion of secondary OM branch on bifurcation level, while CxA was stenosed 70-80%. Management: PCI CxA-OM2 was performed with DES Xience expedition 2,75/33mm in CxA-OM2 with POT 3,25/12mm proximally, then after rewiring and adequate lesion preparation a DEB Sequent please 2,5/25mm in CxA distally was performed. After two months we performed an elective PCI LAD and checked out the result of CxA bifurcation which was optimal.

Conclusion: Using one stent and DCB in true bifurcation lesions still remains questionable. There are no data from a prospective study, while there are data from the observational study which enrolled 130 patients. DCB-only strategy was performed in 54% patients, 34.6% had at least one stent in the main branch, 8.5% had at least one stent in the side branch and 3.1% at least one stent in the main branch and side branch. Study follow up lasted for 9.8 months. The TLR rate was 4.5%, MACE was 6.1%, and no stent thrombosis was detected. This study suggested that the DCB+one stent, and DCB-only strategy was safe and effective in selected bifurcations, possibly allowing for an abbreviated antiplatelet regimen.

RECEIVED:

February 11, 2020

ACCEPTED:

February 22, 2020



LITERATURE |||||

1. Bruch L, Zadura M, Waliszewski M, Platonic Z, Eränen J, Scheller B, et al. Results From the International Drug Coated Balloon Registry for the Treatment of Bifurcations. Can a Bifurcation Be Treated Without Stents? *J Interv Cardiol.* 2016 Aug;29(4):348-56. <https://doi.org/10.1111/joic.12301>
2. American College of Cardiology. The 15th Meeting of the European Bifurcation Club. Available from: <https://www.acc.org/latest-in-cardiology/articles/2019/11/07/08/36/the-15th-meeting-of-the-european-bifurcation-club> (February 10, 2020).