

Title	A variant within the FTO confers susceptibility to diabetic nephropathy in Japanese patients with type 2 diabetes
Author(s)	Taira, Makiko; Imamura, Minako; Takahashi, Atsushi; Kamatani, Yoichiro; Yamauchi, Toshimasa; Araki, Shin-ichi; Tanaka, Nobue; van Zuydam, Natalie R.; Ahlqvist, Emma; Toyoda, Masao; Umezono, Tomoya; Kawai, Koichi; Imanishi, Masahito; Watada, Hirotaka; Suzuki, Daisuke; Maegawa, Hiroshi; Babazono, Tetsuya; Kaku, Kohei; Kawamori, Ryuzo; Groop, Leif C.; McCarthy, Mark I.; Kadowaki, Takashi; Maeda, Shiro
Citation	PLOS ONE (2018), 13(12)
Issue Date	2018-12-19
URL	<a href="http://hdl.handle.net/2433/250039">http://hdl.handle.net/2433/250039</a>
Right	© 2018 Taira et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.
Type	Journal Article
Textversion	publisher

RESEARCH ARTICLE

# A variant within the *FTO* confers susceptibility to diabetic nephropathy in Japanese patients with type 2 diabetes

Makiko Taira<sup>1</sup>, Minako Imamura<sup>1,2,3</sup>, Atsushi Takahashi<sup>4,5</sup>, Yoichiro Kamatani<sup>4,6</sup>, Toshimasa Yamauchi<sup>7</sup>, Shin-ichi Araki<sup>8</sup>, Nobue Tanaka<sup>9</sup>, Natalie R. van Zuydam<sup>10,11</sup>, Emma Ahlqvist<sup>12</sup>, Masao Toyoda<sup>13</sup>, Tomoya Umezono<sup>13</sup>, Koichi Kawai<sup>14</sup>, Masahito Imanishi<sup>15,16</sup>, Hiroataka Watada<sup>17,18</sup>, Daisuke Suzuki<sup>19</sup>, Hiroshi Maegawa<sup>8</sup>, Tetsuya Babazono<sup>9</sup>, Kohei Kaku<sup>20</sup>, Ryuzo Kawamori<sup>18</sup>, The SUMMIT Consortium<sup>1</sup>, Leif C. Groop<sup>12,21</sup>, Mark I. McCarthy<sup>10,11,22</sup>, Takashi Kadowaki<sup>7,23,24</sup>, Shiro Maeda<sup>1,2,3</sup>\*



OPEN ACCESS

**Citation:** Taira M, Imamura M, Takahashi A, Kamatani Y, Yamauchi T, Araki S-i, et al. (2018) A variant within the *FTO* confers susceptibility to diabetic nephropathy in Japanese patients with type 2 diabetes. PLoS ONE 13(12): e0208654. <https://doi.org/10.1371/journal.pone.0208654>

**Editor:** Yong-Gang Yao, Kunming Institute of Zoology, Chinese Academy of Sciences, CHINA

**Received:** September 4, 2018

**Accepted:** November 20, 2018

**Published:** December 19, 2018

**Copyright:** © 2018 Taira et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All summary statistics data are available in the National Bioscience Database Center website at (<http://humandbs.biosciencedbc.jp/en/>.) The associated accession number(s)/URL for our data is as follows: hum0014.v12.T2DMwN.v1 and URL: <https://humandbs.biosciencedbc.jp/hum0014-v12>.

**Funding:** This work was partly supported by a grant from the Ministry of Education, Culture, Science and Technology, Japan (to SM), and from the Japan Agency of Medical Research and

**1** Laboratory for Endocrinology, Metabolism and Kidney Diseases, RIKEN Center for Integrative Medical Sciences, Kanagawa, Japan, **2** Department of Advanced Genomic and Laboratory Medicine, Graduate School of Medicine, University of the Ryukyus, Okinawa, Japan, **3** Division of Clinical Laboratory and Blood Transfusion, University of the Ryukyus Hospital, Okinawa, Japan, **4** Laboratory for Statistical Analysis, RIKEN Center for Integrative Medical Sciences, Kanagawa, Japan, **5** Department of Genomic Medicine, Research Institute, National Cerebral and Cardiovascular Center, Osaka, Japan, **6** Kyoto-McGill International Collaborative School in Genomic Medicine, Kyoto University Graduate School of Medicine, Kyoto, Japan, **7** Department of Diabetes and Metabolic Diseases, Graduate School of Medicine, The University of Tokyo, Tokyo, Japan, **8** Department of Medicine, Shiga University of Medical Science, Shiga, Japan, **9** Diabetes Center, Tokyo Women's Medical University School of Medicine, Tokyo, Japan, **10** Wellcome Centre for Human Genetics, Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom, **11** Oxford Center for Diabetes, Endocrinology and Metabolism, Radcliffe Department of Medicine, University of Oxford, Oxford, United Kingdom, **12** Department of Clinical Sciences, Diabetes and Endocrinology, Lund University Diabetes Centre, Scania University Hospital, Malmö, Sweden, **13** Division of Nephrology, Endocrinology and Metabolism, Department of Internal Medicine, Tokai University School of Medicine, Kanagawa, Japan, **14** Kawai Clinic, Ibaragi, Japan, **15** Division of Nephrology and Hypertension, Department of Internal Medicine, Osaka City General Hospital, Osaka, Japan, **16** Department of Nephrology, and Hemodialysis Unit, Ishikiriseiki Hospital, Higashi-Osaka, Japan, **17** Department of Metabolism and Endocrinology, Juntendo University Graduate School of Medicine, Tokyo, Japan, **18** Sportology Center, Juntendo University Graduate School of Medicine, Tokyo, Japan, **19** Suzuki Diabetes Clinic, Kanagawa, Japan, **20** Department of Internal Medicine, Kawasaki Medical School, Okayama, Japan, **21** Institute for Molecular Medicine Finland, University of Helsinki, Helsinki, Finland, **22** Oxford NIHR BIOMEDICAL Research Centre, Oxford University Hospitals Trust, Oxford, United Kingdom, **23** Department of Prevention of Diabetes and Lifestyle-Related Diseases, Graduate School of Medicine, The University of Tokyo, Tokyo, Japan, **24** Department of Metabolism and Nutrition, Mizonokuchi Hospital, Faculty of Medicine, Teikyo University, Kanagawa, Japan

☞ These authors contributed equally to this work.

¶ The complete membership of the author group can be found in the Acknowledgments

\* [smaeda@med.u-ryukyu.ac.jp](mailto:smaeda@med.u-ryukyu.ac.jp)

## Abstract

To explore novel genetic loci for diabetic nephropathy, we performed genome-wide association studies (GWAS) for diabetic nephropathy in Japanese patients with type 2 diabetes. We analyzed the association of 5,768,242 single nucleotide polymorphisms (SNPs) in Japanese patients with type 2 diabetes, 2,380 nephropathy cases and 5,234 controls. We further performed GWAS for diabetic nephropathy using independent Japanese patients with type 2 diabetes, 429 cases and 358 controls and the results of these two GWAS were combined with an inverse variance meta-analysis (stage-1), followed by a *de novo* genotyping for the candidate SNP loci ( $p < 1.0 \times 10^{-4}$ ) in an independent case-control study (Stage-2; 1,213

Development (17km0405202h0802) (TK, TY, SM). The research leading to these results has received funding from the European Union's Seventh Framework Programme (FP7/2007-2013) for the Innovative Medicine Initiative under grant agreement n° 115006, the SUMMIT Consortium.

**Competing interests:** The authors have declared that no competing interests exist.

cases and 1,298 controls). After integrating stage-1 and stage-2 data, we identified one SNP locus, significantly associated with diabetic nephropathy; rs56094641 in *FTO*,  $P = 7.74 \times 10^{-10}$ . We further examined the association of rs56094641 with diabetic nephropathy in independent Japanese patients with type 2 diabetes (902 cases and 1,221 controls), and found that the association of this locus with diabetic nephropathy remained significant after integrating all association data ( $P = 7.62 \times 10^{-10}$ ). We have identified *FTO* locus as a novel locus for conferring susceptibility to diabetic nephropathy in Japanese patients with type 2 diabetes.

## Introduction

Diabetic nephropathy is a leading cause of end-stage renal disease (ESRD), and its prevalence is progressively increasing according to the increase of number of patients with diabetes mellitus [1,2]. Prolonged persistent hyperglycemia is a principal cause of diabetic microvascular complications, but it has been shown that only ~30% of all patients with diabetes develop overt nephropathy, whereas most patients develop diabetic retinopathy 20–30 years after the onset of diabetes [3]. In addition, familial clustering of diabetic nephropathy has been observed in both type 1 and type 2 diabetes [4–6], implying that genetic factors are involved in the development and/or progression of diabetic nephropathy. Identification of disease susceptibility loci for many common diseases such as type 2 diabetes, has been achieved by the introduction of genome-wide association studies (GWAS). Worldwide efforts to identify susceptibility loci for diabetic nephropathy, however, have not yet met with success so much. Several susceptibility loci to diabetic nephropathy or its related traits which showed their lowest p-values close to or with a genome-wide significance level have been unveiled; rs2268388 in *ACACB* [7], rs7583877 in *AFF3* locus, rs17709344 in *RGMA-MCTP2* locus [8], rs4972593 in *Sp3-CDCA7* locus [9], rs1801239 in *CUBN* locus [10], rs161740 in *EPO* locus [11]. However, the results have not been conclusive, and thus most susceptibility loci for diabetic nephropathy remain to be un-identified, suggesting heterogeneity of the disease or contribution of non-genetic factors, which have not been taken into account, may produce inconsistent results among the studies.

In this study, to identify novel susceptibility loci to diabetic nephropathy, we performed a GWAS meta-analysis for diabetic nephropathy using existing Japanese GWAS data for patients with type 2 diabetes.

## Materials and methods

### Study subjects

**Discovery stage (Stage-1).** We selected 7,641 individuals having type 2 diabetes registered in Biobank Japan, and divided these patients into two groups (stage-1, set-1); 1) 2,380 nephropathy cases, defined as patients with overt albuminuria or under renal replacement therapy and 2) 5,234 controls with normoalbuminuria and with diabetes duration of 5 years or longer or with diabetic retinopathy. We also used independent patients with type 2 diabetes who were extracted from previously reported GWAS data [12], and performed GWAS for diabetic nephropathy (stage-1 set 2, cases,  $n = 429$ , controls,  $n = 358$ ). There was no overlap between set-1 and set-2. Diabetes was diagnosed according to the World Health Organization

Table 1. Clinical characteristics of participants.

	Stage1				Stage2	
	Set1		Set2		case	control
	case	control	case	control		
n	2,380	5,234	429	358	1,213	1,298
M/F	1,611/769	3,162/2,072	304/125	226/131**	833/380	759/539
Men%	67.6	60.4	70.9	63.1	68.6	58.5
age	65.9 ± 10.5	66.3 ± 9.6	67.2 ± 10.0	66.9 ± 9.2	63.9 ± 11.0	63.9 ± 10.5
BMI	23.9 ± 4.0*	23.6 ± 3.6	23.5 ± 4.0	23.8 ± 3.8	24.1 ± 3.9	23.6 ± 3.8
Diabetes duration	13.1 ± 9.8*	14.4 ± 8.5	13.4 ± 10.2	13.2 ± 10.0	12.0 ± 9.5	10.5 ± 8.8

\* P < 0.01 vs. control

\*\* information of sex for one participant is not available

<https://doi.org/10.1371/journal.pone.0208654.t001>

(WHO) criteria [13], and those who were diagnosed as type 1 diabetes, mitochondrial diseases or maturity-onset diabetes of the young were excluded.

**Validation analysis (Stage-2).** We examined independent 2,511 patients with Japanese patients with type 2 diabetes, 1,213 diabetic nephropathy cases and 1,298 controls, from the BioBank Japan that were not included in the discovery stage.

Clinical characteristics of participants for Stage-1 (set-1, set-2) and Stage-2 are shown in Table 1. Genomic DNA was extracted from peripheral leukocytes using the standard procedure. All individuals provided written informed consent to participate in this study.

### Genotyping, quality control and imputation in the discovery stage

Set-1 individuals in stage 1 were genotyped using the Human Omni Express Exome Bead Chip. There were 628,670 autosomal SNPs that passed quality control, a call rate  $\geq 0.99$ , Hardy-Weinberg equilibrium test  $P \geq 1 \times 10^{-6}$  in controls and minor allele frequency (MAF)  $\geq 0.01$ . Set-2 samples were genotyped using the Illumina Human 610K SNP array, and 504,984 autosomal SNPs passed the quality control described above and used for further analysis. For sample quality control, we evaluated cryptic relatedness for each sample using an identity-by-state method and removed samples that exhibited second-degree or closer relatedness. We performed principal component analysis to select individuals belong to the major Japanese cluster (Hondo cluster, S1 Fig) as reported previously [14], and data for 7,614 individuals (2,380 diabetic nephropathy cases and 5,234 controls) in set-1 and 787 individuals (429 diabetic nephropathy cases and 358 controls) in set-2 were used in subsequent analyses. We performed genotype imputation with MACH and Minimac [15,16] using linkage disequilibrium data in the 1000 Genomes Project (phased JPT, CHB and Han Chinese South data  $n = 275$ , March 2012) as reference populations. To evaluate the potential effect of population stratification, we used a quantile-quantile (qq) plot of the observed P-values.

### De novo genotyping

We genotyped 1,213 individuals with diabetic nephropathy and 1,298 controls (Stage-2) registered as type 2 diabetes in BioBank Japan, who were not included in a discovery stage using a multiplex PCR-invader assay [17] for SNPs with p values  $< 10^{-4}$  in the meta-analysis. We also genotyped additional 2,123 Japanese patients with type 2 diabetes (902 diabetic nephropathy cases and 1,221 controls), who visited out-patient clinics of Tokai university hospital, Shiga university of medical science hospital, Juntendo university hospital, Kawasaki medical school hospital, Tokyo women's medical university hospital, Iwate Medical University, Toride Kyodo

Hospital, Kawai Clinic, Osaka City General Hospital, Chiba Tokusyukai Hospital or Osaka Rosai Hospital.

Genotyping success rates < 95% or concordance rates < 99.9% were excluded from the analyses.

### Statistical analysis

The association between each SNP and diabetic nephropathy was assessed using the logistic regression test with an additive model with or without adjusting for age, sex, and log-transformed body mass index (BMI) using Mach2dat. We combined data from the each GWAS, validation and replication studies using METAL [18] as an inverse variance method. Heterogeneity in effect sizes among the studies was evaluated with a Cochran's Q test. Regional association plots were generated using LocusZoom [19].

### Ethics approval


The protocol of this study conformed to the provisions of the Declaration of Helsinki and was approved by the ethical committees at the RIKEN Yokohama Institute, Tokai university hospital, Shiga university of medical science, Juntendo university, Kawasaki medical school, Tokyo women's medical university and Iwate Medical University, and the institutional review boards at Toride Kyodo Hospital, Osaka City General Hospital, Chiba Tokusyukai Hospital and Osaka Rosai Hospital.

## Results

### A meta-analysis of GWAS for diabetic nephropathy in the Japanese patients with type 2 diabetes

We obtained genotype data for 7,521,074 SNPs by imputation, and among them, 5,768,242 SNPs those passed quality control ( $r^2 > 0.7$ ) in both studies (Stage-1 set-1 and set-2) were evaluated in this meta-analysis (Fig 1).

There was no genomic inflation in the qq plots in the both studies (S2 Fig). We selected 77 loci associated with diabetic nephropathy in the meta-analysis ( $P < 1 \times 10^{-4}$ , S1 Table), and the associations of the lead SNPs from the 77 loci with DN were examined in an independent case-control study (Stage-2, 1,213 cases and 1,298 controls). In stage-2 analyses, we successfully obtained the genotype data for 65 loci by the multiplex-PCR invader assay. After combining all association data (Stage-1 set-1, set-2 and Stage-2) using the inverse-variance fixed-effects meta-analysis, we found that one SNP locus, rs56094641 in *FTO* at chromosome (Chr) 16, 16q12.2, showed a genome-wide significant association with diabetic nephropathy ( $P = 7.74 \times 10^{-10}$ , odds ratio (OR) = 1.23, 95% confidence interval (CI) 1.15–1.31, Table 2 and S2 Table). The association of the *FTO* locus with overt nephropathy was not affected by an adjustment for age, sex and BMI (rs9936385,  $r^2 = 1$  with rs56094641, S3 Table). We additionally identified suggestive evidence for the associations of six SNP loci with diabetic nephropathy ( $5.0 \times 10^{-8} < P < 5.0 \times 10^{-7}$ , Table 2 and S2 Table): rs895157 in *PRCD* (17q25.1,  $P = 7.70 \times 10^{-8}$ , OR = 1.28, 95% CI 1.17–1.41), rs10144968 near *RAD51B* (14q24.1–24.2,  $P = 1.22 \times 10^{-7}$ , OR = 1.35, 95% CI 1.21–1.51), rs13306536 in *LRP8* (1p32.3,  $P = 2.70 \times 10^{-7}$ , OR = 1.33, 95% CI 1.19–1.48), rs7544082 near *TRABD2B* (1p33,  $P = 3.08 \times 10^{-7}$ , OR = 1.17, 95% CI 1.10–1.24), rs11101179 in *CHAT* (10q11.2–21.1,  $P = 3.85 \times 10^{-7}$ , OR = 1.19, 95% CI 1.11–1.28), rs710375 near *CCNH* (5q14-15,  $P = 3.96 \times 10^{-7}$ , OR = 1.21, 95% CI 1.12–1.30). Regional plots for these 7 loci are shown in S3 Fig.



**Fig 1. Outline of this study.** \*  $r^2 > 0.7$  in both studies, SNP; single nucleotide polymorphism, MAF; minor allele frequency, HWE P; Hardy-Weinberg Equilibrium test P, RSQ, r square.

<https://doi.org/10.1371/journal.pone.0208654.g001>

### Replication study

We examined the association of rs56094641 in *FTO* with overt nephropathy in 2,123 Japanese patients with type 2 diabetes (cases  $n = 902$ , controls  $n = 1,221$ ). As shown in Table 3, rs56094641 in *FTO* showed the same direction of effect with the original finding in the discovery stage. The association of rs56094641 in *FTO* with overt nephropathy was still genome-wide significant level after integration of all association data ( $P = 7.62 \times 10^{-10}$ , OR = 1.21, 95% CI 1.14–1.29, Table 3) though the association was not statistically significant in the replication study alone ( $P = 0.258$ , OR = 1.11, 95% CI 0.93–1.32, Table 3).

By *in silico* replication using SUMMIT consortium data for type 2 diabetes, the association of *FTO* variants with diabetic nephropathy was not replicated in European patients with type 2 diabetes (Table 3).

### Evaluation of previously reported loci

We reflected our results (meta-analysis of Stage-1 set-1 and set-2) on previously- reported susceptibility loci to overt nephropathy. There was no overlap between the above seven SNP loci



Table 2. Association of 7 SNP loci with diabetic nephropathy in Japanese patients with type 2 diabetes.

SNP ID Gene Chromosome	Alleles	study	RAF	OR (95%CI)	P	Phet
rs56094641	G/A	Stage-1, set-1	0.247/0.211	1.22 (1.13–1.32)	$1.19 \times 10^{-6}$	
<i>FTO</i>		Stage-1, set-2	0.236/0.207	1.19 (0.93–1.52)	0.164	
Ch16		Stage-2	0.252/0.209	1.27 (1.11–1.46)	$3.76 \times 10^{-4}$	
		<b>Combined</b>		<b>1.23 (1.15–1.31)</b>	<b><math>7.74 \times 10^{-10}</math></b>	0.85
rs895157	C/A	Stage-1, set-1	0.130/0.108	1.28 (1.14–1.43)	$2.85 \times 10^{-5}$	
<i>PRCD</i>		Stage-1, set-2	0.126/0.113	1.14 (0.83–1.58)	0.422	
Ch17		Stage-2	0.140/0.109	1.35 (1.13–1.60)	$7.02 \times 10^{-4}$	
		<b>Combined</b>		<b>1.28 (1.17–1.41)</b>	<b><math>7.70 \times 10^{-8}</math></b>	0.67
rs10144968	G/T	Stage-1, set-1	0.078/0.062	1.27 (1.11–1.46)	$3.75 \times 10^{-4}$	
<i>RAD51B</i>		Stage-1, set-2	0.083/0.060	1.41 (0.95–2.10)	$8.48 \times 10^{-2}$	
		Stage-2	0.078/0.051	1.58 (1.25–1.99)	$1.15 \times 10^{-4}$	
		<b>Combined</b>		<b>1.35 (1.21–1.51)</b>	<b><math>1.22 \times 10^{-7}</math></b>	0.28
rs13306536	T/C	Stage-1, set-1	0.089/0.068	1.27 (1.11–1.46)	$3.75 \times 10^{-4}$	
<i>LRP8</i>		Stage-1, set-2	0.068/0.075	0.87 (0.58–1.30)	0.496	
Ch1		Stage-2	0.076/0.058	1.35 (1.08–1.69)	$1.15 \times 10^{-4}$	
		<b>Combined</b>		<b>1.33 (1.19–1.48)</b>	<b><math>2.70 \times 10^{-7}</math></b>	0.1
rs7544082	A/C	Stage-1, set-1	0.525/0.492	1.18 (1.09–1.27)	$3.22 \times 10^{-5}$	
<i>TRABD2B</i>		Stage-1, set-2	0.550/0.514	1.15 (0.77–1.72)	0.164	
Ch1		Stage-2	0.552/0.514	1.16 (1.04–1.31)	$8.70 \times 10^{-3}$	
		<b>Combined</b>		<b>1.17 (1.10–1.24)</b>	<b><math>3.08 \times 10^{-7}</math></b>	0.97
rs11101179	C/T	Stage-1, set-1	0.231/0.197	1.22 (1.12–1.33)	$2.23 \times 10^{-6}$	
<i>CHAT</i>		Stage-1, set-2	0.205/0.205	1.00 (0.77–1.31)	0.977	
Ch10		Stage-2	0.226/0.198	1.17 (1.02–1.34)	$2.10 \times 10^{-2}$	
		<b>Combined</b>		<b>1.19 (1.11–1.28)</b>	<b><math>3.85 \times 10^{-7}</math></b>	0.37
rs710375	C/T	Stage-1, set-1	0.201/0.169	1.24 (1.14–1.36)	$2.23 \times 10^{-6}$	
<i>CCNH-TMEM161B</i>		Stage-1, set-2	0.197/0.154	1.38 (1.04–1.77)	0.977	
Ch5		Stage-2	0.187/0.176	1.08 (0.93–1.24)	$2.10 \times 10^{-2}$	
		<b>Combined</b>		<b>1.21 (1.12–1.30)</b>	<b><math>3.96 \times 10^{-7}</math></b>	0.16

<https://doi.org/10.1371/journal.pone.0208654.t002>

and 21 loci from previously-reported GWAS for susceptibility to overt nephropathy (S4 Table). None of the SNPs in the original reports showed significant association with overt nephropathy in the discovery stage ( $P > 2.38 \times 10^{-3} = 0.05/21$ ). Regional lead SNPs within the *ELMO1*, *RGMA-MCTP2*, *ERBB4* and *SP3-CDCA7* in the discovery stage attained the threshold of the correction of multiple testing error.

## Discussion

From a result of GWAS meta-analysis for diabetic nephropathy followed by validation studies in Japanese patients with type 2 diabetes, which comprising in 4,022 cases and 6,890 controls, we identified significant association between rs56094641 in *FTO* and susceptibility to diabetic nephropathy in Japanese patients with type 2 diabetes.

*FTO* locus has been repeatedly reported to be associated with obesity and/or adiposity [20], and, in this study, we have shown that the risk allele for obesity was significantly associated with susceptibility to diabetic nephropathy in Japanese patients with type 2 diabetes.

Smemo *et al.* mentioned that obesity-associated noncoding sequences within *FTO* are functionally connected, at megabase distances, with the homeobox gene *IRX3* [21]. *IRX3* is a

Table 3. Replication studies for 7 loci associated with diabetic nephropathy.

SNP ID gene	Alleles	Ethnicity	RAF	OR (95%CI)	P	Phet
rs56094641	G/A	Japanese	Discovery	1.23 (1.15–1.31)	$7.74 \times 10^{-10}$	
<i>FTO</i>			Replication	1.11 (0.93–1.32)	0.258	
			Combined	1.21 (1.14–1.29)	$7.62 \times 10^{-10}$	0.67
		Europeans		1.06 (1.19–0.95)	0.300	
rs895157	C/A	Japanese	Discovery	1.28 (1.17–1.41)	$7.70 \times 10^{-8}$	
<i>PRCD</i>			Replication	0.96 (0.77–1.21)	0.748	
			Combined	1.23 (1.13–1.34)	$1.15 \times 10^{-6}$	0.11
		Europeans		0.92 (0.80–1.06)	0.195	
rs10144968	G/T	Japanese	Discovery	1.35 (1.21–1.51)	$1.22 \times 10^{-7}$	
<i>RAD51B</i>			Replication	1.04 (0.76–1.41)	0.808	
			Combined	1.31 (1.18–1.45)	$4.23 \times 10^{-7}$	0.17
		Europeans		0.91 (0.80–1.03)	0.289	
rs13306536	T/C	Japanese	Discovery	1.33 (1.19–1.48)	$2.70 \times 10^{-7}$	
<i>LRP8</i>			Replication	0.98 (0.72–1.34)	0.914	
			Combined	1.30 (1.16–1.42)	$1.38 \times 10^{-6}$	0.05
		Europeans		N/A	N/A	
rs7544082	A/C	Japanese	Discovery	1.17 (1.10–1.24)	$3.08 \times 10^{-7}$	
<i>TRABD2B</i>			Replication	0.99 (0.86–1.15)	0.946	
			Combined	1.16 (1.04–1.31)	$2.58 \times 10^{-6}$	0.24
		Europeans		N/A	N/A	
rs11101179	C/T	Japanese	Discovery	1.19 (1.11–1.28)	$3.85 \times 10^{-7}$	
<i>CHAT</i>			Replication	0.92 (0.77–1.10)	0.365	
			Combined	1.15 (1.08–1.23)	$9.36 \times 10^{-6}$	0.03
		Europeans		N/A	N/A	
rs710375	C/T	Japanese	Discovery	1.21 (1.12–1.30)	$3.96 \times 10^{-7}$	
<i>CCNH</i>			Replication	0.85 (0.70–1.03)	$9.86 \times 10^{-2}$	
			Combined	1.16 (1.08–1.24)	$3.02 \times 10^{-5}$	0.002
		Europeans		1.12 (0.999–1.12)	$3.58 \times 10^{-2}$	

N/A, data is not available

<https://doi.org/10.1371/journal.pone.0208654.t003>

member of the Iroquois homeobox gene family and plays a role in an early step of neural development [22]. Members of this family including *IRX3* and *IRX5* appear to play multiple roles during pattern formation of vertebrate embryos [23]. However, there is no evidence showing functional link between *IRX3/IRX5* and diabetic nephropathy.

Obesity has been recognized as an important risk factor for chronic kidney disease (CKD) [24,25]. The incidence of obesity has actualized an increase the number of patients with obesity-related glomerulopathy and proteinuria [26]. Furthermore, obesity has been reported to be an independent risk factor for ESRD by a large historical cohort study of 177,570 adults after adjustment for multiple epidemiologic and clinical conditions including diabetes [27]. Moreover, a *FTO* variant, rs17817449, which are in absolute linkage disequilibrium to rs56094641, was shown to be associated with ESRD [28], suggesting *FTO* variants confer susceptibility to CKD/ESRD through the mechanisms mediated by obesity/adiposity.

In European patients with type 1 or type 2 diabetes, a genetic risk score constructed from confirmed obesity related SNP loci was associated with diabetic nephropathy [29–31], although *FTO* variant alone did not have significant effect on any of renal phenotype.



Therefore, our report is the first to show a robust association of *FTO* variant with diabetic nephropathy. In addition, adjustment for BMI did not influence the association of the SNP with diabetic nephropathy in this study, suggesting that the SNP gets involved in susceptibility to diabetic nephropathy through the mechanism other than increasing BMI. Values of BMI used in this study, however, were obtained after making diagnosis of type 2 diabetes and some therapeutic interventions might affect the BMI values used in this study. Therefore, we might underestimate the effects of BMI on susceptibility to diabetic nephropathy.

We also obtained six SNP loci, which have shown borderline associations with overt nephropathy at the discovery stage (Table 1:  $5 \times 10^{-8} < \text{combined } P \leq 5 \times 10^{-7}$ ), but all candidate SNPs did not show significant effects in the replication stage in an independent Japanese case-control study (Table 3). In this study, we could not replicate almost all of variants identified by meta-analysis of stage 1 and stage 2 in the additional dataset with a comparable sample size with stage 2 GWAS. Although genome-wide SNPs data are not available for individuals in the replication stage, all samples for the replication stage were collected in the Japan main-island, and it has been shown individuals living in Japan main-island are genetically homogeneous [14]; therefore, we think there is no evidence for population heterogeneity between the discovery stage and the replication stage.

We further performed a GWAS meta-analysis for diabetic nephropathy including patients with microalbuminuria as cases, and found that rs56094641 in *FTO* was significantly associated with diabetic nephropathy also in this analysis ( $P = 2.40 \times 10^{-9}$ , OR = 1.18, 95% CI 1.15–1.31, S5 and S6 Tables). However, the association was not stronger than that in the analysis using overt nephropathy as cases, suggesting the *FTO* variant was associated with advanced stages of diabetic nephropathy.

Our study has some limitations. Firstly, the detail clinical information, i.e. blood pressure, Hemoglobin A1c, lipid profiles, prescription of antihypertensive treatments, was not available in many participants in Stage-1 samples. It means that we cannot exclude the possibility that the association of the SNP with diabetic nephropathy is mediated by some risk factors except age, sex and BMI. Second, by *in silico* replication using SUMMIT consortium data for type 2 diabetes, the association of *FTO* variants with diabetic nephropathy was not replicated in European patients with type 2 diabetes (Table 3, S7 Table), and the association of *FTO* variants with diabetic nephropathy showed a genome-wide significant association only in a joint analysis for the discovery stage; therefore, these observations may not be in line with standards for GWA studies; further study, such as a large-scaled longitudinal study, is required to elucidate the association of *FTO* locus with diabetic nephropathy.

## Conclusions

We performed GWAS for diabetic nephropathy in Japanese patients with type 2 diabetes. One SNP locus, *FTO* locus, showed a significant association with susceptibility to diabetic nephropathy, and the association attained a genome-wide significant level. Further studies are required to confirm the association of this locus with diabetic nephropathy.

## Supporting information

**S1 Fig. Results of principal component analysis in stage-1 and stage-2.**  
(PDF)

**S2 Fig. Quantile-quantile plot.** A: Stage-1, set-1, B: Stage-1, set-2.  
(PDF)

**S3 Fig. Regional plot of each candidate locus.** Results of stage-1 GWAS meta-analysis are shown. Red, diamond-shaped plots indicate the most significant variants in each locus after combining stage 1 and stage 2 data.  $r^2$ , linkage disequilibrium coefficient; chr., chromosome. (PDF)

**S1 Table. SNPs with p values  $< 10^{-4}$  in the discovery stage.**

(PDF)

**S2 Table. Candidate SNP loci for overt diabetic nephropathy in the discovery stage.**

(PDF)

**S3 Table. Effects of the adjustment for age, sex and BMI on association of the *FTO* locus with susceptibility to diabetic nephropathy.**

(PDF)

**S4 Table. Association of previously reported loci with overt diabetic nephropathy in the stage 1 analysis.**

(PDF)

**S5 Table. Six SNP loci associated with diabetic nephropathy including microalbuminuria (Meta-analysis in Discovery Stage,  $P < 5 \times 10^{-7}$ ) in Japanese patients with type 2 diabetes.**

(PDF)

**S6 Table. Six SNP loci associated with diabetic nephropathy including microalbuminuria (Meta-analysis in Discovery Stage,  $P < 5 \times 10^{-7}$ ) in Japanese patients with type 2 diabetes.**

(PDF)

**S7 Table. In silico replication of candidate loci for Diabetic Nephropathy in the data from the SUMMIT Consortium.**

(PDF)

**S8 Table. Contributors for SUMMIT consortium.**

(PDF)

**S1 File. Meta-analysis-on-genetic-association-studies-form.**

(DOCX)

## Acknowledgments

The authors thank all participating doctors, staff, and patients from the collaborating institutes for providing DNA samples. We also thank Mr. Takashi Morizono, Drs. Michiaki Kubo, Momoko, Horikoshi, RIKEN Center for Integrative Medical Sciences for their support of data managements, and technical staffs of the Laboratory for Endocrinology, Metabolism and Kidney diseases, RIKEN Center for Integrative Medical Sciences for performing de novo genotyping.

Members of the SUMMIT consortium.

Michael Mark, Markus Albertini, Carine Boustany, Alexander Ehlgen, Martin Gerl, Jochen Huber, Corinna Schölch, Heike Zimdahl-Gelling at Boehringer-Ingelheim, Ingelheim, Germany

Leif Groop, Elisabet Agardh, Emma Ahlqvist, Tord Ajanki, Nibal Al Maghrabi, Peter Almgren, Jan Apelqvist, Eva Bengtsson, Lisa Berglund, Harry Björckbacka, Ulrika Blom-Nilsson, Mattias Borell, Agneta Burström, Corrado Cilio, Magnus Cinthio, Karl Dreja, Pontus Dunér, Daniel Engelbertsen, Joao Fadista, Maria Gomez, Isabel Goncalves, Bo Hedblad, Anna Hultgårdh, Martin E. Johansson, Cecilia Kennbäck, Jasmina Kravic, Claes Ladvall, Åke

Lernmark, Eero Lindholm, Charlotte Ling, Holger Luthman, Olle Melander, Malin Neptin, Jan Nilsson, Peter Nilsson, Tobias Nilsson, Gunilla Nordin Fredriksson, Marju Orho-Melander, Emilia Ottoson-Laakso, Annie Persson, Margaretha Persson, Mats-Åke Persson, Jacqueline Postma, Elisabeth Pranter, Sara Rattik, Gunnar Sterner, Lilian Tindberg, Maria Wigren, Anna Zetterqvist, Mikael Åkerlund, Gerd Östling at Lund University Clinical Research Centre, Malmö, Sweden.

Timo Kanninen, Anni Ahonen-Bishopp, Anita Eliasson, Timo Herrala, Päivi Tikka-Kleemola at Biocomputing Platforms, Espoo, Finland.

Anders Hamsten, Christer Betsholtz, Ami Björkholm, Ulf de Faire, Fariba Foroogh, Guillem Genové, Karl Gertow, Bruna Gigante, Bing He, Karin Leander, Olga McLeod, Maria Nastase-Mannila, Jaako Patrakka, Angela Silveira, Rona Strawbridge, Karl Tryggvason, Max Vikström, John Öhrvik, Anne-May Österholm at Karolinska Institute, Stockholm, Sweden.

Barbara Thorand, Christian Gieger, Harald Grallert, Tonia Ludwig, Barbara Nitz, Andrea Schneider, Rui Wang-Sattler, Astrid Zierer at Helmholtz Centre, Munich, Germany.

Giuseppe Remuzzi, Ariela Benigni, Maria Domenica Lesti, Marina Noris, Norberto Perico, Annalisa Perna, Rossella Piras, Piero Ruggenenti, Erica Rurali at Mario Negri Institute for Pharmacological Research, Bergamo, Italy.

David Dunger, Ludo Chassin, Neil Dalton, John Deanfield, Jane Horsford, Clare Rice, James Rudd, Neil Walker, Karen Whitehead, Max Wong at University of Cambridge, UK.

Helen Colhoun, Fiona Adams, Tahira Akbar, Jill Belch, Harshal Deshmukh, Fiona Dove, Angela Ellingford, Bassam Farran, Mike Ferguson, Gary Henderson, Graeme Houston, Faisal Khan, Graham Leese, Yiyuan Liu, Shona Livingstone, Helen Looker, Margaret McCann, Andrew Morris, David Newton, Colin Palmer, Ewan Pearson, Gillian Reekie, Natalie Smith at University of Dundee, Scotland.

Angela Shore, Kuni Aizawa, Claire Ball, Nick Bellenger, Francesco Casanova, Tim Frayling, Phil Gates, Kim Gooding, Andrew Hattersley, Roland Ling, David Mawson, Robin Shandas, David Strain, Clare Thorn at Peninsula Medical School, Exeter, UK.

Ulf Smith, Ann Hammarstedt, Hans Häring, Oluf Pedersen, Georgio Sesti at University of Gothenburg, Sweden.

Per-Henrik Groop, Emma Fagerholm, Carol Forsblom, Valma Harjutsalo, Maikki Parkkonen, Niina Sandholm, Nina Tolonen, Iiro Toppila, Erkka Valo at Folkhälsan, Helsinki, Finland.

Veikko Salomaa, Aki Havulinna, Kati Kristiansson, Pia Okamo, Tomi Peltola, Markus Perola, Arto Pietilä, Samuli Ripatti, Marketta Taimi at The National Institute for Health and Welfare, Helsinki, Finland.

Seppo Ylä-Herttua, Mohan Babu, Marika Dijkstra, Erika Gurzeler, Jenni Huusko, Ivana Kholová, Markku Laakso, Mari Merentie, Marja Poikolainen at University of Eastern Finland, Kuopio, Finland.

Mark McCarthy, Chris Groves, Thorhildur Juliusdottir, Fredrik Karpe, Vasiliki Lagou, Andrew Morris, Will Rayner, Neil Robertson, Natalie van Zuydam at University of Oxford, UK.

Claudio Cobelli, Barbara Di Camillo, Francesca Finotello, Francesco Sambo, Gianna Toffolo, Emanuele Trifoglio at University of Padova, Italy.

Riccardo Bellazzi, Nicola Barbarini, Mauro Bucalo, Christiana Larizza, Paolo Magni, Alberto Malovini, Simone Marini, Francesca Mulas, Silvana Quaglioni, Lucia Sacchi, Francesca Vitali at University of Pavia, Italy.

Ele Ferrannini, Beatrice Boldrini, Michaela Kozakova, Andrea Mari, Carmela Morizzo, Lucrecia Mota, Andrea Natali, Carlo Palombo, Elena Venturi, Mark Walker at University of Pisa, Italy.

Carlo Patrono, Francesca Pagliaccia, Bianca Rocca at Catholic University of Rome, Italy.  
Pirjo Nuutila, Johanna Haukkala, Juhani Knuuti, Anne Roivainen, Antti Saraste at University of Turku, Finland.

Paul McKeague, Norma Brown, Marco Colombo at University of Edinburgh, Scotland.  
Birgit Steckel-Hamann, Krister Bokvist, Sudha Shankar, Melissa Thomas at Eli Lilly.

Li-ming Gan, Suvi Heinonen, Ann-Cathrine Jönsson-Rylander, Remi Momo, Volker Schneck, Robert Unwin, Anna Walentinsson, Carl Whatling at AstraZeneca.

Everson Nogoceke, Gonzalo Durán Pacheco, Ivan Formentini, Thomas Schindler at Roche.

Piero Tortoli, Luca Bassi, Enrico Boni, Alessandro Dallai, Francesco Guidi, Matteo Lenge, Riccardo Matera, Alessandro Ramalli, Stefano Ricci, Jacopo Viti at University of Florence, Italy

Bernd Jablonka, Dan Crowther, Johan Gassenhuber, Sibylle Hess, Thomas Hübschle, Hans-Paul Juretschke, Hartmut Rütten, Thorsten Sadowski, Paulus Wohlfart at Sanofi-aventis.

Julia Brosnan, Valerie Clerin, Eric Fauman, Craig Hyde, Anders Malarstig, Nick Pullen, Mera Tilley, Theresa Tuthill, Ciara Vangjeli, Daniel Ziemek at Pfizer.

Lead authors of the SUMMIT consortium are MM ([mark.mccarthy@drl.ox.ac.uk](mailto:mark.mccarthy@drl.ox.ac.uk)) and LG ([leif.groop@med.lu.se](mailto:leif.groop@med.lu.se)).

## Author Contributions

**Conceptualization:** Shiro Maeda.

**Data curation:** Atsushi Takahashi.

**Formal analysis:** Makiko Taira, Minako Imamura, Atsushi Takahashi, Yoichiro Kamatani.

**Funding acquisition:** Toshimasa Yamauchi, Takashi Kadowaki, Shiro Maeda.

**Investigation:** Makiko Taira, Minako Imamura, Shiro Maeda.

**Methodology:** Shiro Maeda.

**Project administration:** Shiro Maeda.

**Resources:** Shin-ichi Araki, Nobue Tanaka, Masao Toyoda, Tomoya Umezono, Koichi Kawai, Masahito Imanishi, Hirotaka Watada, Daisuke Suzuki, Hiroshi Maegawa, Tetsuya Babazono, Kohei Kaku, Ryuzo Kawamori.

**Supervision:** Toshimasa Yamauchi, Takashi Kadowaki, Shiro Maeda.

**Validation:** Shin-ichi Araki, Nobue Tanaka, Natalie R. van Zuydam, Emma Ahlqvist, Masao Toyoda, Tomoya Umezono, Leif C. Groop, Mark I. McCarthy, Shiro Maeda.

**Writing – original draft:** Makiko Taira, Shiro Maeda.

**Writing – review & editing:** Minako Imamura.

## References

1. Saran R, Robinson B, Abbott KC, Agodoa LYC, Bhave N, Bragg-Gresham J, et al. US Renal Data System 2017 Annual Data Report: Epidemiology of Kidney Disease in the United States. *Am J Kidney Dis.* 2018; 71: S1–S672.
2. Yokoyama H, Kawai K, Kobayashi M; Japan Diabetes Clinical Data Management Study Group. Microalbuminuria is common in Japanese type 2 diabetic patients: a nationwide survey from the Japan Diabetes Clinical Data Management Study Group (JDDM 10). *Diabetes Care.* 2007; 30: 989–992. <https://doi.org/10.2337/dc06-1859> PMID: 17392559

3. Krolewski AS, Warram JH, Rand LI, Kahn CR. Epidemiologic approach to the etiology of type I diabetes mellitus and its complications. *New Engl J Med*. 1987; 317: 1390–1398. <https://doi.org/10.1056/NEJM198711263172206> PMID: 3317040
4. Quinn M, Angelico MC, Warram JH, Krolewski AS. Familial factors determine the development of diabetic nephropathy in patients with IDDM. *Diabetologia*. 1996; 39: 940–945. PMID: 8858216
5. Pettitt DJ, Saad MF, Bennett PH, Nelson RG, Knowler WC. Familial predisposition to renal disease in two generation of Pima Indians with type 2 (non-insulin-dependent) diabetes mellitus. *Diabetologia*. 1990; 33: 438–443. PMID: 2401399
6. Fava S, Azzopardi J, Hattersley AT, Watkins PJ. Increased prevalence of proteinuria in diabetic sibs of proteinuric type 2 diabetic subjects. *Am J Kidney Dis*. 2000; 35: 708–712. PMID: 10739793
7. Maeda S, Kobayashi MA, Araki S, Babazono T, Freedman BI, Bostrom MA, et al. A single nucleotide polymorphism within the acetyl-coenzyme A carboxylase beta gene is associated with proteinuria in patients with type 2 diabetes. *PLoS Genet*. 2010; 6: e1000842. <https://doi.org/10.1371/journal.pgen.1000842> PMID: 20168990
8. Sandholm N, Salem RM, McKnight AJ, Brennan EP, Forsblom C, Isakova T, et al. New susceptibility loci associated with kidney disease in type 1 diabetes. *PLoS Genet*. 2012; 8: e1002921. <https://doi.org/10.1371/journal.pgen.1002921> PMID: 23028342
9. Sandholm N, McKnight AJ, Salem RM, Brennan EP, Forsblom C, Harjutsalo V, et al. Chromosome 2q31.1 associates with ESRD in women with type 1 diabetes. *J Am Soc Nephrol*. 2013; 24: 1537–1543. <https://doi.org/10.1681/ASN.2012111122> PMID: 24029427
10. Böger CA, Chen MH, Tin A, Olden M, Köttgen A, de Boer IH, et al. *CUBN* is a gene locus for albuminuria. *J Am Soc Nephrol*. 2011; 22: 555–570. <https://doi.org/10.1681/ASN.2010060598> PMID: 21355061
11. Tong Z, Yang Z, Patel S, Chen H, Gibbs D, Yang X, et al. Promoter polymorphism of the erythropoietin gene in severe diabetic eye and kidney complications. *Proc Natl Acad Sci USA*. 2008; 105: 6998–7003. <https://doi.org/10.1073/pnas.0800454105> PMID: 18458324
12. Hara K, Fujita H, Johnson TA, Yamauchi T, Yasuda K, Horikoshi M, et al. Genome-wide association study identifies three novel loci for type 2 diabetes. *Hum Mol Genet*. 2014; 23: 239–246. <https://doi.org/10.1093/hmg/ddt399> PMID: 23945395
13. Alberti KG, Zimmet PZ. Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: diagnosis and classification of diabetes mellitus provisional report of a WHO consultation. *Diabet Med*. 1998; 15: 539–553. [https://doi.org/10.1002/\(SICI\)1096-9136\(199807\)15:7<539::AID-DIA668>3.0.CO;2-S](https://doi.org/10.1002/(SICI)1096-9136(199807)15:7<539::AID-DIA668>3.0.CO;2-S) PMID: 9686693
14. Yamaguchi-Kabata Y, Nakazono K, Takahashi A, Saito S, Hosono N, Kubo M, et al. Japanese population structure, based on SNP genotypes from 7003 individuals compared to other ethnic groups: effects on population-based association studies. *Am J Hum Genet*. 2008; 83: 445–456. <https://doi.org/10.1016/j.ajhg.2008.08.019> PMID: 18817904
15. Li Y, Willer CJ, Ding J, Scheet P, Abecasis GR. MaCH: using sequence and genotype data to estimate haplotypes and unobserved genotypes. *Genet Epidemiol*. 2010; 34: 816–834. <https://doi.org/10.1002/gepi.20533> PMID: 21058334
16. Howie B1, Fuchsberger C, Stephens M, Marchini J, Abecasis GR. Fast and accurate genotype imputation in genome-wide association studies through pre-phasing. *Nat Genet*. 2012; 44: 955–959. <https://doi.org/10.1038/ng.2354> PMID: 22820512
17. Ohnishi Y, Tanaka T, Ozaki K, Yamada R, Suzuki H, Nakamura Y. A high-throughput SNP typing system for genome-wide association studies. *J Hum Genet*. 2001; 46: 471–477. <https://doi.org/10.1007/s100380170047> PMID: 11501945
18. Willer CJ1, Li Y, Abecasis GR. METAL: fast and efficient meta-analysis of genome wide association scan. *Bioinformatics*. 2010; 26: 2190–2191. <https://doi.org/10.1093/bioinformatics/btq340> PMID: 20616382
19. Pruim RJ1, Welch RP, Sanna S, Teslovich TM, Chines PS, Gliedt TP, et al. LocusZoom: regional visualization of genome-wide association scan results. *Bioinformatics*. 2010; 26: 2336–2337. <https://doi.org/10.1093/bioinformatics/btq419> PMID: 20634204
20. Loos RJ, Yeo GS. The bigger picture of *FTO*: the first GWAS-identified obesity gene. *Nat Rev Endocrinol*. 2014; 10: 51–61. <https://doi.org/10.1038/nrendo.2013.227> PMID: 24247219
21. Smemo S, Tena JJ, Kim KH, Gamazon ER, Sakabe NJ, Gómez-Marín C, et al. Obesity-associated variants within *FTO* form long-range functional connections with *IRX3*. *Nature*. 2014; 507: 371–375. <https://doi.org/10.1038/nature13138> PMID: 24646999

22. Bellefroid EJ1, Kobbe A, Gruss P, Pieler T, Gurdon JB, Papalopulu N. *Xiro3* encodes a *Xenopus* homolog of the *Drosophila* Iroquois genes and functions in neural specification. *EMBO J*. 1998; 17: 191–203. <https://doi.org/10.1093/emboj/17.1.191> PMID: 9427753
23. Lewis MT, Ross S, Strickland PA, Snyder CJ, Daniel CW. Regulated expression patterns of IRX-2, an Iroquois-class homeobox gene, in the human breast. *Cell Tissue Res*. 1999; 296: 549–554. PMID: 10370142
24. Stenvinkel P, Zoccali C, Ikizler TA. Obesity in CKD—what should nephrologists know? *J Am Soc Nephrol*. 2013; 24: 1727–1736. <https://doi.org/10.1681/ASN.2013040330> PMID: 24115475
25. Wickman C., Kramer H. Obesity and kidney disease: potential mechanisms. *Semin Nephrol*. 2013; 33: 14–22. <https://doi.org/10.1016/j.semnephrol.2012.12.006> PMID: 23374890
26. D'Agati VD, Chagnac A, de Vries AP, Levi M, Porrini E, Herman-Edelstein M, et al. Obesity-related glomerulopathy: clinical and pathologic characteristics and pathogenesis. *Nat Rev Nephrol*. 2016; 12: 453–471. <https://doi.org/10.1038/nrneph.2016.75> PMID: 27263398
27. Hsu CY, McCulloch CE, Iribarren C, Darbinian J, Go AS. Body mass index and risk factor for end-stage renal disease. *Ann Intern Med*. 2006; 144: 21–28. PMID: 16389251
28. Hubacek JA, Viklicky O, Dlouha D, Bloudickova S, Kubinova R, Peasey A, et al. The *FTO* gene polymorphism is associated with end-stage renal disease: two large independent case-control studies in a general population. *Nephrol Dial Transplant*. 2012; 27: 1030–1035. <https://doi.org/10.1093/ndt/gfr418> PMID: 21788373
29. Sandholm N, Van Zuydam N, Ahlqvist E, Juliusdottir T, Deshmukh HA, Rayner NW, et al. The Genetic Landscape of Renal Complications in Type 1 Diabetes. *J Am Soc Nephrol*. 2017; 28: 557–574. <https://doi.org/10.1681/ASN.2016020231> PMID: 27647854
30. Todd JN, Dahlström EH, Salem RM, Sandholm N, Forsblom C, McKnight AJ, et al. Genetic Evidence for a Causal Role of Obesity in Diabetic Kidney Disease. *Diabetes*. 2015; 64: 4238–4246. <https://doi.org/10.2337/db15-0254> PMID: 26307587
31. van Zuydam NR, Ahlqvist E, Sandholm N, Deshmukh H, Rayner NW, Abdalla M, et al. A Genome-Wide Association Study of Diabetic Kidney Disease in Subjects With Type 2 Diabetes. *Diabetes*. 2018; 67:1414–1427. <https://doi.org/10.2337/db17-0914> PMID: 29703844